

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/335638340>

A new species of the genus *Synixais Aurivillius*, 1911 (Coleoptera: Cerambycidae) from Mindoro island, Philippines

Article in *Baltic Journal of Coleopterology* · September 2019

CITATIONS

0

READS

45

1 author:



Arvīds Barševskis

Daugavpils University

121 PUBLICATIONS 428 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



IUCN assessment of legally protected Invertebrata species for Latvia [View project](#)



Cerambycidae of Philippines [View project](#)

A new species of the genus *Synixais* Aurivillius, 1911 (Coleoptera: Cerambycidae) from Mindoro island, Philippines

Arvīds Barševskis

Barševskis A. 2019. A new species of the genus *Synixais* Aurivillius, 1911 (Coleoptera: Cerambycidae) from Mindoro island, Philippines. *Baltic J. Coleopterol.*, 19(1): 51 - 56.

Synixais mindoroensis sp. n. (Coleoptera: Cerambycidae) from Mindoro island (Philippines) is described and illustrated. An updated catalogue of the genus *Synixais* Aurivillius, 1911 is proposed. The genus *Synixais* in the world fauna is now represented by nine species.

Key words: taxonomy, new species, long-horned beetles, *Synixais*, Lamiinae, Pteropliini, Philippines, Mindoro

Arvīds Barševskis. Daugavpils University, Institute of Life Sciences and Technology, Coleopterological Research Center, Vienības Str. 13, Daugavpils, LV-5401, Latvia; email: arvids.barsevskis@du.lv

INTRODUCTION

The genus *Synixais* Aurivillius, 1911 (Coleoptera: Cerambycidae) belongs to the subfamily Lamiinae Latreille, 1825 and tribe Pteropliini Thomson, 1861. This insufficiently studied genus in the world fauna contains eight species, distributed in the Oriental Region: Laos, Indonesia, Malaysia and Philippines. Species of *Synixais* are characterized by relatively flattened and elongated body, smoothed and variable in shapes elytral spots, and by densely setose antenna with many long setae.

Recently, Vives (2015) described *S. apoensis* Vives, 2015, the first species of the genus from the Philippines archipelago, and three years later, Barševskis (2018) described *S. willietorresi* Barševskis, 2018. Both these species were described from Mindanao island:

S. apoensis from Kabatung (Bukidnon) and *S. willietorresi* from Gutalac vicinity (Zamboanga).

This article presents description of a new species of the genus from Mindoro Island (Baco), the third species known from the Philippines and the first from Mindoro. The genus *Synixais* in the world fauna is now represented by nine species.

MATERIAL AND METHODS

The following abbreviations of museum collections were used in the present study: BMNH - The Natural History Museum, London, UK; CEV - Private Collection of Eduard Vives, Barcelona, Spain;

DUBC - Daugavpils University Beetles Collection, Ilgas, Daugavpils Distr., Latvia;
MNHN - Muséum National d'Histoire Naturelle, Paris, France;
NMNH - United States National Museum, Washington, USA;
NRS - Naturhistoriska Riksmuseet Stockholm, Stockholm, Sweden;
UNIMAS - University of Malaysia, Sarawak, Malaysia.

Examination of the specimen were made under a Nikon SMZ745T binocular stereomicroscope, NIS-Elements 6 D software. Photographs were taken with a Canon EOS 6D camera and Canon MP-E 65 mm macrolens, and processed using Helicon Focus auto montage computer software and subsequently was edited with Photoshop CS6 Extended. The maximum body length was measured from anterior margin of labrum to apex of elytra, and maximum width of the body was measured at level of basal portion of the elytra between shoulders.

The holotype was deposited in DUBC, Daugavpils University Coleopterological Research Center „ILGAS”, Daugavpils Distr., Latvia.

***Synixais mindoroensis* sp. n.**
(Fig. 1)

Type material. Holotype, female: PHILIPPINES: Mindoro isl. /Baco, 03.2019. / Local collector leg. [white handwritten label]; HOLOTYPUS: / *Synixais / mindoroensis* sp. n. / A.Barševskis descr. 2019 [red handwritten label].

General distribution: Philippines: Mindoro island.

Description. Body elongated, parallel-sided, flattened, brown, with variable in size, larger and smaller smoothed spots between grey pubescence. Length: 11.5 mm, width: 3.7 mm.

Head quadrangular, flattened, covered by dense grey pubescence and coarse punctures, with thin elongate line on middle wide impression between antennae. Apical margin of head concaved. Eyes bilobate, not protruding. Both lobes under antennal bases connected by very thin line. Basal elevation of antennae protruding. Cheeks wide, with very dense yellow-grey pubescence. Clypeus indistinctly visible, yellow-brown. Labrum yellow-brown, covered by yellow-grey pubescence. Mandibles narrow, sharp; laterobasal portion of mandibles covered by sparse yellow-grey pubescence. Basal antennomere dark-brown in basal and black in apical portion, antennomere 2 dark-brown, antennomeres 3 - 4 black; antennomeres 1-4 shiny, with pale, sparse pubescence, remaining antennomeres dark-brown, with dense pubescence; all antennomeres with long setae between sparse pale pubescence. Labial and maxillary palpi yellow-brown.

Pronotum almost cylindrical, flattened, slightly widened before middle, yellow-brown, covered by yellow-grey pubescence, with variable in size smoothed spots between it. Pronotum with very fine microsculpture between sparse, coarse punctures. Basal angles of pronotum in distinct, small. Scutellum rounded apically, covered by pubescence. *Pars stridens* visible under basal margin of prothorax, bilobate apically, with very fine transverse microsculpture.

Elytra slightly impressed dorsally before middle, yellow-brown, shiny, covered by dense yellow-grey pubescence, with smoothed spots between it. Lateral portions of elytra behind shoulders, middle and before apical portions with oblique smoothed spots, and with similar spots on dorsal surface of elytra in about middle and apical portions. Middle portion of elytra along suture with many small smoothed spots. Basal portion of elytra with distinctly denser and coarser setiferous punctures as that in middle portion, covered by dark setae. Apex of elytra rounded, densely setose.



Fig. 1. Holotype of *Synixais mindoroensis* sp. n.

Underside of body with very dense yellow-grey pubescence, with sparse, small, smoothed spots between it.

Legs short and robust, dark-brown, covered by dense yellow-grey pubescence and many long setae. Tarsomeres black.

Male unknown.

Differential diagnosis. Based on the general shape of the body, the new species is similar to

S. willietorresi (Fig. 2), but differs from it by the following: 1) body brown, more paler than that in *S. willietorresi*; 2) antennae slender, almost unicolor, with very indistinct transition between brown to black, while coloration of antennae of *S. willietorresi* is distinctly bicolor; 3) dorsal surface of each elytron with oval large smoothed spot before apex, while apical portion of elytra of *S. willietorresi* with two spots, oval and elongated, located along margin; 4) different shape of dorsal spot behind the middle of elytra.



Fig. 2. Holotype of *Synixais willietorresi* Barševskis, 2018

Etymology. Toponymic. The name of species is derived from the name of island, where the holotype was collected.

A CATALOGUE OF SPECIES OF SYNIXAIS AURIVILLIUS, 1911

1. *S. apoensis* Vives, 2015

Distribution: Philippines: Mindanao island

Type deposited: CEV

References: Vives, 2015: 54; Barševskis 2018: 308

2. *S. argentea* Breuning, 1961

Distribution: Malaysia: Pahang

Type deposited: BMNH

References: Breuning, 1961: 18; Barševskis 2018: 307

3. *S. banksi* Breuning, 1938

Distribution: Malaysia: Borneo island (Sarawak)

Type deposited: UNIMAS

References: Breuning, 1938: 246; Barševskis 2018: 307

4. *S. fuscomaculata* Aurivillius, 1911

Distribution: Indonesia/Malaysia (?): Borneo island

Type deposited: NRS

References: Aurivillius, 1911: 211; Barševskis 2018: 308

5. *S. mindoroensis* Barševskis, sp. n.

Distribution: Philippines: Mindoro island

Type deposited: DUBC

6. *S. notaticollis* Breuning, 1965

Distribution: Laos: Xaignaboury region

Type deposited: BPBM

References: Breuning, 1965: 42; Randon & Breuning, 1970: 411; Barševskis 2018: 308

7. *S. strandi* Breuning, 1940

Distribution: Laos.

Type deposited: USNM

References: Breuning, 1940: 421; Randon & Breuning, 1970: 411; Lingafelter & al., 2014: 325; Barševskis 2018: 308

8. *S. sumatrensis* Breuning, 1982

Distribution: Indonesia: Sumatra island

Type deposited: MNHN

References: Breuning, 1982: 17; Barševskis 2018: 308

9. *S. willietorresi* Barševskis, 2018

Distribution: Philippines: Mindanao island

Type deposited: DUBC

References: Barševskis 2018: 306

ACKNOWLEDGEMENTS

I wish to express my gratitude to my colleagues Dr. Alexey Shavrin (Daugavpils, Latvia) for editorial comments on the manuscript, and Anita Rukmane (Daugavpils, Latvia) for help in the preparation of photographs and the laboratory assistance.

REFERENCES

Aurivillius Ch. 1911. Neue oder wenig bekannte Coleoptera Longicornia. 12. *Arkiv för Zoologi, Uppsala* 7 (19): 187-228, figs 49-57.

Barševskis A. 2018. A new species of the genus *Synixais* Aurivillius, 1911 (Coleoptera: Cerambycidae) from the Philippines. *Baltic J. Coleopterol.*, 18(2): 305-308.

Breuning S. 1938. Novae species Cerambycidae VI. *Festschrift zum 60. Geburtstag von Professor Dr. Embrik Strand, Riga* 4 [1937]: 180-392.

Breuning, 1940. Novae species Cerambycidae X. *Folia Zoologica et Hydrobiologica, Riga* 10 (2): 407-437.

Breuning S. 1961. Nouvelles formes de Lamiaires. *Bulletin de l'Institut Royal des*

- Sciences Naturelles de Belgique, Bruxelles* 37 (20): 1-44.
- Breuning S., 1965. Contribution à la connaissance des Lamiens du Laos (Coll. Céramb.) 13^{ème} Partie. *Bulletin de la Société Royale des Sciences Naturelles du Laos* 14: 31-62.
- Breuning S., 1982. Diagnoses préliminaires de nouveaux Lamiinae du Muséum National d'Histoire Naturelle de Paris [Coleoptera, Cerambycidae]. *Annales de la Société Entomologique de France, Paris (N. S.)* 18 (1): 9-29.
- Lingafelter S. Wayne, Nearn Eu. H., Tavakilian G. L., Monne M. Á. & Biondi M., 2014. Longhorned Woodboring Beetles (Coleoptera: Cerambycidae and Disteniidae) Primary Types of the Smithsonian Institution. *Smithsonian Institution Scholarly Press, Washington D.C.: v-xviii + 1-390.*
- Rondon J.A, Breuning S., 1970. Lamiines du Laos. *Pacific Insects Monograph* 24: 315-571.
- Vives E. 2015. New or interesting Cerambycidae from Philippines (XII). (Coleoptera, Cerambycidae, Lamiinae). *Boletín de la S. E. A.* 56 (1): 49-60.

Received: 14.06.2019.

Accepted: 15.07.2019.

Published: 28.08.2019.