

Far Eastern Entomologist

Дальневосточный энтомолог

Journal published by Far East Branch
of the Russian Entomological Society
and Laboratory of Entomology,
Institute of Biology and Soil Science,
Vladivostok

Number 156: 1-11

ISSN 1026-051X

December 2005

THE GROUND BEETLES (COLEOPTERA: CARABIDAE) ASSEMBLAGES ON THE MURAVJEV-AMURSKY PENINSULA, RUSSIAN FAR EAST

**S. K. Kholin¹⁾, S. Yu. Storozhenko¹⁾, V. S. Sidorenko¹⁾,
G. Sh. Lafer¹⁾ and S. Tanabe²⁾**

1) Institute of Biology and Soil Science, Vladivostok-22, 690022, Russia

2) Kanazawa University, Kanazawa, 920-1192, Japan

Six assemblages of the ground beetles are recognized and described in the natural and anthropogenic ecosystems of the Muravjev-Amursky peninsula. The ecological differentiation of carabid beetles assemblages becomes apparent in the number of species and composition of species, and remains permanent for a studied period. The species diversity of assemblages does not tend to decrease under anthropogenic press.

KEY WORDS: beetles, Carabidae, assemblage, anthropogenic and natural ecosystems, Russian Far East.

С. К. Холин¹⁾, С. Ю. Стороженко¹⁾, В. С. Сидоренко¹⁾, Г. Ш. Лафер¹⁾, С. Танабе²⁾. Ассамблеи жуужелиц (Coleoptera: Carabidae) полуострова Муравьев-Амурский, Дальний Восток // Дальневосточный энтомолог. 2005. N 156. С. 1-11.

Выявлены и описаны шесть ассамблей жуужелиц в естественных и нарушенных биотопах п-ова Муравьев-Амурский. Экологическая дифференциация комплексов жуужелиц проявляется как в количестве видов, так и в видовом составе, и сохраняется на протяжении ряда лет. Показано, что видовое богатство ассамблей не всегда снижается при антропогенном воздействии.

1) Биолого-почвенный институт, Дальневосточное отделение Российской академии наук, Владивосток-22, 690022, Россия.

2) Университет Канадзавы, Канадзава, 920-1192, Япония.

INTRODUCTION

The ground beetles are one of the most common objects to study the spatial distribution, structure and diversity of the soil animal communities in the boreal zone (Chernov, 1975; Thiele, 1977; Striganova & Porjadina, 2005). A check-list of 208 species from 52 genera of the families Cicindelidae and Carabidae of the local fauna of the Muravjev-Amursky peninsula has been published recently (Lafer, 2005), but the composition and structure of carabid communities in different biotops are still unknown. Therefore the main purpose of the present paper is to recognize and to describe the ground beetles assemblages in the natural and anthropogenic ecosystems of the Muravjev-Amursky peninsula.

The present investigation has been supported by grants of the RFBR N 05-04-49544 and Far Eastern Branch of the Russian Academy of Sciences N 04-1-OBH-100.

MATERIAL AND METHODS

The study was carried out on the Muravjev-Amursky peninsula, which is located in the southern part of Primorskii krai (Russia). This area lies in the temperate zone and covered mostly with mixed broadleaved and coniferous forest (so called "blackfir" forest) (Kurentzova, 1968; Krestov, 2003). The trees' species composition in forests is highly variable and depends on local conditions. The large Russian city Vladivostok is situated on the peninsula, therefore the vegetation is prone to anthropogenic damage.

In 2003-2005 13 sites were established in the suburbs of Vladivostok (Fig. 1). These sites included different habitat types of primary forests, oak forests, mixed deciduous forests, shores of small rivers and grasslands, among which one grassland was situated in a village (so called "dacha region"), and one mixed deciduous forest was situated in a city park (Table 1).




Fig. 1. The location of the study area and sampling sites on the Muravjev-Amursky peninsula. For more details see Table 1.

Table 1. Characteristics of the study localities

Locality	Year	Biotop	Ground
1	2003	Wet grassland	Clay, dense grass
1a	2004		
2	2003	Deciduous forest edge	Loam with gravel, tree waste
2a	2004		
3	2003	Deciduous forest after fire forcing on the top of hill	Humus soil with small stones, tree waste
3a	2004		
4	2003	Mixed coniferous-deciduous forest on the slope of hill	Humus soil, tree waste, no grass
4a	2004		
5	2003	Mixed coniferous-deciduous forest on the slope of hill	Humus soil, tree waste, no grass
5a	2004		
6	2003	Oak mixed deciduous forest	Thin layer of humus with small stones, loam, tree waste
6a	2004		
7	2003	City park with deciduous trees	Humus soil, litter, dense grass
7a	2004		
8	2004	Shore of small river in mixed coniferous-deciduous forest	Gravelly and stony
8a	2005		
9	2004	Manufactured grassland in the village with horticultural crops	Thick layer of humus
9a	2005		
10	2004	City park with deciduous trees	Dry and very dense ground without humus
11	2005	Glade in mixed deciduous forest with power line	Humus soil, litter, grass and shrubs
12	2005	Mixed deciduous forest on the bottom of valley	Humus soil, tree waste
13	2005	Shore of small river in mixed deciduous forest	Gravelly and stony

The pitfall-traps were used as a sampling method. Plastic jars (diameter 65 mm, volume 200 ml, depth 9.5 cm) were partially filled with water and detergent, and fifteen traps (1.5-2 m apart) were placed in row on each sampling site. Sampling period on every site was 24 hours. The sampling was carried out monthly from May up October on the same day for all study sites. All the collected beetles have been determined by G. Sh. Lafer and deposited in the Institute of Biology and Soil Science, Far East Branch of the Russian Academy of Sciences (Vladivostok).

The obtained data was analysed by using the Margalef's (1), Shannon's (2), and evenness (3) indices of biological diversity, and a principal coordinate analysis based on the Jaccard's coefficient for similarity estimations (4) was used to differentiate ground beetle assemblages (Legendre & Legendre, 1983):

$$D_{Mg} = (S-1) / \ln N \quad (1),$$

$$H' = -\sum P_i \ln P_i \quad (2),$$

$$\text{Var}(H') = (\sum P_i \ln^2 P_i - (H')^2) / N + (S-1) / 2N^2,$$

$$J' = H' / \ln S \quad (3),$$

$$\text{Var}(J') = \text{Var}(H') / \ln^2 S,$$

$$D_J = a / (a+b+c) \quad (4)$$

where S is number of species in sample, N – number of specimens in sample, P_i – the proportion of species i in sample, a – the number of species common to both samples and b and c are the number of species occurring on each of the site, D_{Mg} – index of species diversity by Margalef, H' – Shannon's index, J' – evenness index, Var – variance of estimation, D_J – Jaccard's coefficient.

RESULTS

A total of 2074 specimens were caught in the different localities (Tables 2, 3). A hundred species of carabid beetles were collected in the all sites during 2003-2005, comprising about 50 % of the species diversity of Muravjev-Amursky peninsula. The number of species on studied sites varied from 4 to 31 in different years. The species diversity was most significant for samplings of wet grassland (31 species), edge of deciduous forest (30 species) and manufactured grassland (31 species), thus, the Margalef's index of species diversity for these sites is very high too (Table 3).

Five dominant species, those that comprise 42% of the total individuals in the whole sampling, were the following: *Bembidion poppii captivorum*, *Agonum mandli*, *Carabus billbergi*, *Pterostichus vladivostokensis*, *Nebria coreica*. First species, *B. poppii captivorum*, was dominant only in the river shores (45-70%), *A. mandli* on the deciduous forest edge (37-45%), *C. billbergi* in the oak forest (48-62%), *P. vladivostokensis* in the mixed deciduous forest (41-54%) and *N. coreica* in the wet grassland (40%). Remainder species had low abundances in all localities (Fig. 2).




Fig. 2. Ranked species-abundance curve for total sample of ground beetles.




Fig. 3. The sampling localities plotted on the 1st, 2nd and 3rd axis of the principal coordinate analysis according to their species composition.

The major environmental factors appearing to determine the distribution of ground beetle assemblages are the substratum type, disturbance and soil moisture, all of which are also important determinants of the distribution of plant communities (Thiele, 1977).

The points of each samples (localities) are plotted against the three PCA axes in Fig. 3. The localities from similar habitats tend to form six clusters. Axis 1 tended to have the different type forests with the lowest scores, and the "grassland" localities with highest: number of species would seem to be important to this axis (a positive correlation between number of species and localities scores: $r=0.76$, $p<0.00001$). Axis 2 was possible related to degree anthropogenic pressure, with the city park localities at the most lower points on this axis. Axis 3 distinguished river shore samples in particular and may be related to substrat characteristics: most other localities were not gravelly or stony. These points lie towards the lower end of this axis.

Table 2. Number of carabid beetles sampled in sampling localities on the Murajev-

Species	Sampling						
	1	1a	2	2a	3	3a	4
<i>Notiophilus impressifrons</i>	0	0	0	0	0	1	0
<i>Leistus niger</i>	0	0	0	1	0	0	2
<i>Nebria subdilata</i>	0	0	0	0	0	0	0
<i>Nebria coreica</i>	44	9	27	10	0	0	0
<i>Nebria ochotica</i>	0	0	0	0	0	0	0
<i>Carabus schrencki</i>	0	0	0	0	0	0	1
<i>Carabus canaliculatus careniger</i>	0	0	0	0	0	0	0
<i>Carabus granulatus ussuriensis</i>	2	10	1	1	0	0	0
<i>Carabus smaragdinus</i>	0	0	0	0	0	1	0
<i>Carabus arvensis</i>	0	0	0	0	0	0	0
<i>Carabus venustus</i>	0	0	0	0	12	0	19
<i>Carabus vietinghoffi</i>	0	0	0	0	0	0	0
<i>Carabus hummeli</i>	0	0	0	0	0	0	0
<i>Carabus billbergi</i>	0	0	4	6	0	0	0
<i>Diplous depressus</i>	0	0	0	0	0	0	0
<i>Patrobus septentrionis</i>	4	1	0	0	0	0	0
<i>Asaphidion semilucidum</i>	0	1	0	0	0	0	0
<i>Asaphidion ussuriense</i>	0	0	1	1	0	3	0
<i>Bembidion mandli</i>	0	0	0	0	0	0	0
<i>Bembidion (Diplocampa) sp.</i>	0	3	0	0	0	0	0
<i>Bembidion elevatum lamprosimile</i>	0	0	0	1	0	4	1
<i>Bembidion semipunctatum</i>	0	0	0	0	0	0	0
<i>Bembidion grapei</i>	0	0	0	0	0	0	0
<i>Bembidion poppii captivorum</i>	0	0	0	0	0	0	0
<i>Bembidion serorum</i>	2	0	0	0	0	0	0
<i>Bembidion tetraporum</i>	0	0	0	0	0	0	0
<i>Bembidion gebleri persuasum</i>	0	0	0	0	0	0	0
<i>Bembidion infuscatipenne</i>	0	0	0	0	0	0	0
<i>Epaphius arsenjevi</i>	1	0	0	0	0	0	0
<i>Epaphius dorsistriatus</i>	1	27	5	0	0	0	0
<i>Trechus apicalis</i>	0	0	0	0	0	0	0
<i>Trechus sp. (? sikhotealinensis Ueno et Lafer)</i>	0	0	0	1	0	0	0
<i>Blemus discus</i>	2	15	14	4	0	0	0
<i>Poecilus encopoleus</i>	0	0	0	0	0	0	0
<i>Poecilus reflexicollis</i>	0	1	1	0	0	1	0
<i>Pterostichus sulcitaris</i>	0	1	0	0	0	0	0
<i>Pterostichus aberrans</i>	0	21	0	2	0	0	0
<i>Pterostichus neglectus</i>	0	0	0	0	0	1	0
<i>Pterostichus subovatus</i>	0	0	0	0	0	0	0
<i>Pterostichus prolongatus</i>	5	17	2	1	0	0	0
<i>Pterostichus sutschanensis</i>	0	0	0	2	0	0	16
<i>Pterostichus vladivostokensis</i>	0	0	7	10	5	1	42
<i>Pterostichus microps</i>	0	0	0	0	0	0	0
<i>Pterostichus nigrita</i>	4	3	2	2	0	0	0
<i>Pterostichus rotundangulus</i>	3	25	1	6	0	0	0
<i>Pterostichus interruptus</i>	0	0	0	0	0	0	2
<i>Pterostichus eobius</i>	0	0	0	0	0	0	1
<i>Pterostichus jankowskii</i>	0	0	0	4	0	0	0
<i>Pterostichus longinquus</i>	0	1	0	0	0	0	0
<i>Pterostichus eschscholtzi fortis</i>	0	0	0	0	0	0	0

Amursky peninsula (order of species are given after Lafer, 2005)

locality														
4a	5	5a	6	6a	7	7a	8	8a	9	9a	10	11	12	13
0	1	0	0	6	0	0	0	0	0	0	0	0	0	0
1	3	0	0	0	0	0	0	1	0	0	0	0	1	0
0	0	0	0	0	0	0	14	7	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	1	3	0	0	0	0
0	0	0	0	0	0	0	45	11	0	0	0	0	0	1
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	2	0	0	0	0	0	0	0	0	0	0	0
0	0	0	1	2	1	4	0	0	7	0	2	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	8	1	0	0	0	0	0	0	0	0	0	0
4	12	1	0	0	0	0	0	0	0	0	0	1	0	0
0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
0	4	0	0	0	0	0	2	0	0	0	0	0	0	0
1	0	0	42	43	21	5	0	0	0	0	1	2	7	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	3
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	1	1	0	1	12	3	0	0	0	0	4	1	0	0
0	0	0	5	15	0	0	1	0	0	0	0	0	2	0
0	0	0	0	0	0	0	0	2	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
0	0	0	0	1	0	0	1	0	0	0	0	0	0	4
0	0	0	0	0	0	0	98	67	0	0	0	0	0	93
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	5	2	0	0	0	0	0	0
0	0	0	0	0	0	0	1	7	0	0	0	0	0	1
0	0	0	0	0	0	0	8	6	0	0	0	0	0	14
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	15	4	0	0	0	0
0	0	1	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	1	0
0	0	0	0	0	0	0	0	2	1	0	0	0	0	0
0	0	0	0	0	0	0	0	0	6	1	0	0	0	0
0	0	0	0	0	0	2	0	1	2	0	1	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	3	0	1	0	0	1	1	0	0	0	1
0	0	0	0	0	2	0	0	0	22	3	0	0	0	0
9	14	9	0	0	0	0	0	0	0	0	0	0	0	0
16	32	22	7	5	0	0	1	3	0	0	0	0	4	0
0	0	1	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	1	0	1	0	0	0	1	0
1	0	0	1	1	0	0	0	2	0	0	0	1	1	1
0	0	1	0	0	0	0	1	1	0	0	0	1	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	1	0	0	0	0
0	0	0	0	0	2	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	2	1	0	0	0	1

Table 2 (continued)

Species	Sampling						
	1	1a	2	2a	3	3a	4
<i>Pterostichus niger</i>	0	0	0	0	0	0	0
<i>Pterostichus microcephalus</i>	0	0	0	0	0	0	0
<i>Pterostichus aereipennis</i>	0	0	3	3	4	5	15
<i>Pterostichus alacer</i>	0	0	0	2	0	0	0
<i>Pterostichus orientalis</i>	0	3	26	32	0	0	0
<i>Agonum gracilipes</i>	0	0	0	0	0	0	0
<i>Agonum jankowskii</i>	0	0	0	2	0	0	0
<i>Agonum mandli</i>	15	12	84	95	0	0	0
<i>Agonum sculptipes</i>	6	2	0	3	0	0	0
<i>Agonum bellicum</i>	0	1	0	21	0	0	0
<i>Agonum gracile</i>	0	0	7	0	0	0	0
<i>Agonum fallax</i>	0	1	1	4	0	0	0
<i>Agonum assimile</i>	0	1	0	25	0	0	0
<i>Calathus halensis</i>	1	6	1	0	0	0	0
<i>Synuchus melantho</i>	0	0	0	0	0	0	0
<i>Synuchus intermedius</i>	0	0	0	0	1	0	0
<i>Synuchus orbicollis</i>	1	3	0	0	0	0	0
<i>Synuchus agonus</i>	0	0	0	0	0	0	0
<i>Synuchus chinensis</i>	0	0	0	0	0	0	0
<i>Synuchus congruus</i>	0	0	0	0	0	0	0
<i>Synuchus nivalis</i>	0	0	0	0	0	0	0
<i>Amara communis</i>	0	1	0	1	0	0	0
<i>Amara familiaris</i>	0	0	0	0	0	0	0
<i>Amara lunicollis</i>	0	0	0	0	0	0	0
<i>Amara magnicollis</i>	0	0	0	0	0	0	0
<i>Amara nitida orienticola</i>	0	0	0	1	0	3	0
<i>Amara ovata</i>	0	0	0	0	0	1	0
<i>Amara similata</i>	0	0	0	0	0	0	0
<i>Amara ussuriensis</i>	0	0	0	0	0	0	0
<i>Amara amplipennis</i>	0	5	0	0	0	0	0
<i>Amara aurichalcea</i>	1	0	0	0	0	0	0
<i>Amara mikae</i>	1	1	0	0	0	0	0
<i>Amara pseudosimplicidens</i>	1	0	1	1	0	0	0
<i>Amara plebeja</i>	0	0	0	0	0	0	0
<i>Curtonotus gonioderus</i>	0	0	0	0	0	0	0
<i>Anisodactylus signatus</i>	0	0	0	0	0	0	0
<i>Harpalus latus ussuricus</i>	0	0	0	1	0	0	0
<i>Pseudoophonus capito</i>	1	2	0	0	0	0	0
<i>Pseudoophonus eous</i>	0	0	0	0	0	0	0
<i>Pseudoophonus jureceki</i>	0	0	0	0	0	0	0
<i>Pseudoophonus tridens</i>	0	0	0	0	0	0	0
<i>Pseudoophonus ussuriensis</i>	15	28	0	7	0	0	0
<i>Trichotichnus coruscus</i>	1	1	0	3	0	0	0
<i>Trichotichnus nishioi</i>	0	0	0	0	0	0	0
<i>Bradycellus glabratus</i>	0	0	1	0	0	1	1
<i>Bradycellus curtulus</i>	0	0	0	0	0	0	0
<i>Lioholus jedlickai</i>	0	1	0	0	0	0	0
<i>Chlaenius pallipes</i>	1	2	0	0	0	0	0
<i>Lachnocrepis prolixus</i>	0	1	0	0	0	0	0
<i>Oodes integer</i>	0	0	0	0	0	0	0

locality														
4a	5	5a	6	6a	7	7a	8	8a	9	9a	10	11	12	13
1	3	3	0	6	0	0	0	2	0	0	0	0	5	0
0	2	0	0	0	0	0	0	0	0	0	0	0	0	0
0	3	0	1	4	0	0	1	0	0	0	0	0	1	0
0	0	0	0	0	0	1	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	13	2	0	0	0	0	1	2
0	0	0	0	0	0	0	1	0	1	0	0	0	0	0
0	0	1	0	1	0	6	9	2	2	1	0	12	1	0
0	0	0	0	0	5	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	1	0	0	0	0	0
0	0	0	0	0	0	0	7	0	0	0	0	0	0	0
0	0	0	0	0	2	0	0	0	4	1	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	4	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	2	0	0	0	0
5	0	0	0	0	0	0	0	0	0	0	0	0	1	1
0	0	0	0	0	0	0	0	0	0	0	0	0	1	0
0	0	0	0	0	1	0	2	0	2	0	0	4	2	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
0	0	0	0	0	0	2	2	0	5	0	2	0	0	0
0	0	0	0	0	0	1	0	0	1	0	0	0	0	0
0	0	0	0	0	1	20	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
1	0	1	0	0	7	12	1	0	2	0	4	8	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	1	2	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	1	0	0	1	0	1
0	0	0	0	0	2	0	1	1	2	1	1	0	0	0
0	0	0	0	0	0	0	1	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	3	0	0	0	1
0	0	0	0	0	1	0	0	0	2	0	0	0	0	0
0	0	0	0	0	3	14	0	0	0	0	0	0	0	0
0	0	0	0	0	1	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	1	0	0	0	0	0
0	0	0	0	1	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	1	0	0	0
0	0	0	0	0	0	0	0	0	2	1	0	0	0	0
0	0	0	0	0	5	8	5	0	2	0	0	0	0	5
0	0	0	1	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	1	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	1	2	0	0	0	0	0
0	0	0	0	0	0	0	0	0	5	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	1	0	0	0	0	0
0	0	0	0	0	0	0	0	0	3	2	0	0	0	0
0	0	0	0	0	6	3	0	0	5	7	2	0	0	0
0	0	0	0	0	0	0	0	0	5	1	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	2

Table 3. Characteristics of the samples from different localities on the Muravjev-Amursky peninsula

Locality	P*	S	N	D _{Mg}	H'	Var (H')	J'	Var (J')
1	60	21	112	4.239	2.173	0.015	0.714	0.002
1a	75	31	206	5.631	2.783	0.005	0.810	0.0004
2	60	19	189	3.434	1.910	0.009	0.649	0.001
2a	75	30	253	5.241	2.361	0.008	0.694	0.001
3	60	4	22	0.971	1.118	0.022	0.806	0.011
3a	75	11	22	3.235	2.174	0.030	0.906	0.005
4	60	10	100	1.954	1.598	0.009	0.694	0.002
4a	75	9	39	2.457	1.671	0.024	0.760	0.005
5	60	10	75	2.085	1.724	0.013	0.749	0.003
5a	75	10	41	2.424	1.492	0.035	0.648	0.007
6	60	9	68	1.896	1.327	0.021	0.604	0.004
6a	75	14	90	2.889	1.809	0.017	0.686	0.002
7	60	18	74	3.950	2.371	0.015	0.820	0.002
7a	60	15	84	3.160	2.307	0.009	0.852	0.001
8	75	23	221	4.075	1.954	0.009	0.623	0.001
8a	45	17	118	3.354	1.739	0.018	0.614	0.002
9	75	31	109	6.395	2.955	0.011	0.860	0.001
9a	45	17	34	4.537	2.590	0.022	0.914	0.003
10	60	9	18	2.768	2.043	0.029	0.930	0.006
11	45	11	33	2.860	1.879	0.033	0.784	0.006
12	45	15	33	4.004	2.420	0.023	0.894	0.003
13	45	17	133	3.272	1.295	0.018	0.457	0.002
Total	1365	100	2074	12.963	3.563	0.001	0.770	0.00004

* - number of pitfall traps per day.

Six assemblages were identified due to analysis of the ground beetles species composition of 13 sites on the Muravjev-Amursky peninsula, as follows: 1) different type of forest, 2) shore of small river usually shaded by trees, 3) forest edge, 4) glade, 5) city park, 6) wet and manufactured grassland (Fig. 3). The ecological differentiation of carabid beetles assemblages becomes apparent in the number of species and species composition. Carabid assemblage structure did not tend to change significantly between years. The species diversity of assemblages does not always decrease under anthropogenic press. For example, the number of species in the different types of forest is 2-3 times less than in the city park or manufactured grassland. Moreover, the most diversified carabid fauna is found in the forest edge and wet grasslands (Table 3). Originality of the river shore assemblage is defined by the riparian *Bembidion* species.

The species collected in this study provides a rich database for more detailed research on the ecology and life history of individual species. Moreover, this data can be used to distinguish and classify major habitat groups, however more extensive collecting is needed to produce a detailed interpretable classification within major habitats.

REFERENCES

- Chernov, Yu.N. 1975. Prirodnaya zonalnost' i zivotnyi mir sushi. [The natural zoning and overland fauna]. Moscow: Mysl' Publ. 222 p. (In Russian).
- Krestov, P.V. 2003. Forest Vegetation of Easternmost Russia (Russian Far East). – In: Kolbek J., Srutek, M. & Box, E. (eds.). Forest vegetation of Northeast Asia. Kluwer, Dordrecht: 93-180.
- Kurentzova, G.E. 1968. Rastitelnost' Primorskogo kraya. [The vegetation of Primorye region]. Vladivostok. 192 p. (In Russian).
- Lafer, G.Sh. 2005. A check-list of ground beetles (Coleoptera: Cicindelidae, Carabidae) of the Muravjev-Amursky peninsula, Primorsky kray. – Far Eastern Entomologist 151: 1-8.
- Legendre, L. & Legendre, P. 1983. Numerical ecology. Developments in Environmental Modelling. 3. Amsterdam; Oxford; New York. 419 p.
- Striganova, B.R. & Porjadina, N.M. 2005. Soil animal population in boreal forests of West-Siberian Plain. Moscow: KMK Sci. Press Ltd. 234 p.
- Thiele, H.-U. 1977. Carabid beetles in their environments. – Springer Verlag. Berlin. 369 p.

SHORT COMMUNICATION

A. L. Ozerov. FIRST RECORD OF THE FAMILY NERIIDAE (INSECTA: DIPTERA) IN RUSSIA. – Far Eastern Entomologist. 2005. N 156: 12.

А. Л. Озеров. Первая находка мух-нерIID (Insecta: Diptera: Neriidae) в России // Дальневосточный энтомолог. 2002. N 156. С. 12.

Up to now the representatives of the family Neriidae were not known from Russia. One specimen of the genus *Nerius* Fabricius has been found in the collection of Institute of Biology and Soil Science, Far Eastern Branch of the Russian Academy of Sciences (Vladivostok). The majority of species of the genus *Nerius* Fabricius, 1805 are Neotropical, but one species is known from Palaearctic Region.

Nerius femoratus Coquillett, 1898

MATERIAL. Russia: Primorskii krai, Vladivostok, Akademgorodok, 17.VIII 1967, 1 ♀ (V.A. Trjapitzin).

DISTRIBUTION. Russia (new record); Japan [1].

1. Soós, Á. 1984. Fam. Neriidae. – In: Soós Á. & Papp, L. (eds). Catalogue of Palaearctic Diptera. Vol. 9. Micropezidae-Agromyzidae. Budapest: 24-25.

Author's address:

Zoological Museum,
Moscow Lomonosov State University,
Bol'shaya Nikitskaya 6,
Moscow, 125009, Russia

SHORT COMMUNICATION

V. N. Kuznetsov¹⁾ & S. Shiyake²⁾. *COCCINELLA EXPLANATA* MIYATAKE, 1963 IS NEWLY RECORDED SPECIES OF LADY BEETLES (COLEOPTERA, COCCINELLIDAE) FROM RUSSIA. – Far Eastern Entomologist. 2005. N 156: 13.

В. Н. Кузнецов¹⁾, С. Сияке²⁾. *Coccinella explanata* Miyatake, 1963 – новый для России вид кокциnellид (Coleoptera, Coccinellidae) // Дальневосточный энтомолог. 2002. N 156. С. 13.

A new for the Russian fauna species of the genus *Coccinella* Linnaeus, 1758 is recorded below.

Coccinella explanata Miyatake, 1963

MATERIAL. Russia: Primorskii krai, Lazovskii District, the mouth of Kievka River, sandy seashore, in aphides colonies on the grasses *Glehnia littoralis* Fr. Schmidt ex Miq. and *Oenothera beinnis* L., 3.VIII 2005, 53 specimens (V. Kuznetsov, S. Shiyake); Khasanskii District, 1 km SE Ryazanovka, seashore, 12.VII 2002, 8 specimens (S. Shiyake).

DISTRIBUTION. Russia (new record); South Korea [1], Japan (Hokkaido, Honshu) [2].

1. Park, H.C. 1993. Systematics and Ecology of Coccinellidae (Insecta: Coleoptera) in Korea. Seoul. 299 p. [In Korean].

2. Sasaji, H. 1971. Fauna Japonica. Coccinellidae (Insecta: Coleoptera). Tokyo. 345 p.

Author's address:

1)Institute of Biology and Soil Science,
Vladivostok-22, 690022, Russia

2) Osaka Museum of Natural History,
Nagai Park 1-23,
Osaka, 546-0034, Japan

FAR EASTERN ENTOMOLOGIST
2005

CONTENTS

	N of issue	Pages	Date of issue
A. S. Ryabukhin. A new species of <i>Lathrobium</i> Gravenhorst, 1802 (Coleoptera: Staphylinidae: Paederinae) from Kamchatka Peninsula	144	1-4	Jan.
E. V. Mikhailjova. Description of <i>Diplomaragna provecta</i> sp. n., with the new records of two congeners from the Russian Far East (Diplopoda, Chordeumatida, Diplomaragnidae)	145	1-4	Feb.
Z. A. Fedotova and V. S. Sidorenko. New species of gall midges of the supertribe Oligotrophidi (Diptera, Cecidomyiidae) from the Russian Far East	146	1-12	Apr.
E. P. Nartshuk and V. A. Korneyev. Data on the fauna of Pyrgotidae (Diptera, Cyclorrhapha) of the Russian Far East	147	1-10	May
V. S. Sidorenko. Additions to the fauna of Drosophilidae (Diptera) of Russia and Ukraine	147	11-12	May
A. V. Gorochov. New taxa of Tettigoniidae (Orthoptera) from Vietnam	148	1-12	June
A. I. Zaitzev. New data on Keroplatidae (Diptera, Sciaroidea) from Russian Far East	149	1-4	July
Yu. A. Pesenko. Contributions to the halictid fauna of the Eastern Palaearctic Region: genus <i>Halictus</i> Latreille (Hymenoptera: Halictidae, Halictinae)	150	1-12	Aug.
G. Sh. Lafer. A check-list of ground beetles (Coleoptera: Cicindelidae, Carabidae) of the Muravjev-Amursky Peninsula, Primorskii krai	151	1-8	Sep.
Yu. A. Pesenko. Contributions to the halictid fauna of the Eastern Palaearctic Region: subfamily Nomioidinae (Hymenoptera: Halictidae)	152	1-12	Sep.
M. Yu. Proshchalykin and A. N. Kupianskaya. The bees (Hymenoptera, Apoidea) of the northern part of the Russian Far East	153	1-39	Oct.
S. A. Shabalin. On distribution of <i>Apogonia nigroolivacea</i> Heyden (Coleoptera, Scarabaeidae) in Russia	153	40	Oct.
Yu. V. Astafurova and Yu. A. Pesenko. Contributions to the halictid fauna of the Eastern Palaearctic Region: subfamily Nomiinae (Hymenoptera: Halictidae)	154	1-16	Nov.
E. A. Beljaev and M. G. Ponomarenko. New lepidopterological finds (Lepidoptera: Gelechiidae, Tortricidae, Geometridae) in south of Russian Far East	155	1-11	Nov.

	N of issue	Pages	Date of issue
S. Yu. Storozhenko. <i>Gampsocleis schelkovnikovae</i> Adelung, 1916 is newly recorded katydid species (Orthoptera, Tettigoniidae) from Ukraine	155	12	Nov.
S. K. Kholin, S. Yu. Storozhenko, V. S. Sidorenko, G. Sh. Lafer and S. Tanabe. The ground beetles (Coleoptera: Carabidae) assemblages on the Muravjev- Amursky peninsula, Russian Far East	156	1-11	Dec.
A. L. Ozerov. First record of the family Neriidae (Insecta: Diptera) in Russia	156	12	Dec.
V. N. Kuznetsov and S. Shiyake. <i>Coccinella explanata</i> Miyatake, 1963 is newly recorded species of lady beetles (Coleoptera, Coccinellidae) from Russia	156	13	Dec.

INSTRUCTIONS FOR AUTHORS

Far Eastern Entomologist is journal publishing original papers on entomology, including taxonomy, systematic, morphology, phylogeny, as well biology, ecology and biogeography. Reviews, comprehensive or revisionary studies of the insects thought other East Asia are especially welcome and will be given first priority for publication. Faunistic papers based on materials from the Russian Far East may be submitted also. Submission of a manuscript to Far Eastern Entomologist implies that the report is original, unpublished and is not being considered for publication elsewhere. Papers in languages other than English are not accepted. Articles should be concise and the number of tables and figures limited to what is strictly necessary. Manuscripts should not exceed 16 pages (including figures and tables); additional printed pages are at the expense of the author(s).

Manuscripts should be prepared in accordance with the style and format of recent issues. (Current issues of Far Eastern Entomologist should be checked for style and format). An abstract should be followed by Key Words (2-7) and include no more than 100 words totally. Cite the author and year of publication of genera and species on first mention. The names of genera and species should be underlined. New description must confirm with the current edition of the Code of Zoological Nomenclature. If a new taxon is described, the institution or museum where the type material is deposited must be indicated. The description of new taxa on types deposited in personal collection will not be accepted.

References in the text, as follows: “Bey-Bienko (1932) states...” or “Bey-Bienko (1932: 25) states...” when the author wishes to refer to a specific page, or “(Bey-Bienko, 1932)” as the author of a statement. Joint authors must be connected by “&” in both the text and the references. When there are more than two authors use “et al.” (Bey-Bienko et al., 1932) in the text. If journal names are not spelled out completely they should follow a consistent and accepted format.

Illustration should be numbered in a single series throughout in Arabic numerals. Tables and legends must be typed on separate sheets and should be self-explanatory.

The following transliterations of Russian alphabet should be used:

A - a	E - e	K - k	П - p	Ф - f	Щ - shch
Б - b	Ж - zh	Л - l	Р - r	Х - kh	Ы - y
В - v	З - z	М - m	С - s	Ц - ts	Э - e
Г - g	И - i	Н - n	Т - t	Ч - ch	Ю - yu
Д - d	Й - i	О - o	У - u	Ш - sh	Я - ya

The editors reserve the right to make minor textual corrections that do not alter the author's meaning. Twenty reprints of each article are provided free of charge to the first author. An order form and prices for additional reprints will be sent with the proofs.

Inquiries regarding content, subscription, manuscripts and copies should be sent to editor: S.Yu. Storozhenko, Institute of Biology and Soil Science, Vladivostok, 690022, Russia.

© **Far Eastern entomologist (Far East. entomol.)** Journal published since October 1994.

Editor-in-Chief: S.Yu. Storozhenko

Editorial Board: A.S. Lelej, V.S. Sidorenko, N.V. Kurzenko

Address: Institute of Biology and Soil Science, Far East Branch of Russian Academy of Sciences, 690022, Vladivostok-22, Russia.

E-mail: entomol@ibss.dvo.ru

FAX: (4232) 310 193