

ISSN 0132-8069

Russian Entomological Journal

Русский энтомологический журнал

**Vol. 24. No. 3
Том 24. Вып. 3**

Москва ❖ Moscow

2015

EDITOR-IN-CHIEF

Dr. K.G. Mikhailov

Zoological Museum, Moscow State University, Bolshaya Nikitskaya str. 6, Moscow 125009 Russia.

EDITORIAL BOARD

Dr. A.V. Antropov

Zoological Museum, Moscow State University, Bolshaya Nikitskaya str. 6, Moscow 125009 Russia.

Dr. V.E. Gokhman

Botanical Garden, Moscow State University, Vorobyevy Gory, Moscow 119899 Russia.

Prof. Dr. R.D. Zhantiev

Chair of Entomology, Biological Faculty, Moscow State University, Vorobyevy Gory, Moscow 119899 Russia.

Dr. G.Yu. Lyubarsky

Zoological Museum, Moscow State University, Bolshaya Nikitskaya str. 6, Moscow 125009 Russia.

Dr. K.V. Makarov

Moscow State Pedagogical University, Department of Zoology & Ecology, Kibalchicha str. 6 build. 5, Moscow 129278 Russia.

Dr. A.V. Matalin

Moscow State Pedagogical University, Department of Zoology & Ecology, Kibalchicha str. 6 build. 5, Moscow 129278 Russia.

Dr. L.N. Medvedev

Institute of Ecology and Evolution, Russian Academy of Sciences, Leninsky prosp. 33, Moscow 117071 Russia.

Dr. A.L. Ozerov

Zoological Museum, Moscow State University, Bolshaya Nikitskaya str. 6, Moscow 125009 Russia.

Prof. A.P. Rasnitsyn

Palaeontological Institute, Russian Academy of Sciences, Profsoyuznaya str. 123, Moscow 117321 Russia.

Prof. J. Serrano

Catedrático de Zoología, Facultad de Veterinaria, Campus de Espinardo, Universidad de Murcia, Murcia 30071 Spain.

Dr. A.V. Sviridov

Zoological Museum, Moscow State University, Bolshaya Nikitskaya str. 6, Moscow 125009 Russia.

Dr. S. Venn

Dept. of Geography, Physicum A134, P.O. Box 64, University of Helsinki, Helsinki, 00014 Finland.

ГЛАВНЫЙ РЕДАКТОР

К.Г. Михайлов

Зоологический музей МГУ, ул. Большая Никитская 6, Москва 125009 Россия.

РЕДАКЦИОННАЯ КОЛЛЕГИЯ

А.В. Антропов

Зоологический музей МГУ, ул. Большая Никитская 6, Москва 125009 Россия.

В.Е. Гохман

Ботанический сад МГУ, Воробьевы горы, Москва 119899 Россия.

проф. Р.Д. Жантиев

Кафедра энтомологии, Биологический факультет МГУ, Воробьевы горы, Москва 119899 Россия.

Г.Ю. Любарский

Зоологический музей МГУ, ул. Большая Никитская 6, Москва 125009 Россия.

К.В. Макаров

Московский педагогический государственный университет, кафедра зоологии и экологии, ул. Кибальчича, д.6, корп 5, Москва 129278 Россия.

А.В. Маталин

Московский педагогический государственный университет, кафедра зоологии и экологии, ул. Кибальчича, д.6, корп 5, Москва 129278 Россия.

Л.Н. Медведев

Институт проблем экологии и эволюции РАН, Ленинский просп. 33, Москва 117071 Россия.

А.Л. Озеров

Зоологический музей МГУ, ул. Большая Никитская 6, Москва 125009 Россия.

проф. А.П. Расницын

Палеонтологический институт РАН, Профсоюзная ул. 123, Москва 117321 Россия.

проф. Х. Сerrano

Catedrático de Zoología, Facultad de Veterinaria, Campus de Espinardo, Universidad de Murcia, Murcia 30071 Spain.

А.В. Свиридов

Зоологический музей МГУ, ул. Большая Никитская 6, Москва 125009 Россия.

С. Венн

Dept. of Geography, Physicum A134, P.O. Box 64, University of Helsinki, Helsinki, 00014 Finland.

D.E. Shcherbakov Permian and Triassic ancestors of webspinners (Embiodea) Д.Е. Щербakov Пермские и триасовые предки эмбий (Embiodea)	187–200
R.D. Zhantiev, O.S. Korsunovskaya Acoustic signals of the bush-crickets of tribe Barbitistini (Orthoptera: Tettigoniidae: Phaneropterinae) from Eastern Europe and Caucasus. П. <i>Leptophyes</i> Fieber, 1853, <i>Euconocercus</i> Bey-Bienko, 1950, <i>Barbitistes</i> Charpentier, 1825, <i>Polysarcus</i> Fieber, 1853 Р.Д. Жантиев, О.С. Корсуновская кустические сигналы кузнечиков трибы Barbitistini (Orthoptera: Tettigoniidae: Phaneropterinae) восточной Европы и Кавказа. П. <i>Leptophyes</i> Fieber, 1853, <i>Euconocercus</i> Bey-Bienko, 1950, <i>Barbitistes</i> Charpentier, 1825, <i>Polysarcus</i> Fieber, 1853	201–210
S.V. Kazantsev New species of <i>Metriorrhynchus</i> Gemminger et Harold, 1869 and <i>Porrostoma</i> Laporte, 1838 from New Guinea (Coleoptera: Lycidae) С.В. Казанцев Новые виды <i>Metriorrhynchus</i> Gemminger et Harold, 1869 и <i>Porrostoma</i> Laporte, 1838 из Новой Гвинеи (Coleoptera: Lycidae)	211–233
A.I. Miroshnikov A new species of the genus <i>Mimosebasmia</i> Pic, 1946 (Coleoptera: Cerambycidae) from Vietnam А.И. Мирошников Новый вид рода <i>Mimosebasmia</i> Pic, 1946 (Coleoptera: Cerambycidae) из Вьетнама	235–238
A.I. Miroshnikov The longicorn beetle genus <i>Anaglyptus</i> Mulsant, 1839 (Coleoptera: Cerambycidae) in Vietnam, with descriptions of two new species Жуки-дровосеки рода <i>Anaglyptus</i> Mulsant, 1839 (Coleoptera: Cerambycidae) фауны Вьетнама с описанием двух новых видов	239–242
S.N. Myartseva Three new species of the genus <i>Coccobius</i> Ratzeburg (Hymenoptera: Aphelinidae) from Mexico С.Н. Мярцева Три новых вида рода <i>Coccobius</i> Ratzeburg (Hymenoptera: Aphelinidae) из Мексики	243–246



© “*Russian Entomological Journal*”
составление, редактирование
compiling, editing

На титуле: Byrsax kimurai Miyatake, 1970
Фото К.В. Макарова

Отдел подписки: *К.Г. Михайлов*
Distribution manager: *K.G. Mikhailov*
E-mail: mikhailov2000@gmail.com

The journal is issued quarterly since 1992.

Address for correspondence:
Dr. K.G. Mikhailov,
Zoological Museum of the Moscow State University,
Bolshaya Nikitskaya Street 6, Moscow 125009 Russia.

**Index of contents of vol.1–24, instruction for authors, exact publication dates of former issues
are available in our web-site: <http://kmkjournals.com>**

Printed in September 2015
Отпечатано в сентябре 2015

Издание осуществляется при содействии Зоологического музея МГУ

A new species of the genus *Mimosebasmia* Pic, 1946 (Coleoptera: Cerambycidae) from Vietnam

Новый вид рода *Mimosebasmia* Pic, 1946 (Coleoptera: Cerambycidae) из Вьетнама

Alexandr I. Miroshnikov^{1, 2}
А.И. Мирошников^{1, 2}

¹ Russian Entomological Society, Krasnodar, Russia. E-mail: miroshnikov-ai@yandex.ru

² Sochi National Park, Moskovskaya str., 21, Sochi, Krasnodar region 354002, Russia.

¹ Русское энтомологическое общество, Краснодар, Россия

² Сочинский национальный парк, ул. Московская, 21, Сочи, Краснодарский край 354002, Россия.

KEY WORDS: Coleoptera, Cerambycidae, Cerambycini, *Mimosebasmia*, new species, Vietnam.

КЛЮЧЕВЫЕ СЛОВА: Coleoptera, Cerambycidae, Cerambycini, *Mimosebasmia*, новый вид, Вьетнам.

ABSTRACT. A new species, *Mimosebasmia fedorenkoi* sp.n., is described from Vietnam. Its differences not only from *M. purpurea* Pic, 1946, but also from members of the genus *Gibbocerambyx* Pic, 1923 are listed. The structural features of *M. fedorenkoi* sp.n. notably extend the diagnosis of the genus *Mimosebasmia* Pic, 1946 and increase the morphological similarity of the latter to the genus *Gibbocerambyx*. In this connection, the taxonomic status of *Mimosebasmia* requires revision.

РЕЗЮМЕ. Описывается новый вид *Mimosebasmia fedorenkoi* sp.n. из Вьетнама. Показаны его отличия не только от *M. purpurea* Pic, 1946, но и от представителей рода *Gibbocerambyx* Pic, 1923. Отмечено, что особенности строения *M. fedorenkoi* sp.n. значительно расширяют диагноз рода *Mimosebasmia* Pic, 1946 и усиливают морфологическое сходство последнего с родом *Gibbocerambyx*. В связи с этим указана необходимость уточнения таксономического статуса *Mimosebasmia* при дальнейших исследованиях.

Introduction

The genus *Mimosebasmia* Pic, 1946 has hitherto included a single species, *M. purpurea* Pic, 1946, described from Tonkin (now part of northern Vietnam) [Pic, 1946]. This species is only known to me from the original description and some specimens taken from southern China, Myanmar, Vietnam, Laos, Thailand, and Cambodia. My repeated attempts to relocate the type specimen of *M. purpurea* in the Muséum national d'Histoire naturelle, Paris, France (MNHN), including directly in the Pic Collection, have failed even despite an active help of several colleagues. The type thus seems to be kept in some other place. Moreover, to the best of my knowledge, there seems to be no other publication which would treat this little-known species.

When studying a rich material collected this year in Vietnam, Kon Tum Province, I have discovered a new species of the genus *Mimosebasmia* which is described in this paper. The holotype of this new species is to be deposited in the Zoological Institute of the Russian Academy of Sciences, St. Petersburg, Russia (ZISP).

Mimosebasmia fedorenkoi Miroshnikov, sp.n.
Figs 1–5


MATERIAL. Holotype ♀ (ZISP), Vietnam, Kon Tum Prov., Kon Plong Distr., Dak Khe River, 14°43'20"N / 108°18'58"E, 1030 m, 8.–23.IV.2015, at light, leg. D. Fedorenko.

Paratypes*, 2 ♂♂ 1 ♀ (the author's collection), Vietnam, Da Nang, Bana Mt., 1400 m, 15°59'N / 107°59'E, V.2015, leg. Toi Nguyen.

DIAGNOSIS. The new species resembles *M. purpurea*, but differs very clearly in many characters, including the slenderer body, the more strongly elongated elytra, the mostly distinctly coarser sculpture and the less strongly developed setation of the pronotum, the clearly less strongly elevated elytra at the very base on the sides of the scutellum, the presence of teeth at their apex, the poorly (not strongly) impressed scutellum, the larger eyes with larger ocelli and a comparatively less deep emargination, the shorter genae, the much less strongly developed antennal tubercles, a distinguished shape of the submentum with a clearly smaller puncturation, the longer and slenderer antennae (at least so in the female), the peculiar shape of the apical external angle of antennomeres 7–10, the absence of a tubercle at the apex of the prosternal process, the slenderer legs, the clearly sparser setation of the elytra, but a noticeably denser setation on the venter which especially strongly obscures the sculpture of the metasternum and visible sternites, as well as the denser

* After completion of the paper, I have received from my colleague, Dr. Evgeny S. Koshkin (Khabarovsk, Russia) several quality photographs of further three specimens, 2 males and 1 female, which belong to *Mimosebasmia fedorenkoi* sp.n. These are included in the type series, even though the beetles have not reached me yet.

The males, 25 mm in length, are very similar to the female, but the antennae are barely longer, and the elytra more or less distinctly narrowed from base towards the apex.



Figs 1–5. *Mimosebasmia fedorenkoi* sp.n., holotype, female: 1 — habitus, dorsal view; 2 — habitus, ventral view; 3 — head and pronotum, dorsal view; 4 — head, ventral view; 5 — apex of elytra.

Рис. 1–5. *Mimosebasmia fedorenkoi* sp.n., голотип, самка: 1 — общий вид, сверху; 2 — общий вид, снизу; 3 — голова и переднеспинка, сверху; 4 — голова, снизу; 5 — вершина надкрылий.

pubescence of the antennae and legs.

The similarity of *M. fedorenkoi* **sp.n.** to members of the genus *Gibbocerambyx* Pic, 1923 is also noteworthy (see Remarks). This, to varying degrees, is strongly shown in habitus, the structure of the antennae, some details of the head, the shape of the pronotum, the character of its sculpture and setation, the structure of the elytra, including their base and the teeth at the apex, the structure of the legs, as well as in a number of the other features. However, it differs clearly from all species of that genus at least by a red coloration of the setation of the dorsum, while from most species it differs in a monochrome (not striped) setation of the elytra. In addition, *M. fedorenkoi* **sp.n.**, due to a peculiar complex of its characters (except for the above features of setation) described below, is well distinguished from each species of *Gibbocerambyx* individually.

DESCRIPTION. Female. Body length 23.0 mm, humeral width 5.0 mm. Coloration of integument combines mostly black and red tones; head mainly in area of neck, eyes entirely, antennae mostly, pronotum partly at apex, at base and on coarsely sculptured partitions, scutellum mostly, prosternum partly, meso- and metasterna to a small part, coxae partly, femora and tibiae completely, tarsi almost entirely black; head mostly except for neck, pronotum mainly in spaces between partitions formed by a coarse sculpture, elytra completely except for lighter epipleura, many of antennomeres partly, but most clearly in the area of their apical external angle, venter mostly dark red with this or that colour intensity.

Head with weakly developed antennal tubercles (in *M. purpurea*, these tubercles usually very well-developed); genae about equal in length to last segment of maxillary palpi and much shorter than transverse diameter of eye (in *M. purpurea*, at least in female, genae much longer than last segment of maxillary palpi and about equal in length to transverse diameter of eye); eyes large, rather strongly convex, deeply emarginate, with moderately large ocelli (but clearly larger than in *M. purpurea*); submentum trapezoidal in shape, apex 2.5 times as wide as long in the middle (Fig. 4), with a heterogeneous, irregular puncturation (in *M. purpurea*, submentum apex more than 4 times as wide as long in the middle, with a coarse, irregular, mostly confluent puncturation); antennae noticeably longer than body, extended behind apex of elytra by penultimate antennomere (Figs 1–2) (in *M. purpurea*, antennae of female usually way or another shorter than body, sometimes only reaching apex of elytra); antennomere 2 barely transverse; length ratio of antennomeres 1–11, 41: 9: 55: 35: 62: 64: 62: 53: 52: 47: 57; apical external angle of apical antennomeres sharpened, most strongly in antennomeres 7–10 (but unlike *M. purpurea* not incurved at the very apex).

Pronotum subequal in length and width near middle, at base 1.26 times as long as width at apex and there with a clear constriction; with a complex, patterned sculpture somewhat resembling that in *M. purpurea*, predominantly very coarse, the most coarse on disk and there forming high partitions, mainly longitudinal, mostly strongly to very strongly sinuous, and a transverse, median, broad, somewhat sinuous and highest partition (Fig. 3); with an almost smooth surface both of partitions proper and spaces between them, only with a gentle microsculpture.

Scutellum longitudinal, at apex rounded, slightly impressed (in *M. purpurea*, scutellum very strongly impressed).

Elytra strongly elongated, 3.44 times as long as humeral width; in basal half about parallel-sided, in apical part mostly slightly narrowed towards apex and only before apex strongly narrowed towards it; at the very base on sides of scutellum clearly, but moderately tuberculiform elevated; apical sutural angle extended into a well-expressed, flattened, almost verti-

cally arranged, ensiform tooth, on external side mostly hidden under dense setation, apical external angle forming a short, but clear tooth completely hidden under dense setation dorsally (Fig. 5); with a clear, very dense and confluent puncturation, gradually weakened to apex, forming a finely cellulate, relatively gentle sculpture (in *M. purpurea*, elytra more or less clearly broadened towards apex, in female often broadened more noticeably than in male, but sometimes even in female almost parallel-sided; 2.8–2.93 times as long as humeral width regardless of sex; at the very base lateral to scutellum tuberculiform, very strongly elevated so that scutellum appearing sunken between these elevations; neither apical sutural nor external angle with a tooth, thereby external angle usually broadly rounded).

Prosternum with a heterogeneous, mostly rough and coarse, predominantly rugose sculpture, thereby in apical one-third mainly with transverse folds, in middle part, before coxae, partly with longitudinal folds; prosternal process very strongly broadened at apex; mesosternal process about 2 times as broad as prosternal process between coxae; metasternum and visible sternites with small, dense punctures, more large ones near the middle suture of metasternum; last (visible) sternite at apex broadly rounded, with a very weak emargination, without impression in apical part.

Legs slender, long; femora not claviform; metatarsomere 1 subequal in length to next two metatarsomeres combined.

Head dorsally in most part, pronotum due to two very broad, longitudinal, paramedian stripes, to two narrow stripes best developed in basal half near middle, and to an apical shortened fascia clothed with dense, recumbent, bright red setae, as in Fig. 3 (in *M. purpurea*, pronotum mostly with similar setae, but forming no pattern of stripes and fasciae); elytra with same, but much sparser setae barely obscuring puncturation (in *M. purpurea*, elytra clothed with rather dense, recumbent, red setae strongly hiding puncturation); pronotum on sides due to separate spots, venter almost entirely, legs mostly with recumbent, greyish setae, the most dense on metasternum and visible sternites (Fig. 2); protibiae ventrally and on inner side predominantly in apical part, meso- and metatibiae at apex mostly on inner side, as well as tarsi partly with golden yellow setae; long, erect, light setae mainly developed on head, on sides of pronotum and on abdomen apically; the most abundant, erect, but relatively short, light setation covering apical part of prosternum.

REMARKS. Taking into account the structural features of *M. fedorenkoi* **sp.n.** and, accordingly, the additional new traits which characterize the genus *Mimosebasmia*, its clear similarity to the genus *Gibbocerambyx* becomes apparent. In this connection, in the future the taxonomic status of *Mimosebasmia* seems to require revision.

ETYMOLOGY. The new species honours my colleague and friend, Dr. Dmitry N. Fedorenko (Institute for Problems of Ecology and Evolution, Russian Academy of Sciences, Moscow, Russia), who collected the holotype.

ACKNOWLEDGEMENTS. I am most grateful to Dmitry N. Fedorenko, who was funded by the Russia-Vietnam Tropical Center, for the valuable material he rendered to me for study. My sincere thanks also go to Thierry Deuve and Gérard L. Tavakilian (MNHN) for the opportunity to study the museum material while Denis G. Kasatkin (Rostov-on-Don, Russia) and Francesco Vitali (Luxembourg) provided the specimens of *M. purpurea* from their private collections. I am deeply indebted to Sergey V. Murzin (Moscow, Russia) who has arranged the receipt of some material for study, to Kirill V. Makarov (Moscow Pedagogical State University, Russia) for his help in the preparation

of photographs, again to Gérard L. Tavakilian and Sergey V. Murzin for having kindly performed a thorough search for the type of *M. purpurea* in the MNHN collection, as well as to Evgeny S. Koshkin who shared a valuable information.

References

- Pic M. 1946. Coléoptères du globe (suite) // L'Échange. Revue Linnéenne. Ann.62. No.504. P.5-8.