

FOLIA HEYROVSKYANA

**Check-list of Czechoslovak Insects IV
(Coleoptera)**

Seznam československých brouků



Supplementum 1

Praha 1993

Check-list of Czechoslovak Insects IV

(Coleoptera)

Seznam československých brouků

**Prepared cooperatively by specialists
on the various groups of *Coleoptera*
under the direction of**

JOSEF JELÍNEK

FOLIA HEYROVSKYANA

Supplementum 1

Praha, 1993

Explanations

All names of the taxa of family and genus groups are listed according to the system and distinguished by different type:

Superfamily **TENEBRIONOIDEA**

Family	Meloidae
Subfamily	Meloinae
Tribus	MELOINI
Genus	Meloe
Subgenus	Lampronmeloe

Within each genus and subgenus, the names of species are listed alphabetically and indented. All synonyms are printed in brevier and introduced by the mark of equation (=).

Abbreviations and symbols used in the text:

B	- Bohemia
M	- Moravia incl. Silesia
S	- Slovakia
?B ?M ?S	- dubious or not document records
'B 'M 'S	- species extinct in a particular region
?B ?M ?S	- adventive species
?B ?M ?S	- adventive species acclimatized in a particular region
⁴ (before the name of species)	- species only synanthropic in Czechoslovakia
*B *M *S	- unpublished record for a particular region

Species listed without letters (B, M, S) are listed according to the data in the literature (mostly foreign) pertaining to Czechoslovakia without specification of particular region.

Vysvětlivky

Všechna jména taxonů skupiny čeledi a rodu jsou řazena podle systému a odlišena typem písma:

Nadčeleď **TENEBRIONOIDEA**

Čeleď	Meloidae
Podčeleď	Meloinae
Tribus	MELOINI
Rod	Meloe
Podrod	Lampronmeloe

V rámci rodu či podrodu jsou jména druhů a poddruhů řazena abecedně a vytiskena při levém okraji textu. Synonyma všech taxonů jsou uvozena značkou "=".

Zkratky a symboly použité v textu:

B	- Čechy
M	- Morava včetně Slezska
S	- Slovensko
?B ?M ?S	- pochybný nebo nedoložený údaj
'B 'M 'S	- druh vyhynulý na uvedeném území
?B ?M ?S	- adventivní druh
?B ?M ?S	- druh aklimatizovaný na uvedeném území
⁴ (před jménem druhu)	- druh v Československu pouze synantropní
*B *M *S	- nález dosud nepublikovaný z uvedené oblasti

Druhy uváděné bez výše uvedených písmen (B, M, S) jsou založeny na literárních (především zahraničních) údajích, vztahujících se k Československu bez bližšího upřesnění.

SAPERDINI*Saperda* Fabricius, 1775*Saperda* s. str.*octopunctata* (Scopoli, 1772)*perforata* (Pallas, 1773)*populnea* (Linnaeus, 1758)*punctata* (Linnaeus, 1767)*scalaris* (Linnaeus, 1758)*Anarea* Mulsant, 1839*carcharias* (Linnaeus, 1758)*similis* (Laicharting, 1784)*Menesia* Mulsant, 1856*bipunctata* (Zoubkov, 1829)*Stenostola* Dejean, 1835*dubia* (Laicharting, 1784)*ferrea ferrea* (Schrank, 1776)

B M S

B M S

B M S

B M S

B M S

B M S

B M S

B M S

B M S

B M S

B M S

B M S

B M S

OBEREINI*Oberea* Dejean, 1835*Oberea* s. str.*linearis* (Linnaeus, 1761)*oculata* (Linnaeus, 1758)*pupillata* (Gyllenhal, 1817)

B M S

B M S

B M S

Amaurostoma* J. Müller, 1906erythrocephala* (Schrank, 1776)

B M S

euphorbiae (Germar, 1813)

M S

moravica Kratochvíl, 1989

M S

PHYTOECIINI*Phytoecia* Dejean, 1835*Pilemia* Fairmaire, 1864*hirsutula hirsutula* (Frölich, 1793)

?M S

Cardoria Mulsant, 1863*scutellata* (Fabricius, 1792)

M S

Musaria J. Thomson, 1864*argus* (Frölich, 1793)

M S

nigripes nigripes (Voët, 1778)

M S

=affinis (Harrer, 1784)*Phytoecia* s. str.*coerulea coerulea* (Scopoli, 1772)

M S

cylindrica (Linnaeus, 1758)

B M S

icterica (Schaller, 1783)

B M S

nigricornis (Fabricius, 1781)

B M S

pustulata (Schrank, 1776)

B M S

virgula (Charpentier, 1825)

M S

Opsilia Mulsant, 1863*coerulescens* (Scopoli, 1763)

B M S

molybdaena (Dalman, 1817)

M S

uncinata (W. Redtenbacher, 1842)

B M S

Chrysomelidae

(J. STREJČEK)

The present list is based on data in the literature as well as on the extensive faunal research carried out by the author during last twenty years. The manuscript has been complemented also by further specialists on the family *Chrysomelidae*, namely R. Farion, F. Kantner and M. Zúber. The list contains all species found on the territory of Czechoslovakia by 1992. The species known from Czechoslovakia only from data in the literature have been included only in the case that their occurrence there comports with their known geographic distribution. Species, the occurrence of which is quite improbable were omitted.

Dubious remains the taxonomic status of *Chrysolina lapidaria* Bechyně. It is tentatively listed as a not fully resolved synonym of *Ch. rufa* Duftschmid.

Seznam je založen na literárních údajích a na rozsáhlém faunistickém výzkumu, prováděném autorem v posledních dvaceti letech. Na doplnění rukopisu se podíleli další naši specialisté na tuto čeleď, a to kolegové ing. R. Farion, F. Kantner a M. Zúber, jimž na tomto místě děkuji.

Seznam zahrnuje druhy zjištěné na území Československa do r. 1992, druhy uváděné pouze v literatuře jen pokud jejich výskyt zde odpovídá přirozenému geografickému rozšíření a současným faunistickým poznatkům. Druhy pro území Československa zcela nepravděpodobné již uváděny nejsou.

Problematickým druhem je *Chrysolina lapidaria* Bechyně, která je zde předběžně zařazena jako ne zcela dořešené synonymum k druhu *Ch. rufa* Duftschmid.

D o n a c i i n a e**PLATEUMARINI**

Plateumaris C. G. Thomson, 1859
 =*Euplateumaris* Iablokov-Khnzorian, 1966
 =*Julisiana* Reitter, 1920

- braccata* (Scopoli, 1772) B M S
consimilis (Schrank, 1781) B M S
 =*tenuicornis* (Balthasar, 1934)
rustica (Kunze, 1818) B M S
 =*affinis* Kunze, 1818
 =*sulcifrons* Weise, 1900
sericea (Linnaeus, 1761) B M S
 =*discolor* Panzer, 1795
 =*discolor kratochvili* Bechyně, 1945
 =*lacordairii* Perris, 1864
 =*obsoleta* Jacobson, 1894

DONACIINI

Sominella Jacobson, 1908
 =*Pseudodonacia* Reitter, 1920
 =*Plateumaroides* Iablokov-Khnzorian, 1962
reticulata (Gyllenhal, 1817) B M
 =*appendiculata* Ahrens, 1810

Donaciella Reitter, 1920
cinerea (Herbst, 1784) B M S
 =*hydrocharis* (Fabricius, 1801)
clavipes (Fabricius, 1793)
tomentosa (Ahrens, 1810)

Donacia Fabricius, 1775
Donaciomima Medvedev, 1973

aquatica (Linnaeus, 1758)
 =*dentipes* (Fabricius, 1793)

bicolor Zschach, 1788

brevicornis Ahrens, 1810

brevitarsis C. G. Thomson, 1884

dentata Hoppe, 1795

impressa Paykull, 1799

malinowskyi Ahrens, 1810

marginata Hoppe, 1795

=*lemnæ* (Fabricius, 1801)

=*limbata* (Panzer, 1796)

obscura Gyllenhal, 1813

semicuprea Panzer, 1796

simplex Fabricius, 1775

sparganii Ahrens, 1810

springeri G. Müller, 1916

thalassina Germar, 1811

versicolorea (Brahm, 1790)

vulgaris Zschach, 1788

Donacia s. str.

=*Eudonacia* Haupt, 1956

crassipes Fabricius, 1775

HAEMONIINI

Macroplea Samouelle, 1819
 =*Apelma* Billberg, 1820
 =*Haemona* Dejean, 1821
appendiculata (Panzer, 1794) B M
 =*zosterae* (Fabricius, 1801)

Orsodacninae
Orsodacne Latreille, 1802
cerasi (Linnaeus, 1758) B M S
lineola (Panzer, 1795) B M S

Zeugophorinae
Zeugophora Kunze, 1818
flavicollis (Marsham, 1802) B M S
frontalis Suffrian, 1840 B ?M ?S
scutellaris Suffrian, 1840 B M S
subspinosa (Fabricius, 1781) B M S
turneri Power, 1863 M

Criocerinae**CRIOCERINI**

Lilioceris Reitter, 1912

lili (Scopoli, 1763) B M S
merdigera (Linnaeus, 1758) B M S
tibialis (A. et G. B. Villa, 1838) ?M
Crioceris O. F. Müller, 1764
asparagi asparagi (Linnaeus, 1758) B M S
duodecimpunctata (Linnaeus, 1758) B M S
quatuordecimpunctata (Scopoli, 1763) M S
quinquepunctata (Scopoli, 1763) M S

LEMINI

Lema Fabricius, 1798

cyanella (Linnaeus, 1758) B M S
 =*puncticollis* (Curtis, 1830)

Oulema Des Gozis, 1886

dufschmidi (Redtenbacher, 1874) B M S
erichsoni (Suffrian, 1841) B M S
gallaeciana (Heyden, 1870) B M S
 =*lichenis* auct. nec (Voët, 1806)
 =*cyanella* auct. nec (Linnaeus, 1758)
melanopus (Linnaeus, 1758) B M S
rufocyanea (Suffrian, 1847) B
septentrionis (Weise, 1880) B M S
tristis (Herbst, 1786) B M S

Clytrinae

Labidostomis Chevrolat in Dejean, 1837

cyanicornis (Germar, 1817) B M S
humeralis (Schneider, 1792) B M S
longimana (Linnaeus, 1761) B M S
lucida axillaris (Lacordaire, 1848) B M S

<i>pallidipennis</i> (Gebler, 1830)	?B	?M	S	<i>Asiopus</i> Lopatin, 1965	?B	?M	S
<i>tridentata</i> (Linnaeus, 1758)	B	M	S	<i>apicalis</i> Gebler, 1830			S
<i>Lachnaia</i> Chevrolat, 1837				<i>gamma</i> Herrich-Schäffer, 1829			
= <i>Lachnaea</i> Agassiz, 1846				<i>quatuordecimmaculatus</i> Schneider, 1792	B	M	S
<i>sexpunctata</i> (Scopoli, 1763)	B	M	S	<i>Lamellosus</i> Tomov, 1979			
<i>Clytra</i> Laicharting, 1781				<i>laevicollis</i> Gebler, 1830			M S
= <i>Clytira</i> Fabricius, 1798				<i>Cryptocephalus</i> s. str.			
<i>appendicina</i> Lacordaire, 1848	B	M	S	<i>aureolus aureolus</i> Suffrian, 1847	B	M	S
<i>laeviuscula</i> Ratzeburg, 1837	B	M	S	<i>aureolus illyricus</i> Franz, 1949			S
<i>quadripunctata</i> (Linnaeus, 1758)	B	M	S	<i>biguttatus</i> (Scopoli, 1763)	B	M	S
<i>Smaragdina</i> Chevrolat, 1837				= <i>bipustulatus</i> (Fabricius, 1775)			
= <i>Cyaniris</i> L. Redtenbacher, 1845				<i>bipunctatus</i> (Linnaeus, 1758)	B	M	S
= <i>Gynandropthalma</i> Lacordaire, 1848				<i>coerulescens</i> C. R. Sahlberg, 1839	B	M	S
= <i>Monrosia</i> Medvedev, 1971				<i>cordiger</i> (Linnaeus, 1758)	B	M	S
<i>affinis</i> (Illiger, 1794)	B	M	S	<i>coryli</i> (Linnaeus, 1758)	B	M	S
<i>aurita</i> (Linnaeus, 1766)	B	M	S	<i>decennamaculatus</i> (Linnaeus, 1758)	B	M	S
<i>diversipes</i> (Letzner, 1839)	?B	M		<i>distinguendus</i> Schneider, 1792	B	M	S
<i>flavicollis</i> (Charpentier, 1825)	B	M	S	<i>elongatus</i> Germar, 1824	B	M	S
<i>salicina</i> (Scopoli, 1763)	B	M	S	<i>flavipes</i> Fabricius, 1781	B	M	S
= <i>cyanea</i> (Fabricius, 1775)				<i>frenatus</i> Laicharting, 1781	B	M	S
<i>xanthaspis</i> (Germar, 1824)	M	S		<i>gridellii</i> Burlini, 1950			S
<i>Cheilotoma</i> Chevrolat, 1837				<i>hypocoeridis</i> (Linnaeus, 1758)	B	M	S
= <i>Chilotoma</i> Agassiz, 1846				= <i>cristula</i> Dufour, 1843			
<i>erythrosoma</i> Faldermann, 1837	M			<i>imperialis</i> Laicharting, 1781	B	M	S
<i>musciformis</i> (Goeze, 1777)	B	M	S	<i>janthinus</i> Germar, 1824	B	M	S
<i>Antipus</i> De Geer, 1778				<i>laetus</i> Fabricius, 1792			M S
= <i>Antipa</i> auct.				<i>marginatus</i> Fabricius, 1781	B	M	S
<i>Tituboea</i> Lacordaire, 1848				<i>moraei</i> (Linnaeus, 1758)	B	M	S
<i>macropus</i> (Illiger, 1800)	M	S		<i>nitidulus</i> Fabricius, 1787	B	M	S
				<i>nitidus</i> (Linnaeus, 1758)	B	M	S
				<i>octacosmus</i> Bedel, 1891	B	M	S
				= <i>octoguttatus</i> Schneider, 1792			
				= <i>sexpustulatus</i> Rossi, 1790			
				<i>octomaculatus</i> Rossi, 1790			S
				= <i>quinquepunctatus</i> Harrer, 1784 nec Scopoli, 1763			
				<i>octopunctatus</i> Scopoli, 1763	B	M	S
				= <i>variabilis</i> Schneider, 1791			
				<i>pallifrons</i> Gyllenhal, 1813			M
				= <i>pallidifrons</i> Clavareau, 1913			
				<i>parvulus</i> O. F. Müller, 1776	B	M	S
				<i>primarius</i> Harold, 1872	B		
				<i>punctiger</i> Paykull, 1799	B	M	S
				<i>quadriguttatus</i> Richter, 1820	B	M	S
				<i>quadripustulatus</i> Gyllenhal, 1813	B	M	S
				<i>quinquepunctatus</i> Scopoli, 1763	B	M	S
				= <i>interruptus</i> Suffrian, 1847			
				= <i>signatus</i> Laicharting, 1781			
				<i>sericeus</i> (Linnaeus, 1758)	B	M	S
				<i>sexpunctatus</i> (Linnaeus, 1758)	B	M	S
				<i>signatifrons</i> Suffrian, 1847	B		S
				<i>transiens</i> Franz, 1949			M S
				<i>trimaculatus</i> Rossi, 1790	?B	?M	S
				<i>variegatus</i> Fabricius, 1781	B	M	S

<i>violaceus</i> Laicharting, 1781	B	M	S	<i>Chrysomelinae</i>
<i>virens</i> Suffrian, 1847	B	M	S	<i>TIMARCHINI</i>
<i>vittatus</i> Fabricius, 1775	B	M	S	<i>Timarcha</i> Latreille, 1829
<i>Burlinius</i> Lopatin, 1965	B	M	S	<i>Timarcha</i> s. str.
<i>bilineatus</i> (Linnaeus, 1767)	B	M	S	<i>goettingensis</i> <i>goettingensis</i> (Linnaeus, 1758) B M S
<i>carpathicus</i> Frivaldszky, 1883		M	S	= <i>coraria</i> (Laicharting, 1781)
<i>chrysopus</i> Gmelin, 1788	B	M	S	<i>goettingensis</i> <i>carpathica</i> Bechyně, 1946 S
= <i>huebneri</i> Fabricius, 1792				<i>pratensis</i> Duftschmid, 1825 S
<i>connexus</i> Olivier, 1807	B	M	S	<i>rugulosa</i> <i>rugulosa</i> Herrich-Schäffer, 1838 M S
<i>elegantulus</i> Gravenhorst, 1807	B	M	S	= <i>rugulosa</i> <i>stictai</i> Bechyně, 1947
<i>exiguus</i> Schneider, 1792	B	M	S	<i>rugulosa</i> <i>lomnickii</i> Miller, 1867 ?S
= <i>wasastjernii</i> Gyllenhal, 1827				<i>tenebricosa</i> <i>moravica</i> Bechyně, 1945 ?M S
<i>frontalis</i> Marsham, 1802	B	M	S	<i>tenebricosa</i> <i>submontana</i> Bechyně, 1945 ?B
<i>fulvus</i> Goeze, 1777	B	M	S	<i>Metallotimarcha</i> Motschulsky, 1860
<i>labiatus</i> (Linnaeus, 1761)	B	M	S	<i>metallica</i> <i>metallica</i> (Laicharting, 1781) B M S
<i>macellus</i> Suffrian, 1860		M	S	
<i>ocellatus</i> Drapiez, 1819	B	M	S	
= <i>geminus</i> Gyllenhal, 1827		M		
<i>ochroleucus</i> Stephens, 1834		B	M	<i>CHRYSOMELINI</i>
<i>populi</i> Suffrian, 1848	B	M	S	<i>Leptinotarsa</i> Chevrolat in Dejean, 1837
<i>pusillus</i> Fabricius, 1777	B	M	S	<i>decemlineata</i> (Say, 1824) 2B 2M 2S
= <i>minutus</i> Herbst, 1783				<i>Crosita</i> Motschulsky, 1860
<i>pygmaeus</i> Fabricius, 1792	B			<i>salviae</i> (Germar, 1824) ?S
<i>querceti</i> Suffrian, 1848	B	M	S	<i>Chrysolina</i> Motschulsky, 1860
<i>rufipes</i> Goeze, 1777	B	M	S	= <i>Chrysomela</i> auct. nec Linnaeus, 1758
<i>saliceti</i> Zebe, 1855	B	M	S	<i>Threnosoma</i> Motschulsky, 1860
<i>strigosus</i> Germar, 1823	B	M	S	<i>fimbrialis</i> <i>fimbrialis</i> (Küster, 1845) M S
<i>vittula</i> Suffrian, 1848	B	M	S	<i>Ovostoma</i> Motschulsky, 1860
				<i>globipennis</i> <i>globipennis</i> (Suffrian, 1851) M S
				<i>globipennis</i> <i>slovaca</i> Bechyně, 1946 S
				<i>olivieri</i> <i>slovaca</i> Bechyně, 1946 M S
				= <i>coerulea</i> <i>slovaca</i> Bechyně, 1946
<i>Lamprosoma</i> oomorphus Curtis, 1831				<i>Menhastrilla</i> Bechyně, 1950
= <i>Lamprosoma</i> auct. nec Kirby, 1818				<i>herbacea</i> (Duftschmid, 1825) B M S
<i>concolor</i> (Sturm, 1807)	B	M	S	<i>Synerga</i> Weise, 1900
= <i>kolbei</i> (Scholz, 1926)				<i>coeruleans</i> <i>coeruleans</i> (L. G. Scriba, 1791) B M S
				<i>Euchrysolina</i> Bechyně, 1950
				<i>graminis</i> <i>graminis</i> (Linnaeus, 1758) B M S
<i>Eumolpini</i>				<i>Taeniochrysea</i> Bechyně, 1950
<i>Eumolpus</i> Illiger, 1798				<i>americana</i> (Linnaeus, 1758) ?M
= <i>Chrysochus</i> Chevrolat in Dejean, 1837				<i>Heliostola</i> Motschulsky, 1860
<i>aslepiadeus</i> (Pallas, 1776)	B	M	S	<i>carpathica</i> <i>gabrieli</i> (Weise, 1903) ?M S
				<i>lichenis</i> <i>lichenis</i> (Richter, 1820) B
				<i>lichenis</i> <i>moraviaca</i> (Weise, 1882) ?M S
				<i>schniederi</i> (Weise, 1882) S
<i>ADOXINI</i>				<i>Erythrochrysa</i> Bechyně, 1950
<i>Eupales</i> Lefevre, 1885				<i>polita</i> <i>polita</i> (Linnaeus, 1758) B M S
<i>ulema</i> (Germar, 1813)		S		<i>Chrysolina</i> s. str.
<i>Adoxus</i> Kirby, 1837				<i>staphylea</i> <i>staphylea</i> (Linnaeus, 1758) B M S
= <i>Bromius</i> Chevrolat in Dejean, 1837				<i>Chrysomorpha</i> Motschulsky, 1860
<i>obscurus</i> <i>obscurus</i> (Linnaeus, 1758)	B	M	S	<i>cerealis</i> <i>cerealis</i> (Linnaeus, 1767) B
<i>obscurus</i> <i>villosulus</i> (Schrank, 1781)		M	S	<i>cerealis</i> <i>plorans</i> Bechyně, 1948 B M S
<i>Pachnephorus</i> L. Redtenbacher, 1845	B	M	S	
<i>pilosus</i> (Rossi, 1790)				
<i>tessellatus</i> (Duftschmid, 1825)	M	S		
<i>vilosus</i> (Duftschmid, 1825)	M	S		

<i>Colaphoptera</i> Motschulsky, 1860		<i>Chalcoidea</i> Motschulsky, 1860	
<i>crassimargo ominosa</i> Bechyně, 1950	B	<i>analis</i> (Linnaeus, 1767)	B M S
<i>marcasitica turgida</i> (Weise, 1882)	B M S	<i>carnifex carnifex</i> Fabricius, 1792	B M S
<i>purpurascens purpurascens</i> (Germar, 1882)	B M S	<i>marginata marginata</i> (Linnaeus, 1758)	B M S
<i>rufa bohemiae</i> Bechyně, 1950	B	<i>Craspeda</i> Motschulsky, 1860	
<i>rufa diminuta</i> Bechyně, 1950	M S	<i>limbata limbata</i> (Fabricius, 1775)	B M
= <i>dahlii</i> Suffrian, 1851 nec Matzek, 1843		<i>limbata findeli</i> (Suffrian, 1870)	?M S
=? <i>lapidaria</i> Bechyně, 1950		<i>limbata kavani</i> Bechyně, 1950	S
<i>rufa squalida</i> Bechyně, 1950	B M	<i>Fastuolina</i> Motschulsky, 1860	
<i>rufa staphylaeoides</i> Bechyně, 1950	B	= <i>Dlochrysa</i> auct.	
<i>umbratilis</i> (Weise, 1887)	B M S	<i>fastuosa</i> (Scopoli, 1763)	B M S
= <i>olivacea</i> (Suffrian, 1851)		<i>Oreina</i> Chevrolat in Dejean, 1837	
<i>Minckia</i> E. Strand, 1935		= <i>Chrysochloa</i> Hoppe, 1840	
<i>chalcites</i> (Germar, 1824)	S	= <i>Orina</i> Agassiz, 1846	
<i>oricalcia</i> (O. F. Müller, 1776)	B M S	<i>Oreina</i> s. str.	
= <i>oricalcea</i> auct.		= <i>Romalarina</i> Weise in Heyden, Reitter et Weise, 1906	
<i>rufoaenea</i> (Suffrian, 1851)	S	<i>alpestris alpestris</i> (Schummel, 1844)	M S
<i>Colaphodes</i> Motschulsky, 1860		<i>alpestris polonina</i> Bechyně, 1958	S
<i>haemoptera haemoptera</i> (Linnaeus, 1758)	B M S	<i>alpestris umbrosa</i> (Weise, 1884)	?B
<i>Allocrysolina</i> Bechyně, 1850		<i>bifrons decora</i> (Richter, 1820)	?B M
<i>fuliginosa galii</i> (Weise, 1884)	B	<i>bifrons obenbergeri</i> Marchand, 1939	S
<i>Colaphosoma</i> Motschulsky, 1860		<i>intricata intricata</i> (Germar, 1824)	B M S
<i>sturmi</i> (Westhoff, 1882)	B M S	<i>viridis</i> (Duftschmid, 1825)	S
= <i>diversipes</i> (Bedel, 1892)		<i>Allorina</i> Weise, 1902	
= <i>goettingensis</i> (Linnaeus, 1761) nec (Linnaeus, 1758)		<i>bidentata</i> Bontems, 1981	?S
= <i>violacea</i> (Weise, 1916)		= <i>tristis</i> auct.	
<i>Taeniosticha</i> Motschulsky, 1860		<i>caerulea</i> (Olivier, 1790)	S
<i>reitteri</i> (Weise, 1884)	?S	= <i>rugulosa</i> (Suffrian, 1851)	
= <i>lurida</i> (Linnaeus, 1767) nec (Scopoli, 1763)		<i>Protorina</i> Weise, 1894	
<i>Stichoptera</i> Motschulsky, 1860		<i>plagiata</i> (Suffrian, 1861)	?M S
<i>gypsophilae gypsophilae</i> (Küster, 1845)	B M S	<i>Chrysochloa</i> Hoppe, 1840	
<i>kuesteri kuesteri</i> (Helligsen, 1912)	B M S	<i>cacaliae marani</i> (Fassati, 1961)	B
= <i>marginalis</i> auct.		= <i>cacaliae</i> var. <i>bohemica</i> (Weise, 1889)	
= <i>sanguinolenta</i> (Küster, 1845) nec (Linnaeus, 1758)		<i>cacaliae senecionis</i> (Schummel, 1843)	B M S
<i>rossia</i> (Illiger, 1802)	M S	<i>speciosissima speciosissima</i> (Scopoli, 1763)	B M S
<i>sanguinolenta</i> (Linnaeus, 1758)	B M S	<i>speciosissima ignita</i> (Letzner, 1852)	B ?M
<i>Anopachys</i> Motschulsky, 1860		<i>speciosissima juncorum</i> (Suffrian, 1851)	?M S
<i>aurichalcea bohemica</i> G. Müller, 1948	B M S	<i>virgulata</i> (Germar, 1824)	?B M S
<i>perplexa</i> (Breit, 1920)	?S	<i>Colaphus</i> Dahl, 1823	
<i>Pleurosticha</i> Motschulsky, 1860		= <i>Colaphellus</i> Weise, 1916	
<i>morio</i> (Krynický, 1832)	S	= <i>Colaspidema</i> L. Redtenbacher, 1845	
= <i>susterai</i> Bechyně, 1950		<i>sophiae</i> (Schaller, 1783)	B M S
<i>Sphaeromela</i> Bedel, 1899		<i>Gastrophysa</i> Chevrolat in Dejean, 1837	
<i>varians</i> (Schaller, 1783)	B M S	= <i>Gastroidea</i> Agassiz, 1846	
= <i>lichenis havelkai</i> Bechyně, 1949		<i>polygoni</i> (Linnaeus, 1758)	B M S
<i>Hypericia</i> Bedel, 1899		<i>viridula</i> (De Geer, 1775)	B M S
<i>cuprina cuprina</i> (Duftschmid, 1825)	B M S	<i>Phaedon</i> Dahl, 1823	
<i>didymata didymata</i> (L. G. Scriba, 1791)	?S	<i>Phaedon</i> s. str.	
<i>geminata</i> (Paykull, 1799)	B M S	<i>armoraciae</i> (Linnaeus, 1758)	B M S
= <i>cuprina</i> auct. nec (Duftschmid, 1825)		<i>cochleariae</i> (Fabricius, 1792)	B M S
<i>hyperici hyperici</i> (Forster, 1771)	B M S	<i>laevigatus</i> (Duftschmid, 1825)	?B M S
		<i>Neophaedon</i> Jacobson, 1901	
		<i>pyritosus</i> (Rossi, 1792)	B M S

<i>Sternoplatus</i> Motschulsky, 1860				<i>Spartoxena</i> Motschulsky, 1860			
<i>segnis</i> (Weise, 1884)	?B	M	S	= <i>Spartomena</i> Reitter, 1912			
<i>Sclerophaedon</i> Weise, 1892	B	M	S	<i>fornicata</i> (Brüggemann, 1873)	B	M	S
<i>carniolicus</i> (Germar, 1824)				<i>Phratora</i> Chevrolat, 1843			
<i>carpathicus</i> (Weise, 1875)			S	= <i>Phyllodecta</i> Kirby, 1837			
<i>orbicularis</i> (Suffrian, 1851)	B	M	S	<i>Phratora</i> s. str.			
<i>Hydrothassa</i> C. G. Thomson, 1866	B	M	S	<i>atrovirens</i> Cornelli, 1857	B	M	S
<i>glabra</i> (Herbst, 1783)				<i>laticollis</i> Suffrian, 1851	B	M	S
= <i>aucta</i> (Fabricius, 1787)				<i>tibialis tibialis</i> (Suffrian, 1851)	B	M	S
<i>hannoverana</i> (Fabricius, 1775)		M	S	<i>tibialis cornelli</i> Weise, 1882	B	M	S
<i>marginella</i> (Linnaeus, 1758)	B	M	S	= <i>viennensis</i> Weise, 1882			
<i>Prasocuris</i> Latreille, 1802	B	M	S	<i>vitellinae</i> (Linnaeus, 1758)	B	M	S
<i>juncii</i> (Brahm, 1790)				<i>Chaeroceta</i> Mroczkowski, 1990			
<i>phelandrii</i> (Linnaeus, 1758)	B	M	S	<i>vulgatissima</i> (Linnaeus, 1758)	B	M	S
<i>Plagioderma</i> Chevrolat in Dejean, 1837	B	M	S	<i>Entomoscelis</i> Chevrolat, 1843			
<i>versicolora</i> (Laicharting, 1781)				<i>adonis</i> (Pallas, 1771)	B	M	S
<i>Chrysomela</i> Linnaeus, 1758				<i>sacra</i> (Linnaeus, 1758)	?M	S	
= <i>Lina</i> L. Redtenbacher, 1849							
= <i>Melasoma</i> Stephens, 1834							
<i>Chrysomela</i> s. str.							
<i>populi</i> Linnaeus, 1758	B	M	S	<i>G a l e r u c i n a e</i>			
<i>saliceti</i> (Weise, 1884)	B	M	S	<i>GALERUCINI</i>			
<i>tremulae</i> Fabricius, 1787	B	M	S	<i>Galerucella</i> Crotch, 1873			
<i>Pachylina</i> Medvedev, 1969	B	M	S	<i>Galerucella</i> s. str.			
<i>collaris</i> Linnaeus, 1758				<i>fergussoni</i> Fowler, 1910	B		
<i>Strickerus</i> R. Lucas, 1920	B	M	S	<i>grisescens</i> Joannis, 1866	?B		
= <i>Microdera</i> Stephens, 1839 nec Eschscholz, 1831	B	M	S	<i>nymphaeae</i> (Linnaeus, 1758)	B	M	S
<i>cuprea</i> Fabricius, 1775				<i>sagittariae</i> (Gyllenhal, 1813)	B	?M	S
<i>lapponica</i> Linnaeus, 1788	B	M	S	<i>Neogalerucella</i> Chujô, 1962			
<i>viginipunctata</i> (Scopoli, 1763)	B	M	S	<i>calmariensis</i> (Linnaeus, 1767)	B	M	S
<i>Linaeidea</i> Motschulsky, 1860	B	M	S	<i>lineola</i> (Fabricius, 1781)	B	M	S
<i>aenea</i> (Linnaeus, 1758)				<i>pusilla</i> (Duftschmid, 1825)	B	M	S
<i>Gonioctena</i> Chevrolat, 1843				<i>tenella</i> (Linnaeus, 1761)	B	M	S
= <i>Phytodecta</i> Kirby, 1837				<i>Pyrrhalta</i> Joannis, 1866			
<i>Gonioctena</i> s. str.				<i>viburni</i> (Paykull, 1799)	B	M	S
<i>decemnotata</i> (Marsham, 1802)	B	M	S	<i>Xanthogaleruca</i> Laboissière, 1922			
= <i>rufipes</i> (De Geer, 1775) nec (Linnaeus, 1758)				<i>luteola</i> (O. F. Müller, 1776)	M	S	
<i>flavicornis</i> (Suffrian, 1851)	B	M	S	<i>Lochmaea</i> Weise, 1883			
<i>kaufmanni</i> (Miller, 1881)			S	<i>capreae</i> (Linnaeus, 1758)	B	M	S
<i>linnaeana</i> (Schrank, 1781)	B	M	S	<i>crataegi</i> (Forster, 1771)	B	M	S
<i>nivosa</i> (Suffrian, 1851)	?B			<i>suturalis</i> (C. G. Thomson, 1866)	B	M	S
<i>vininalis</i> <i>vininalis</i> (Linnaeus, 1758)	B	M	S	<i>Galeruca</i> Geoffroy, 1762			
<i>Spartophila</i> Stephens, 1834	B	M	S	= <i>Adimonia</i> Laicharting, 1781			
<i>olivacea</i> (Forster, 1771)				= <i>Ademonia</i> Agassiz, 1846			
<i>Goniomena</i> Motschulsky, 1860	B	M	S	<i>Galeruca</i> s. str.			
<i>intermedia</i> Helliessen, 1911	B			<i>dahli dahli</i> Joannis, 1866		S	
<i>interposita</i> (Franz et Palmén, 1950)	B	M	S	<i>interrupta interrupta</i> Illiger, 1802	?B	S	
<i>pallida carpathica</i> (Bechyně, 1947)				<i>interrupta circumdata</i> Duftschmid, 1825	B	M	S
<i>pallida nigricolor</i> (Reitter, 1914)			S	<i>laticollis laticollis</i> C. R. Sahlberg, 1837	B	M	S
<i>pallida sudetica</i> (Bechyně, 1947)	B	M		<i>pomonae pomonae</i> (Scopoli, 1763)	B	M	S
<i>pallida vulgaris</i> (Bechyně, 1947)	B	M		<i>tanaceti tanaceti</i> (Linnaeus, 1758)	B	M	S
<i>quinquepunctata</i> (Fabricius, 1787)	B	M	S	<i>Haptoscelis</i> Weise, 1886			
				<i>melanocephala melanocephala</i> Ponza, 1805	B		S
				<i>Emarhopa</i> Weise, 1886			
				<i>rufa</i> (Germar, 1824)	?B	M	S

SERMYLASSINI

=*Sermylini*
=*Agelasini*

Sermyllassa Reitter, 1912

=*Sermyla* Chapuis, 1875 nec H. et A. Adams, 1854
halensis (Linnaeus, 1767)

B M S

*AGELASTICINI**Agelastica* Chevrolat, 1837
alni (Linnaeus, 1758)

B M S

*LUPERINI**Phylllobrotica* L. Redtenbacher, 1845*adusta* (Creutzer, 1799)

M S

quadrimaculata (Linnaeus, 1758)

B M S

Euluperus Weise, 1886*xanthopus* (Duftschmid, 1825)

S

Calomicrus Stephens, 1834*circumfusus* Marsham, 1802

B M S

pinicola (Duftschmid, 1825)

B M S

Luperus Geoffroy, 1762=*Lyperus* Billberg, 1820*flavipes* (Linnaeus, 1767)

B M S

longicornis (Fabricius, 1781)

B M S

lyperus (Sulzer, 1776)

B M S

saxonicus (Gmelin, 1790)

B

viridipennis (Germar, 1824)

B M S

xanthopoda (Schrank, 1781)

B M S

*Alticinae**Phyllotreta* Stephens, 1836*aerea* Allard, 1859

B M S

armoraciae (Koch, 1803)

B M S

astrachanica Lopatin, 1977

B M S

=*diademata astrachanica* Lopatin, 1977*atra* (Fabricius, 1775)

B M S

austriaca Heikertinger, 1909

B M S

christinae Heikertinger, 1941

B S

cruciferae (Goeze, 1777)

B M S

diademata Foudras, 1859

?B ?M S

dilatata C. G. Thomson, 1866

B M S

exclamationis (Thunberg, 1784)

B M S

flexuosa (Illiger, 1794)

B M S

hochetlingeri Fleischer, 1917

S

nemorum (Linnaeus, 1758)

B M S

nigripes (Fabricius, 1775)

B M S

nodicornis (Marsham, 1802)

B M S

ochripes (Curtis, 1837)

B M S

procera (L. Redtenbacher, 1849)

B M S

scheuchi Heikertinger, 1941

B M S

striolata (Fabricius, 1803)

B M S

=*sinuata* (L. Redtenbacher, 1849)

B M S

=*vittata* (Fabricius, 1801) nec (Fabricius, 1775)

B M S

tetrastigma (Cornolli, 1837)

B M S

undulata Kutschera, 1860

B M S

vittula (L. Redtenbacher, 1849)

B M S

Aphtona Chevrolat, 1842*abdominalis* (Duftschmid, 1825)

M S

aeneomicans Allard, 1875

S

=*metallica* Weise, 1889*atrovirens* Förster, 1849

B M S

=*orientalis* Roubal, 1936 nec Rey, 1873*pieninensis* Roubal, 1936*beckeri* Jacobson, 1895

S

cyanella (L. Redtenbacher, 1849)

B M S

cyparissiae (Koch, 1803)

B M S

czwalinai czwalinai Weise, 1888

B M S

euphorbiae (Schrank, 1781)

B M S

flava Guillebeau, 1894

S

herbigrada (Curtis, 1837)

B M S

lacertosa Rosenhauer, 1847

M S

lutescens lutescens (Gyllenhal, 1808)

B M S

=*sternalis* Rey, 1873*nigriscutis* Foudras, 1859

B M S

nonstriata (Goeze, 1777)

B M S

=*coerulea* (Geoffroy, 1785)*ovata* Foudras, 1859

B M S

pallida (Bach, 1856)

B M S

pygmaea pygmaea Kutschera, 1861

B M S

semicyanea semicyanea Allard, 1859

B M S

venustula venustula Kutschera, 1861

B M S

violacea (Koch, 1803)

M S

Longitarsus Latreille, 1825*absynthii* Kutschera, 1862

B M S

albineus (Foudras, 1859)

S

anchusae (Paykull, 1799)

B M S

apicalis (Beck, 1817)

B M S

=*analis* (Duftschmid, 1825)*atricillus* (Linnaeus, 1761)

B M S

ballotae (Marsham, 1802)

B M S

brunneus (Duftschmid, 1825)

B M S

cerinthes (Schrank, 1798)

B M S

curtus (Allard, 1860)

B M S

echii (Koch, 1803)

B M S

exoletus exoletus (Linnaeus, 1758)

B M S

exoletus rufulus (Foudras, 1859)

S

ferrugineus (Foudras, 1859)

B M S

foudrasi Weise, 1893

B M S

fowleri Allen, 1967

S

fulgens (Foudras, 1859)

B M S

fuscoaeneus fuscoaeneus

L. Redtenbacher, 1849

M S

fuscoaeneus polonicus

Warchałowski, 1967

?M ?S

ganglbaueri Heikertinger, 1912

B M S

gracilis Kutschera, 1864

B M S

<i>holsticus</i> (Linnaeus, 1758)	B	M	S	<i>Altica</i> Geoffroy, 1762			
<i>jacobaeae</i> (Waterhouse, 1858)	B	M	S	= <i>Haltica</i> Koch, 1803			
<i>kutscherae</i> Rye, 1872	B	M		<i>aenescens</i> (Weise, 1888)	B	M	
<i>languidus</i> Kutschera, 1863	B	M	S	<i>brevicollis brevicollis</i> (Foudras, 1859)	M	S	
<i>lateripunctatus personatus</i> Weise, 1893	M	S		<i>brevicollis coryletorum</i> Král, 1964	M	S	
<i>linnaei</i> (Duftschmid, 1825)	S			= <i>quercetorum</i> (Weise, 1888) nec (Foudras, 1859)			
<i>longipennis</i> Kutschera, 1863	B	M	S	<i>carduorum cirsii</i> (Israelson, 1956)	?B	S	
<i>longiseta</i> Weise, 1889	B	M	S	<i>chamaenerii</i> (H. Lindberg, 1926)		S	
<i>luridus luridus</i> (Scopoli, 1763)	B	M	S	<i>cornivorax</i> Král, 1969		S	
<i>lycopi lycopi</i> (Foudras, 1859)	B	M	S	<i>impressicollis</i> (Reiche, 1862)	M	S	
<i>medvedevi</i> Shapiro, 1956	B	M	S	<i>lythri</i> Aubé, 1843	B	M	S
<i>melanocephalus melanocephalus</i> (De Geer, 1775)	B	M	S	<i>oleracea oleracea</i> (Linnaeus, 1758)	B	M	S
<i>membranaceus</i> (Foudras, 1859)	B	M	S	<i>oleracea breddini</i> (Mohr, 1958)	B	M	S
<i>minimus</i> Kutschera, 1864	B	M	S	<i>palustris</i> (Weise, 1888)	B	M	S
<i>minusculus</i> (Foudras, 1859)	B	M	S	<i>pusilla</i> (Duftschmid, 1825)	B	M	S
<i>nanus</i> (Foudras, 1859)	B	M	S	<i>quercetorum quercetorum</i> Foudras, 1859		S	
<i>nasturtii</i> (Fabricius, 1792)	B	M	S	<i>quercetorum saliceti</i> (Weise, 1888)	B	M	S
<i>niger</i> (Koch, 1803)	M	S		<i>tamaricis tamaricis</i> (Schrank, 1785)	B	M	S
<i>nigerrimus</i> (Gyllenhal, 1827)	B			<i>Hermeophaga</i> Foudras, 1859			
<i>nigrofasciatus nigrofasciatus</i> (Goeze, 1777)	B	M		<i>mercurialis</i> (Fabricius, 1792)	B	M	S
<i>noricus</i> Leonardi, 1976	B	M	S	<i>Batophila</i> Foudras, 1859			
<i>obliteratus</i> (Rosenhauer, 1847)	B	M	S	<i>aerata</i> (Marsham, 1802)		?S	
<i>ochroleucus</i> (Marsham, 1802)	B	M	S	<i>rubi</i> (Paykull, 1790)	B	M	S
<i>pallidicornis</i> Kutschera, 1863	M	S		<i>Ochrosis</i> Foudras, 1859			
= <i>hubenthali</i> Wanka, 1917				<i>ventralis ventralis</i> (Illiger, 1807)	?B	M	S
<i>parvulus</i> (Paykull, 1799)	B	M	S	<i>Lythraria</i> Bedel, 1897			
<i>pellucidus</i> (Foudras, 1860)	B	M	S	<i>salicariae</i> (Paykull, 1800)	B	M	S
<i>pinguis</i> Weise, 1888	S			<i>Asiorestia</i> Jacobson, 1925			
<i>pratensis</i> (Panzer, 1794)	B	M	S	= <i>Crepidodera</i> auct. nec Chevrolat, 1837			
<i>pulmonariae</i> Weise, 1893	B	M	S	<i>brevicollis</i> (J. Daniel, 1904)	B	S	
<i>quadriguttatus</i> (Pontoppidan, 1765)	B	M	S	<i>corpulenta</i> (Kutschera, 1860)		S	
= <i>quadripustulatus</i> (Fabricius, 1775)				<i>cyanescens</i> (Duftschmid, 1825)	M	S	
= <i>quadripunctatus</i> (Fourcroy, 1785)				<i>cyanipennis puncticollis</i> (Reitter, 1897)	S		
<i>reichei</i> (Allard, 1860)	M	S		= <i>picea</i> (Petri, 1907)			
<i>rubellus</i> (Foudras, 1859)	M	S		= <i>alpina marani</i> (Roubal, 1939)			
<i>rubiginosus</i> (Foudras, 1859)	B	M	S	<i>femorata</i> (Gyllenhal, 1813)	B	M	S
<i>salviae</i> Gruev, 1975	B	M	S	<i>ferruginea</i> (Scopoli, 1763)	B	M	S
= <i>obliteratus</i> auct. nec (Rosenhauer, 1847)				<i>melanostoma</i> (L. Redtenbacher, 1849)		S	
<i>scutellaris</i> (Rey, 1847)	B	M	S	<i>nigritula</i> (Gyllenhal, 1813)	B	M	S
<i>substriatus</i> Kutschera, 1863	M	S		<i>transsilvanica</i> Fuss, 1864		S	
<i>succineus succineus</i> (Foudras, 1859)	B	M	S	<i>sublaevis</i> (Motschulsky, 1859)	B	M	S
<i>suturalis suturalis</i> (Marsham, 1802)	S			<i>transversa</i> (Marsham, 1802)	B	M	S
<i>suturellus</i> (Duftschmid, 1825)	B	M	S	<i>Orestia</i> Gemar, 1845			
<i>sympyti</i> Heikertinger, 1912	B	M	S	<i>aubei arcuata</i> Miller, 1868		S	
<i>tabidus</i> (Fabricius, 1775)	B	M	S	<i>carpathica</i> Reitter, 1880		S	
<i>tristis</i> Weise, 1888	M			<i>Derocrepis</i> Weise, 1886			
= <i>pannonicus</i> Kaszab, 1962				<i>rufipes</i> (Linnaeus, 1758)	B	M	S
<i>waterhousei</i> Kutschera, 1864	B	M	S	<i>Hippuriphila</i> Foudras, 1859			
				<i>modeeri</i> (Linnaeus, 1761)	B	M	S

<i>isatidis</i> Heikertinger, 1912	B	M	S	<i>nebulosa</i> Linnaeus, 1758	B	M	S
<i>luteola</i> (O. F. Müller, 1776)	B	M	S	<i>pannonica</i> Suffrian, 1844	B	M	S
= <i>pallida</i> (Stephens, 1831)				<i>panzeri</i> Weise, 1907	B	M	S
<i>napi</i> (Fabricius, 1792)	B	M	S	<i>prasina prasina</i> Illiger, 1798	B		S
= <i>rapae</i> (Illiger, 1807)	B	M	S	= <i>chloris</i> Suffrian, 1844			
<i>picina picina</i> (Marsham, 1802)	B	M	S	<i>rubiginosa</i> O. F. Müller, 1776	B	M	S
= <i>picea</i> L. Redtenbacher, 1849				<i>rufovirens</i> Suffrian, 1844	B	M	S
<i>sophiae</i> (Illiger, 1807)	B	M	S	<i>sanguinolenta</i> O. F. Müller, 1776	B	M	S
= <i>cyanoptera</i> L. Redtenbacher, 1849				<i>sanguinosa</i> Suffrian, 1844	B	M	S
<i>subaenea subaenea</i> Kutschera, 1864		S		<i>seladonia</i> Gyllenhal, 1827		M	
<i>thlaspis</i> Foudras, 1859	B	S		<i>stigmatica</i> Suffrian, 1844	B	M	S
<i>toelgi</i> Heikertinger, 1914	B			<i>vibex</i> Linnaeus, 1767	B	M	S
<i>vindobonensis</i> Heikertinger, 1914	M	S		<i>Cassiduella</i> E. Strand, 1928			
H i s p i n a e				<i>nobilis</i> Linnaeus, 1758	B	M	S
<i>Hispa</i> Linnaeus, 1767				<i>vittata</i> Villers, 1789	B	M	S
= <i>Hisrella</i> Chapuis, 1875				<i>Pseudocassida</i> Desbrochers, 1891			
<i>atra</i> Linnaeus, 1767	B	M	S	<i>murræa murræa</i> Linnaeus, 1767	B	M	S
<i>Diaclispa</i> Gestro, 1897				<i>Mionycha</i> Weise, 1891			
<i>testacea</i> (Linnaeus, 1767)		?S		<i>azurea</i> Fabricius, 1801	?B	M	S
C a s s i d i n a e				<i>margaritacea</i> Schaller, 1783	B	M	S
<i>Cassida</i> Linnaeus, 1758				<i>subreticulata</i> Suffrian, 1844	M	S	
<i>Cassida</i> s. str.				= <i>splendidula</i> Suffrian, 1844			
<i>atrata</i> Fabricius, 1787	?B	M	S	<i>Lordiconia</i> Reitter, 1926			
<i>aurora</i> Weise, 1907		M	S	<i>canaliculata</i> Laicharting, 1781	?B	M	S
<i>berolinensis</i> Suffrian, 1844	B	M	S	= <i>austriaca</i> Fabricius, 1792			
<i>denticollis</i> Suffrian, 1844	B	M	S	<i>Odontionycha</i> Weise, 1891			
<i>ferruginea</i> Goeze, 1777	B	M	S	<i>viridis</i> Linnaeus, 1758	B	M	S
<i>flaveola</i> Thunberg, 1794	B	M	S	<i>Mionychella</i> Spaeth, 1952			
<i>inquinata</i> Brullé, 1832		M	S	<i>hemisphaerica</i> Herbst, 1799	B	M	S
<i>lineola</i> Creutzer, 1799	B	M	S				

Bruchidae

(J. STREJČEK)

The list corresponds with the recent paper on *Bruchidae* of Czechoslovakia by Strejček (1991). Following species were previously reported from Czechoslovakia. Some of them were erroneously identified, the other ones are not documented with relevant collection specimens and their occurrence in Czechoslovakia does not comport with their geographic distributions: *Bruchus emarginatus* Allard, *B. laticollis* Boh., *Bruchidius annulicornis* (Allard), *B. bimaculatus* (Ol.), *B. nanus* (Germar).

Seznam koresponduje s nedávným zpracováním československých druhů čeledi *Bruchidae* (Strejček, 1991). V literatuře jsou z území Československa uváděny i některé další druhy, u kterých byla při revizi zjištěna chybná determinace. Dále jsou to druhy, u kterých nebyl zjištěn žádný doklad a jejichž výskyt neodpovídá jejich známému rozšíření. Takovéto druhy byly ze seznamu vypuštěny. Jsou to *Bruchus emarginatus* Allard, *B. laticollis* Boh., *Bruchidius annulicornis* (Allard), *B. bimaculatus* (Ol.) a *B. nanus* (Germar).