

Recent advances in the knowledge of Mexican Alticinae (Coleoptera, Chrysomelidae)¹

David G. Furth¹

¹ Department of Entomology, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20560, USA

Corresponding author: David G. Furth (furthd@si.edu)

Academic editor: M. Schmitt | Received 23 July 2017 | Accepted 20 September 2017 | Published 11 December 2017

<http://zoobank.org/142AC5A7-981F-417F-B04E-02304FBEE985>

Citation: Furth DG (2017) Recent advances in the knowledge of Mexican Alticinae (Coleoptera, Chrysomelidae). In: Chaboo CS, Schmitt M (Eds) Research on Chrysomelidae 7. ZooKeys 720: 23–46. <https://doi.org/10.3897/zookeys.720.17790>

Abstract

The present study updates previously published biodiversity/faunistics of the flea beetles of Mexico published by the author after examination of 6132 specimens from 8 institutional collections. The following 9 genera were selected as indicators of the effects of known diversity only through examination of museum specimens (i.e., “indoor collecting”): *Alagoasa* Bechyné; *Asphaera* Chevrolat; *Capraita* Bechyné; *Disonycha* Chevrolat; *Kuschelina* Bechyné; *Omophoita* Chevrolat; *Prasona* Baly; *Systema* Chevrolat; and *Waltherianella* Bechyné. From the specimens examined in these genera from the 8 collections, there were 394 new records for Mexican states of the 287 new species records representing 47% new records of the species recorded from those states. Total new state records 287 from 80 species. States with most new records: Chiapas (32); Nayarit (27); Sinaloa (24). 80 spp. (47%) with new state records. *Systema oberthuri* Baly is reported from Mexico for the first time. The current total of Alticinae in Mexico is 90 genera/626 species. The difficulties of the generic boundaries between *Systema* and *Prasona*, *Alagoasa* and *Kuschelina*; as well as the specific boundaries between *A. jacobiana* and *A. decemguttatus* and the specific level pattern variation in *Disonycha glabrata* and *Alagoasa decemguttatus* are discussed. *Kuschelina semipurpurea*, formerly placed in *Alagoasa*, is placed is considered as a new combination.

Keywords

Mexico state records, Flea Beetles, indoor collecting, faunistics, new combinations, *Systema/Prasona* and *Alagoasa/Kuschelina* confusion, *Alagoasa decemguttatus*, *Omophoita octomaculata*, Mexican biodiversity

¹ Contribution to the 9th International Symposium on the Chrysomelidae, Orlando, FL, USA, September 28, 2016

Introduction

For about 30 years I have been studying the biodiversity/faunistics of the Central American Alticinae (Furth and Savini 1996, 1998, Furth et al. 2003), especially the Mexican Alticinae (Furth 2004, 2006, 2009, 2013). The current project is a continuation of research about the biodiversity/faunistics, distribution, and biogeography of the Alticinae (Flea Beetles) of Mexico. The author has conducted extensive field work in the majority of the 32 Mexican states as well as examination and determination of thousands of Mexican specimens from many museums in North America. Two previous published surveys by the author were of individual selected states. The first one from Chihuahua and Sonora (Furth 2009) revealed 26 genera with 70 species from Chihuahua, including 44 species new to Chihuahua, 9 species new to Sonora, 10 species new to Mexico, and two species new to science. The second from Oaxaca (Furth 2013) discovered 68 genera with 275 species (113 species known only from Oaxaca) in that state alone elevating the total recorded Alticinae fauna of Mexico to 90 genera and 625 species. This Oaxaca study demonstrated a very strong biogeographic affinity to the Neotropical Region. Some of the species in this Oaxaca study were at the time unidentifiable morpho-species and some either far from their known distribution or even new to science. The last comprehensive list of the entire Mexican Alticinae fauna was over 10 years ago and a lot of new specimen data has become accessible. There are still many more specimens in the remaining Alticinae genera currently being studied by the author from the institutions mentioned in the current study as well as others in a few other North American and Mexican institutions. Some details of the Mexican Alticinae fauna as well as about Mexican biogeography were included in Furth (2006) which recorded 89 genera and 524 species of Alticinae in Mexico.

Some of the author's recent publications (mentioned above) concerning the Mexican fauna stressed the value of a combination of strategies for biodiversity research beginning with a comprehensive search of the literature examination, combined with fieldwork and extensive search of institutional collections. The current study concentrates on the latter of these strategies to uncover additional specimens and localities in order to better understand Mexican flea beetle diversity.

Methods

During the past 10+ years the author has visited the institutions listed below and has pulled out all Alticinae from Mexico and borrowed them. For the current study nine genera, including the genera of the “Oedionychini/a” tribe/subtribe (*Alagoasa*, *Asphaera*, *Capraita*, *Kuschelina*, *Omophoita*, *Walterianella*), as well as a few other randomly selected genera (*Disonycha*, *Prasona*, *Systema*) were studied and determined to species. There were 6625 adult specimens studied and determined to species based on morphology and using literature and reference collections. The term species refers also to all taxa, including subspecies recorded from Mexico.

The institutional collections studied are: American Museum of Natural History (New York, New York, USA) [AMNH]; Monte L. Bean Museum of Brigham Young University (Provo, Utah, USA) [BYU]; Natural History Museum (London, United Kingdom) [NHM]; California Academy of Sciences (San Francisco, California, USA) [CAS]; California Department of Food and Agriculture, Sacramento, California, USA [CDEA]; University of California Berkeley (Berkeley, California, USA) [UCB]; University of California Davis, Davis, California, USA [UCD]; U.S. National Museum/NMNH (Washington, D.C., USA) [USNM]; and a few specimens donated to the author by R. Wills Flowers (Florida State University, Tallahassee, Florida, USA) [RWF]. Specimens were studied and determined using a Leica MZ-APO dissecting stereomicroscope, as well as specimen reference collections at the U. S. National Museum of Natural History, Smithsonian Institution, Washington, D.C. USA [USNM/NMNH], relevant literature in the author's library and certain online references, e.g., MCZ Type Database.

The habitus photographic images were taken using the Visionary Digital BK Lab Imaging system outfitted with the Canon EOS 5D and a MP-E 65 mm 1–5× Canon macrolens. Stacked images were processed in part with Helicon Focus; final editing was done with Adobe PhotoShop. Specimens will be returned to their original institutional collections with some vouchers deposited at the USNM.

Images of adult species for Figures 4–6, 8–9 were taken randomly from the Internet by searching for the species name. Figures 1–3 were taken by the author; Figure 7 was taken by K. Darrow.

In Table 2 (totaled in Table 1) there are some new state records from multiple institutional collections, but each new state record is only counted once (e.g., for *Alagoasa acutangula* Nayarit is a new state record and was recorded from AMNH, CAS, CDEA, and UCB, but it is only counted as a single state record). The taxa in Table 2 are referred to in the text as species, but a few are subspecies names.

The author follows a less popular classification, i. e., Alticinae rather than Alticini, as explained in several publications, e.g., Furth and Lee (2000), Furth and Suzuki (1998), Mohamedsaid and Furth (2011).

Table 1. Indoor Collecting. Numbers of specimens examined, by institutional collection. For Specimens Examined, the number in parentheses are those not determined to species.

Collection	Specimens Examined	New State Records/Collection
AMNH	374 (9)	30
NHM	120 (14)	1
BYU	309 (33)	29
CAS	1412 (93)	105
CDEA	374 (20)	27
UCB	3131 (223)	150
UCD	846 (97)	46
USNM	59 (4)	6
TOTALS	6625 (493)	394/287*

*Repeats/Actual

Table 2. List of determined species by institution with state records (new state records in bold print).

Genus	Species	Author	Distribution	AMINH	BMINH	BYU	CAS	CDEFA	NHIM	UCB	UCD	RWF	USNM	New Records	No. New Records
	<i>acutangula</i>	(Jacoby)	CHIS, COL, DGO, GRO, JAL, MEX, MOR, NL, OAX, VER	NAY		GRO, JAL,	CHIS, GRO, MOR, NAY, SIN, SLP	JAL, MICH, NAY	VER	CHIS, DGO, GRO, JAL, MICH, MOR, NAY, NL, PUE, SIN, VER	GRO, MOR, PUE, SIN		GRO, QRO	MICH, NAY, PUE, QRO, SIN, SLP	6
	<i>bipunctata</i>	(Chevrolat)	CHIS, DF, OAX, SLP, VER, YUC	CHIS, SLP, QRO, TAMPS, VER, YUC		PUE, SLP	VER		MOR, VER	CHIS, QRO, SLP, VER	SLP			MOR, PUE, QRO, TAMPS	5
	<i>chevrolati</i>	(Baly)	OAX, VER, YUC				VER		VER						
	<i>elypata</i>	(Jacoby)	CHIS, DGO, FGO, MICH, OAX, TAB, VER			VER		CHIS, VER	VER	CHIS, DGO, OAX, VER	CHIS, VER			DGO	1
	<i>donckieri</i>	(Jacoby)	GRO					PUE						PUE	1
	<i>extrema</i>	(Harold)	MOR, OAX, TAB, VER	CHIS, VER			CHIS		VER	VER	VER		YUC	CHIS, YUC	2
	<i>fimbriata</i>	(Forster)	GRO, MICH, MOR, OAX				MICH								
	<i>hoegeri</i>	(Jacoby)	OAX, VER				VER								
	<i>inconspicua</i>	(Jacoby)	DGO, JAL	JAL		JAL	SIN			DGO, JAL, SIN				SIN	1
	<i>jaobiana</i> * [comb. n.]	(Horn)		CHIH, CHIS, DGO, JAL, MOR, NAY, OAX, SON, SIN, YUC		GRO, SON	CHIS, MOR, NAY, SIN, SON	COL, JAL, NAY, SIN	MEX	CHIH, CHIS, DGO, GRO, JAL, NAY, OAX, SIN, SON	CHIH, JAL, MOR, SIN			SIN, YUC	2
	<i>lateralis</i>	(Jacoby)	COL, GRO, JAL, MEX, MICH, MOR, NAY, OAX	NAY		PUE	CHIH, COL, MOR, NAY, NL, SIN, VER	COL, JAL, NAY, SIN	VER	GRO, MOR, NAY, SIN, VER	CHIS, MOR			CHIH, CHIS, COL, NL, SIN, VER	6
	<i>longicollis</i>	(Jacoby)	OAX				VER							VER	1
	<i>semipurpurea</i> **	(Jacoby)	VER											CHIH	1
	<i>seriata</i>	(Baly)	GRO, MOR, OAX, PUE, VER	MOR			MOR, SLP, TAMPS, VER			CHIS, GRO				CHIS, SLP, TAMPS	3

Alagausa

Genus	Species	Author	Distribution	AMNH	BMINH	BYU	CAS	CDEA	NHM	UCB	UCD	RWF	USNM	New Records	No. New Records
<i>Diomphya</i>	<i>antennata</i>	Jacoby	COL, DGO, GRO, JAL, MEX, MICH, MOR, OAX, VER		MEX		MICH, MOR, NAV, SIN, VER	SIN		CHIS, GRO, JAL, MICH, NAV, OAX, SIN, VER	CHIH, MICH, SIN, VER			CHIH, CHIS, NAV, SIN	4
	<i>arizonae</i>	Casey	CHIH, DGO?, GRO, MOR							CHIH, COAH, NL, SON	CHIH, NL			COAH, NL, SON	3
	<i>barberi</i>	Blake	GRO, SIN, VER				SIN	MICH		NAV, VER	SIN			MICH, NAV	2
	<i>brevilineata</i>	Jacoby	DGO, GRO, JAL, MOR, OAX		MEX		CHIS, MOR	JAL		CHIS, JAL, NAV				CHIS, MEX, NAV	3
	<i>brunneofasciata</i>	Jacoby	GRO, PUE, SLP							OAX, SIN				OAX, SIN	2
	<i>collata</i>	(Fabricius)	CHIH, COAH, DE, DGO, GTO, JAL, MEX, MICH, MOR, OAX, PUE, TAB, VER, YUC				DGO, JAL, MEX, MOR, NAV, PUE, SIN, SON	SIN, VER		DE, JAL, NAV, SIN	NAV, SIN			NAV, SIN, SON	3
	<i>dorsata</i>	Harold	MOR, OAX, TAB, VER, YUC			VER	CHIS, JAL, MOR, NAV, VER	JAL		CHIS, NAV, OAX, SLP, VER				CHIS, JAL, MOR, NAV, SLP	5
	<i>figurata</i>	Jacoby	AGS, CHIH, CHIS, COAH, COL, DE, DGO, GRO, GTO, JAL, MEX, MICH, MOR, NAV, OAX, SIN, TAB, VER, YUC	CHIH	MEX	CHIH, JAL, SON	CHIH, MOR, NAV, SIN	JAL, MOR, SIN		CHIH, CHIS, COL, DE, DGO, GRO, GTO, JAL, MICH, NAV, OAX, SIN, SON, TAMPS, VER	CHIH, CHIS, JAL, MICH, NAV, SIN, ZAC			SON, TAMPS, ZAC	3
	<i>finnata finnata</i>	LeConte	BC, CHIH, CHIS, DGO, GRO, HGO, JAL, MEX, MICH, MOR, NL, OAX, PUE, SLP, SON, TAB, VER, ZAC	COL, TAB	MEX	JAL, MEX	CHIS, GRO, NAV, PUE, SIN, SON, ZAC	JAL, MOR, SIN		BCS, CHIS, COL, DGO, GRO, JAL, MICH, NAV, NL, PUE, SIN, SON, VER	CHIH, JAL			BCS, COL, NAV, SIN	4
	<i>glabrata</i>	(Fabricius)	BC, BCS, CAMP, CHIS, COL, DGO, GRO, JAL, MOR, NAV, OAX, PUE, SIN, SON, TAB, TAMPS, YUC, VER	COL, TAB	JAL, MOR	SON	BCS, CHIH, CHIS, JAL, MEX, MOR, NAV, NL, SON, SIN, SON, TAMPS, VER	COL, JAL, NAV, SIN, SON		CAMP, CHIH, CHIS, COL, GRO, GTO, JAL, MICH, MOR, NAV, NL, OAX, QROO, SIN, SON, TAB, VER, YUC	GRO, JAL, MOR, NAV, SIN, SLP, SON, VER			CHIH, GTO, MEX, MICH, NL, QROO, SLP	7
	<i>guatemalensis</i>	Jacoby	CHIS, GRO, MOR, OAX, VER				CHIS			CHIS, GRO, JAL, TAB, VER	MOR, TAMPS			JAL, TAB, TAMPS	3
	<i>jalopensis</i>	Blake	VER				CHIS			CHIS, COL, SIN	VER			CHIS, COL, SIN	3

Genus	Species	Author	Distribution	AMNH	BMINH	BYU	CAS	CDEA	NHM	UCB	UCD	RWF	USNM	New Records	No. New Records
	<i>knabi</i>	Blake	GRO							OAX, VER				OAX, VER CHIH, CHIS,	2
	<i>leptolineata texana</i>	Schaeffer	DGO, GRO, JAL, MOR, NL, OAX, QROO, TAMPS, YUC		MEX		NAY, SON			CHIH, CHIS, COL, MICH, OAX, VER	CHIS, MOR			COL, MEX, MICH, NAY, OAX, SON, VER	9
	<i>limbata</i>	Jacoby	DGO, MEX, MOR							DGO					
	<i>maculipes</i>	Jacoby	CHIS, VER	SON			BCS, DGO, SON	COAH		CHIS, TAMPS				BCS, COAH, DGO, SON, TAMPS	5
	<i>melanocephala</i>	Jacoby	VER				CHIS			CHIS, VER				CHIS	1
	<i>mexicana</i>	Jacoby	TAB, SLP, VER, YUC			TAMPS	VER			BC, CHIS, DGO, NAY, OAX, PUE, SON, TAB, VER			BC, CHIS, DGO, NAY, OAX, PUE, SON, TAMPS	8	
	<i>militaris</i>	Jacoby	TAB, VER, YUC			CHIS	CHIS			CHIS				CHIS	1
	<i>nigrita</i>	Jacoby	OAX (Furth, 2013)					COL, NAY		NAY				COL, NAY	2
	<i>pluriligata</i>	LeComte	BC, CHIH, DGO, JAL, NAY, SIN, SLP, SON, VER		JAL	CHIH, JAL, SLP, SON	CHIS, JAL, NAY, SLP, SON, TAMPS, VER			CHIS, DGO, OAX, VER	CAMP, CHIH, DGO, MEX, MICH, SIN, SON			CAMP, CHIS, MEX, MICH, OAX, TAMPS	6
	<i>politula</i>	Horn	AGS, CAMP, CHIH, DF, DGO, GRO, GTO, HGO, JAL, MEX, MOR, OAX, PUE, QRO, SLP, SON, TAMPS, VER, ZAC	VER	DGO	CHIH	AGS, CHIS, DGO, JAL, NAY, SON			AGS, CHIH, DGO, GRO, GTO, HGO, JAL, MICH, NAY, NL, PUE, SLP, SON, ZAC	CHIH, DGO, HGO, JAL, PUE, ZAC			CHIS, MICH, NAY, NL	4
	<i>proena</i>	Casey	NAY	NAY			NAY								

Disonycha

Genus	Species	Author	Distribution	AMNH	BMINH	BYU	CAS	CDEA	NHM	UCB	UCD	RWF	USNM	New Records	No. New Records
	<i>quinquevittata</i>	(Larrelle)	CHIS, COL, GRO, OAX, QROO, TAB, TAMPS, VER				JAL	JAL		VER				JAL	1
	<i>reticulolis</i>	(Jacoby)	NAY, VER				VER				VER				
	<i>sallei</i>	(Baly)	VER												
<i>Disonychia</i>	<i>scriptipennis</i>	(Jacoby)	CHIS, COL, DGO, GRO, MOR, NAY, OAX, YUC				CHIS, NAY			CHIS, JAL, MICH, SIN	GRO, JAL, NAY			JAL, MICH, SIN	3
	<i>subaenea</i>	Jacoby	DGO, GRO, MOR, OAX							JAL				JAL	1
	<i>tenicornis</i>	Horn	CHIH, DGO, HGO				CHIS, SIN	HGO, JAL, SIN		CHIH, DGO, HGO, NL, SIN				CHIS, JAL, NL, SIN	4
<i>Kuschelina</i>	<i>trifasciata</i>	Jacoby	CHIS				CHIS			CHIS					
	<i>laeta</i>	(Terbosc)	TAMPS, VER	COL, NAY		CHIH	BC, MOR, NAY, NL, VER			BCS, DGO, GTO, JAL, NAY, SIN, SON				BC, BCS, CHIH, COL, DGO, GTO, JAL, MOR, NAY, NL, SIN, SON	12
	<i>modesta</i>	(Jacoby)	CHIH, CHIS, DE, DGO, GRO, GTO, HGO, MEX, MOR, OAX, PUE, SLP, TLAX, VER			CHIH	PUE			MEX	CHIH				
<i>Omophoita</i>	<i>semipurpurea</i> ***	(Jacoby)		CHIH										CHIH	1
	<i>aequinoctialis</i>	(Linnaeus)	Mexico						TAB	TAB, VER				TAB, VER	2
	<i>aequinoctialis affinis</i> ?	(Jacoby)	Mexico ?							CHIS				CHIS	1
<i>Omophoita</i>	<i>cinetipennis</i>	(Chevrolat)	JAL, OAX, PUE, SLP, VER				NL, SLP			PUE, SLP, VER	VER			NL	1
	<i>guanipennis</i>														
	<i>ocotomaculata</i> **** Crotch (= some <i>aequinoctialis</i> & <i>punctulata</i> Bechyne & Bechyne)	(Crotch)	OAX, TAB, TAMPS, VER	CHIS, NAY, SLP, TAB, TAMPS, VER, YUC		NL, SLP, TAMPS	CHIS, MEX, NL, SLP, VER	VER	CHIS, TAB, VER	CAMP, CHIS, COAH, OAX, SLP, VER	SLP, TAMPS, VER			CAMP, CHIS, COAH, MEX, NAY, NL, SLP, YUC	8
<i>quadrinotata centraliamericana</i>	Bechyne	OAX, TAB, VER	TAB, VER			CHIS, MEX, VER		TAB, VER	CHIS, TAB, VER	VER	VER		CHIS, MEX	2	

Genus	Species	Author	Distribution	AMNH	BMINH	BYU	CAS	CDEA	NHM	UCB	UCD	RWF	USNM	New Records	No. New Records
	<i>recticollis</i>	(Baly)	CHIS, HGO, OAX, TAB, TAMPS, VER				VER			CHIS, VER					
	<i>violacea</i>	Jacoby	GRO	NAY			NAY			CHIS, JAL, MICH, NAY, SON	SIN			CHIS, JAL, MICH, NAY, SIN, SON	6
<i>Prasoma</i>	<i>viridis</i>	Baly	VER			PUE	VER	JAL?		PUE, VER				JAL, PUE	2
	<i>abbreviata</i>	Jacoby	PUE			MOR	PUE, VER			PUE				MOR, VER	2
	<i>bitaeniata</i>	LeConte	CHIH			CHIH	BC				VER			BC, VER	2
	<i>blanda</i>	Melsheimer	BC, CHIH, JAL, MICH, NL, SIN, SLP, SON, TAB, VER			CHIH, JAL	NAY, NL, PUE, SLP, SON			CHIS, JAL, MOR, NAY, SIN, SLP	CHIH, COAH, JAL, SIN, SON			CHIS, COAH, MOR, NAY, PUE, SLP	6
<i>Systema</i>	<i>contigua</i>	Jacoby	CHIS, GRO, GTO, HGO, OAX, SON, TAMPS, VER, ZAC			CHIS, HGO, SLP	CHIS, VER	JAL		NAY, NL, PUE, QRO, SIN, SON, VER				JAL, NAY, NL, PUE, QRO, SIN, SLP, SON, VER	9
	<i>discicollis</i>	Clark	CAMP, CHIH, DE, DGO, GTO, JAL, MEX, MICH, TAB, TAMPS, VER, ZAC	MICH		DGO	DE, JAL			DGO, HGO, MEX, MICH	JAL, MICH, ZAC			HGO	1

Genus	Species	Author	Distribution	AMNH	BMINH	BYU	CAS	CDEA	NHM	UCB	UCD	RWF	USNM	New Records	No. New Records
	<i>gracilentata</i>	Blake	NL	CHIH		CHIH, CHIS, SLP, SON, TAMPS	MOR, SLP, TAMPS, VER, ZAC	CHIS		CHIH, GRO, PUE, SIN, SLP, VER	CHIH, HGO, SLP, VER		QROO, ZAC	CHIH, GRO, HGO, MOR, PUE, QROO, SIN, SLP, SON, TAMPS, VER, ZAC	13
	<i>lucis</i>	Blake					BC							BC	1
	<i>marginata</i>	Jacoby	PUE, VER				CHIS							CHIS	1
	<i>nigropilgata</i>	Jacoby	AGS, CHIH, DE, DGO, GTO, GRO, JAL, MICH, MOR, OAX, PUE, VER			CHIH, CHIS, GRO, JAL, MOR, OAX, SLP, SON, VER	MOR, PUE, SIN, TAMPS, VER			DGO, GRO, GTO, JAL, MICH, MOR, PUE, SIN	CHIH, JAL, MOR, SIN			CHIS, SIN, SLP, SON, TAMPS	5
	<i>oberthuri</i>	Baly					CHIS							CHIS	1
	<i>pectoralis</i>	Clark	CHIS, GTO, OAX, VER				VER			CHIS, PUE, VER	GRO			GRO, PUE	2
	<i>s-litena</i>	(Linnaeus)	CHIS, GTO, TAB, VER			VER	VER	VER		CHIS, TAB, VER	VER				
	<i>sabini</i>	Jacoby	CHIS				CHIS				CHIS				
	<i>semitittata</i>	Jacoby	BCS, GRO, GTO, HGO, MEX, MOR, NL, OAX, SIN	CHIH		CHIH, SON	PUE	MOR		CHIH, CHIS, NAV, SON	CHIH, CHIS, JAL, NAV, PUE, SON, ZAC			CHIH, CHIS, JAL, NAV, PUE, SON, ZAC	7
	<i>subcostata</i>	Jacoby	MICH, MOR, VER				GRO	MICH		GRO, JAL, NAV, VER	MICH			GRO, JAL, NAV	3
	<i>subrigosa</i>	Jacoby	GTO, MICH, MOR				TAB	GRO		GRO, PUE	ZAC			GRO, PUE, TAB, ZAC	4
	<i>sulphurea</i>	Jacoby	CHIH, DGO, GRO, MOR, OAX							CHIH, CHIS, PUE, SON	GRO, MOR, PUE			CHIS, PUE, SON	3

Systema

Genus	Species	Author	Distribution	AMNH	BMINH	BYU	CAS	CDEA	NHM	UCB	UCD	RWF	USNM	New Records	No. New Records
<i>Systema</i>	<i>thoracica</i>	Jacoby	CAMP, HGO, PUE, QROO, TAB, VER	CHIS, MEX, VER			JAL, NAY, SLP, VER			CHIS, JAL, NAY, SIN, TAB, VER	JAL, VER			CHIS, JAL, MEX, NAY, SIN, SLP	6
	<i>undulata</i>	Jacoby	AGS, CAMP, GRO, GTO, MOR, VER			MOR	MEX			GRO, MICH				GRO, MEX, MICH, MOR	4
	<i>variabilis</i>	Jacoby	CHIH, CHIS, COL, DGO, GRO, GTO, MICH, MOR, NAY, OAX, VER		GRO, MOR, PUE, SLP, TAMPS, VER		CHIH, JAL, MOR, NAY, SIN, VER	JAL, SIN		CHIH, JAL, MICH, MOR, NAY, QRO, SIN, VER	CHIH, MICH, SIN, VER			JAL, PUE, QRO, SIN, SLP, TAMPS	6
<i>Waterianella</i>	<i>biarvata</i>	(Chevrolat)	CHIS, VER				NAY, VER		VER	VER				NAY	1
	<i>dwangoensis</i>	(Jacoby)	CHIH, DGO			SON	NAY, SON			DGO, JAL	CHIH, SIN		DGO	JAL, NAY, SIN, SON	4
	<i>inscripta</i>	(Jacoby)	OAX, SLP, VER			SLP	CHIS	VER		CHIS, VER				CHIS	1
	<i>oculata?</i>	(Fabricius)	VER				VER			VER					
	<i>signata</i>	(Jacoby)	CHIS, JAL, TAB, TAMPS, VER, YUC	QROO, VER, YUC			CHIS, MOR ?VER		VER	CHIS	SLP?			MOR, QROO, SLP?	2
	<i>sublimeata</i>	(Jacoby)	OAX, TAB, VER, YUC	QROO, VER			CHIS, VER			CHIS, SLP, VER				CHIS, QROO, SLP	3
<i>Waterianella</i>	<i>tenuicincta</i>	(Jacoby)	SLP, TAB, VER			SLP	CHIS, NAY, SLP, VER							CHIS, NAY	2
	<i>tenuistula</i>	(Schauffuss)	CHIS, COL? GRO, JAL, MICH, MOR, NAY, QROO, TAMPS, VER, YUC	VER, YUC		NAY, SLP	COL, JAL, MOR, NAY, OAX, SON	JAL, MICH, NAY	JAL	JAL, SON	MOR, NAY, SIN, VER	CHIS		COL, NAY, OAX, SIN, SLP, SON	6
TOTAL															287

*This species has apparently been confused with *A. 10-guttatus* (Fabricius); therefore, most *A. 10-guttatus* records are actually *A. jacobiana* (originally described and only known from USA: AZ, TX).

** *Alagoasa semipurpurea* = comb. n., this species should be placed in *Kuschelina*.

*** *Kuschelina semipurpurea* = comb. n., see *Alagoasa*.

**** *aequinoctialis aequinoctialis* (= *cyanipennis octomaculata* Crotch & *punctulata* Bechyne & Bechyne). BUT, re Blake 1931 *O. aequinoctialis* (s.s.) has a black metasternum & black metafemora. Indeed there seems to be some differences consistent with this & an elytral pattern different where *aequinoctialis* (s.s.) has the median/central spots more rounded & only slightly angled whereas for *O. cyanipennis* 8-maculata they are more distinctly angled and slender.

Results

Although 6625 Mexican Alticinae specimens from nine institutional collections were studied (see list in the Methods section above) some of these (493) that could not be determined reliably to recorded species (Table 1); therefore, a total of 6132 specimens were determined to species. Table 1 also demonstrates that a total of 394 new state records were found in these 8 collections, but 107 were repeated so that actually there are 287 new state records. The specimens studied belonged to a somewhat random assortment of genera, including the subtribe Oedionychina (*Alagoasa*, *Asphaera*, *Capraita*, *Kuschelina*, *Omophoita*, *Walterianella*) and *Disonycha*, *Prasona*, *Systema*. The genera with the most specimens belonged to *Alagoasa*, *Asphaera*, *Omophoita*, *Disonycha*, and *Systema*. Table 2 lists only the species in these genera with the new state records as discovered in the current study with those new state records in bold type; the full distribution of each species can be determined by combining these with the distributions in Furth (2006, 2009, 2013). Map 1 demonstrates the number of species by state as of Furth (2013) in comparison to the current study illustrating in Map 2 only the new species records by states and in Map 3 the total species per state.

As indicated in Table 2 the nine genera studied have a high percentage of new state records based only on examination of these institutional collections, as follows: *Alagoasa* (13 of 18 species with new state records of the 44 species recorded from Mexico) [see also Fig. 1]. However, the two apparent new state records (SIN and YUC) for *A. jacobiana* are not included as new records because of its confusion with *A. decemguttata*; *Asphaera* (4 of 4 species of the 10 species recorded from Mexico) [see also Fig. 2]; *Capraita* (2 of 2 species of the 4 species recorded from Mexico) [see also Fig. 3]; *Disonycha* (27 of 33 species of the 49 species recorded from Mexico) [see also Fig. 4]; *Kuschelina* (2 of 3 species of the 8 species recorded from Mexico) [see also Fig. 5]; *Omophoita* (6 of 7 species of the 13 species record from Mexico) [see also Fig. 6]; *Prasona* (1 of 1 species of the 1 species recorded from Mexico) [see also Fig. 7]; *Systema* (18 of 20 species of the 31 species recorded from Mexico) [see also Fig. 8]; and *Walterianella* (7 of 8 species of the 10 species recorded from Mexico) [see also Fig. 9]. Thus, there are 80 species of the 97, or almost 83% of the species examined from the nine institutional collections with new state records, and this is 47% of the total 170 species in these genera recorded from Mexico (Fig. 10).

Of special note in Table 2 are the confirmed species determinations that verify some questionable state records indicated in Furth (2006). They are as follows: *Disonycha guatemalensis* from Veracruz; *Omophoita affinis* from Mexico, a state record for a species only recorded previously as from the country of Mexico; *Systema bitaeniata* from Veracruz; *Systema blanda* from San Luis Potosi; *Systema contigua* from Sonora and Veracruz; *Systema undulata* from Guerrero and Morelos; *Walterianella venustula* from Colima.

Systema oberthuri Baly is reported for the first time from Mexico (Table 2; Fig. 12).

From the current study, it is evident that the numbers of recorded species have changed, in some cases significantly (see Maps 1, 2, 3). Map 2 shows these new records clearly (the details are in Table 2). The Mexican states with the most new species records are Chiapas (32), Nayarit (27), and Sinaloa (24).




Figure 1. *Alagoasa* Bechyné new state records versus previously recorded state records.




Figure 2. *Asphaera* Chevrolat new state records versus previously recorded state records.




Figure 3. *Capraita* Bechyné new state records versus previously recorded state records.




Figure 4. *Disonycha* Chevrolat new state records versus previously recorded state records.




Figure 5. *Kuschelina* Bechyné new state records versus previously recorded state records.




Figure 6. *Omophoita* Chevrolat new state records versus previously recorded state records.



Figure 7. *Prasona* Baly new state records versus previously recorded state records.




Figure 8. *Systema* Chevrolat new state records versus previously recorded state records.




Figure 9. *Walterianella* Bechyné new state records versus previously recorded state records.




Figure 10. Summary of all new records for selected genera.





Figure 11. Summary of all genera with species percentages.




Systema laevis Blake

Systema oberthuri Baly
NEW TO MEXICO

Prasona viridis Baly

Figure 12. *Systema* and *Prasona* generic confusion.

Melanism only known from Chihuahua, Nayarit, Sonora, Sinaloa




Disonycha glabrata (Fabricius)
normal form

Disonycha glabrata (Fabricius)
partially melanistic form

Disonycha glabrata (Fabricius)
melanistic form

Figure 13. *Disonycha glabrata* (Fabricius) species color forms.



Ex Biologia Centrali Americana, Jacoby, 1887
(Figures 15, 18, 22)



Biologia Centrali Americana specimen
Panama: V. de Chiriqui. Champion

Figure 14. *Alagoasa decemguttata* (Fabricius) intraspecific variation.




Figure 15. *Alagoasa decemguttata* intraspecific variation (ex Bechyné, 1955).



Figure 16. *Alagoasa jacobiana* (Horn) species confusion and intraspecific variation.




Figure 17. *Alagoasa* and *Kuschelina* generic confusion.

División Política Estatal

Records of Oaxaca Study (Furth, 2013)



Map 1. Species numbers by states from Furth (2013).

División Política Estatal

New Records



Map 2. New species records by states from the current study.



Map 3. Total species records by states as of the current study.

Taxonomic problems

Generic level confusion

Systema versus *Prasona* (Fig. 12). There has long been some confusion concerning the genus *Prasona* Baly as to whether it is synonymous with *Systema* Chevrolat. *Prasona viridis* Baly from Mexico is the type species (Baly 1861) and the only one recorded from Central America (Furth and Savini 1996) and eight other species from various parts of South America (Bechyné 1971). *Prasona* seems to have the primary characteristics of *Systema*, but seems only to differ in being much larger in size than species of *Systema*. *Prasona* was placed near *Systema* and *Cyrsylus* in the “catalog phylogeny” (Furth and Suzuki 1998) of Bechyné (1971).

Alagoasa versus *Kuschelina* (Fig. 17). *Kuschelina* Bechyné differs from *Alagoasa* Bechyné by having moniliform antennal segments; smaller eyes (diameter less than 3 times as small as width of frons); head (frons/vertex) rugosely punctured; male terminal sternite ventrally without depression; epipleura bent/slanted downward, i.e., visible in lateral view; elytral pleura narrow, not grooved or explanate; antero-lateral corners of pronotum only slightly protruding (not appearing to surround eyes/head; body shape elongate oval (not very rounded). The first placement of North American *Oedionychus* Berthold or *Oedionychis* Latreille (most *Alagoasa* were placed in one of these generic names previously) into *Kuschelina* was by Balsbaugh and Hays (1972) [for 14 spp.]. In Riley et al. (2003) there were 28 species of *Oedionychis* Latreille listed (following Wil-

cox 1975), but only one (*K. scripticollis* (Say) was noted as a new combination; however, many of the others are also technically new combinations, e.g., *K. amplivittata* (Blake), *K. barberi* (Blake), *K. flavocyanea* (Crotch), *K. jacobiana* (Horn), etc. Based on the above characters the author considers *K. semipurpurea* (Jacoby), formerly placed in *Alagoasa*, to be a new combination.

Intra-specific variation:

Disonycha glabrata (Fabricius) (Fig. 13). Because of melanization of the elytra there are intermediate and dark forms; known only from Chihuahua, Nayarit, Sonora, and Sinaloa.

Alagoasa decemguttatus (Fabricius) (Figs 14, 15). Intra-specific variation in this species is well-documented in the literature (Jacoby 1886; Bechyné 1955).

Species confusion (see Table 2):

Alagoasa decemguttatus versus *A. jacobiana* (Horn) (Figs 14, 16). *Alagoasa decemguttatus*: Some confusion with this species. According to Bechyné (1971) *A. decemguttatus* is only from South America and, therefore, most *A. decemguttatus* recorded from Mexico are probably *A. jacobiana*; therefore, some records for *A. jacobiana* may be new records because of this confusion.

Alagoasa duodecimmaculata (Jacoby): There is some confusion of this species with *A. trifasciata escuintla* Bechyné & probably most specimens are *A. trifasciata escuintla*. Its occurrence in Mexico is still somewhat in question (Furth 2006).

Omophoita cyanipennis octomaculata Crotch or *O. octomaculata*: There has historically been confusion about the exact identity of *O. aequinoctialis aequinoctialis*. In this study, the author considers records of *O. a. aequinoctialis* to consist of both *O. cyanipennis octomaculata* Crotch and *O. punctulata* (Bechyné & Bechyné). However, according to Blake (1931) *O. aequinoctialis* (s.s.) has a black metasternum and black metafemora. Indeed, there seems to be some consistent differences between these and an elytral pattern different where *aequinoctialis* (s.s.) has the median/central spots more rounded and only slightly angled, whereas for *O. cyanipennis octomaculata* they are more distinctly angled and slender.

Discussion

Interesting that this study did not reveal new genera and only discovered one new species for Mexico (Furth 2004, 2006, 2009, 2013). This may indicate that the number of described species found in Mexico may be reaching the level of being relatively well known. Only one new species record for Mexico (*Systema oberthuri*) was discovered in this study. However, based on the author's previous and current research, there are certainly many additional undescribed species living within Mexico; probably as many as another 300–400

species. Few other Central American countries have been surveyed for Chrysomelidae. Only Costa Rica is relatively well-known with 350 species in 89 genera (Furth et al. 2003) with only 113 species in 43 genera known previously from the literature. Panama is poorly-known and has 270 species in 70 genera recorded (Furth and Savini 1996, 1998).

Indoor Collecting (Table 1):

The author is not aware of any references in the literature referring to “indoor collecting”, especially in the meaning used in this study. One interview in 2015 of Dr. Art Evans referred to this term for picking up a beetle collection, something the author of this study has been doing for 50 years. However, the meaning for the current study refers to visiting institutional or private scientific collections for the purposes of scientific research, e.g., systematic revisions, faunistics, biodiversity, biogeographical, etc. Most biologists interested in nature, prefer the fun of “outdoor collecting” coupled with subsequent study of material and data from this back home, “indoor collecting” can be just as fun and usually even more productive scientifically; as evidenced from the current study. This kind of work is certainly nothing new, all students and professional biologists conduct this kind of work because the wealth of information “hiding” in such collections is phenomenal, vast, full of valuable information that can help answer many scientific questions and enhance most studies, and for the most part except data already published (at least in entomology), not available elsewhere. The current study is an example of “indoor collecting” where a large percentage (47%) of new Mexican state records were discovered.

Although the choice of the 9 genera and 8 collections for this study was rather random they still provide a very good example of the scientific value of collections. The author plans to continue such research on the Mexican Flea Beetle fauna based on much more specimen material he has borrowed from the same (and eventually other) collections. The 8 collections sampled represent a large percentage of the collections not yet studied by the author likely to have material from Mexico, notable exceptions are the collections of the Los Angeles County Museum, Texas A. & M. University, and, of course, the Universidad Nacional Autónoma de Mexico.

There is no particular pattern to the states with the most new records (Map 2). This is probably only an artifact of the historical collecting of individuals whose material is associated with the nine collections sampled. One of the 3 states with the most new records is Chiapas that is one of several southern states with strong tropical biogeographical affinities (Furth 2013). As evident in Map 2, this study produced new state records in all but three states (Aguascalientes, Distrito Federal, Tlaxcala) and there were new records with 10 or more in 13 states, i.e., over 30%; this is a testament to the value of “indoor collecting”. In previous studies of the Mexican Alticinae fauna some records were questionable because of unclear label data, unclear assignment in the literature, etc. (Table 2), but in the current study of the specimens in these 8 collections some of these were confirmed or enhanced for 7 species (see results above) for eight of the state, including one for the country.

In the Results section above the author has pointed out and attempted to clarify the taxonomic confusion at the generic and specific levels that came to light during the current study. There is no real need to elaborate in detail about these. Resolution

of the confusion between *Systema* and *Prasona* requires considerably more study, but it is quite possible that *Prasona* will become a synonym of *Systema*. The confusion within the “Oedionychina” of *Alagoasa* and *Kuschelina* has caused some problems in faunistic studies in the Nearctic and Neotropical Regions, e. g., Riley et al. (2003). In the Results section above the author attempts to explain his interpretation of these two genera and to clarify the morphological differences, as well as to point out some new combinations created by this confusion.

Therefore, there are 80 species of the 96 species found in this study from 8 collections, or 83%, with new state records, and this is 47% of the total 170 species in these genera recorded from Mexico (Fig. 10). Figure 11 illustrates the nine genera in this study and their species percentages of the total.

In conclusion, the author hopes that not only does this study of a relatively few collections and genera significantly increase the knowledge of the Mexican Flea Beetle fauna, but also that it demonstrates the value of “indoor collecting” as an integral part of any biodiversity and faunistic research.

Acknowledgements

Lee Herman (AMNH), Sharon Shute (NHM), Shawn Clark (BYU), Norm Penny† (CAS), Charles Bellamy† (C DFA), Cheryl Barr (UCB), Steve Heydon (UCD), Dr. R. Wills Flowers (RWF). For professional assistance with imaging I am very grateful to Karolyn Darrow (NMNH). For help with producing the generic charts of new/old state records I would like to thank Dr. Floyd Shockley (NMNH). For access to the NMNH Chrysomelidae collection I am indebted to Dr. Alexander Konstantinov (Systematic Entomology Laboratory, NMNH). I also thank Dr. Raymond Gagne (NMNH) for assistance in translation of the French description of *Kuschelina* Bechyne. Finally, I am indebted to an anonymous reviewer and the Editor (M. Schmitt) for helpful comments.

References

- Balsbaugh E, Hays K (1972) Leaf beetles of Alabama. Agricultural Experiment Station, Auburn University, 223 pp.
- Baly JS (1861) Descriptions of new genera and species of Phytophaga. *Journal of Entomology* 1(4): 275–302.
- Bechné J (1955) Troisième note sur les Chrysomeloidea Néotropicaux des collections de l'Institut Royal des Sciences Naturelles de Belgique. *Institut Royal des Sciences naturelles de Belgique* 31(5): 1–23.
- Bechné J (1955) Troisième note sur les Chrysomeloidea Néotropicaux des collections de l'Institut Royal des Sciences Naturelles de Belgique (Col. Phytophaga) Deuxième Partie (1). 31(19): 1–28.
- Bechné J (1971) Unpublished catalogue of the Neotropical Alticinae (4 parts). <http://chrysomelidae.miza-ucv.org.ve/content/cat%C3%A1logos-in%C3%A9ditos>

- Blake DH (1931) Notes on West Indian and Central American flea-beetles (Halticinae). Bulletin of the Brooklyn Entomological Society 26: 76–83.
- Furth DG, Lee JE (2000) Comparative study of larvae and adults of some species of the *Blepharida* Chevrolat-group (Coleoptera: Chrysomelidae: Alticinae). Journal of the New York Entomological Society 108(1–2): 26–51.
- Furth DG, Savini V (1996) Checklist of the Alticinae of Central America, including Mexico (Coleoptera: Chrysomelidae). Insecta Mundi 10(1–4): 45–68.
- Furth DG, Savini V (1998) Corrections, clarifications, and additions to the checklist of the Alticinae of Central America, including Mexico (Coleoptera: Chrysomelidae). Insecta Mundi 12(1–2): 133–138.
- Furth DG, Suzuki K (1998) Studies on Oriental and Australian Alticinae genera, based on the comparative morphology of the metafemoral spring, genitalia, and hind wing venation. Proceedings of the Fourth International Symposium on the Chrysomelidae. Atti Museo Regionale Scienze Naturali Torino 1998: 91–124.
- Furth DG, Longino JT, Paniagua M (2003) Survey and quantitative assessment of flea beetle diversity in a Costa Rican rainforest (Coleoptera: Chrysomelidae: Alticinae). In: Furth DG (Ed.) 16 Special Topics in Leaf Beetle Biology: Proceedings of the Fifth International Symposium on the Chrysomelidae. Pensoft Publishers, Sofia-Moscow, 1–23.
- Furth DG (2004) Alticinae (Coleoptera: Chrysomelidae). In: Llorente Bousquets J, Morrone JJ, Yáñez O, Vargas I (Eds) Biodiversidad, Taxonomía y Biogeografía de Artrópodos de México: Hacia una síntesis de su conocimiento. Volumen IV. Universidad Nacional Autónoma de México, México, D. F, 669–684.
- Furth DG (2006) The current status of knowledge of the Alticinae of Mexico (Coleoptera: Chrysomelidae). Proceedings of the Sixth International Symposium on the Chrysomelidae. Bonner Zoologische Beiträge 54(4): 197–225
- Furth DG (2009) Flea beetle diversity of the Sierra Tarahumara, Copper Canyon, Mexico (Chrysomelidae: Alticinae). In: Jolivet P, Schmitt M, Santiago-Blay J (Eds) Research on Chrysomelidae, Volume 2. Koninklijke Brill, Leiden, 131–151.
- Furth D (2013) Diversity of Alticinae in Oaxaca, Mexico: A preliminary study (Coleoptera, Chrysomelidae). In: Jolivet P, Santiago-Blay J, Schmitt N (Eds) Research on Chrysomelidae 4. ZooKeys 332: 1–32. <https://doi.org/10.3897/zookeys.332.4790>
- Jacoby M (1884–1886) Biologia Centrali-Americana, Insecta, Coleoptera, Galerucidae, vol. 6, pt. 1, London, 263–625.
- Jacoby M (1891) Biologia Centrali-Americana, Insecta, Coleoptera, Supplement to Phytophaga, vol. 6, pt. 1, London, 233–312.
- Mohamedsaid MS, Furth DG (2011) Secondary sexual characteristics in the Galerucinae (*sensu stricto*) (Coleoptera: Chrysomelidae). International Scholarly Research Network, Zoology 2011: 1–60.
- Riley E, Clark S, Seeno TN (2003) Catalog of Leaf Beetles North of Mexico (Coleoptera: Megalopodidae, Orsodacnidae and Chrysomelidae, excluding Bruchinae). The Coleopterists Society, Special Publication, 1, 290 pp.
- Wilcox JA (1975) Leaf Beetles (Red Version). Checklist of the beetles of Canada, United States, Mexico, and the West Indies. BRIA (Biol. Res. Insts. Amer), NABFP 1(7): 1–166.