


Igor K. Lopatin  
Oleg R. Aleksandrovich  
Alexander S. Konstantinov



# CHECK LIST OF LEAF-BEETLE

*Chrysomelidae (Coleoptera)*

OF THE EASTERN EUROPE AND NORTHERN ASIA

**MANTIS**

**CHECK LIST OF LEAF-BEETLES**  
**(*COLEOPTERA, CHRYSOMELIDAE*)**  
**OF THE EASTERN EUROPE AND NORTHERN ASIA**

---

---






Fig. 1. P. S. Pallas



Fig. 2. E. Ménétriés




Fig. 3. V. I. Motschulsky



Fig. 4. S. M. Solsky



Fig. 5. G. G. Jacobson



Fig. 6. D. A. Ogloblin

Igor K. Lopatin  
Oleg R. Aleksandrovich  
Alexander S. Konstantinov

**CHECK LIST OF LEAF-BEETLES**  
*(COLEOPTERA, CHRYSOMELIDAE)*  
**OF THE EASTERN EUROPE AND NORTHERN ASIA**

Olsztyn



---

On the cover: *Donacia semicuprea* PANZER  
Phot. Piotr DYNOWSKI

© 2004 WYDAWNICTWO MANTIS, Olsztyn

ISBN 83-918125-6-1

Date of publication: 10. XI. 2004

Publisher's address:  
Wydawnictwo Mantis  
ul. Słowicza 11, 11-041 Olsztyn, Poland  
e-mail: mantis@mgt.pl

## Contens

Introduction	7
Brief historical account	7
Taxonomic structure of the fauna	11
Species diversity in separate regions of the Eastern Europe and Northern Asia	13
Acknowledgments	15
Abstract	15
Check list	17
References	231
Index	291

## Introduction

The purpose of this work is to provide a complete list of names proposed for leaf beetles (*Chrysomelidae*) of the Eastern Europe and Northern Asia, the area that was previously known as the Soviet Union (USSR). On the west this territory is limited by the western borders of Russia, the Baltic countries, Belarus, Ukraine, and Moldova. The southern border follows those of Ukraine, the Transcaucasian and Central Asian countries and Russia (Map 1). The northern and the Eastern borders are those of Eurasia.

The list contains the currently accepted valid names (in bold, at the beginning of each entry) followed by all proposed names and combinations that apply to this taxon. The flea beetle part of the list also mentions erroneous attributions introduced by Gruev and Döberl (1997) in the most recent account of the flea beetle species names in the Palearctic region. The list was compiled based on all available literature, extensive material collected over the years by the authors in all regions of this vast territory, and museums, private collections and scientific institutions of the former USSR, Hungary, Czech Republic, Bulgaria, Germany, Austria, Denmark and Switzerland.

## Brief historical account

The following list is a result of more than 200 years of exploration, collecting and research by many generations of professional researchers and amateurs interested in leaf beetles. The history of these studies deserves special attention; here we mention only the most influential figures in this history.

The earliest data on leaf beetles of the Russian Empire date back to the second half of the XVIII century to the expeditions organized by the Russian Academy of Sciences. The first expeditions involved in significant natural history collecting were the ones headed by P. S. Pallas (Fig. 1). He was invited to come to Russia by tsarina Catherine the Second, when he already was a highly respected researcher in Europe. Pallas's first expedition in southern Russia, Kazakhstan, and Siberia lasted from 1768 to 1774 and led him and his small group (including V. Zuev, N. Sokolov, and A. Valter, taxidermist Shumsky, illustrator Dmitriev, an unknown hunter and a cook) from St. Petersburg to Tobol'sk and back. Pallas's second and last expedition to southern Russia and Crimea (1793–1794) was made at his own expense, but with permission of Catherine the Second.


Significant leaf beetle material was collected in the early 1800's by F. Gebler. He was a medical doctor in Barnaul (Siberia) and traveled extensively in Altai, southern Siberia and Kazakhstan. E. Ménétriés (Fig. 2) (1802–1861) was the first professional entomologist in Russia. Born in Paris, he was a student of Cuvier and Latreille. In 1826 he was invited to take a position of a conservator at the "Kunstkamera" (which contained the original natural history collection founded by Peter the Great in 1714, before it was transferred to the Zoological Museum (later Institute) of the Russian Academy of Sciences). Ménétriés made





Map 1. Eastern Europe and Northern Asia.





Map 2. Russian Empire with names dates and trip roots of early collectors (after Motschulsky, 1860)

extensive collections in Caucasus, Siberia and Central Asia. His first expedition (1829–1830) was to the Caucasus, from the environs of Mount Elbrus in the northwestern part of the mountains to the Lenkoran' area in the southeast. The expedition was funded by the Russian Academy of Sciences. It accompanied Russian troops headed by general Emmanuel to fight in the Caucasian war. The group of researchers included physicist A. Ya. Kupfer, geographer E. F. Lents, botanists K. A. Meier and Ménériés, who was in charge of making zoological collections. Important leaf beetle material was also collected in the Caucasus, particularly its southeastern parts, by F. Faldermann who published two volumes describing his rich findings that included several new species (Faldermann 1837, 1838).

Undoubtedly, the most famous Russian coleopterists of the second half of the XIX century was V. I. Motschulsky (Fig. 3). He was a military officer, who later in his career received the rank of colonel and traveled extensively in Russia and also in the USA and India. In addition to many short trips he conducted several long trips to Europe (1836), Russian Caucasus, Caspian Sea, and Siberia, which reached as far as Lake Baikal (1839–1840), Kirgiz Steppes (1847), Europe, which also extended to Egypt and India (1850–1851). In 1853 Motschulsky traveled to the United States, visiting mostly the Western part of the country. This collecting resulted in tremendous amount of material. Motschulsky also worked up, identified and described the gigantic beetle material collected by L. Schrenck (1854–1867) in the Russian Far East, and I. G. Vosnesensky in Siberia, Kamchatka, Alaska and northern California. He identified material collected by many other collectors whose names, dates and trip routes can be found on the map (Map 2) of the Russian Empire attached to the "Reisen und forschungen im Amur-Lande" (Motschulsky 1860). Until approximately 1861, Motschulsky was closely associated with the Zoological Museum in St. Petersburg. Along with a few other amateurs encouraged by Ménériés, he actively worked with the Museum collection. He mounted and identified beetles streaming to the Museum from collectors, acquisitions and other activities, keeping a number of them for his growing personal collection. After Ménériés death, the new director of the Museum, F. F. Brandt changed the Museum rules attempting to regulate Motschulsky's activity. That effectively ended Motschulsky's collaboration with the Museum. Later he moved to Moscow and left his tremendous collection to the Zoological Museum of Moscow University.

At the end of the nineteenth century S. M. Solsky (Fig. 4) (1881) contributed substantially to the leaf beetles studies of the Russian Empire. Like Motschulsky, he was a military officer, devoting all his free time to the study of beetles. Solsky was an active member of the Russian Entomological Society, serving as scientific secretary, librarian, editor, and later vice president. Like all his predecessors, Solsky was not interested in leaf beetles in particular, he described them as a part of his general *Coleoptera* studies.

G. G. Jacobson (Fig. 5) was the first Russian entomologist for whom leaf beetles were the favorite group (Jacobson 1892–1927). He was born in St. Petersburg in 1871, graduated from St. Petersburg University in 1893, and in 1895 accepted the position of zoologist at the Zoological Museum of the Russian Academy of Sciences where he worked until his death in 1926. The most famous of Jacobson's works, "Beetles of Russian Empire, Western Europe and neighboring countries" (1905–1916) was only partially published and the published part does not include leaf beetles. The single copy of complete manuscript was lost in a fire which destroyed the publishing house in the fall of 1917, however the plates containing illustrations of many endemic genera and species of leaf beetles described by Jacobson



were already printed. Although Jacobson did not revise any substantial group of leaf beetles, he laid the necessary foundation for future revisionary and faunistic work. Jacobson practically built the leaf beetle collection at the Zoological Museum. He also identified thousands of leaf beetle specimens for other collections, researchers and amateurs, many of which were published later documenting many local faunas at the turn of the century.

D. A. Ogloblin (Fig. 6) made the first serious attempt to revise a large group of leaf beetles of the territory under consideration. He was born in Borisoglebsk (Tambov gubernia) in 1893 and graduated from the Natural History department of Kiev University in 1916. First he worked in several positions in the Poltava Museum and later (1930) in Leningrad at the Institute of Plant Protection. From 1933 he was employed by the Zoological Institute where he became a chair of the *Coleoptera* Department in 1938. Ogloblin further developed and improved the leaf beetle collection of the Institute, adding many, especially exotic, species. He published the only volume of the Fauna of the USSR devoted to leaf beetles (Ogloblin 1936) which is still an invaluable source of information on this important group. Ogloblin was a wonderful scientific illustrator. He left many beautifully illustrated unpublished manuscripts, some of which were published posthumously (Ogloblin and Medvedev 1971). Ogloblin died tragically in 1942 in Leningrad during the siege of the city.

Studies of the leaf beetles of Eastern Europe and Northern Asia were significantly stimulated by publication of World (Clavareau 1913–1914) and Palearctic (Winkler 1929) catalogues. In addition to numerous faunistic papers, major works providing keys for leaf beetles species of large territories also appeared (Central Asia – Lopatin 1977; Ukraine – Brovdij 1977, 1983; Kazakhstan – Lopatin and Kulenova 1986; Belarus, Lithuania, Latvia and Estonia – Lopatin 1986; Siberia – Medvedev and Dubeshko 1992; and Far East – Medvedev 1992; European part of former USSR – Medvedev and Shapiro 1965, Bienkovski 1999).

Several monographic works treating the leaf beetles of neighboring countries and larger territories which include the one under consideration are also important for understanding the leaf beetle fauna of Eastern Europe and Northern Asia (Borowiec 1999, Doguet 1994, Gressitt and Kimoto 1961, 1963, Gruev and Döberl 1997, Gruev and Tomov 1984, 1986, Kimoto and Takizawa 1994, 1997, Konstantinov 1998, Konstantinov and Vandenberg 1996, Mohr 1966, Scherer 1969, Warchałowski 1985, 1985a, 1991, 1993, 1994, 1995, 1998, 2000). The attempt to catalog all the flea beetle species of the Palearctic region (Gruev and Döberl 1997) incorrectly included numerous new combinations incorrectly attributed to authors who never proposed them. We indicate these incorrect attributions in the following checklist.

### Taxonomic structure of the fauna

The checklist follows the traditional classification of leaf beetles accepted in early works on this fauna (Lopatin 1977, Medvedev 1992). For *Donaciinae* we follow Askevold (1990). The list contains 16 subfamilies, 22 tribes, 167 genera and 1826 species and subspecies (Tab. 1).

Four subfamilies contain the greatest number of the genera: *Alticinae*, *Chrysomelinae*, *Galerucinae* and *Eumolpinae*. The highest species diversity occurs in the subfamilies *Alticinae*, *Chrysomelinae*, and *Cryptocephalinae*, each of which contains more than 300

species. Both *Galerucinae* and *Clytrinae* contain more than 100 species. The other subfamilies have from 1 to 26 genera and from 1 to 73 species.

Within the largest subfamilies most of the species are concentrated in a few genera. Although the fauna of the former USSR contains a large number of species, its generic diversity is not as great as those of neighboring faunas. For example, in the subfamily *Clytrinae* most of the species are placed in the genera *Labidostomis* (49 species) and *Smaragdina* (31 species). In the *Cryptocephalinae* the largest genus is *Cryptocephalus* (224 species), in the subfamily *Chrysomelinae* – *Chrysolina* (202), in the subfamily *Alticinae* – *Longitarsus* (130), *Aphthona* (61 species), *Phyllotreta* (72), *Psylliodes* (53), and *Chaetocnema* (47), and in the subfamily *Cassidinae* – *Cassida* (56).

Table 1.

Subfamily	Tribe	Genera	Species
<i>Donaciinae</i>	<i>Donaciini</i>	2	45
	<i>Haemoniini</i>	2	4
	<i>Plateumarini</i>	1	13
<i>Criocerinae</i>	<i>Criocerini</i>	2	15
	<i>Lemiini</i>	2	14
<i>Zeugophorinae</i>		2	10
<i>Orsodacninae</i>		1	2
<i>Megalopodinae</i>		3	2
<i>Synetinae</i>		2	3
<i>Clytrinae</i>	<i>Clytrini</i>	2	114
<i>Cryptocephalinae</i>	<i>Cryptocephalini</i>	3	228
	<i>Pachybrachini</i>	3	119
	<i>Stylosomini</i>	1	10
<i>Chlamisinae</i>		1	1
<i>Lamprosomatinae</i>	<i>Lamprosomatini</i>	2	2
<i>Eumolpinae</i>	<i>Adoxini</i>	16	42
	<i>Colasposomini</i>	1	1
	<i>Eumolpini</i>	3	9
	<i>Nodinini</i>	6	14
<i>Chrysomelinae</i>	<i>Chrysomelini</i>	31	396
	<i>Timarchini</i>	1	8
<i>Galerucinae</i>	<i>Agelasticini</i>	4	8
	<i>Galerucini</i>	12	69
	<i>Luperini</i>	15	74
	<i>Sermvlassini</i>	1	1
<i>Alticinae</i>		34	539
<i>Hispinae</i>		6	10
<i>Cassidinae</i>	<i>Aspidimorphini</i>	1	2
	<i>Cassidini</i>	7	71
Total	22	167	1826

These genera also contain the majority of species in the region. The remaining 14 subfamilies are represented by a few genera and species (1 or 2 genera are known for *Orsodacninae*, *Megalopodinae*, *Synetinae*, *Chlamisinae*, and *Lamprosomatinae*). Some subfamilies are generally small, the others are more tropical in nature and the majority of their species occur outside of Eastern Europe and Northern Asia

### Species diversity in separate regions of the Eastern Europe and Northern Asia

The territory of Eastern Europe and Northern Asia occupies most of the Palearctic region and contains a diversity of geographic zones from tundra to desert with different climatic and soil conditions and a tremendous variety of habitats. In addition to the vast plains, several major mountain systems occur in the west and south (Carpathians, Caucasus, Ural, mountains of Central Asia and southern Siberia). The diversity of habitats, proximity of important centers of speciation, and long history of the fauna defines leaf beetle diversity of Eastern Europe and Northern Asia. As a whole this fauna is truly Palearctic, with a small addition of Oriental and Afrotropical elements in the south and southeast.

There are 5 major regions recognized in Eastern Europe and Northern Asia: Eastern Europe including European part of the former USSR; Carpathians and Caucasus; Siberia; Kazakhstan; Central Asia; and Far East. The number of leaf beetle species in each region is presented in the Table 2.

Table 2.

Region	Number of species
Eastern Europe	639
Siberia	403
Far East	435
Kazakhstan	590
Central Asia	766

The leaf beetles of Eastern Europe are much better known than the leaf beetles of other regions. For example, for the last 10 years no previously unrecorded species were discovered in Lithuania, Latvia, Estonia, European part of Russia, Belarus, and Moldova. The leaf beetles of the Caucasus and Crimea are much less known. Some, particularly alpine, groups of leaf beetles, are in need of revision, which will undoubtedly result in the discovery of previously unknown taxa. The leaf beetles of Central Asia are even less known. Despite years of explorations and extensive research in Kirghizstan, Kazakhstan, Tajikistan, and Uzbekistan there are still vast mountain ranges where exploration was brief and fragmentary. Recent attempts to revise certain taxa resulted in great numbers of species descriptions and changes in taxonomic status (Konstantinov and Lopatin 1992, 2000).

The leaf beetles of Eastern Europe are typically European with a small addition of Mediterranean and Near Eastern species. The endemics are not numerous and occur only in the southern parts of the region (Crimea and Caucasus). Among the endemics *Cecchiniola platyscelidina* Jebb. is particularly important since it is the single known generic endemic in



this region. The other examples of endemics are *Luperus kiritschenkoi* Ogl., *Cryptocephalus floralis* Kryn., *Chrysolina rosti* Wse., *Ch. pliginskii* Reitt., *Ch. abchasica* Wse., *Ch. apsilaena* Silf. The relatively small number of endemics can be easily understood since the fauna is comparatively young, as the most of the region was under a glacial cover during the Pleistocene.

Siberia, with a severe climate and relative monotony of vegetation, has the poorest fauna of *Chrysomelidae*. Nevertheless there are a number of endemics, some of which are not necessarily alpine. These are local subspecies of widely distributed species: *Labidostomis metallica imitatrix* Ogl., and *Clytra atraphaxidis punctata* Wse. Most of the endemics are species of *Cryptocephalus* (*pilosellus* Suff., *agnus* Wse., *korotyaevi* L. Medv., *miczechi* Tapp., *bivulneratus* Fald., *bidenulus* Suff., *biguttulatus* Gebl.), *Pachybrachis* (*tuvensis* L. Medv., *sericans* Suff., *transbaicalicus* L. Medv.), and *Chrysolina* (*instabilis* Mäkl., *difficilis* Motsch., *haemochlora* Gebl., *jakovlevi* Wse., *spectabilis* Motsch., *schewyrewi* Jcbs., *ordinata* Gebl. etc.). Quite remarkable is *Oreothassa martjanovi* Jcbs., a monotypic endemic genus. *Sternoplatys clementzi* Jcbs., *Colaphellus alpinus* Gebl., *C. foveolatus* Gebl., *Diorhabda rybakovi* Wse., *Luperus anthracinus* Ogl., *Altica sajanica* Csiki, *Aphthona chekanovskii* Konst. and *Pentamesa duodecimpunctata* (Motsch.) are limited in their distribution to the mountainous parts of the region: Altai and Sayan.

The Far East consists of two parts. The northeastern part is an extension of East Siberia, and has a relatively poor fauna. The southern part, Cisamuria and Maritime Province, have much higher species diversity. However, the percent of endemics is not large and the majority of species occur both in Korea and China. The species of oriental origin join with Palearctic fauna only in this region. These are members of the following genera: *Clythraxeloma* and *Temnaspis* (*Megalopodinae*), *Melixanthus* (*Cryptocephalinae*), *Oomorphoides* (*Lamprosomatinae*), *Abiromorphus*, *Basilepta*, *Cleoporus*, *Colasposoma*, *Lypesthes* (*Eumolpinae*), *Ambrostoma*, *Paropsides*, *Gastrolinoides* (*Chrysomelinae*), *Clitena*, *Euliroetis*, *Cneorane*, *Morphosphaera*, *Atrachya*, *Galerucida* (*Galerucinae*), *Sangariola*, *Liprus*, *Phygasia*, *Argopistes*, *Hemipyxis* (*Alticinae*), *Cassidispia*, *Dactylispia*, *Rhadinosa* (*Hispiniae*), *Aspidimorpha*, *Thlaspida*, and *Glyphocassis* (*Cassidinae*).

The leaf beetle faunas of Kazakhstan and Central Asia are more complicated than the previous ones. Northern and central Kazakhstan are populated by a large number of leaf beetles closely associated with steppe and semidesert landscapes. Less common are species associated with forests which occur close to the Altai Mountains (*Oulema lichenis* Voet., *O. tristis* Hbst., *Lema decempunctata* Gebl., *Cryptocephalus distinguendus* Schnd., *C. octopunctatus* Scop., *Chrysolina gypsophilae* Küst., *Ch. reitteri* Wse. (*lurida* L.), *Ch. hyrcana* Wse., *Chrysomela vigintipunctata* L., *Ch. cuprea* L.). Widespread polyzonal, Euro-Siberian species are also common.

There are two main landscapes in southern Kazakhstan: deserts and mountains. The leaf beetle fauna contains more endemics and a number of species distributed also in Central Asia. The latter occupies the territory from the east coast of the Caspian Sea to the Pamirs and Dzhungaria. Deserts and mountains define the landscapes of this region. The leaf beetle fauna of Central Asia is the richest and most unique among the regional faunas of Eastern Europe and Northern Asia. The following genera (some of which are represented by many species) are endemic: *Exaesiognatha*, *Jaxartiolus*, *Aphilenia*, *Caspiana*, *Parnops*, *Atomyria*, *Bedelia*, *Andosia*, *Anidania*, *Adoxinia*, *Callipta*, *Gnathomela*,

*Neophaedon*, *Xenomela*, *Pallasiola*, and *Chiridula*. In many subfamilies endemics reach 60–75 % of total number of species. It is particularly characteristic for *Cryptocephalinae*. For example, almost all species of the subgenus *Asionus* of the genus *Cryptocephalus* have extremely limited ranges (*dilectus* Wse., *spurius* Lop., *medvedevi* Lop., *iskanderi* Lop., *irenae* Lop., *oxianus* Lop., *pavlovskii* Lop., *heydeni* Wse.). The majority of *Pachybrachis* and *Acolastus* species are endemics. In *Chrysomelinae* most species of *Chrysolina* (*nigrovittata* Ball., *tianshanica* Jebs., *octocosta* Jebs., *ballioni* Lop., *kryzhanovskii* Lop., *kaikana* Lop., *tastavica* Lop., *dolini* Lop., *sogdiorum* Wse., *koenigi* Jebs., *kiritschenkoi* Lop.) and all *Oreomela* are endemics. The only two subfamilies without endemics are *Orsodacninae* and *Hispinae*. In *Alticinae* many particularly alpine and ephemeral species of *Phyllotreta*, *Aphthona*, *Longitarsus* are endemics as well. Among them some species have extremely narrow ranges covering only one or two neighboring mountain ridges. The ranges of some species extend into Iran, Afghanistan and China.

### Acknowledgments

We are most grateful to Prof. G. S. Medvedev (Zoological Institute of Russian Academy of Sciences, St. Petersburg, Russia) and Prof. V. G. Dolin (Institute of Zoology, Kiev, Ukraine) for encouragement, support and friendship over the years. We thank Dr. A. Norrbom (Systematic Entomology Laboratory, USDA, ARS) and C. Staines (Smithsonian Institution, Washington DC, USA) for their thorough reviews of this manuscript and numerous constructive comments. We sincerely thank Dr. B. A. Korotyaev (Zoological Institute of Russian Academy of Sciences, St. Petersburg, Russia) for providing photographs of Jacobson and Ménétrés and his useful suggestions. Prof. V. N. Tanasijchuk (Zoological Institute of Russian Academy of Sciences, St. Petersburg, Russia), Prof. L. Borowiec (University of Wrocław, Poland) and Dr. V. Korneev (Institute of Zoology, Kiev, Ukraine) helped to obtain important publications.

### Abstract

The checklist of all known leaf beetle (*Coleoptera*, *Chrysomelidae*) taxa of the Eastern Europe and Northern Asia is provided. It contains 16 subfamilies, 167 genera and 1826 species and subspecies. The replacement name *Longitarsus silbursaiicus* Lopatin, new name, is provided for *Longitarsus seriepunctatus* Lopatin, 1966 (non Roubal, 1943). The history of the leaf beetle studies in the Eastern Europe and Northern Asia is briefly outlined. Zoogeographic features of the fauna, particularly an endemism, are also discussed.

## Check list

Family: *Chrysomelidae* LATREILLE, 1802

Subfamily: *Donaciinae* KIRBY, 1837

Tribe: *Donaciini* KIRBY, 1837

Genus: *Donacia* FABRICIUS, 1775

*Donaciella* REITTER, 1920

*Pseudodonacia* REITTER, 1920

*Donaciomima* L. MEDVEDEV, 1973

Subgenus: *Donacia (Donacia)* FABRICIUS, 1775

*Donacia (Donacia) aequidorsis* JACOBSON, 1894

Distribution: SE European Russia, W Kazakhstan.

*Donacia (Donacia) antiqua* KUNZE, 1818

*Donacia gracilis* SUFFRIAN, 1845

*Donacia simplicifrons* LACORDAIRE, 1845

Distribution: N, Central and northern part of E Europe.

*Donacia (Donacia) bactriana bactriana* J. WEISE, 1887

Distribution: S Kazakhstan, Central Asia.

*Donacia (Donacia) bactriana sahlbergi* JACOBSON, 1901

Distribution: Central Asia.

*Donacia (Donacia) breviuscula* JACOBSON, 1899

Distribution: Asian Russia (Amurland).

*Donacia (Donacia) crassipes* FABRICIUS, 1775

*Donacia striata* PANZER, 1794

*Donacia micans* HOPPE, 1795

*Donacia micans* ab. *anthracina* EVERTS, 1903

Distribution: Europe, Kazakhstan, Asian Russia (Siberia east to Baikal).

*Donacia (Donacia) fedtschenkoae* JACOBSON, 1899

Distribution: Uzbekistan.

*Donacia (Donacia) flemola* GOECKE, 1944

*Donacia jacobsoniana* SHAVROV, 1946

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea.



***Donacia (Donacia) gracilicornis* JACOBSON, 1892***Donacia transcaucasica* SUMAKOV, 1901

Distribution: Transcaucasia, W Iran, ?Turkmenistan.

***Donacia (Donacia) kirgizkaisaka* JACOBSON, 1925**

Distribution: Siberia (Irtysh River).

***Donacia (Donacia) knipovitschi* JACOBSON, 1927**

Distribution: Asian Russia (S Maritime Prov.), NE China.

***Donacia (Donacia) koenigi* JACOBSON, 1899**

Distribution: Transcaucasia.

***Donacia (Donacia) kraatzi* J. WEISE, 1882***Donacia fastuosa* KHNZORIAN, 1962

Distribution: Armenia, N Turkey.

***Donacia (Donacia) mistschenkoi* JACOBSON, 1910**

Distribution: Caucasus.

***Donacia (Donacia) ochroleuca* J. WEISE, 1882***Donacia flavidula* REITTER, 1913

Distribution: Asian Russia (Transbaikalia, Amurland, Maritime Prov.), China.

***Donacia (Donacia) polita* KUNZE, 1818**

Distribution: S Europe, Moldavia.

***Donacia (Donacia) ussuriensis* L. MEDVEDEV, 1973**

Distribution: Asian Russia (S Maritime Prov.).

**Subgenus: *Donacia (Donaciella)* REITTER, 1920*****Donacia (Donaciella) cinerea* HERBST, 1784***Donacia tarsata* PANZER, 1796*Donacia hydrochaeridis* FABRICIUS, 1801

Distribution: Europe, Transcaucasia, Kazakhstan, W Siberia.

***Donacia (Donaciella) clavipes clavipes* FABRICIUS, 1793***Donacia menyanthidis* AHRENS, 1810

Distribution: All over the Palaearctic except for the extreme North.

***Donacia (Donaciella) clavipes glabrata* SOLSKY, 1872**

Distribution: Asian Russia (E Siberia, Maritime Prov.), Mongolia.

***Donacia (Donaciella) tomentosa* AHRENS, 1810**

Distribution: N and Central Europe, Azerbaijan, E Kazakhstan, Uzbekistan, Asian Russia (S Siberia, Altai).

**Subgenus: *Donacia (Donaciomima)* L. MEDVEDEV, 1973*****Donacia (Donaciomima) aquatica* (LINNAEUS, 1758)***Leptura aquatica* LINNAEUS, 1758*Leptura coccineo-fasciata* HARRER, 1784*Donacia fasciata* HERBST, 1784

*Donacia dentipes* FABRICIUS, 1792  
*Donacia aquatica* ab. *concinna* J. WEISE, 1898  
*Donacia aquatica* ab. *chalybaea* J. SAHLBERG, 1900  
*Donacia aquatica* ab. *cyanicollis* OLSUFJEV, 1903  
*Donacia aquatica* ab. *waldaica* OLSUFJEV, 1903  
 Distribution: Europe, Asian Russia (Siberia), N China, Japan.

***Donacia (Donaciomima) bicolora bicolora* ZSCHACH, 1788**

*Donacia sagittariae* FABRICIUS, 1792  
*Donacia collaris* PANZER, 1794  
*Donacia aurea* HOPPE, 1795  
 Distribution: Europe except for the South, Asian Russia (W Siberia), Central Asia.

***Donacia (Donaciomima) brevicornis* AHRENS, 1810**

*Donacia platysterna* C. G. THOMSON, 1866  
*Donacia brevicornis* ab. *noeskei* HÄNEL, 1912  
 Distribution: N and NE Europe.

***Donacia (Donaciomima) brevitarsis* C. G. THOMSON, 1884**

Distribution: N and NE Europe, Alps.

***Donacia (Donaciomima) clavareani* JACOBSON, 1906**

Distribution: Asian Russia (Maritime Prov.), China, Japan.

***Donacia (Donaciomima) dentata* HOPPE, 1795**

*Donacia angustata* KUNZE, 1818  
*Donacia phellandrii* C. R. SAHLBERG, 1839  
*Donacia dentata* ab. *indentata* KOLOSSOW, 1925  
*Donacia dentata* ab. *chrysochlora* WESTHOFF, 1882  
*Donacia dentata* ab. *autumnalis* WESTHOFF, 1882  
*Donacia dentata* ab. *diabolus* WESTHOFF, 1882  
 Distribution: Europe except for the extreme North, Kazakhstan, Asian Russia (W Siberia).

***Donacia (Donaciomima) fennica* (PAYKULL, 1800)**

*Rhagium fennicum* PAYKULL, 1800  
 Distribution: N, northeast of Central and E Europe, Asian Russia.

***Donacia (Donaciomima) impressa* PAYKULL, 1799**

*Donacia impressa* ab. *armillata* SHAYROV, 1948  
 Distribution: Europe except for the outermost North and South. Kazakhstan, Asian Russia.

***Donacia (Donaciomima) malinovskiyi* AHRENS, 1810**

*Donacia Silesiaca* WEIGEL, 1806 [nomen nudum]  
*Donacia arundinis* AHRENS, 1810  
*Donacia malinovskiyi* ab. *ladonensis* THIERRIAT, 1946  
*Donacia malinovskiyi* ab. *arvensis* THIERRIAT, 1946  
*Donacia malinovskiyi* ab. *mequignoni* THIERRIAT, 1946  
*Donacia malinovskiyi* ab. *caroli* THIERRIAT, 1946  
 Distribution: Central and E Europe, Kazakhstan, Asian Russia (W Siberia).

***Donacia (Donaciomima) marginata marginata* HOPPE, 1795***Donacia limbata* PANZER, 1796*Donacia vittata* PANZER, 1796*Donacia lemnae* FABRICIUS, 1801*Donacia limbata* var. *unicolor* WESTHOFF, 1882

Distribution: Europe except for the North, Caucasus, Kazakhstan, Uzbekistan, Asian Russia (W Siberia).

***Donacia (Donaciomima) obscura obscura* GYLLENHAL, 1813***Donacia impressa* AHRENS, 1810, non PAYKULL, 1799*Donacia obscura* ab. *barovskyi* JACOBSON, 1922

Distribution: Europe except for the Mediterranean.

***Donacia (Donaciomima) obscura splendens* JACOBSON, 1894**

Distribution: Asian Russia (W and Middle Siberia, Altai, Transbaikalia, Yakutia, Amurland, Maritime Prov.).

***Donacia (Donaciomima) semicuprea* PANZER, 1796***Donacia semicuprea* ab. *iris* WESTHOFF, 1882*Donacia semicuprea* ab. *tenebrans* WESTHOFF, 1882*Donacia semicuprea* ab. *concolor* WESTHOFF, 1882

Distribution: W and Central Europe, Asian Russia (Siberia east to Baikal).

***Donacia (Donaciomima) simplex* FABRICIUS, 1775***Donacia linearis* HOPPE, 1795*Donacia linearis* var. *pulcherrima* HUMMEL, 1825*Donacia simplex* var. *aeruginosa* WESTHOFF, 1882*Donacia simplex* var. *atrocoerulea* WESTHOFF, 1882*Donacia simplex* var. *aurichalcea* WESTHOFF, 1882*Donacia simplex* var. *sanguinea* WESTHOFF, 1882

Distribution: W and Central Europe, Asian Russia (S Siberia, Transbaikalia), Mongolia.

***Donacia (Donaciomima) sparganii sparganii* AHRENS, 1810***Donacia sparganii* ab. *tridens* JACOBSON, 1892*Donacia sparganii* ab. *coelestis* J. WEISE, 1893

Distribution: N, Central and E Europe, Asian Russia (W Siberia).

***Donacia (Donaciomima) sparganii gracilipes* JACOBY, 1885***Donacia gracilipes* JACOBY, 1885*Donacia tschitscherini* SEMENOV, 1895

Distribution: Asian Russia (Transbaikalia, NE Siberia, Kamchatka, Amurland, Maritime Prov., Sakhalin, Kurile Is.), Japan.

***Donacia (Donaciomima) thalassina intermedia* JACOBSON, 1899***Donacia thalassina rufovariegata* JACOBSON, 1900*Donacia semenovi* JACOBSON, 1907

Distribution: Asian Russia (Siberia, Maritime Prov.), Mongolia.

***Donacia (Donaciomima) thalassina thalassina* GERMAR, 1811***Donacia thalassina* var. *porphyrogenita* WESTHOFF, 1882*Donacia thalassina* ab. *viridis* ŠAVROV, 1948*Donacia thalassina* ab. *extensa* ŠAVROV, 1948

*Donacia thalassina* ab. *inermipes* ŠAVROV, 1948

Distribution: Europe except for the extreme North, Asian Russia (S Siberia, Transbaikalia, Tuva), Mongolia.

***Donacia (Donaciomima) versicoloreae* (BRAHM, 1790)**

*Donacia versicoloreae* BRAHM, 1790 [nomen nudum]

*Leptura versicoloreae* BRAHM, 1790

*Donacia bidens* OLIVIER, 1791

*Donacia cincta* GERMAR, 1810

*Donacia clavipes* AHRENS, 1810, non FABRICIUS, 1792

*Donacia clavipes* ab. *suffriani* WESTHOFF, 1882

*Donacia clavipes* ab. *lusatica* HÄNEL, 1911

Distribution: Europe, Kazakhstan, Asian Russia (SW Siberia).

***Donacia (Donaciomima) vulgaris issykensis* JACOBSON, 1900**

Distribution: Kazakhstan, Central Asia.

***Donacia (Donaciomima) vulgaris vulgaris* ZSCHACH, 1788**

*Donacia typhae* AHRENS, 1810

*Donacia typhae* ab. *viridula* J. SAHLBERG, 1871

*Donacia typhae* var. *concolor* WESTHOFF, 1882

Distribution: All over the Palaearctic.

**Subgenus: *Donacia (Cyphogaster)* GOECKE, 1934**

***Donacia (Cyphogaster) lenzi* SCHÖNFELDT, 1888**

*Donacia aeraria* KOLBE, 1886, non BALY, 1865

Distribution: Asian Russia (S Maritime Prov.), China, Korea, Japan.

***Donacia (Cyphogaster) provosti* FAIRMAIRE, 1885**

*Donacia nitidicollis* J. WEISE, 1898

*Donacia brevicollis* J. WEISE, 1898

*Donacia sicanga* GRESSITT, 1942

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

**Genus: *Sominella* JACOBSON, 1908**

***Sominella macrocnemia* (FISCHER-WALDHEIM, 1823)**

*Donacia macrocnemia* FISCHER-WALDHEIM, 1823

Distribution: Asian Russia (Transbaikalia, Yakutia, Amurland, Khabarovsk Prov., Maritime Prov.), China.

**Tribe: *Haemoniini* CHEN, 1941**

**Genus: *Macrolea* SAMOUELLE, 1819**

*Haemonia* DEJEAN, 1821

***Macrolea appendiculata* (PANZER, 1794)**

*Donacia appendiculata* PANZER, 1794

*Donacia mucronata* HOPPE, 1795

*Donacia equiseti* FABRICIUS, 1798

*Haemonia appendiculata* ab. *lineata* CHEVROLAT, 1844

*Haemonia appendiculata* ab. *chevrolati* LACORDAIRE, 1845

*Haemonia mosellae* BEDEL, 1869

*Haemonia appendiculata* ab. *rugipennis* J. SAHLBERG, 1870

*Haemonia appendiculata* ab. *flavicollis* BELLEVOYE, 1870

Distribution: Europe except for the North and South, W Siberia, ? N Africa.

***Macroplea mutica mutica* (FABRICIUS, 1792)**

*Rhagium muticum* FABRICIUS, 1792

*Donacia zosteræ* FABRICIUS, 1801

*Donacia ruppiae* GERMAR, 1831

*Haemonia curtisii* LACORDAIRE, 1845

*Haemonia gyllenhalii* LACORDAIRE, 1845

*Haemonia sahlbergii* LACORDAIRE, 1845

Distribution: Coastal parts of the North, White, Baltic, Mediterranean, Black and Caspian Sea.

***Macroplea mutica japana* (JACOBY, 1885)**

*Haemonia japana* JACOBY, 1885

Distribution: Asian Russia (E Siberia), Japan.

**Genus: *Neohaemonia* SZEKÉSSY, 1941**

***Neohaemonia voronovae* L. MEDVEDEV, 1977**

Distribution: Asian Russia (Transbaikalia), Mongolia.

**Tribe: *Plateumarini* ASKEVOLD, 1990**

**Genus: *Plateumaris* C. G. THOMSON, 1866**

*Juliusina* Reitter, 1920

***Plateumaris amurensis* J. WEISE, 1898**

*Plateumaris hirashimai* KIMOTO, 1963

Distribution: Asian Russia (Yakutia, Amurland, Maritime Prov., Sakhalin, Kurile Is.), Japan.

***Plateumaris braccata* (SCOPOLI, 1772)**

*Prionus braccatus* SCOPOLI, 1772

*Leptura violacea* PALLAS, 1773

*Donacia palustris* HERBST, 1784

*Donacia nigra* FABRICIUS, 1792

*Donacia?* *adominalis* OLIVIER, 1795

*Donacia braccata* ab. *fairmairei* LEGRAND, 1861

Distribution: Europe except for the North and South, Caucasus, SE Kazakhstan, Asian Russia (S Siberia).

***Plateumaris consimilis consimilis* (SCHRANK, 1781)**

*Leptura consimilis* SCHRANK, 1781

*Donacia rufipes* OLIVIER, 1791

*Donacia discolor* HOPPE, 1795

*Donacia variabilis* KUNZE, 1818

*Plateumaris consimilis* ab. *aerea* BECHYNÉ, 1942

*Plateumaris consimilis* ab. *coerulea* BECHYNÉ, 1942

*Plateumaris consimilis* ab. *flavipes* BECHYNÉ, 1942

*Plateumaris consimilis* ab. *nigripes* BECHYNÉ, 1942

*Plateumaris consimilis* ab. *violacea* BECHYNÉ, 1942

*Plateumaris consimilis* ab. *viridis* BECHYNÉ, 1942

Distribution: Europe except for the North and South, Urals, Asian Russia (Siberia).

***Plateumaris consimilis orientalis* SHAVROV, 1948**

Distribution: Asian Russia (S Maritime Prov.).

***Plateumaris constricticollis kurilensis* L. MEDVEDEV, 1978**

Distribution: Asian Russia (S Kurile Is.: Kunashir).

***Plateumaris obsoleta* JACOBSON, 1894**

Distribution: Asian Russia (S Maritime Prov.).

***Plateumaris roscida* J. WEISE, 1912**

*Donacia annularis* REITTER, 1920

Distribution: Asian Russia (Transbaikalia, Yakutia, Amurland, Maritime Prov.).

***Plateumaris rustica* (KUNZE, 1818)**

*Donacia rustica* KUNZE, 1818

*Donacia affinis* KUNZE, 1818

*Donacia planicollis* KUNZE, 1818

*Donacia pallipes* KUNZE, 1818

*Donacia rustica* ab. *picipes* J. WEISE, 1898

*Donacia rustica* ab. *forojulensis* GÖRIANI, 1906

Distribution: Europe except for the South, Asian Russia (W Siberia), N Algeria.

***Plateumaris sachalinensis* L. MEDVEDEV, 1973**

Distribution: Asian Russia (S Sakhalin).

***Plateumaris sericea sericea* (LINNAEUS, 1758)**

*Leptura sericea* LINNAEUS, 1758

*Donacia caerulea* ZSCHACH, 1788

*Donacia festucae* FABRICIUS, 1792

*Donacia discolor* PANZER, 1795

*Donacia violacea* HOPPE, 1795

*Donacia micans* PANZER, 1796

*Donacia armata* PAYKULL, 1799

*Donacia nymphaeae* FABRICIUS, 1793

*Donacia proteus* KUNZE, 1818

*Donacia asiatica* FALDERMANN, 1837

*Donacia palustris* SCHILLING, 1838

*Donacia comari* SUFFRIAN, 1846

*Donacia geniculata* C. G. THOMSON, 1866

*Plateumaris sericea* ab. *atropurpurea* WESTHOFF, 1882

*Plateumaris sericea* ab. *tenebricola* WESTHOFF, 1882

*Plateumaris sericea* ab. *luctuosa* WESTHOFF, 1882



*Plateumaris discolor* ab. *nigrita* SCHILSKY, 1908  
*Plateumaris sericea* ab. *intermedia* APFELBECK, 1916  
*Plateumaris discolor* ab. *tatica* BALTHASAR, 1934  
*Plateumaris sericea* ab. *viridis* CSIKI, 1953  
*Plateumaris sericea* ab. *laevigata* CSIKI, 1953  
*Plateumaris discolor* ab. *isolacordairei* BECHYNÉ, 1945  
*Plateumaris discolor* ab. *cupraria* BECHYNÉ, 1945  
*Plateumaris discolor* ab. *purpuricena* BECHYNÉ, 1945  
*Plateumaris discolor* ab. *pseudoviolacea* BECHYNÉ, 1945  
*Plateumaris discolor* ab. *coelicolor* BECHYNÉ, 1945  
 Distribution: Europe except for the South, W and Central Siberia.

***Plateumaris sericea caucasica* ZAITZEV, 1930**

Distribution: NW Caucasus (Teberda, Bakuriani).

***Plateumaris sericea sibirica* (SOLSKY, 1872)**

*Donacia sericea* var. *sibirica* SOLSKY, 1872

*Plateumaris socia* CHEN, 1941

*Plateumaris nipponensis* NAKANE, 1963

Distribution: Asian Russia (Siberia, Amurland, Maritime Prov., Kamchatka, Sakhalin), Mongolia, China, Japan.

***Plateumaris weisei* DUVIVIER, 1885**

*Plateumaris mongolica* SEMENOV, 1895

*Plateumaris morimotoi* KIMOTO, 1963

Distribution: N and C Europe, Asian Russia (W Siberia, Tuva, Transbaikalia, Amurland, Maritime Prov., Sakhalin), Mongolia, Japan.

Subfamily: *Orsodacninae* J. THOMSON, 1866

Genus: *Orsodacne* LATREILLE, 1802

*Orsodacna* LATREILLE, 1804

***Orsodacne cerasi* (LINNAEUS, 1758)**

*Chrysomela cerasi* LINNAEUS, 1758

*Crioceris cantharoides* FABRICIUS, 1775

*Crioceris chlorotica* OLIVIER, 1791

*Crioceris melanura* FABRICIUS, 1792

*Crioceris glabrata* PANZER, 1794

*Orsodacna limbata* OLIVIER, 1808

*Orsodacna lineola* LACORDAIRE, 1845

*Orsodacne cerasi* var. *duftschmidti* J. WEISE, 1891

*Orsodacne cerasi* ab. *Horváthi* LACZO, 1912

*Orsodacne cerasi* var. *lacordairei* PIC, 1913

*Orsodacne cerasi* ab. *theresae* PIC, 1913

*Orsodacne cerasi* ab. *theresae* PIC, 1913

*Orsodacne cerasi* ab. *baudii* ROUBAL, 1914

*Orsodacne cerasi* ab. *suturalis* JACOBSON, 1922

Distribution: Europe, Caucasus, Asian Russia (W Siberia, Altai, Cisbaikalia).

***Orsodacne lineola* (PANZER, 1795)**

*Crioceris lineola* PANZER, 1795  
*Orsodacna humeralis* LAIRELLE, 1804  
*Orsodacne nigriceps* LAIRELLE, 1807  
*Crioceris nigricollis* OLIVIER, 1808  
*Orsodacne coerulescens* DUFESCHMIDT, 1825  
? *Orsodacna lineola* ab. *mespili* LACORDAIRE, 1845  
*Orsodacne lineola* ab. *croatica* J. WEISE, 1883  
*Orsodacne lineola* ab. *marginata* CSIKI, 1899  
*Orsodacne lineola* ab. *brancsiki* LACZO, 1909  
*Orsodacne lineola* ab. *hispanica* PIC, 1913  
*Orsodacne lineola* ab. *kraatzii* PIC, 1913

Distribution: Europe except for the North, Mediterranean.

Subfamily: *Megalopodinae* CLAVAREAU, 1913Genus: *Temnaspis* LACORDAIRE, 1845***Temnaspis bonneuili* PIC, 1947**

Distribution: Asian Russia (Amurland, Maritime Prov.).

Genus: *Clythraeloma* KRAATZ, 1879***Clythraeloma cyanipennis* KRAATZ, 1879**

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea.

Subfamily: *Synetinae* EDWARDS, 1953Genus: *Syneta* LACORDAIRE, 1845***Syneta adamsi* BALY, 1877**

Distribution: Asian Russia (Amurland, Khabarovsk Prov., Maritime Prov., Sakhalin, Kurile Is.), China, Korea.

***Syneta betulae betulae* (FABRICIUS, 1792)**

*Crioceris betulae* FABRICIUS, 1792  
Distribution: N and northern parts of E Europe, Asian Russia (Siberia east to Transbaikalia), Mongolia.

***Syneta betulae amurensis* PIC, 1901**

*Syneta ussurica* BRANCSIK, 1914  
Distribution: Asian Russia (Amurland, Khabarovsk Prov., Maritime Prov., Sakhalin).

Subfamily: *Zeugophorinae* CHŪJŌ, 1952Genus: *Zeugophora* KUNZE, 1818*Auchenia* THUNBERG, 1792 (Decision of ICZN N 1382, 9.06.1986)Subgenus: *Zeugophora (Zeugophora)* KUNZE, 1818*Zeugophora (Zeugophora) bimaculata* KRAATZ, 1879

Distribution: Asian Russia (Amurland, Maritime Prov.).

*Zeugophora (Zeugophora) flavicollis* (MARSHAM, 1802)*Auchenia flavicollis* MARSHAM, 1802*Zeugophora flavicollis* var. *australis* J. WEISE, 1881

Distribution: N, Central and E Europe, Kazakhstan, Asian Russia (W Siberia, Tuva, Yakutia).

*Zeugophora (Zeugophora) frontalis* SUFFRIAN, 1841

Distribution: Europe, Asian Russia (Siberia, Tuva, Sakhalin).

*Zeugophora (Zeugophora) hozumii* CHŪJŌ, 1953

Distribution: Russian Far East, Japan.

*Zeugophora (Zeugophora) scutellaris* SUFFRIAN, 1841

Distribution: Central and E Europe, Ciscaucasia, Kazakhstan, Tajikistan, Kirghizstan, Asian Russia (southern W Siberia, Altai).

*Zeugophora (Zeugophora) subspinosa* (FABRICIUS, 1781)*Crioceris subspinosa* FABRICIUS, 1781

Distribution: All over the Palaearctic except for the Central Asia.

*Zeugophora (Zeugophora) turneri* POWER, 1863*Zeugophora rufo-testacea* KRAATZ, 1872*Zeugophora turneri* ab. *australis* J. WEISE, 1881*Zeugophora turneri* ab. *notatipes* PIC, 1925

Distribution: Central and East Europe, Asian Russia (southern W Siberia, Transbaikalia), Mongolia.

*Zeugophora (Zeugophora) weisei* REITTER, 1889

Distribution: Armenia, N Iran.

Subgenus: *Zeugophora (Pedrillia)* WESTWOOD, 1864*Zeugophora (Pedrillia) annulata* BALY, 1873*Zeugophora biguttata* KRAATZ, 1879

Distribution: Asian Russia (Amurland, Khabarovsk Prov., Maritime Prov., Sakhalin, Kurile Is.), China, Korea, Japan.

*Zeugophora (Pedrillia) bicolor* KRAATZ, 1879

Distribution: Asian Russia (Amurland, Maritime Prov., Kurile Is.), China.

Subfamily: *Criocerinae* LATREILLE, 1807

Tribe: *Criocerini* LATREILLE, 1807

Genus: *Liliocerus* REITTER, 1912

***Liliocerus lili* (SCOPOLI, 1763)**

*Attelabus lili* SCOPOLI, 1763

*Chrysomela stercoraria* LINNAEUS, 1767

*Criocerus merdigera* FABRICIUS, 1775

*Criocerus liliorum* THOMSON, 1866

*Criocerus regeli* BALLION, 1878

*Liliocerus amurensis* PIC, 1916

*Liliocerus lili* ab. *schepmani* PIC, 1934

Distribution: Europe except for the North, SE Kazakhstan (Dzhungar Alatau), Asian Russia (W and Middle Siberia, Amurland), N Africa.

***Liliocerus merdigera* (LINNAEUS, 1758)**

*Chrysomela merdigera* LINNAEUS, 1758

*Criocerus rufipes* HERBST, 1783

*Criocerus brunnea* FABRICIUS, 1792

*Criocerus merdigera* ab. *suffriani* W. SCHMIDT, 1842

*Criocerus merdigera* ab. *medicana* LACORDAIRE, 1845

*Criocerus merdigera* ab. *collaris* LACORDAIRE, 1845

*Criocerus laticollis* REITTER, 1893

*Criocerus wagneri* JACOBSON, 1893

Distribution: All over the Palaearctic except for the extreme North, S and Central Asia, N America.

***Liliocerus ruficollis* (BALY, 1865)**

*Criocerus ruficollis* BALY, 1865

*Criocerus sieversi* HEYDEN, 1887

Distribution: Asian Russia (S Maritime Prov.), China, Korea.

***Liliocerus rugata* (BALY, 1865)**

*Criocerus rugata* BALY, 1865

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

***Liliocerus scapularis* (BALY, 1859)**

*Criocerus scapularis* BALY, 1859

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

Genus: *Criocerus* GEOFFROY, 1762

***Criocerus asparagi* (LINNAEUS, 1758)**

*Chrysomela asparagi* LINNAEUS, 1758

*Chrysomela campestris* LINNAEUS, 1767

*Lema pupillata* AHRENS, 1812

*Lema maculipes* GEBLER, 1834

*Criocerus asparagi* ab. *limaei* PIC, 1895

*Criocerus asparagi* ab. *anticeconjuncta* PIC, 1900

*Criocerus asparagi* ab. *apiceconjuncta* PIC, 1900

- Crioceris asparagi* ab. *impupillata* PIC, 1900  
*Crioceris asparagi* ab. *incrucifer* PIC, 1900  
*Crioceris asparagi* ab. *trifasciata* SCHUSTER, 1905  
*Crioceris asparagi* ab. *quadripunctata* SCHUSTER, 1905  
*Crioceris asparagi* ab. *pici* HEYDEN, 1906  
*Crioceris asparagi* ab. *schusteri* HEYDEN, 1906  
*Crioceris asparagi turkestanica* L. MEDVEDEV, 1955

Distribution: Europe except for the North, Mediterranean (except for Apennines Peninsula, Sicily, Corsica, Sardinia, Crete), Caucasus, Transeaucasia, Turkmenistan.  
 Introduced to N America and Argentina.

***Crioceris bicrucjata* (C. R. SAHLBERG, 1823)**

- Lema bicrucjata* C. R. SAHLBERG, 1823  
*Crioceris distincta* LACORDAIRE, 1845  
*Crioceris decorata* MORAWITZ, 1860

Distribution: SE European Russia (Volgograd Distr.), Caucasus, Greece, Turkey, Central Asia.

***Crioceris duodecimpunctata duodecimpunctata* (LINNAEUS, 1758)**

- Chrysomela 12-punctata* LINNAEUS, 1758  
*Lema dodecastigma* SUTERIAN, 1841  
*Crioceris duodecimpunctata* ab. *bisbiconnexa* PIC, 1919  
*Crioceris duodecimpunctata* ab. *delagrangei* PIC, 1919  
*Crioceris duodecimpunctata* ab. *reyyi* KASZAB, 1962  
*Crioceris duodecimpunctata* ab. *gisellae* KASZAB, 1962  
*Crioceris duodecimpunctata* ab. *ruffi* KASZAB, 1962  
*Crioceris duodecimpunctata* ab. *guranyii* KASZAB, 1962  
*Crioceris duodecimpunctata* ab. *bisquadripunctata* KASZAB, 1962  
*Crioceris duodecimpunctata* ab. *steinmanni* EODOR, 1965

Distribution: All over the Palaearctic except the North.

***Crioceris duodecimpunctata hypopsila* JACOBSON, 1907**

- Crioceris demokidovi* SEMENOV, 1908

Distribution: S Kazakhstan, Central Asia.

***Crioceris duodecimpunctata orientalis* (JACOBY, 1885)**

- Crioceris orientalis* JACOBY, 1885  
*Crioceris duodecimpunctata tshingisana* LOPATIN, 1966

Distribution: Asian Russia (Amurland, Khabarovsk Prov., Maritime Prov.), China, Korea, Japan.

***Crioceris iliensis* J. WEISE, 1900**

Distribution: Kazakhstan, Turkmenistan.

***Crioceris oshanini* DOHRN, 1884**

- Crioceris egregia* J. WEISE, 1900  
*Crioceris koltzei* J. WEISE, 1892

Distribution: Kazakhstan, Central Asia.

***Crioceris quatuordecimpunctata quatuordecimpunctata* (SCOPOLI, 1763)**

- Attelabus 14-punctatus* SCOPOLI, 1763  
*Crioceris quatuordecimpunctata* ab. *russica* PIC, 1916  
*Crioceris quatuordecimpunctata* ab. *moravica* ROUBAL, 1946

*Crioceris quatuordecimpunctata* ab. *parvipunctata* ROUBAL, 1949  
*Crioceris quatuordecimpunctata* ab. *deficiens* ROUBAL, 1949  
*Crioceris quatuordecimpunctata* ab. *internepunctata* CSIKI, 1953  
*Crioceris quatuordecimpunctata* ab. *externepunctata* CSIKI, 1953  
*Crioceris quatuordecimpunctata* ab. *lichtneckerti* KASZAB, 1962  
*Crioceris quatuordecimpunctata* ab. *csikii* KASZAB, 1962  
*Crioceris quatuordecimpunctata* *tshingisana* ab. *imitatrix* LOPATIN, 1967  
 Distribution: Central and SE Europe.

***Crioceris quatuordecimpunctata sibirica* J. WEISE, 1887**

Distribution: Asian Russia (Amurland, Khabarovsk Prov., Maritime Prov.), NE China, Korea, Japan.

***Crioceris quinquepunctata* (SCOPOLI, 1763)**

*Attelabus 5-punctatus* SCOPOLI, 1763  
*Crioceris quinquepunctata* ab. *thoracica* J. WEISE, 1893  
*Crioceris quinquepunctata* ab. *kaufmanni* KASZAB, 1962  
 Distribution: Mountains of Central and East Europe, Dinars Mts., S Belarus, Ukraine, S European Russia.

Tribe: *Lemiini* GYLLENHAL, 1813

Genus: *Oulema* DES GOZIS, 1886

*Oulema* BEDEL, 1889

***Oulema dilutipes* (FAIRMAIRE, 1885)**

*Lema dilutipes* FAIRMAIRE, 1885  
*Lema kajimurai* OHNO, 1962  
 Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

***Oulema duftschmidi* (L. REDTENBACHER, 1874)**

*Lema duftschmidi* L. REDTENBACHER, 1874  
 Distribution: Europe, Kazakhstan, Central Asia.

***Oulema erichsonii* (SUFFRIAN, 1841)**

*Lema erichsonii* SUFFRIAN, 1841  
*Lema erichsonii* var. *lipperti* GREDLER, 1866  
*Lema weisei* SEIDLITZ, 1891  
*Lema sapporoensis* KUWAYAMA, 1932  
*Lema viridula* GRESSITT, 1942  
 Distribution: Europe except for the extreme North and South, Asian Russia (Siberia, Maritime Prov.), Mongolia, Japan.

***Oulema gallaeciana* (HEYDEN, 1870)**

*Lema gallaeciana* HEYDEN, 1870  
*Crioceris cyanella* PAYKULL, 1799, non *Chrysomela cyanella* LINNAEUS, 1758  
*Chrysomela lichenis* VOET, 1806  
*Crioceris obscura* STEPHENS, 1829, non *Lema obscura* FABRICIUS, 1801  
*Lema lichenis* J. WEISE, 1881  
*Lema cyanella* PAYKULL, 1799, non LINNAEUS, 1758



*Lema nigra* DELHERM DE LARCENE, 1885

Distribution: All over Palaearctic except for the Kazakhstan and Central Asia.

***Oulema melanopus* (LINNAEUS, 1758)**

*Chrysomela melanopus* LINNAEUS, 1758

*Chrysomela melanopa* LINNAEUS, 1761

*Lema cyanella* var. *atrata* WALT., 1835

Distribution: Europe, Kazakhstan, Asian Russia (S Siberia east to Mongolia).

***Oulema oryzae* (KUWAYAMA, 1929)**

*Lema oryzae* KUWAYAMA, 1929

*Lema suvorovi* JACOBSON, 1931

Distribution: Asian Russia (Maritime Prov.), China, Japan.

***Oulema pygmaea* KRAATZ, 1879**

Distribution: Asian Russia (Amurland, Khabarovsk Prov., Maritime Prov.), China, Korea.

***Oulema septentrionis* J. WEISE, 1880**

Distribution: Europe except for the extreme North and South.

***Oulema tristis* (HERBST, 1786)**

*Crioceris tristis* HERBST, 1786

*Lema flavipes* W. SCHMIDT, 1841

Distribution: Europe, Asian Russia (S Siberia, Amurland), Kazakhstan, Korea, Japan.

**Genus: *Lema* FABRICIUS, 1798**

**Subgenus: *Lema* (*Lema*) FABRICIUS, 1798**

***Lema* (*Lema*) *cyanella* (LINNAEUS, 1758)**

*Chrysomela cyanella* LINNAEUS, 1758

*Crioceris puncticollis* CURTIS, 1830

*Lema rugicollis* SUFFERIAN, 1841

*Lema cirsicola* CHIJÓ, 1959

Distribution: All over Palaearctic except for the Central Asia.

***Lema* (*Lema*) *diversa* BALY, 1873**

*Lema lewisi* BALY, 1873

Distribution: Asian Russia (S Maritime Prov.), Korea, China, Japan.

***Lema* (*Lema*) *scutellaris* (KRAATZ, 1879)**

*Oulema scutellaris* KRAATZ, 1879

Distribution: Asian Russia (Amurland, Maritime Prov.), Korea, China, Japan.

**Subgenus: *Lema* (*Microlema*) PIC, 1932**

***Lema* (*Microlema*) *decempunctata* GEBLER, 1830**

Distribution: Asian Russia (Altai, Maritime Prov.), Mongolia, N China, Japan, Kazakhstan.

***Lema* (*Microlema*) *quadrinaculata* GEBLER, 1845**

*Lema akinini* HEYDEN, 1887

*Lema margelanica* J. WEISE, 1894

Distribution: Kazakhstan, Uzbekistan, Kirghizstan.

Subfamily: *Clytrinae* KIRBY, 1837

Tribe: *Clytrini* KIRBY, 1837

Genus: *Labidostomis* GERMAR, 1822

Subgenus: *Labidostomis (Labidostomis)* GERMAR, 1822

***Labidostomis (Labidostomis) amurensis* HEYDEN, 1884**

*Labidostomis orientalis* CHŪJŌ, 1940

Distribution: Asian Russia (Amurland, Khabarovsk Prov., Maritime Prov.), China, Korea.

***Labidostomis (Labidostomis) arcuata* PIC, 1920**

Distribution: S Kazakhstan, Kirghizstan, NW China.

***Labidostomis (Labidostomis) arnoldii* L. MEDVEDEV, 1962**

Distribution: Northern Caucasus.

***Labidostomis (Labidostomis) asiatica* FALDERMANN, 1837**

Distribution: Transcaucasia.

***Labidostomis (Labidostomis) beckeri* J. WEISE, 1882**

Distribution: S and SE parts of European Russia, Ukraine, Crimea, Moldavia, Ciscaucasia, E Kazakhstan, Asian Russia (W Siberia).

***Labidostomis (Labidostomis) bipunctata bipunctata* (MANNERHEIM, 1825)**

*Clytra bipunctata* MANNERHEIM, 1825

Distribution: Asian Russia (Altai, Tuva, Sayan, Transbaikalia, Russian Far East), Mongolia.

***Labidostomis (Labidostomis) bipunctata indigiriensis* L. MEDVEDEV, 1978**

Distribution: Asian Russia (Yakutia).

***Labidostomis (Labidostomis) brevipennis* FALDERMANN, 1837**

*Labidostomis armeniaca* LACORDAIRE, 1848

Distribution: Transcaucasia, Turkey.

***Labidostomis (Labidostomis) chinensis* LEFÈVRE, 1887**

Distribution: Asian Russia (Transbaikalia, Amurland, Maritime Prov.), Mongolia, China, Korea.

***Labidostomis (Labidostomis) crebrecollis* L. MEDVEDEV, 1962**

Distribution: Asian Russia (Transbaikalia, Amurland, Maritime Prov.), Mongolia, Korea.

***Labidostomis (Labidostomis) curtus* LOPATIN, 1961**

Distribution: Kazakhstan Plains, S Uzbekistan, S Tajikistan.

***Labidostomis (Labidostomis) cyanicornis* GERMAR, 1822**

Distribution: SE and southern parts of Central and E Europe, Ciscaucasia, N Kazakhstan, Asian Russia (W Siberia, Baikal Area, Tuva, Sayan, Yakutia), Mongolia.

***Labidostomis (Labidostomis) decipiens* FALDERMANN, 1837**

Distribution: Balkans, Transcaucasia, Turkey, Syria, N Iran.

***Labidostomis (Labidostomis) elegans* LEFÈVRE, 1876***Labidostomis reitteri* J. WEISE, 1885

Distribution: Transcaucasia, Turkey, N Iran.

***Labidostomis (Labidostomis) fedtschenkoi* LOPATIN, 1963**

Distribution: Uzbekistan, Tajikistan (Mountains).

***Labidostomis (Labidostomis) glasunovi* JACOBSON, 1893**

Distribution: Tajikistan (Mountains), Pamirs.

***Labidostomis (Labidostomis) humeralis* (D. H. SCHNEIDER, 1792)***Clytra humeralis* D. H. SCHNEIDER, 1792

Distribution: S, Central and the southern part of E Europe, N Kazakhstan, Turkey.

***Labidostomis (Labidostomis) kaszabi* L. MEDVEDEV, 1962**

Distribution: Turkey.

***Labidostomis (Labidostomis) lepida* LEFÈVRE, 1872**

Distribution: Latvia, Lithuania, Belarus, Ukraine, W European Russia.

***Labidostomis (Labidostomis) lipskyi* LOPATIN, 1961**

Distribution: Tajikistan.

***Labidostomis (Labidostomis) longimana longimana* (LINNAEUS, 1761)***Chrysomela longimana* LINNAEUS, 1761*Labidostomis longimana* ab. *obscura* TRELLA, 1930

Distribution: W, Central and E Europe, Kazakhstan, Asian Russia (S Siberia, Altai).

***Labidostomis (Labidostomis) longimana dalmatina* (LACORDAIRE, 1848)***Clythra dalmatina* LACORDAIRE, 1848*Clythra uralensis* LACORDAIRE, 1848*Labidostomis glycyrrhizae* BAKER, 1880*Labidostomis caucasica* L. MEDVEDEV, 1962

Distribution: S Europe, Crimea, Caucasus, Turkey.

***Labidostomis (Labidostomis) lucida lucida* (GERMAR, 1824)***Clytra lucida* GERMAR, 1824*Labidostomis fulgida* LACORDAIRE, 1848

Distribution: W, Central, and western parts of East Europe.

***Labidostomis (Labidostomis) lucida axillaris* (LACORDAIRE, 1848)***Clythra axillaris* LACORDAIRE, 1848*Labidostomis sareptana* KRAATZ, 1872

Distribution: Steppe zone of Palaeartic, W Iran.

***Labidostomis (Labidostomis) maculipennis* LEFÈVRE, 1870***Labidostomis kindermanni* KRAATZ, 1872

Distribution: Transcaucasia, Turkey.

***Labidostomis (Labidostomis) medvedevi* WARCHALOWSKI, 1985***Labidostomis montana* L. MEDVEDEV, 1970, non COBOS, 1954

Distribution: SE Azerbaijan (Talysh), W Turkmenistan, NW Iran.

***Labidostomis (Labidostomis) mesopotamica peregrina* J. WEISE, 1900***Labidostomis escherichi* J. WEISE, 1897*Labidostomis frontalis* J. WEISE, 1900*Labidostomis iconiensis* J. WEISE, 1900

Distribution: Daghestan, Transcaucasia, E Turkey.

***Labidostomis (Labidostomis) metallica metallica* LEFÈVRE, 1872**

Distribution: SE part of European Russia, Daghestan, W Kazakhstan.

***Labidostomis (Labidostomis) metallica centrisculpta* PIC, 1920**

Distribution: S Kazakhstan, Kirghizstan, SE Tajikistan.

***Labidostomis (Labidostomis) metallica dzhungarica* L. MEDVEDEV, 1980**

Distribution: Mongolia, NW China.

***Labidostomis (Labidostomis) metallica imitatrix* L. MEDVEDEV, 1971**

Distribution: Asian Russia (Yakutia).

***Labidostomis (Labidostomis) metallica mongolica* L. MEDVEDEV, 1980**

Distribution: Asian Russia (Tuva), Mongolia.

***Labidostomis (Labidostomis) metallica ornatipennis* L. MEDVEDEV, 1971**

Distribution: S Kazakhstan, Kirghizstan (W Tian-Shan).

***Labidostomis (Labidostomis) metallica steppensis* L. MEDVEDEV, 1971**

Distribution: Asian Russia (Altai), N and E Kazakhstan.

***Labidostomis (Labidostomis) oertzeni villosula* J. WEISE, 1889***Labidostomis heteromalla* LEFÈVRE, 1889

Distribution: Transcaucasia, E Turkey.

***Labidostomis (Labidostomis) ornatus* L. MEDVEDEV, 1962***Labidostomis signatipennis* LOPATIN, 1963

Distribution: S Kazakhstan (W Tian-Shan).

***Labidostomis (Labidostomis) pachysoma* L. MEDVEDEV, 1965**

Distribution: SE of European Russia, Steppe of Crimea Peninsula, Daghestan, Georgia.

***Labidostomis (Labidostomis) pallidipennis* (GEBLER, 1830)***Clythra pallidipennis* GEBLER, 1830*Clythra chalybeicornis* BRULLÉ, 1832*Labidostomis longipennis* GEBLER, 1841*Clythra distinguenda* ROSENHAUER, 1847*Clythra pilicollis* LACORDAIRE, 1848.

Distribution: S and southern parts of E Europe, Caucasus, Transcaucasia, Turkey, Kazakhstan, Asian Russia (Siberia).

***Labidostomis (Labidostomis) propinqua* FALDERMANN, 1837**

Distribution: Balkans, Ukraine (Crimea), Caucasus, Transcaucasia, Turkey.

***Labidostomis (Labidostomis) rugicollis* LEFÈVRE, 1872**

Distribution: Tajikistan, S Turkmenistan (Kopetdagh Mountains), N Iran.

***Labidostomis (Labidostomis) schneideri* J. WEISE, 1900***Labidostomis polygoni* LOPATIN, 1961

Distribution: Uzbekistan, Tajikistan.

***Labidostomis (Labidostomis) senicula* KRAATZ, 1872***Labidostomis nitida* BALLIION, 1878*Labidostomis albida* JACOBSON, 1894*Labidostomis turcomanica* LOPATIN, 1954

Distribution: SE of European Russia, Kazakhstan, Central Asia, Mongolia, NW China.

***Labidostomis (Labidostomis) sibirica sibirica* (GERMAR, 1824)***Clythra sibirica* GERMAR, 1824

Distribution: Asian Russia (Siberia), N Mongolia, N China.

***Labidostomis (Labidostomis) sibirica tjutschewi* JACOBSON, 1900**

Distribution: Asian Russia (Tuva, Transbaikalia), N Mongolia.

***Labidostomis (Labidostomis) sibirica transitoria* JACOBSON, 1900**

Distribution: Asian Russia (Altai, S Siberia).

***Labidostomis (Labidostomis) stenostoma* J. WEISE, 1900***Labidostomis alaiensis* PIC, 1920

Distribution: E Uzbekistan, Tajikistan, S Kirghizstan.

***Labidostomis (Labidostomis) subfasciata* J. WEISE, 1885***Labidostomis trigemina* J. WEISE, 1898

Distribution: Azerbaijan, S Turkmenia, N and W Iran.

***Labidostomis (Labidostomis) sulcicollis* (LACORDAIRE, 1848)***Clythra sulcicollis* LACORDAIRE, 1848

Distribution: Armenia, Turkey.

***Labidostomis (Labidostomis) tridentata* (LINNAEUS, 1758)***Chrysomela tridentata* LINNAEUS, 1758*Clythra kluczyckii* M. NOWICKI, 1872*Labidostomis leithneri* L. REDTENBACHER, 1874

Distribution: Europe, Asian Russia (S Siberia, Russian Far East), Mongolia, N China.

**Genus: *Antipus* DEGEER, 1778***Antipa* auct.**Subgenus: *Antipus (Tituboea)* LACORDAIRE, 1848*****Antipus (Tituboea) macropus* ILLIGER, 1800***Tituboea macropus* ab. *binaeva* REITTER, 1912*Tituboea macropus* ab. *armeniaca* PIC, 1918

Distribution: S Europe, Caucasus, Turkey, Syria, Iran, Turkmenistan, Uzbekistan.

***Antipus (Tituboea) nigriventris* LEFÈVRE, 1872**

Distribution: S Europe, Caucasus, Turkey, Syria, Iran, Turkmenistan, Uzbekistan.

***Antipus (Tituboea) puncticollis* L. MEDVEDEV, 1962**

Distribution: N Iran, Turkmenistan.

***Antipus (Tituboea) rugulosa* J. WEISE, 1898**

Distribution: S Transcaucasia.

***Antipus (Tituboea) silensis* J. WEISE, 1894**

Distribution: Turkmenistan, Uzbekistan, Tajikistan, S Kazakhstan, E Iran.

**Genus: *Lachnaia* CHEVROLAT in DEJEAN 1837***Lachnaea* Agassiz, 1846***Lachnaia sexpunctata* (SCOPOLI, 1763)***Buprestis sexpunctata* SCOPOLI, 1763*Cryptocephalus longipes* FABRICIUS, 1775*Lachnaea sexpunctata* ab. *pontica* J. WEISE, 1884

Distribution: W and S Europe, S Ukraine, Turkey.

**Genus: *Clytra* LAICHARTING, 1781***Clythra* FABRICIUS, 1798**Subgenus: *Clytra (Clytraria)* SEMENOV, 1903***Clytrella* L. MEDVEDEV, 1961*Clitrella* L. MEDVEDEV, 1961***Clytra (Clytraria) atraphaxidis atraphaxidis* (PALLAS, 1773)***Chrysomela atraphaxidis* PALLAS, 1773

Distribution: S Europe, the southern part of E Europe, Caucasus, N and E Kazakhstan, Mountains of Central Asia, Turkey, W Iran.

***Clytra (Clytraria) atraphaxidis asiatica* CHŪJŌ, 1941***Clytra atraphaxidis* var. *asiatica* CHŪJŌ, 1941

Distribution: Asian Russia (Amurland), China, Korea.

***Clytra (Clytraria) atraphaxidis maculifrons* ZUBKOV, 1833**

Distribution: SE of European Russia, N and Middle Kazakhstan, Plains of Central Asia, N Iran.

***Clytra (Clytraria) atraphaxidis punctata* J. WEISE, 1890**

Distribution: Asian Russia (Altai), E Kazakhstan, Mongolia, NW China.

***Clytra (Clytraria) jacobsoni* SEMENOV, 1903***Clytra opaca* JACOBSON, 1898, non ROSENHAUER, 1847*Clytra bucharica* PIC, 1915

Distribution: Tajikistan.

***Clytra (Clytraria) novempunctata* Olivier, 1808***Clytra persica* PIC, 1920

Distribution: Balkans, Daghestan, Transcaucasia, Turkey, W Iran, Iraq, Turkmenistan, NE Africa.

***Clytra (Clytraria) popovi* L. MEDVEDEV, 1961**

Distribution: Tajikistan.

***Clytra (Clytraria) rufina* (SOLSKY, 1882)***Coptocephala rufina* SOLSKY, 1882

Distribution: S Kazakhstan, Deserts of Central Asia.

***Clytra (Clytraria) valerianae valerianae* MÉNÉTRIÉS, 1832***Clythra valerianae* MÉNÉTRIÉS, 1832

Distribution: Balkans, southern E Europe, Caucasus, Transcaucasia, Turkey, NW Iran.

***Clytra (Clytraria) valerianae tetrastigma* (W. SCHMIDT, 1841)***Clythra valerianae tetrastigma* W. SCHMIDT, 1841*Clytra valerianae iranica* L. MEDVEDEV, 1961*Clytra valerianae taurica* L. MEDVEDEV, 1961*Clytra valerianae opacipennis* Lopatin, 1962

Distribution: S Ukraine, Crimea, NW Iran.

**Subgenus: *Clytra (Clytra)* LAICHARTING, 1781*****Clytra (Clytra) arida* J. WEISE, 1889**Distribution: Asian Russia (W Siberia, Tuva, Sayan, Transbaikalia, Russian Far East),  
Mongolia, China, Korea.***Clytra (Clytra) laeviuscula* (RATZEBURG, 1837)***Clythra fasciata* RATZEBURG, 1837, non FABRICIUS, 1801*Clythra laeviuscula* RATZEBURG, 1837*Clythra laeviuscula* var. *connexa* FRICKEN, 1888*Clythra laeviuscula* ab. *antistita* J. WEISE, 1889*Clythra laeviuscula* ab. *biinterrupta* REITTER, 1912Distribution: Central, S and southern parts of E Europe, Ciscaucasia, Transcaucasia,  
Turkey, N and E Kazakhstan, E Kirghizstan, Afghanistan, N Tajikistan, NW China.***Clytra (Clytra) quadripunctata quadripunctata* (LINNAEUS, 1758)***Chrysomela 4-punctata* LINNAEUS, 1758*Clytra quadripunctata* var. *dissimilis* J. WEISE, 1889*Clytra messae* G. MÜLLER, 1921Distribution: Europe, Caucasus, Transcaucasia, Turkey, N Iran, Kazakhstan, Asian  
Russia (southern W and Middle Siberia, Amurland).***Clytra (Clytra) quadripunctata appendicina* (LACORDAIRE, 1848)***Clythra appendicina* LACORDAIRE, 1848

Distribution: Southern part of Central and W Europe, Kazakhstan, Central Asia.

***Clytra quadripunctata turfanica* LOPATIN, 1962**

Distribution: China (Turfan).

**Genus: *Smaragdina* CHEVROLAT in DEJEAN, 1837***Cyaniris* CHEVROLAT in DEJEAN, 1837, non DALMAN, 1816 (*Lepidoptera*)*Gynandrophthalma* LACORDAIRE, 1848*Gynandrophthalma* BACH, 1856



**Subgenus: *Smaragdina (Calyptorhina)* LACORDAIRE, 1848**

***Smaragdina (Calyptorhina) aeneoviridis* LOPATIN, 1975**

Distribution: S Tajikistan.

***Smaragdina (Calyptorhina) biornata* LEFÈVRE, 1872**

Distribution: Transcaucasia, Turkey, W Iran.

***Smaragdina (Calyptorhina) chloris caucasica* L. MEDVEDEV, 1971**

Distribution: SW Ciscaucasia, Abkhazia.

***Smaragdina (Calyptorhina) golda* JACOBSON, 1925**

Distribution: Asian Russia (S Amurland, Maritime Prov.), Korea, Japan

***Smaragdina (Calyptorhina) limbata* (STEVEN, 1806)**

*Clytra limbata* STEVEN, 1806

*Clytra dorsalis* OLIVIER, 1808

Distribution: Balkans, Romania, Hungary, E Ciscaucasia, Transcaucasia, Turkey, Near and Middle East.

***Smaragdina (Calyptorhina) unipunctata* (OLIVIER, 1808)**

*Clytra unipunctata* OLIVIER, 1808

*Clytra menetriesi* FALDERMANN, 1837

Distribution: Transcaucasia, Turkey, Middle East, Turkmenistan.

***Smaragdina (Calyptorhina) viridis viridis* (KRAATZ, 1882)**

*Gynandrophthalma viridis* KRAATZ, 1882

Distribution: S Kazakhstan, Turkey.

***Smaragdina (Calyptorhina) viridis comans* (JACOBSON, 1925)**

*Calyptorhina comans* JACOBSON, 1925

Distribution: SE Azerbaijan, Turkmenistan, N Iran.

**Subgenus: *Smaragdina (Nanosmaragdina)* LOPATIN et KULENOVA, 1985**

*Smaragdinella* L. MEDVEDEV, 1971

***Smaragdina (Nanosmaragdina) macilenta* (J. WEISE, 1887)**

*Gynandrophthalma macilenta* J. WEISE, 1887

Distribution: Kazakhstan (Chu-Ili Mountains), Uzbekistan, W Tajikistan.

**Subgenus: *Smaragdina (Monrosiana)* L. MEDVEDEV, 1992**

*Monrosia* L. MEDVEDEV, 1971

***Smaragdina (Monrosiana) affinis* (ILLIGER, 1794)**

*Clytra affinis* ILLIGER, 1794

*Smaragdina aurita* ab. *noesskei* SCHATZMAYR, 1928

*Smaragdina aurita* ab. *trimaculata* PORTA, 1934

Distribution: S, Central and E Europe, Turkey, N Africa.

***Smaragdina (Monrosiana) aurita aurita* (LINNAEUS, 1767)**

*Chrysomela aurita* LINNAEUS, 1767

Distribution: Europe except for the North, Turkey.

***Smaragdina (Monrosiana) aurita auritoides* ACHARD, 1923**

Distribution: Caucasus, NW Iran, NE Turkey.

***Smaragdina (Monrosiana) aurita hammarstroemi* (JACOBSON, 1901)***Gynandrophthalma aurita* var. *hammarstroemi* JACOBSON, 1901

Distribution: Asian Russia (Siberia, Russian Far East), Mongolia.

***Smaragdina (Monrosiana) aurita nigrocyanea* (MOTSCHULSKY, 1866)***Gynandrophthalma nigrocyanea* MOTSCHULSKY, 1866

Distribution: Asian Russia (Sakhalin, Kurile Is.), Japan.

***Smaragdina (Monrosiana) collaris collaris* (FABRICIUS, 1781)***Cryptocephalus collaris* FABRICIUS, 1781*Clytra lateralis* GEBLER, 1848

Distribution: Kazakhstan, Asian Russia (Tuva, E Siberia), Mongolia, China.

***Smaragdina (Monrosiana) collaris lateralis* GEBLER, 1848**

Distribution: E Kazakhstan, Asian Russia (Altai, Transbaikalia, Yakutia).

***Smaragdina (Monrosiana) discolor discolor* (SOLSKY, 1881)***Gynandrophthalma discolor* SOLSKY, 1881

Distribution: Kazakhstan, Central Asia.

***Smaragdina (Monrosiana) discolor rufilabris* (J. WEISE, 1887)***Gynandrophthalma rufilabris* J. WEISE, 1887

Distribution: Kazakhstan (Chu-Ili Mountains).

***Smaragdina (Monrosiana) discolor viridiceps* (J. WEISE, 1892)***Gynandrophthalma viridiceps* J. WEISE, 1892*Cyaniris ignorata* LOPATIN, 1954

Distribution: Mountains of the SE Kazakhstan and Central Asia (except for the Turkmenistan).

***Smaragdina (Monrosiana) flavicollis* (CHARPENTIER, 1825)***Clytra flavicollis* CHARPENTIER, 1825*Clytra diversipes* LETZNER, 1839*Gynandrophthalma flavicollis* ab. *picticollis* J. WEISE, 1889

Distribution: Europe except for the extreme North and South, Turkey, Algeria.

***Smaragdina (Monrosiana) hypocrita* (LACORDAIRE, 1848)***Clytra hypocrita* LACORDAIRE, 1848

Distribution: Serbia, Bulgaria, Ukraine (S Crimea), Caucasus, Turkey.

***Smaragdina (Monrosiana) labilis sahlbergi* (JACOBSON, 1901)***Gynandrophthalma sahlbergi* JACOBSON, 1901*Smaragdina discithorax* PIC, 1902*Smaragdina discalis* CHŪJŌ, 1941

Distribution: Asian Russia (Transbaikalia, Amurland, Maritime Prov.), Mongolia, Korea.

***Smaragdina (Monrosiana) obscuripes* (J. WEISE, 1887)***Gynandrophthalma obscuripes* J. WEISE, 1887*Smaragdina compresipennis* PIC, 1927*Smaragdina obscuricolor* PIC, 1927

Distribution: Asian Russia (Transbaikalia, Maritime Prov.), N China.

***Smaragdina (Monrosiana) potanini* L. MEDVEDEV, 1970**

Distribution: Asian Russia (S Maritime Prov.), China.

***Smaragdina (Monrosiana) punctatissima* J. WEISE, 1892**

Distribution: S Kazakhstan, Kirghizstan.

***Smaragdina (Monrosiana) salicina* (SCOPOLI, 1763)***Buprestis salicina* SCOPOLI, 1763*Cryptocephalus cyaneus* FABRICIUS, 1775*Smaragdina salicina* ab. *notaticollis* ROUBAL, 1922*Smaragdina salicina* ab. *nigripennis* TRELIA, 1930

Distribution: Europe, Asian Russia (W Siberia), Caucasus, Turkey, E Kazakhstan.

***Smaragdina (Monrosiana) semiauratiaca* FAIRMAIRE, 1888**

Distribution: Asian Russia (S Amurland, S Maritime Prov.), China, Korea, Japan.

***Smaragdina (Monrosiana) stenroosi* (JACOBSON, 1901)***Gynandrophthalma stenroosi* JACOBSON, 1901

Distribution: S Kazakhstan, Kirghizstan.

***Smaragdina (Monrosiana) thoracica thoracica* FISCHER-WALDHEIM, 1842**

Distribution: Asian Russia (Altai, Siberia), Mountains of Kirghizstan, Tajikistan, SE Kazakhstan.

***Smaragdina (Monrosiana) thoracica dzungarica* L. MEDVEDEV, 1980**

Distribution: NW China.

***Smaragdina (Monrosiana) xanthaspis* (GERMAR, 1824)***Clytra xanthaspis* GERMAR, 1824*Cyaniris thoracica* KÜSTER, 1846

Distribution: SE and southern parts of Central Europe, Transcaucasia, Turkey.

**Genus: *Cheilotoma* CHEVROLAT in DEJEAN, 1837***Chilotoma* Agassiz, 1846**Subgenus: *Cheilotoma (Cheilotoma)* CHEVROLAT in DEJEAN, 1837*****Cheilotoma (Cheilotoma) erythrostroma* (FALDERMANN, 1837)***Cheilotoma erythrostroma* CHEVROLAT in DEJEAN, 1836, 1837 [nomen nudum]*Cheilotoma erythrostroma* FALDERMANN, 1837

Distribution: SE Europe, NW Caucasus, Transcaucasia, Iran.

***Cheilotoma (Cheilotoma) musciformis* (GOEZE, 1777)***Chrysomela musciformis* GOEZE, 1777*Clytra bucephala* SCHALLER, 1783*Cheilotoma musciformis* ab. *orientalis* JACOBSON, 1901*Cheilotoma musciformis* ab. *leosinii* LEONI, 1906*Cheilotoma musciformis* ab. *siegae* G. MÜLLER, 1921

Distribution: S Europe, Caucasus, Kazakhstan, Asian Russia (southern parts of W and Middle Siberia, Transbaikalia).

**Subgenus: *Cheilotoma (Exaesiognatha)* JACOBSON, 1923*****Cheilotoma (Exaesiognatha) ivanovi* JACOBSON, 1923***Cheilotoma ivanovi turcomanica* L. MEDVEDEV, 1971

Distribution: Plains of Central Asia from W Turkmenistan to Fergana Valley and S Tajikistan.

**Genus: *Coptocephala* CHEVROLAT in DEJEAN, 1837****Subgenus: *Coptocephala (Coptocephala)* CHEVROLAT in DEJEAN, 1837*****Coptocephala (Coptocephala) chalybaea chalybaea* (GERMAR, 1824)***Clytra chalybaea* GERMAR, 1824

Distribution: SE Europe, NW and Minor Caucasus, Daghestan.

***Coptocephala (Coptocephala) chalybaea apicalis* LACORDAIRE, 1848***Coptocephala apicalis* LACORDAIRE, 1848

Distribution: Steppe zone from S Ukraine and Crimea to Central Kazakhstan, Ciscaucasia.

***Coptocephala (Coptocephala) gebleri* GEBLER, 1841***Coptocephala gebleri* CHEVROLAT in DEJEAN, 1837 [nomen nudum]*Coptocephala gebleri* GEBLER, 1841

Distribution: SE Balkan, S Ukraine (Crimea) and European Russia, Caucasus, Transcaucasia, Kazakhstan, Kirghizstan, Asian Russia (southern parts of W Siberia, Altai).

***Coptocephala (Coptocephala) hissarica* LOPATIN, 1963***Coptocephala tadhika* L. MEDVEDEV, 1965

Distribution: SE Uzbekistan, W Tajikistan.

***Coptocephala (Coptocephala) orientalis* BALY, 1873***Coptocephala freija* REITTER, 1900*Coptocephala asiatica* CHŪJŌ, 1940

Distribution: Asian Russia (S Siberia, Amurland, Maritime Prov.), Mongolia, China, Japan.

***Coptocephala (Coptocephala) rubicunda rubicunda* (LAICHARTING, 1781)***Clytra rubicunda* LAICHARTING, 1781*Coptocephala tetradyma* KÜSTER, 1845*Coptocephala rubicunda* ab. *subobliterata* PIC, 1901*Coptocephala rubicunda* ab. *massiliensis* PIC, 1914*Coptocephala rubicunda* ab. *dalmatina* PIC, 1918*Coptocephala rubicunda* ab. *vittata* PETRI, 1926*Coptocephala rubicunda* f. *bimaculata* POLENTZ, 1939 [nomen nudum]*Coptocephala rubicunda* ab. *didymata* ROUBAL, 1948*Coptocephala rubicunda* ab. *connexa* CSIKI, 1953*Coptocephala rubicunda* ab. *efasciata* CSIKI, 1953*Coptocephala rossica* L. Medvedev, 1977

Distribution: SW, Central and southern E Europe, Ciscaucasia, Georgia, N Kazakhstan, Asian Russia (southern parts of W and Middle Siberia, Cisbaikalia).

***Coptocephala (Coptocephala) unifasciata unifasciata* (SCOPOLI, 1763)**

- Egyprestis unifasciata* SCOPOLI, 1763  
*Pyrosometa 4-maculata* LINNAEUS, 1767  
*Coptocephala femoralis* KÜSTER, 1845  
*Coptocephala quadrimaculata* LACORDAIRE, 1848, non LINNAEUS, 1767  
*Coptocephala quadrimaculata* fa. *maculiceps* KRAATZ, 1872  
*Coptocephala unifasciata* var. *fallax* J. WEISE, 1887  
*Coptocephala unifasciata* ab. *nigripes* JACOBSON, 1915

Distribution: Europe except for N, Transcaucasia, Turkey, Syria, E Kazakhstan, W Pamirs, Asian Russia (southern parts of W Siberia).

***Coptocephala (Coptocephala) unifasciata australis* L. MEDVEDEV, 1965**

Distribution: S Kazakhstan, Central Asia (Tian-Shan and Hissar-Darvaz Mountains).

***Coptocephala (Coptocephala) unifasciata deserta* L. MEDVEDEV, 1965**

Distribution: Desert and semi-desert areas of Kazakhstan and Central Asia.

***Coptocephala (Coptocephala) unifasciata destinoi* FAIRMAIRE, 1884**

*Coptocephala intremedia* REINECK, 1908

Distribution: Transcaucasia, Turkey, Near East.

Subfamily: *Cryptocephalinae* GYLLENHAL, 1813

Tribe: *Cryptocephalini* GYLLENHAL, 1813

Genus: *Cryptocephalus* GEOFFROY, 1762

*Cryptocephalus* O. F. MÜLLER, 1764

Subgenus: *Cryptocephalus (Protophysus)* CHEVROLAT in DEJEAN, 1837

*Protophysus* L. Redtenbacher, 1845

***Cryptocephalus (Protophysus) schaefferi* SCHRANK, 1789**

- Cryptocephalus haemorrhoidalis* OLIVIER, 1791  
*Cryptocephalus unicolor* OLIVIER, 1791  
*Cryptocephalus notatus* D. H. SCHNEIDER, 1792  
*Cryptocephalus lobatus* FABRICIUS, 1792  
*Cryptocephalus schaefferi* ab. *hungaricus* CSIKI, 1901  
*Cryptocephalus schaefferi* ab. *sabaudus* PIC, 1908  
*Cryptocephalus schaefferi* ab. *gynaecoides* ROUBAL, 1942  
*Cryptocephalus schaefferi* ab. *aubryi* TIBERGHEN, 1971  
*Cryptocephalus schaefferi* ab. *pulcher* TIBERGHEN, 1971

Distribution: SE and southern parts of Central Europe, Caucasus, Asian Russia (S Siberia east to Transbaikalia), Kazakhstan.

Subgenus: *Cryptocephalus (Disopus)* CHEVROLAT in DEJEAN, 1837

***Cryptocephalus (Disopus) pini difformis* (JACOBY, 1885)**

*Cryptocephalus difformis* JACOBY, 1885

Distribution: Asian Russia (E Siberia, Sakhalin), Japan.



***Cryptocephalus (Disopus) pini pini* (LINNAEUS, 1758)***Chrysomela pini* LINNAEUS, 1758*Cryptocephalus abietis* SUFFRIAN, 1847*Cryptocephalus porrii* SUFFRIAN, 1847*Cryptocephalus pini* ab. *apiceflavus* G. MÜLLER, 1948

Distribution: Europe except for the South, Asian Russia (W Siberia).

**Subgenus: *Cryptocephalus (Asionus)* LOPATIN, 1988***Asiopus* LOPATIN, 1965, non SPIX, 1825***Cryptocephalus (Asionus) altaicus* HAROLD, 1872***Cryptocephalus agnus* J. WEISE, 1898

Distribution: Asian Russia (Altai, Tuva, Sayan, Transbaikalia), Mongolia, China.

***Cryptocephalus (Asionus) apicalis* GEBLER, 1850***Cryptocephalus flavoguttatus* SUFFRIAN, 1847*Cryptocephalus apicalis* ab. *eburatus* J. WEISE, 1882*Cryptocephalus apicalis* ab. *rinskii* JACOBSON, 1907*Cryptocephalus apicalis* ab. *holtzi* PIC, 1909*Cryptocephalus apicalis* ab. *hortobagyanus* PIC, 1909*Cryptocephalus apicalis* ab. *delicatissimus* ROUBAL, 1940*Cryptocephalus apicalis* ab. *csepelensis* CSIKI, 1953*Cryptocephalus apicalis* ab. *limbipennis* CSIKI, 1953*Cryptocephalus apicalis* ab. *marginatipennis* CSIKI, 1953

Distribution: S Ukraine (Crimea), Moldavia, Ciscaucasia, Kazakhstan, Asian Russia (S Siberia).

***Cryptocephalus (Asionus) arnoldii arnoldii* L. MEDVEDEV, 1956**

Distribution: S Kazakhstan (W Tian-Shan Mountains), Uzbekistan (Baisuntau Mountain Range), SW Tajikistan.

***Cryptocephalus (Asionus) arnoldii sogdianus* LOPATIN, 1963**

Distribution: N Tajikistan (Zeravshan Mountain Range), N Uzbekistan (Nuratau Mountain Range).

***Cryptocephalus (Asionus) atriplicis* LOPATIN, 1967**

Distribution: Central and E Kazakhstan, Azerbaijan (Sumghait).

***Cryptocephalus (Asionus) balassogloi* JACOBSON, 1895**

Distribution: Kazakhstan, Uzbekistan, N Tajikistan.

***Cryptocephalus (Asionus) beckeri* MORAWITZ, 1860***Cryptocephalus suffriani* DOHRN, 1859, non SUFFRIAN, 1849

Distribution: SE Russian Plain (Povolzhie), NW Kazakhstan.

***Cryptocephalus (Asionus) bidentatus* L. MEDVEDEV, 1956**

Distribution: Kazakhstan (Talas Alatau), Kirghizstan (W Tian-Shan Mountain Ranges, Fergan Mountain Range), N Tajikistan (Kuramin Mountain Range).

***Cryptocephalus (Asionus) bidentulus bidentulus* SUFFRIAN, 1854**

Distribution: Asian Russia (Tuva, S Transbaikalia), N and E Mongolia.

- Cryptocephalus (Asionus) bidentulus klementzi* OGLOBLIN, 1956**  
Distribution: Asian Russia (Tuva), NW Mongolia.
- Cryptocephalus (Asionus) bidentulus subdesertus* L. MEDVEDEV, 1980**  
Distribution: S Mongolia.
- Cryptocephalus (Asionus) bitaeniatus* SOLSKY, 1876**  
Distribution: Uzbekistan, S Tajikistan, Turkmenistan (Amu-Darya Valley).
- Cryptocephalus (Asionus) bivulneratus* FALDERMANN, 1835**  
Distribution: Asian Russia (Tuva, Transbaikalia), Mongolia.
- Cryptocephalus (Asionus) bohemi* DRAPIEZ, 1819**  
*Cryptocephalus böhmi* GERMAR, 1824  
*Cryptocephalus semenovi* J. WEISE, 1889  
*Cryptocephalus tentator* J. WEISE, 1889  
Distribution: SE Europe, N and Central Kazakhstan.
- Cryptocephalus (Asionus) borochoensis* PIC, 1907**  
*Cryptocephalus kulenovae* LOPATIN, 1967  
Distribution: E Kazakhstan, NW China.
- Cryptocephalus (Asionus) crux* GEBLER, 1848**  
*Cryptocephalus equestris* GEBLER, 1830  
*Cryptocephalus manchuricus* GRESSITT et KIMOTO, 1961  
Distribution: E Kazakhstan, Asian Russia (S Siberia, Yakutia, Tuva, Russian Far East), Mongolia.
- Cryptocephalus (Asionus) curda* JACOBSON, 1897**  
Distribution: Transcaucasia.
- Cryptocephalus (Asionus) diadochus* LOPATIN, 1963**  
Distribution: Tajikistan (Zeravshan Mountain Range).
- Cryptocephalus (Asionus) dilectus* J. WEISE, 1894**  
Distribution: S Kazakhstan (W Tian-Shan), Kirghizstan, E Tajikistan.
- Cryptocephalus (Asionus) ergenensis* MORAWITZ, 1863**  
*Cryptocephalus beckeri* SUFFRIAN, 1864  
*Cryptocephalus ergenensis georgicus* L. MEDVEDEV et SEPTELADZE, 1962  
Distribution: NE Ciscaucasia (Ergeni), Georgia, S Urals, W and Central Kazakhstan.
- Cryptocephalus (Asionus) fasciatointerruptus* BERTI et RAPILLY, 1979**  
Distribution: Iran.
- Cryptocephalus (Asionus) flavicollis* FABRICIUS, 1781**  
Distribution: Steppe of S Russian Plain, Kazakhstan, Kirghizstan.
- Cryptocephalus (Asionus) flexuosus* KRYNICKI, 1834**  
Distribution: Steppe of Ukraine, Crimea, Asian Russia (SW Altai), Kazakhstan, Kirghizstan.
- Cryptocephalus (Asionus) floralis* KRYNICKI, 1834**  
*Cryptocephalus halophilus* SUFFRIAN, 1847  
Distribution: Crimea.

***Cryptocephalus (Asionus) fulmenifer* REITTER, 1889***Cryptocephalus undatus* SUFFRIAN, 1847*Cryptocephalus nigrinus* BERTI et RAPILLY, 1979

Distribution: Transcaucasia.

***Cryptocephalus (Asionus) gamma* HERRICH-SCHÄFFER, 1829***Cryptocephalus sesquistriatus* KRYNICKI, 1832*Cryptocephalus maculipes* ZOUBKOFF, 1833*Cryptocephalus ypsilon* SUFFRIAN, 1847*Cryptocephalus gamma* ab. *semilugens* DUDICH, 1924

Distribution: Steppe of S Russian Plain, Caucasus, Kazakhstan, Asian Russia (S Siberia: Novosibirsk Distr.).

***Cryptocephalus (Asionus) globulus* LOPATIN, 1953**

Distribution: Kazakhstan, Kirghizstan, Tajikistan, Uzbekistan.

***Cryptocephalus (Asionus) halophilus* GEBLER, 1830***Cryptocephalus scopini* LOPATIN, 1961

Distribution: Central Kazakhstan.

***Cryptocephalus (Asionus) hamatus* SUFFRIAN, 1854**

Distribution: Asian Russia (S Siberia, Amurland, Maritime Prov.), Mongolia.

***Cryptocephalus (Asionus) hammadae* LOPATIN, 1961**

Distribution: S Tajikistan.

***Cryptocephalus (Asionus) hauseri* J. WEISE, 1887**

Distribution: S Kazakhstan (Kendyk Tau Mountain Range).

***Cryptocephalus (Asionus) heydeni* J. WEISE, 1886**

Distribution: SE Kazakhstan, Mountains of Central Asia to S Tajikistan (Gazimailik Mountain Range).

***Cryptocephalus (Asionus) hirtipennis* FALDERMANN, 1835**

Distribution: Asian Russia (S Siberia, Yakutia, Russian Far East), Mongolia.

***Cryptocephalus (Asionus) impressipygus* OGLOBLIN, 1956**

Distribution: Steppe of SE Russian Plain, Kazakhstan from Uralsk to Zaysan Lake.

***Cryptocephalus (Asionus) irenae* LOPATIN, 1958**

Distribution: Tajikistan.

***Cryptocephalus (Asionus) iskanderi* LOPATIN, 1961**

Distribution: Central Tajikistan.

***Cryptocephalus (Asionus) jaxarticus* LOPATIN, 1986***Cryptocephalus simulator* LOPATIN, 1971, non GRESSITT, 1942

Distribution: W Kazakhstan.

***Cryptocephalus (Asionus) karakalensis* L. MEDVEDEV, 1955**

Distribution: S Turkmenistan (Kopetdagh Mountain Range), Uzbekistan, Tajikistan.

***Cryptocephalus (Asionus) karatavicus* LOPATIN, 1971**

Distribution: S Kazakhstan.

- Cryptocephalus (Asionus) kerzhneri* LOPATIN, 1968  
Distribution: S Kazakhstan.
- Cryptocephalus (Asionus) klarae* LOPATIN, 1990  
Distribution: SE Kazakhstan.
- Cryptocephalus (Asionus) koltzei* J. WEISE, 1887  
Distribution: Asian Russia (Amurland, Maritime Prov.), Mongolia, N China, Korea.
- Cryptocephalus (Asionus) korotjaevi korotjaevi* L. MEDVEDEV, 1980  
Distribution: Asian Russia (Tuva), Mongolia.
- Cryptocephalus (Asionus) korotjaevi shamoensis* L. MEDVEDEV, 1980  
Distribution: Mongolia.
- Cryptocephalus (Asionus) lateralis* SUFFRIAN, 1863  
Distribution: Steppe of SE Europe, Caucasus, Kazakhstan.
- Cryptocephalus (Asionus) lemniscatus* SUFFRIAN, 1854  
*Cryptocephalus discoideus* FAIRMAIRE, 1888  
*Cryptocephalus confucius* CLAVAREAU, 1914  
*Cryptocephalus hoffmanni* PIC. 1938  
Distribution: Asian Russia (Transbaikalia, Maritime Prov.), Mongolia, N China.
- Cryptocephalus (Asionus) medvedevi* LOPATIN, 1953  
*Cryptocephalus lopatini* IABLOKOFF-KHINZORIAN, 1983, BOB L. MEDVEDEV, 1978  
Distribution: S Turkmenistan (Kopetdagh Mountain Range), Tajikistan, NE Afghanistan.
- Cryptocephalus (Asionus) mniczechi* TAPPES, 1869  
Distribution: Asian Russia (Altai, Tuva, Sayan).
- Cryptocephalus (Asionus) ogloblini* LOPATIN, 1953  
Distribution: S Kazakhstan, Uzbekistan, Kirghizstan.
- Cryptocephalus (Asionus) oxianus* LOPATIN, 1975  
Distribution: S Tajikistan.
- Cryptocephalus (Asionus) pavlovskii* LOPATIN, 1956  
Distribution: S Kazakhstan (W Tian-Shan), Uzbekistan.
- Cryptocephalus (Asionus) personatus* J. WEISE, 1892  
*Cryptocephalus alaiensis* PIC. 1937  
Distribution: Kirghizstan (Kirghiz Alatau, Alai Mountain Range).
- Cryptocephalus (Asionus) prasolovi* LOPATIN, 1992  
Distribution: SE Kazakhstan.
- Cryptocephalus (Asionus) pseudolateralis* L. MEDVEDEV, 1979  
Distribution: E Kazakhstan.
- Cryptocephalus (Asionus) quatuordecimmaculatus* D. H. SCHNEIDER, 1792  
*Cryptocephalus coloratus* FABRICIUS, 1798  
*Cryptocephalus pilleri* SCHRANK, 1789  
*Cryptocephalus quatuordecimmaculatus* ab. *magyaranus* PIC. 1909

*Cryptocephalus quatuordecimmaculatus* ab. *minipennis* PIC, 1909  
*Cryptocephalus quatuordecimmaculatus* ab. *kaszabi* CSIKI, 1953  
*Cryptocephalus quatuordecimmaculatus* ab. *kuthyi* CSIKI, 1953  
*Cryptocephalus quatuordecimmaculatus* ab. *gyorffy* CSIKI, 1953  
*Cryptocephalus quatuordecimmaculatus* ab. *guranyii* CSIKI, 1953  
*Cryptocephalus quatuordecimmaculatus* ab. *stredai* CSIKI, 1953  
 Distribution: SE Europe, Turkey.

***Cryptocephalus (Asionus) richteri* L. MEDVEDEV, 1957**

Distribution: Transcaucasia.

***Cryptocephalus (Asionus) rubi rubi* MÉNÉTRIÉS, 1832**

Distribution: Transcaucasia.

***Cryptocephalus (Asionus) rubi glasunovi* JACOBSON, 1894**

Distribution: S Turkmenistan, S Kazakhstan.

***Cryptocephalus (Asionus) rubi properus* J. WEISE, 1900**

Distribution: SE Kazakhstan, Uzbekistan, S Tajikistan.

***Cryptocephalus (Asionus) sarafschanensis sarafschanensis* SOLSKY, 1882**

*Cryptocephalus placidus* J. WEISE, 1892

Distribution: Tajikistan and Kirghizstan (Turkestan, Zeravshan and N slope of Alai Mountain Ranges).

***Cryptocephalus (Asionus) sarafschanensis betpakdalensis* LOPATIN, 1976**

Distribution: Kazakhstan (Betpakdala desert).

***Cryptocephalus (Asionus) sarafschanensis duvivieri* J. WEISE, 1892**

*Cryptocephalus stenroosi* JACOBSON, 1901

Distribution: Kirghizstan (Kirghiz Alatau Mountain Range), Kazakhstan (Syr-Daryan' Karatau Mountain Range).

***Cryptocephalus (Asionus) sarafschanensis grumi* LOPATIN, 1976**

Distribution: E Kazakhstan, SW Mongolia.

***Cryptocephalus (Asionus) sarafschanensis iliensis* J. WEISE, 1900**

Distribution: Kazakhstan Folding Land (Cisralia and Cisbaikhashia).

***Cryptocephalus (Asionus) sarafschanensis narynensis* LOPATIN, 1976**

Distribution: Kirghizstan (Fergana Mountain Range).

***Cryptocephalus (Asionus) sarafschanensis sugatensis* LOPATIN, 1999**

Distribution: Kazakhstan (Sjughaty Mountains).

***Cryptocephalus (Asionus) sarafschanensis talassicus* LOPATIN, 1976**

Distribution: Uzbekistan and Kirghizstan (Talas and Karzhantau Mountain Ranges).

***Cryptocephalus (Asionus) sarafschanensis tiro* J. WEISE, 1900**

Distribution: Kirghizstan: Central Tian-Shan.

***Cryptocephalus (Asionus) sareptanus* MORAWITZ, 1863**

Distribution: SE of the Russian plain, E Ciscaucasia (Ergeni, Nogaiskaya Steppe), N, Central and E Kazakhstan).

***Cryptocephalus (Asionus) saryarkensis* L. MEDVEDEV et KULENOVA, 1980**

Distribution: Central Kazakhstan.

***Cryptocephalus (Asionus) spurius* LOPATIN, 1956**

Distribution: S Turkmenistan (Kopetdagh Mountain Range).

***Cryptocephalus (Asionus) stackelbergi* LOPATIN, 1971**

Distribution: Central and E Kazakhstan.

***Cryptocephalus (Asionus) stschukini* FALDERMANN, 1835**

Distribution: Asian Russia (Altai, Tuva, Sayan, Transbaikalia, Russian Far East), Mongolia.

***Cryptocephalus (Asionus) tamaricis* SOLSKY, 1867***Cryptocephalus astracanicus* SUFFRIAN, 1867

Distribution: SE of Russian Plain, E Transcaucasia, Kazakhstan, Turkmenistan, Uzbekistan, Tajikistan, Mongolia, NW China.

***Cryptocephalus (Asionus) terminassianae* LOPATIN, 1967**

Distribution: SE Kazakhstan.

***Cryptocephalus (Asionus) undulatus* SUFFRIAN, 1854**

Distribution: Kazakhstan, Turkmenistan, Kirghizstan, Uzbekistan, Tajikistan (except for the Pamirs).

***Cryptocephalus (Asionus) verae* L. MEDVEDEV, 1956**

Distribution: W and S Turkmenistan (Kopetdagh Mountain Range).

***Cryptocephalus (Asionus) volkovitschi* LOPATIN, 1976**

Distribution: S Transcaucasia.

***Cryptocephalus (Asionus) zaitzevi* LOPATIN, 1977**

Distribution: Transcaucasia.

***Cryptocephalus (Asionus) zinovievi* L. MEDVEDEV, 1973**

Distribution: Asian Russia (Amurland, Maritime Prov.), Mongolia.

**Subgenus: *Cryptocephalus (Heterichnus)* WARCHALOWSKI, 1991***Heterodactylus* L. MEDVEDEV, 1963, non SPIX, 1825***Cryptocephalus (Heterichnus) macrodactylus macrodactylus* GEBLER, 1830***Cryptocephalus scapularis* JACOBSON, 1894

Distribution: Asian Russia (S Altai), Kirghizstan, Kazakhstan.

***Cryptocephalus (Heterichnus) macrodactylus levipes* LOPATIN, 1999**

Distribution: Kirghizstan (Central Tian-Shan).

***Cryptocephalus (Heterichnus) macrodactylus persimilis* LOPATIN, 1956***Cryptocephalus (Heterichnus) persimilis* LOPATIN, 1956

Distribution: Kirghizstan (Kavak-Too and Susamyr Mountain Ranges).



***Cryptocephalus (Heterichnus) macrodactylus prosdocetus* JACOBSON, 1898***Cryptocephalus superbus* LOPATIN, 1953

Distribution: Kirghizstan (Fergan and Alai Mountain Ranges)

***Cryptocephalus (Heterichnus) tarsalis tarsalis* J. WEISE, 1887***Cryptocephalus aksuensis* LOPATIN, 1952

Distribution: Mountains of Tajikistan, Kirghizstan and Uzbekistan: from Badakhshan Mountains in the east, up to Karzhantau Mountain Range in the west.

***Cryptocephalus (Heterichnus) tarsalis submontanus* LOPATIN, 1999**

Distribution: S Tajikistan (Gazimailik Mountain Range).

**Subgenus: *Cryptocephalus (Cryptocephalus)* GEOFFROY, 1762*****Cryptocephalus (Cryptocephalus) alborzensis* RAPILLY, 1980**

Distribution: E Azerbaijan, N Iran.

***Cryptocephalus (Cryptocephalus) anticus* SUFFRIAN, 1848***Cryptocephalus nigerrimus* BALIION, 1878*Cryptocephalus sexpustulatus* RUSSL, 1790, non VILLERS, 1789*Cryptocephalus octoguttatus* D. H. SCHNEIDER, 1792, non LINNAEUS, 1767*Cryptocephalus sexpustulatus* ab. *omissus* J. WEISE, 1882*Cryptocephalus sexpustulatus* ab. *oncratus* J. WEISE, 1882*Cryptocephalus (Cryptocephalus) octacosmus* BEDEL, 1891*Cryptocephalus sexpustulatus* ab. *laeniger* PIC, 1913*Cryptocephalus sexpustulatus* ab. *pusztae* KASZAB, 1962*Cryptocephalus sexpustulatus* ab. *vastus* KASZAB, 1962

Distribution: Central and S Europe, Caucasus, Turkey, Kazakhstan, Uzbekistan, Kirghizstan, Asian Russia (southern parts of W Siberia).

***Cryptocephalus (Cryptocephalus) araxicola* IABLOKOFF-KHINZORIAN, 1968**

Distribution: S Armenia.

***Cryptocephalus (Cryptocephalus) aureolus aureolus* SUFFRIAN, 1847***Cryptocephalus aureolus* var. *coeruleus* SCHILSKY, 1888*Cryptocephalus aureolus* ab. *discolor* GIARDI, 1909*Cryptocephalus aureolus* ab. *subcoeruleicolor* PIC, 1912*Cryptocephalus aureolus* ab. *coeruleocinctus* DEPOLI et GORDANICH, 1926*Cryptocephalus aureolus* ab. *lydiae* DEPOLI, 1926*Cryptocephalus aureolus* ab. *aureicollis* DEPOLI, 1940*Cryptocephalus aureolus* ab. *purpureus* CSIKI, 1953

Distribution: Europe except for the extreme North.

***Cryptocephalus (Cryptocephalus) aureolus transcaucasicus* JACOBSON, 1898**

Distribution: Caucasus, Transcaucasia.

***Cryptocephalus (Cryptocephalus) bicolor* ESCHSCHOLTZ, 1818***Cryptocephalus conchinar* SUTCHINS, 1853

Distribution: S and SE Europe, S Ukraine, Caucasus, Turkey, N Iran.

***Cryptocephalus (Cryptocephalus) biguttatus* (SCOPOLI, 1763)***Baryscis biguttata* SCOPOLI, 1763*Cryptocephalus bimaculatus* FABRICE, 1775

*Cryptocephalus biguttatus* ab. *ornatus* ROUBAL, 1909  
*Cryptocephalus biguttatus* ab. *carniolicus* CLAVAREAU, 1913  
 Distribution: Europe, E Kazakhstan, Asian Russia (S Siberia).

***Cryptocephalus (Cryptocephalus) biguttulatus* GEBLER, 1841**

Distribution: Asian Russia (Altai, Tuva, Sayan, E Siberia), Mongolia.

***Cryptocephalus (Cryptocephalus) biguttulus* SUFFRIAN, 1848**

*Cryptocephalus rufilabris* SUFFRIAN, 1853  
 Distribution: S Crimea.

***Cryptocephalus (Cryptocephalus) bipunctatus bipunctatus* (LINNAEUS, 1758)**

*Chrysomela bipunctata* LINNAEUS, 1758  
*Buprestis sanguinolenta* SCOPOLI, 1763  
*Cryptocephalus lineola* FABRICIUS, 1792  
*Cryptocephalus dispar* PAYKULLI, 1798  
*Cryptocephalus quadrinotatus* HERRICH-SCHÄFFER, 1838  
*Cryptocephalus bipunctatus* var. *thomsoni* J. WEISE, 1882  
*Cryptocephalus bipunctatus* var. *clericus* SEIDLITZ, 1891  
*Cryptocephalus bipunctatus* var. *oblitteratipennis* PIC, 1900  
*Cryptocephalus bipunctatus* ab. *immaculatipennis* PIC, 1900  
*Cryptocephalus bipunctatus* ab. *subimmaculatus* PIC, 1900  
*Cryptocephalus bipunctatus* ab. *agnani* PIC, 1908  
*Cryptocephalus bipunctatus* ab. *nebulosonotatus* PIC, 1911  
*Cryptocephalus bipunctatus* ab. *bimaculatus* TRELLA, 1930  
*Cryptocephalus bipunctatus* ab. *bimacula* CSIKI, 1953  
 Distribution: Europe, Dagestan, Kazakhstan, Asian Russia (S Siberia).

***Cryptocephalus (Cryptocephalus) bipunctatus cautus* J. WEISE, 1893**

Distribution: Asian Russia (Transbaikalia, Amurland, Maritime Prov.), NE China, Korea.

***Cryptocephalus (Cryptocephalus) bodungeni bodungeni* JACOBSON, 1905**

Distribution: Kazakhstan (Syr-Darya River Valley), Karzhantau Mountain Range.

***Cryptocephalus (Cryptocephalus) bodungeni tadjhika* JACOBSON, 1925**

Distribution: N Tajikistan and Uzbekistan, (W of Fergana Valley).

***Cryptocephalus (Cryptocephalus) caerulescens* C. R. SAHLBERG, 1839**

*Cryptocephalus flavilabris* GYLLENHAL, 1813, non FABRICIUS, 1787  
*Cryptocephalus pelleti* MARSEUL, 1875  
*Cryptocephalus androgyne* MARSEUL, 1875  
*Cryptocephalus caerulescens* ab. *praehumeralis* PIC, 1908  
*Cryptocephalus caerulescens* ab. *larouzei* PIC, 1908  
*Cryptocephalus caerulescens* fa. *quadripunctata* POLENTZ, 1938  
*Cryptocephalus similis* L. MEDVEDEV, 1956  
*Cryptocephalus caerulescens* ab. *distractus* BURLINI, 1967  
*Cryptocephalus matsumurai* TAKIZAWA, 1971  
 Distribution: N and Central Europe, Asian Russia (southern parts of W Siberia, Amurland, Maritime Prov., Sakhalin).

***Cryptocephalus (Cryptocephalus) coeruleans* MARSEUL, 1875**

*Cryptocephalus coeruleans* SUFFRIAN, 1847  
*Cryptocephalus angustelimbatus* REITTER, 1912

*Cryptocephalus kraatzi* CHŪJŌ, 1935

Distribution: Asian Russia (Cisbaikalia, Amurland, Maritime Prov.), N Mongolia, NE China.

***Cryptocephalus (Cryptocephalus) concolor* SUFFRIAN, 1847**

*Cryptocephalus unicolor* FALDERMANN, 1838

*Cryptocephalus duplicatus persicus* RAPILLY, 1980

Distribution: SE Azerbaijan (Talysh), Turkmenistan (Kopetdagh Mountains), Iran.

***Cryptocephalus (Cryptocephalus) cordiger* (LINNAEUS, 1758)**

*Chrysomela cordigera* LINNAEUS, 1758

*Cryptocephalus cordiger* ab. *vlascanus* PIC, 1910

*Cryptocephalus cordiger* ab. *tihanyensis* SZEKESY, 1943

*Cryptocephalus cordiger* ab. *magistrettii* BURLINI, 1948

*Cryptocephalus cordiger* ab. *nigripes* G. MÜLLER, 1948

*Cryptocephalus cordiger* ab. *talaris* CSIKI, 1953

Distribution: Europe, Kazakhstan, Asian Russia (Tuva, Altais).

***Cryptocephalus (Cryptocephalus) coryli* (LINNAEUS, 1758)**

*Chrysomela coryli* LINNAEUS, 1758

*Cryptocephalus vitis* FABRICIUS, 1775

*Cryptocephalus temesiensis* SUFFRIAN, 1847

*Cryptocephalus coryli* ab. *temesiensis* SUFFRIAN, 1847

*Cryptocephalus coryli* var. *benoiti* PIC, 1900

*Cryptocephalus coryli* ab. *dorsosignatus* REITTER, 1912

*Cryptocephalus coryli* ab. *curtii* MADER, 1927

*Cryptocephalus coryli* ab. *brevecinctus* PIC, 1953

*Cryptocephalus coryli* ab. *weiseanus* PIC, 1953

*Cryptocephalus coryli* ab. *semiconnexus* PIC, 1953

*Cryptocephalus coryli* ab. *combinatus* KASZAB, 1962

*Cryptocephalus coryli* ab. *stilleri* KASZAB, 1962

*Cryptocephalus coryli* ab. *franzi* BURLINI, 1967

*Cryptocephalus coryli* ab. *rubroscutellatus* BURLINI, 1967

Distribution: All over Palaearctic except for the Central Asia, extreme South and North.

***Cryptocephalus (Cryptocephalus) cribratus* SUFFRIAN, 1847**

*Cryptocephalus trapezensis* TAPPES, 1871

Distribution: Caucasus, Turkey, W Turkmenistan (Kopetdagh Mountain Range), N Iran.

***Cryptocephalus (Cryptocephalus) cruciger* HELLÉN, 1922**

Distribution: N Europe.

***Cryptocephalus (Cryptocephalus) curtissimus* PIC, 1907**

Distribution: Sandy Deserts of Kazakhstan and Central Asia, delimited in the north by N beach of Aral Sea and Afghanistan in the south.

***Cryptocephalus (Cryptocephalus) decemmaculatus* (LINNAEUS, 1758)**

*Chrysomela decemmaculata* LINNAEUS, 1758

*Chrysomela bothnica* LINNAEUS, 1758

*Chrysomela barbareae* LINNAEUS, 1761

*Chrysomela decempunctata* LINNAEUS, 1761

*Cryptocephalus frenatus* LAICHARTING, 1781

*Cryptocephalus ornatus* HERBST, 1783

*Cryptocephalus flavescens* D. H. SCHNEIDER, 1792  
*Cryptocephalus betulae nanae* SCHILLING, 1838  
*Cryptocephalus decemmaculatus* var. *scenicus* J. WEISE, 1882  
*Cryptocephalus frenatus* var. *fabricii* J. WEISE, 1882  
*Cryptocephalus frenatus* var. *seminiger* J. WEISE, 1882  
*Cryptocephalus frenatus* ab. *nigricans* TRELLA, 1932  
 Distribution: Europe.

***Cryptocephalus (Cryptocephalus) distinguendus* D. H. SCHNEIDER, 1792**

*Cryptocephalus variegatus* GYLLENHAL, 1813  
*Cryptocephalus alni* MÉNÉTRIÉS, 1836  
*Cryptocephalus distinguendus* ab. *humeralis* SUFFRIAN, 1847  
*Cryptocephalus distinguendus* ab. *bihumeralis* PIC, 1908  
*Cryptocephalus distinguendus* ab. *subdistinguendus* PIC, 1908  
*Cryptocephalus distinguendus* ab. *octephilus* CLAVAREAU, 1913  
*Cryptocephalus distinguendus* ab. *pomeranicus* BURLINI, 1962  
 Distribution: N, Central, and western parts of East Europe, Asian Russia (Siberia, Magadan Distr.), E Kazakhstan.

***Cryptocephalus (Cryptocephalus) duplicatus* SUFFRIAN, 1847**

Distribution: N and W Caucasus, Transcaucasia, Turkey.

***Cryptocephalus (Cryptocephalus) elongatus* GERMAR, 1824**

Distribution: Central and SE Europe, N and Central Kazakhstan, Asian Russia (southern part of W Siberia, Altai, Sayan, S Yakutia).

***Cryptocephalus (Cryptocephalus) eous* LOPATIN, 1952**

Distribution: Asian Russia (Transbaikalia), Mongolia, NE China.

***Cryptocephalus (Cryptocephalus) flavipes* FABRICIUS, 1781**

*Cryptocephalus wydleri* FALDERMANN, 1837  
*Cryptocephalus flavipes* var. *nigrescens* GRADL, 1881  
*Cryptocephalus flavipes* var. *dispar* J. WEISE, 1882  
*Cryptocephalus flavipes* ab. *obscuripes* J. WEISE, 1882  
*Cryptocephalus flavipes* ab. *pumilionis* APFELBECK, 1912  
*Cryptocephalus flavipes* ab. *cordifrons* G. MÜLLER, 1953  
*Cryptocephalus flavipes* ab. *virago* BURLINI, 1967  
 Distribution: All over Palaearctic except for the Central Asia (present at Turkmenistan only), Mongolia, Korea, Japan.

***Cryptocephalus (Cryptocephalus) haroldi* KRAATZ, 1879**

Distribution: Asian Russia (Amurland, Maritime Prov.), NE China.

***Cryptocephalus (Cryptocephalus) hyacinthinus* SUFFRIAN, 1860**

*Cryptocephalus approximatus* BALY, 1873  
*Cryptocephalus fortunatus* BALY, 1873  
 Distribution: Asian Russia (Maritime Prov., S Kurile Is.), N China, Korea, Japan.

***Cryptocephalus (Cryptocephalus) hypochoeridis hypochoeridis* (LINNAEUS, 1758)**

*Chrysomela hypochoeridis* LINNAEUS, 1758  
*Cryptocephalus similis* STEPHENS, 1834  
*Cryptocephalus cristatus* DUFOUR, 1843  
*Cryptocephalus cristula* DUFOUR, 1843

*Cryptocephalus rugulipennis* SUFFRIAN, 1853

*Cryptocephalus cristula* ab. *auratus* GERHARDT, 1909

*Cryptocephalus cristula* ab. *bicolor* GERHARDT, 1909

*Cryptocephalus cristula* ab. *violaceus* GERHARDT, 1909

*Cryptocephalus cristula* ab. *iridescens* TRELLA, 1930

Distribution: Central and E Europe, Asian Russia (Siberia), N and E Kazakhstan.

***Cryptocephalus (Cryptocephalus) hypochoeridis praticola* J. WEISE, 1889**

*Cryptocephalus ruguliventris* FRANZ, 1949

Distribution: Caucasus (Abkhazia, Daghestan, Georgia).

***Cryptocephalus (Cryptocephalus) hypochoeridis transiens* FRANZ, 1949**

Distribution: Switzerland, Yugoslavia, Slovakia, Transcarpathia.

***Cryptocephalus (Cryptocephalus) imperialis* LAICHAITING, 1781**

*Cryptocephalus bistrispunctatus* GERMAR, 1824

*Cryptocephalus imperialis* ab. *posticeinstitutus* PIC, 1900

*Cryptocephalus imperialis* ab. *inexternus* PIC, 1910

Distribution: Europe, S European Russia, Central Georgia, Turkey, Cyprus.

***Cryptocephalus (Cryptocephalus) janthinus* GERMAR, 1824**

*Cryptocephalus megalostomus* FALDERMANN, 1837

*Cryptocephalus rusticus* FALDERMANN, 1837

*Cryptocephalus convexissimus* PIC, 1908

*Cryptocephalus weymarni* GRESSITT et KIMOTO, 1961

*Cryptocephalus janthinus* ab. *melanarius* BURLINI, 1967

Distribution: Southern parts of W and Central Europe, SE Europe, NW Caucasus, Asian Russia (S Siberia, Maritime Prov., Sakhalin).

***Cryptocephalus (Cryptocephalus) kiritschenkiellus* LOPATIN, 1961**

Distribution: Uzbekistan (Fergana Valley).

***Cryptocephalus (Cryptocephalus) krutovskii gebleri* JACOBSON, 1924**

*Cryptocephalus niger* GEBLER, 1830, non GEOFFROY, 1758

Distribution: E Kazakhstan, Asian Russia (Altai, Tuva, Cisbaikalia).

***Cryptocephalus (Cryptocephalus) krutovskii krutovskii* JACOBSON, 1900**

Distribution: Asian Russia (E Altai, Tuva, Cisbaikalia), Mongolia.

***Cryptocephalus (Cryptocephalus) krutovskii triangulifer* JACOBSON, 1926**

*Cryptocephalus karafutonis* KONO et TAMANUKI, 1929

Distribution: Asian Russia (Altai, Middle Siberia, S and Middle Yakutia, Amurland, Kamchatka, Maritime Prov., Sakhalin), Japan.

***Cryptocephalus (Cryptocephalus) kulibini* GEBLER, 1832**

*Cryptocephalus nasutulus* J. WEISE, 1889

Distribution: Asian Russia (Transbaikalia, Amurland, Maritime Prov.), Mongolia, NE China, Korea.

***Cryptocephalus (Cryptocephalus) laetus* FABRICIUS, 1792**

*Cryptocephalus mixtus* D. H. SCHNEIDER, 1792

*Cryptocephalus steveni* ADAMS, 1817

*Cryptocephalus laetus* ab. *miconotatus* PIC, 1908

*Cryptocephalus laetus* ab. *innotatus* CSIKI, 1953  
*Cryptocephalus laetus* ab. *ernoii* KASZAB, 1962  
*Cryptocephalus laetus* ab. *balatonicus* KASZAB, 1962  
*Cryptocephalus laetus* ab. *octosignatus* CSIKI: KASZAB, 1962  
*Cryptocephalus laetus* ab. *bifasciatus* CSIKI: KASZAB, 1962

Distribution: Europe except for the West and North, N Caucasus, Kazakhstan except for the South, Asian Russia (W Siberia, Altai).

***Cryptocephalus (Cryptocephalus) laevicollis* GEBLER, 1830**

*Cryptocephalus laevicollis* ab. *viennensis* J. WEISE, 1882  
*Cryptocephalus rossicus* J. WEISE, 1900  
*Cryptocephalus laevicollis* ab. *nigrofemoratus* BURLINI, 1962  
*Cryptocephalus laevicollis* ab. *pseudoviennensis* BURLINI, 1962

Distribution: E and SE Europe, Crimea, Ciscaucasia, Kazakhstan (except for the South), Asian Russia (southern parts of W Siberia, Altai).

***Cryptocephalus (Cryptocephalus) latimargo* L. MEDVEDEV, 1971**

Distribution: Asian Russia (Cisbaikalia, Transbaikalia, Magadan Distr.), Mongolia.

***Cryptocephalus (Cryptocephalus) limbatipennis* JACOBY, 1885**

*Cryptocephalus yanoi* CHŪJŌ, 1938  
*Cryptocephalus kononovi* L. MEDVEDEV, 1973

Distribution: Asian Russia (Maritime Prov.), E China, Japan.

***Cryptocephalus (Cryptocephalus) luridipennis luridipennis* SUFFRIAN, 1854**

*Cryptocephalus quindecimnotatus* SUFFRIAN, 1854  
*Cryptocephalus instabilis* BALY, 1873  
*Cryptocephalus pustulipes* KIMOTO, 1964

Distribution: Asian Russia (southern S part of W Siberia, Altai, Tuva, S Yakutia, Magadan Distr.), Mongolia, Japan.

***Cryptocephalus (Cryptocephalus) luridipennis pallescens* KRAATZ, 1879**

*Cryptocephalus serinicolor* PIC, 1907

Distribution: Asian Russia (Amurland, Maritime Prov.), Mongolia, WE China, Korea.

***Cryptocephalus (Cryptocephalus) mannerheimi* GEBLER, 1825**

Distribution: Asian Russia (Altai, Tuva, S Yakutia, Amurland, Maritime Prov.), Mongolia, N China, Korea.

***Cryptocephalus (Cryptocephalus) marginatus* FABRICIUS, 1781**

*Cryptocephalus terminatus* GERMAR, 1824  
*Cryptocephalus palliatus* SUFFRIAN, 1861  
*Cryptocephalus marginatus* ab. *crucifer* BURLINI, 1946  
*Cryptocephalus marginatus* ab. *discoidalis* BURLINI, 1946  
*Cryptocephalus marginatus* ab. *devaulxi* MÉQUIGNON, 1948  
*Cryptocephalus marginatus* ab. *pseudoterminalis* BURLINI, 1948  
*Cryptocephalus marginatus* ab. *sabaudus* MÉQUIGNON, 1948  
*Cryptocephalus marginatus* ab. *wellschmeidti* BURLINI, 1961

Distribution: Europe except for the North.

***Cryptocephalus (Cryptocephalus) melanoxanthus melanoxanthus* SOLSKY, 1876**

*Cryptocephalus grombczewskii* JACOBSON, 1895

Distribution: Kazakhstan, Kirghizstan, Tajikistan, Uzbekistan.

***Cryptocephalus (Cryptocephalus) melanoxanthus chotanensis* (JACOBSON, 1895)***Cryptocephalus chotanensis* JACOBSON, 1895

Distribution: Kirghizstan (Chon-Khemine River Valley), NW China.

***Cryptocephalus (Cryptocephalus) melanoxanthus flaviventris* LOPATIN, 1952**

Distribution: Kazakhstan (Lower Ili and Kharatal Rivers), Turkmenistan.

***Cryptocephalus (Cryptocephalus) melanoxanthus variolosus* (JACOBSON, 1894)***Cryptocephalus variolosus* JACOBSON, 1894

Distribution: Kirghizstan (Shores of Issyk-Kul Lake).

***Cryptocephalus (Cryptocephalus) modestus* SUFFRIAN, 1848**

Distribution: SE of European Russia, Kazakhstan, Asian Russia (Altai).

***Cryptocephalus (Cryptocephalus) monilis* J. WEISE, 1890**

Distribution: Kazakhstan, S Mongolia, N China.

***Cryptocephalus (Cryptocephalus) moraei* (LINNAEUS, 1758)***Chrysomela moraei* LINNAEUS, 1758*Cryptocephalus bivittatus* GYLLENHAL, 1813*Cryptocephalus interruptofasciatus* STURM, 1826 [nomen nudum]*Cryptocephalus interrupto-fasciatus* SUFFRIAN, 1847*Cryptocephalus moraei* ab. *vittiger* MARSEUL, 1875*Cryptocephalus moraei* ab. *arquatus* J. WEISE, 1882*Cryptocephalus moraei* ab. *bivittiger* PIC, 1909*Cryptocephalus moraei* ab. *winkleri* PIC, 1913*Cryptocephalus moraei* ab. *alleoni* PORTEVIN, 1934*Cryptocephalus moraei* ab. *marseuli* CSIKI, 1953*Cryptocephalus moraei* ab. *nigellus* BURLINI, 1955

Distribution: Europe, Caucasus, Turkey, N Iran, Urals, Asian Russia (W Siberia, Altai).

***Cryptocephalus (Cryptocephalus) multiplex liothorax* (SOLSKY, 1871)***Cryptocephalus parvicollis* JACOBSON, 1895*Cryptocephalus jacobsoni* CLAVAREAU, 1913*Cryptocephalus chujoe* NAKANE, 1963

Distribution: Asian Russia (southern part of E Siberia, Maritime Prov.), N China, N Korea.

***Cryptocephalus (Cryptocephalus) nitidulus* FABRICIUS, 1787***Cryptocephalus ochrostoma* HAROLD, 1872*Cryptocephalus nitidulus* ab. *splendidulus* CSIKI, 1953*Cryptocephalus nitidulus* ab. *stigmula* BURLINI, 1955*Cryptocephalus nitidulus* ab. *aurithorax* KASZAB, 1962*Cryptocephalus nitidulus* ab. *perfectecinctus* BURLINI, 1962*Cryptocephalus fortunatus borealis* L. MEDVEDEV, 1966

Distribution: N, Central, and E Europe, Urals, Asian Russia (S Siberia, Sakhalin, Kurile Is.), N China, Japan.

***Cryptocephalus (Cryptocephalus) nitidus* (LINNAEUS, 1758)***Chrysomela nitida* LINNAEUS, 1758*Chrysomela nitens* LINNAEUS, 1761*Cryptocephalus flavifrons* FABRICIUS, 1792

Distribution: Europe, Asian Russia (S Siberia).



***Cryptocephalus (Cryptocephalus) nobilis* KRAATZ, 1879**

Distribution: Asian Russia (Amurland), Japan.

***Cryptocephalus (Cryptocephalus) ochroloma* GEBLER, 1830**

Distribution: Asian Russia (Altai, Tuva, E Siberia, Amurland, Maritime Prov.), N Mongolia, N China.

***Cryptocephalus (Cryptocephalus) octomaculatus* ROSSI, 1790**

*Cryptocephalus quinguepunctatus* HARRER, 1784, non SCOPOLI, 1763

*Cryptocephalus 8-maculatus* ROSSI, 1790

*Cryptocephalus 12 punctatus* FABRICIUS, 1792

*Cryptocephalus? octomaculatus* ab. *testaceus* A. VILLA et J. B. VILLA, 1833

*Cryptocephalus? octomaculatus* ab. *stramineus* SUFFRIAN, 1847

*Cryptocephalus octomaculatus* ab. *derivatus* PIC, 1923

*Cryptocephalus octomaculatus* ab. *volviciensis* PIC, 1923

*Cryptocephalus octomaculatus* ab. *horvathianus* KASZAB, 1962

Distribution: S, Central and E Europe, Azerbaijan.

***Cryptocephalus (Cryptocephalus) octopunctatus* (SCOPOLI, 1763)**

*Buprestis octopunctata* SCOPOLI, 1763

*Cryptocephalus variabilis* D. H. SCHNEIDER, 1791

*Cryptocephalus octopunctatus* ab. *diversenotatus* PIC, 1907

*Cryptocephalus octopunctatus* ab. *colmariensis* PIC, 1918

*Cryptocephalus octopunctatus* ab. *multiinterruptus* PIC, 1918

*Cryptocephalus octopunctatus* ab. *sanguinarius* PIC, 1918

*Cryptocephalus octopunctatus* ab. *subhumeralis* PIC, 1918

*Cryptocephalus octopunctatus* ab. *viturati* PIC, 1918

*Cryptocephalus 8-punctatus* ab. *bipunctatus* TRELLA, 1930

*Cryptocephalus 8-punctatus* ab. *10-punctatus* TRELLA, 1930

*Cryptocephalus 8-punctatus* ab. *impunctatus* TRELLA, 1930

*Cryptocephalus 8-punctatus* ab. *postiunctus* TRELLA, 1930

*Cryptocephalus 8-punctatus* ab. *4-punctatus* TRELLA, 1930

*Cryptocephalus 8-punctatus* ab. *6-punctatus* TRELLA, 1930

*Cryptocephalus 8-punctatus* ab. *subinterruptus* PIC, 1939

*Cryptocephalus octopunctatus* ab. *praetermissus* BURLINI, 1948

*Cryptocephalus octopunctatus* ab. *antisejunctus* CSIKI, 1953

*Cryptocephalus octopunctatus* ab. *bijunctus* CSIKI, 1953

*Cryptocephalus octopunctatus* ab. *internotatus* CSIKI, 1953

*Cryptocephalus potaissiae* CSIKI, 1953

*Cryptocephalus octopunctatus* ab. *jaszoensis* KASZAB, 1962

*Cryptocephalus octopunctatus* ab. *scalaris* KASZAB, 1962

Distribution: Europe, Ciscaucasia, W and E Kazakhstan, Asian Russia (southern parts of W Siberia, Altai).

***Cryptocephalus (Cryptocephalus) orotschena* JACOBSON, 1925**

Distribution: Asian Russia (northern part of W Siberia, Yakutia, Magadan Distr.).

***Cryptocephalus (Cryptocephalus) oxysternus* JACOBSON, 1895**

Distribution: Asian Russia (Maritime Prov.), Mongolia, NE China.

***Cryptocephalus (Cryptocephalus) parvulus* O. F. MÜLLER, 1776**

*Cryptocephalus nigrocoeruleus* GOEZE, 1777

*Cryptocephalus flavilabris* FABRICIUS, 1787

*Cryptocephalus coeruleus* OLIVIER, 1791  
*Cryptocephalus fulcratus* GERMAR, 1824  
*Cryptocephalus obliquostratus* MOTSCHULSKY, 1866  
*Cryptocephalus permolestus* BALY, 1873  
*Cryptocephalus inurbanus* HAROLD, 1874  
*Cryptocephalus consolanus* BALY, 1874  
*Cryptocephalus parvulus* ab. *klettei* GERHARDT, 1909  
*Cryptocephalus parvulus* ab. *burlinii* TIBERGHEN, 1971

Distribution: All over Palaearctic except for the Central Asia.

***Cryptocephalus (Cryptocephalus) peliopterus* SOLSKY, 1871**

*Cryptocephalus bivittatus* GEBLER, 1830  
*Cryptocephalus japonus* BALY, 1873  
*Cryptocephalus raddei* KRAATZ, 1879

Distribution: Asian Russia (S Amurland and Maritime Prov.), NE China, Japan.

***Cryptocephalus (Cryptocephalus) prusias* SUFFRIAN, 1853**

*Cryptocephalus delagrangi* PIC, 1898

Distribution: SE Balkan Peninsula, Transcaucasia, Turkey.

***Cryptocephalus (Cryptocephalus) pseudocautus* L. MEDVEDEV, 1973**

Distribution: Asian Russia (Transbaikalia).

***Cryptocephalus (Cryptocephalus) punctiger* PAYKULL, 1799**

*Cryptocephalus punctiger* ab. *reticulaticollis* ROUBAL, 1929

Distribution: N and Central Europe.

***Cryptocephalus (Cryptocephalus) pustulipes* MÉNÉTRIÉS, 1836**

*Cryptocephalus transversalis* SUFFRIAN, 1853

Distribution: Asian Russia (S Siberia, Tuva, S Yakutia, Amurland, Maritime Prov.), Mongolia, NE China, Korea.

***Cryptocephalus (Cryptocephalus) putjatae* JACOBSON, 1895**

Distribution: Asian Russia (S Transbaikalia, Maritime Prov.), Mongolia, NE China.

***Cryptocephalus (Cryptocephalus) quadriguttatus* RICHTER, 1820**

*Cryptocephalus nigribuccis* GEBLER, 1830  
*Cryptocephalus quadriguttatus* ab. *maurus* SUFFRIAN, 1848  
*Cryptocephalus quadriguttatus* ab. *nuntius* J. WEISE, 1898  
*Cryptocephalus quadriguttatus* ab. *latemaculatus* CSIKI, 1953

Distribution: Central and E Europe (North to Moscow-Smolensk), Ciscaucasia, N and E Kazakhstan, Asian Russia (southern part of W Siberia, Altai, Tuva), Mongolia.

***Cryptocephalus (Cryptocephalus) quadripustulatus* GYLLENHAL, 1813**

*Cryptocephalus 4-pustulatus* GYLLENHAL, 1813  
*Cryptocephalus quadripustulatus* ab. *similis* SUFFRIAN, 1853  
*Cryptocephalus rhaeticus* STIERLIN, 1863  
*Cryptocephalus quadripustulatus* ab. *aethiops* J. WEISE, 1882  
*Cryptocephalus quadripustulatus* ab. *bosnicus* APFELBECK, 1912  
*Cryptocephalus quadripustulatus* ab. *trismarginatus* BURLINI, 1967

Distribution: Europe, W and Middle Siberia.

***Cryptocephalus (Cryptocephalus) quinquepunctatus* (SCOPOLI, 1763)**

- Buprestis quinquepunctata* SCOPOLI, 1763  
*Cryptocephalus signatus* LAICHARTING, 1781  
*Cryptocephalus interruptus* SUFFRIAN, 1847  
*Cryptocephalus? quinquepunctatus* ab. *rubellus* J. WEISE, 1881  
*Cryptocephalus? quinquepunctatus* ab. *ravouxi* PIC, 1911  
*Cryptocephalus? quinquepunctatus* ab. *erratus* PIC, 1919  
*Cryptocephalus? quinquepunctatus* ab. *tyrolensis* PIC, 1919  
*Cryptocephalus signatus* ab. *bifasciatus* TRELLA, 1930  
*Cryptocephalus signatus* ab. *unifasciatus* TRELLA, 1930  
*Cryptocephalus? quinquepunctatus* ab. *vallorbensis* PIC, 1939  
*Cryptocephalus? quinquepunctatus* ab. *anticecruciatu*s CSIKI, 1953  
*Cryptocephalus? quinquepunctatus* ab. *bicharicus* KASZAB, 1962  
 Distribution: Central Europe, Carpathians, Transcarpathia, Moldavia, Transcaucasia, S European Russia, S Urals.

***Cryptocephalus (Cryptocephalus) regalis* GEBLER, 1830**

- Cryptocephalus pilosus* BALY, 1873  
*Cryptocephalus mutatus* KRAATZ, 1879  
*Cryptocephalus dauricus* J. WEISE, 1882  
*Cryptocephalus cyanescens* J. WEISE, 1887  
*Cryptocephalus sutschanus* MACHATSCHKE, 1959  
 Distribution: Asian Russia (S Siberia, Altai, Tuva, Amurland, Maritime Prov.), Mongolia, N China, Korea, Japan.

***Cryptocephalus (Cryptocephalus) rufofasciatus* SOLSKY, 1882**

- Cryptocephalus plustschevskiyi* JACOBSON, 1894  
*Cryptocephalus transversalis* REITTER, 1895  
*Cryptocephalus meyeri* REITTER, 1897  
*Cryptocephalus rickmersi* J. WEISE, 1900  
 Distribution: Sandy Deserts of Kazakhstan and Central Asia from Ryn-Peski in the north, up to S Uzbekistan in the south.

***Cryptocephalus (Cryptocephalus) ruralis* J. WEISE, 1887**

- Distribution: Asian Russia (S Cisbaikalia, Yakutia, Amurland, Magadan Distr.).

***Cryptocephalus (Cryptocephalus) semiargenteus* REITTER, 1894**

- Cryptocephalus semiargenteus mujuncumensis* LEBEDEV, 1931  
 Distribution: Deserts of S Central Asia and Kazakhstan, from lower Syr-Darya River, up to S Uzbekistan and Tajikistan.

***Cryptocephalus (Cryptocephalus) sericeus* (LINNAEUS, 1758)**

- Chrysomela sericea* LINNAEUS, 1758  
*Cryptocephalus sericeus* fa. *pratorum* SUFFRIAN, 1847  
*Cryptocephalus robustus* SUFFRIAN, 1853  
*Cryptocephalus bidens* C. G. THOMSON, 1868  
*Cryptocephalus zambanellus* MARSEUL, 1875  
*Cryptocephalus sericeus* var. *intrusus* J. WEISE, 1881  
*Cryptocephalus sericeus* ab. *coeruleus* J. WEISE, 1881  
*Cryptocephalus sericeus* ab. *violaceus* J. WEISE, 1881  
*Cryptocephalus sericeus* ab. *purpurascens* REITTER, 1912  
*Cryptocephalus sericeus* ab. *parellinus* CLAVAREAU, 1913

*Cryptocephalus sericeus* ab. *purpureicolor* PIC, 1914  
*Cryptocephalus sericeus* ab. *semiviridescens* PIC, 1914  
*Cryptocephalus sericeus* ab. *bicolor* DEPOLI, 1926  
*Cryptocephalus sericeus* ab. *viridiaureus* DEPOLI, 1926  
*Cryptocephalus sericeus* fa. *pratonus* PORTEVIN, 1934  
*Cryptocephalus sericeus* ab. *bicolorata* KASZAB, 1962  
*Cryptocephalus sericeus* ab. *funebri* BURLINI, 1962  
*Cryptocephalus sericeus* ab. *mirandus* BURLINI, 1967

Distribution: Europe except for the North and Mediterranean, Caucasus, Transcaucasia, Turkey, N Kazakhstan, Asian Russia (Siberia east to Yakutia).

***Cryptocephalus (Cryptocephalus) sexpunctatus* (LINNAEUS, 1758)**

*Chrysomela 6-punctata* LINNAEUS, 1758  
*Cryptocephalus sexpunctatus* ab. *pictus* SUFFRIAN, 1847  
*Cryptocephalus sexpunctatus* ab. *separandus* SUFFRIAN, 1847  
*Cryptocephalus sexpunctatus* ab. *gyllenhali* J. WEISE, 1881  
*Cryptocephalus sexpunctatus* ab. *thoracicus* J. WEISE, 1881  
*Cryptocephalus sexpunctatus* ab. *simplarius* HEYDEN, 1884  
*Cryptocephalus sexpunctatus* ab. *subnigrofasciatus* PIC, 1918  
*Cryptocephalus sexpunctatus* ab. *trijunctus* TRELLA, 1930  
*Cryptocephalus sexpunctatus* ab. *fortinsi* PIC, 1939  
*Cryptocephalus sexpunctatus* ab. *pedemontanus* PIC, 1939

Distribution: All over the Palaearctic except for the Central Asia.

***Cryptocephalus (Cryptocephalus) signatifrons* SUFFRIAN, 1847**

*Cryptocephalus fulvipes* var. *signatifrons* SUFFRIAN, 1847

Distribution: Europe except for the North, SE Poland, SW Ukraine (Carpathians).

***Cryptocephalus (Cryptocephalus) splendens* KRAATZ, 1879**

*Cryptocephalus zubovskii* JACOBSON, 1894

Distribution: Asian Russia (Altai, Tuva, Sayan, Cisbaikalia, Dauria, S Yakutia, Amurland, Maritime Prov., Sakhalin), Mongolia, NE China.

***Cryptocephalus (Cryptocephalus) surdus* RAPILLY, 1980**

Distribution: Transcaucasia, Turkey.

***Cryptocephalus (Cryptocephalus) tataricus* GEBLER, 1841**

*Cryptocephalus hectastigmus* SOLSKY, 1882

*Cryptocephalus rufus* KRAATZ, 1882

*Cryptocephalus dorsiger* BREIT, 1918

*Cryptocephalus montivagus* LOPATIN, 1954

Distribution: S and E Kazakhstan, Asian Russia (S Cisbaikalia, Tuva), Central Asia, SW Mongolia.

***Cryptocephalus (Cryptocephalus) tetradecaspilotus* BALY, 1873**

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Japan.

***Cryptocephalus (Cryptocephalus) tetratyrus* SOLSKY, 1871**

*Cryptocephalus amurensis* HEYDEN, 1884

Distribution: Asian Russia (S Amurland, Maritime Prov.), Korea.

***Cryptocephalus (Cryptocephalus) tricoloratus* JACOBSON, 1895**

Distribution: Asian Russia (S Maritime Prov.), SW China.

***Cryptocephalus (Cryptocephalus) trimaculatus* ROSSI, 1790***Cryptocephalus salicis* FABRICIUS, 1792*Cryptocephalus sexmaculatus* OLIVIER, 1808*Cryptocephalus trimaculatus* ab. *posticefasciatus* PIC, 1900

Distribution: S and SE Europe, Caucasus, Turkey.

***Cryptocephalus (Cryptocephalus) turangae* LOPATIN, 1961**

Distribution: S Tajikistan (Vahsh and Kyzyl-Su River Valleys).

***Cryptocephalus (Cryptocephalus) umbrosus* LOPATIN, 1956***Cryptocephalus (Cryptocephalus) semiargenteus umbrosus* LOPATIN, 1956

Distribution: Iran.

***Cryptocephalus (Cryptocephalus) violaceus* LAICHARTING, 1781***Cryptocephalus violaceus* ab. *smaragdinus* SUFFRIAN, 1847*Cryptocephalus violaceus* ab. *viridicollis* HÄNEL, 1934*Cryptocephalus violaceus* ab. *morelianus* PIC, 1953*Cryptocephalus violaceus* ab. *sepulchralis* CSIKI, 1953

Distribution: Central and S Europe, Ciscaucasia.

***Cryptocephalus (Cryptocephalus) virens* SUFFRIAN, 1847**

Distribution: Central and SE Europe, Steppe of Kazakhstan, Asian Russia (S Siberia).

***Cryptocephalus (Cryptocephalus) vittatus* FABRICIUS, 1775***Cryptocephalus vittatus* var. *negligens* J. WEISE, 1882*Cryptocephalus vittatus* ab. *optatus* J. WEISE, 1898*Cryptocephalus vittatus* ab. *schrammi* PIC, 1907*Cryptocephalus vittatus* ab. *lineellus* GERHARDT, 1909*Cryptocephalus vittatus* ab. *interruptus* EVERTS, 1921*Cryptocephalus vittatus* ab. *reynosensis* PIC, 1923*Cryptocephalus vittatus* ab. *patens* CSIKI, 1953

Distribution: W, SW and Central Europe.

***Cryptocephalus (Cryptocephalus) zejensis* MIKHAILOV, 1999**

Distribution: Asian Russia (Far East).

**Subgenus: *Cryptocephalus (Burlinius)* LOPATIN, 1965*****Cryptocephalus (Burlinius) appositus* LOPATIN, 1958**

Distribution: S Kirghizstan (Fergan Mountain Range).

***Cryptocephalus (Burlinius) bilineatus* (LINNAEUS, 1767)***Chrysomela bilineata* LINNAEUS, 1767*Pachybrachis armeniacus* FALDERMANN, 1837*Cryptocephalus spitzyi* SUFFRIAN, 1848*Cryptocephalus bilineatus* var. *moestus* J. WEISE, 1882*Cryptocephalus bilineatus* var. *bisbilineatus* PIC, 1904*Cryptocephalus bilineatus* ab. *ehmanni* CSIKI, 1953*Cryptocephalus bilineatus* ab. *gesztelensis* CSIKI, 1953*Cryptocephalus bilineatus* ab. *ujhelyi* CSIKI, 1953*Cryptocephalus bilineatus* ab. *pseudoarmeniacus* BURLINI, 1955*Cryptocephalus bilineatus* ab. *bakonyensis* KASZAB, 1962

Distribution: All over the Palaearctic except for the extreme North and Central Asia.

***Cryptocephalus (Burlinius) carpathicus* FRIVALDSZKY, 1883**

Distribution: Ukrainian Carpathians, Hungary.

***Cryptocephalus (Burlinius) chrysopus* GMELIN, 1787***Chrysomela biguttata* SCHALLER, 1763, non SCOPOLI, 1763*Cryptocephalus hübneri* FABRICIUS, 1792

Distribution: Europe, N Caucasus, Asian Russia (Sayan).

***Cryptocephalus (Burlinius) confusus* SUFFRIAN, 1854***Cryptocephalus discretus* BALY, 1873

Distribution: Asian Russia (S Siberia, Tuva, Yakutia, Russian Far East), Mongolia.

***Cryptocephalus (Burlinius) connexus* OLIVIER, 1807***Cryptocephalus tschimganensis ahngeri* JACOBSON, 1901*Cryptocephalus connexus* ab. *arenarius* J. WEISE, 1882*Cryptocephalus connexus* ab. *subconnexus* J. WEISE, 1882*Cryptocephalus connexus* ab. *fasciatus* PETRI, 1912*Cryptocephalus connexus* ab. *bisbireductellus* KASZAB, 1962*Cryptocephalus connexus* ab. *heterepistaticus* KASZAB, 1962*Cryptocephalus connexus* ab. *lichtmeckerti* KASZAB, 1962*Cryptocephalus connexus* ab. *redutevittatus* KASZAB, 1962*Cryptocephalus connexus* ab. *sajoi* KASZAB, 1962

Distribution: Mediterranean, E Europe, Caucasus, Kazakhstan, Central Asia (Steppe).

***Cryptocephalus (Burlinius) dilutellus dilutellus* JACOBSON, 1901**

Distribution: Kazakhstan, Kirghizstan, N and Central Tajikistan.

***Cryptocephalus (Burlinius) dilutellus badakshanicus* LOPATIN, 1999**

Distribution: E Tajikistan (Badakhshan Mountains).

***Cryptocephalus (Burlinius) elegantulus* GRAVENHORST, 1807***Cryptocephalus tessellatus* GERMAR, 1813*Cryptocephalus jucundus* FALDERMANN, 1837*Cryptocephalus elegantulus* ab. *inadumbratus* PIC, 1904*Cryptocephalus elegantulus* ab. *horvathi* CSIKI, 1953*Cryptocephalus elegantulus* ab. *deubeli* KASZAB, 1962*Cryptocephalus elegantulus* ab. *negrei* BURLINI, 1967

Distribution: Central and SE Europe, Asian Russia (W Siberia), Turkey.

***Cryptocephalus (Burlinius) exiguus* D. H. SCHNEIDER, 1792***Cryptocephalus wasastjernii* GYLLENHAL, 1827*Cryptocephalus amiculis* BALY, 1873*Cryptocephalus exiguus* ab. *adocetus* JACOBSON, 1901

Distribution: Europe. Siberia.

***Cryptocephalus (Burlinius) fausti* J. WEISE, 1882**

Distribution: E Caucasus (Daghestan, Azerbaijan).

***Cryptocephalus (Burlinius) fedtschenkoi* JACOBSON, 1901***Cryptocephalus dilutellus* ab. *fedtschenkoi* JACOBSON, 1901

Distribution: Kazakhstan, Tajikistan, Kirghizstan.

***Cryptocephalus (Burlinius) flavoscutellaris* L. MEDVEDEV, 1973**

Distribution: Asian Russia (Amurland), Korea.

***Cryptocephalus (Burlinius) frontalis* MARSHAM, 1802***Cryptocephalus marginellus* GRAVENHORST, 1807, non OLIVIER, 1791

Distribution: Europe, Asian Russia (Siberia, Maritime Prov., Sakhalin).

***Cryptocephalus (Burlinius) fulvus* (GOEZE, 1777)***Chrysomela fulva* GOEZE, 1777*Cryptocephalus minutus* FABRICIUS, 1792*Cryptocephalus ochraceus* STEPHENS, 1829*Cryptocephalus signaticollis* SUFFRIAN, 1848*Cryptocephalus minutus* fa. *fulvicollis* SUFFRIAN, 1848*Cryptocephalus fulvus* ab. *succinctus* J. WEISE, 1882*Cryptocephalus fulvus* ab. *vittatus* GOZIS, 1907*Cryptocephalus fulvus* ab. *gozisi* PIC, 1908*Cryptocephalus fulvus* ab. *weiseanus* BREIT, 1918*Cryptocephalus fulvus* ab. *obscuricollis* GABRIEL in W. KOLBE, 1927*Cryptocephalus fuscolineatus* CHŪJŌ, 1940*Cryptocephalus fulvus schatzmayri* BURLINI, 1969

Distribution: Europe, Transcaucasia, Near East, Asian Russia (Maritime Prov.), W and Central Kazakhstan, N China, Korea.

***Cryptocephalus (Burlinius) gussakovskii* LOPATIN, 1952**

Distribution: Asian Russia (Maritime Prov.).

***Cryptocephalus (Burlinius) invisus* LOPATIN, 1963***Cryptocephalus polymorphus* ab. *dilecta* LOPATIN, 1956

Distribution: Kirghizstan, Tajikistan.

***Cryptocephalus (Burlinius) kabaki* LOPATIN, 2002**

Distribution: N China (Tzinzian).

***Cryptocephalus (Burlinius) kurentzovi* L. MEDVEDEV, 1966**

Distribution: Asian Russia (S Kurile Is.), Japan.

***Cryptocephalus (Burlinius) labiatus* (LINNAEUS, 1761)***Chrysomela labiata* LINNAEUS, 1761*Cryptocephalus exilis* STEPHENS, 1832*Cryptocephalus digrammus* SUFFRIAN, 1848*Cryptocephalus labiatus* var. *ocularis* HEYDEN, 1863

Distribution: Europe, Asian Russia (W Siberia), Mongolia.

***Cryptocephalus (Burlinius) lederi* J. WEISE, 1889***Cryptocephalus oomorphus* JACOBSON, 1895

Distribution: Transcaucasia, Iran, Afghanistan, Turkmenistan (Kopetdagh Mountains).

***Cryptocephalus (Burlinius) macellus* SUFFRIAN, 1860***Cryptocephalus ochroleucus* SUFFRIAN, 1868*Cryptocephalus macellus* ab. *pseudoconnexus* G. MÜLLER, 1948*Cryptocephalus macellus iranicus* L. MEDVEDEV, 1975

Distribution: S and Central Europe, Transcaucasia, Near and Middle East, N Africa.

***Cryptocephalus (Burlinius) nigrofasciatus* JACOBY, 1885***Cryptocephalus turpis* CHEN, 1942*Cryptocephalus kimotoi* NAKANE, 1963

Distribution: Asian Russia (Amurland, Maritime Prov., Sakhalin), N China, Mongolia, Japan.

***Cryptocephalus (Burlinius) ocellatus* DRAPIEZ, 1819**

*Cryptocephalus geminus* GYLLENHAL, 1827

*Cryptocephalus ochropezus* SUFFRIAN, 1853

*Cryptocephalus ocellatus* var. *nigrifrons* BEDEL, 1899

*Cryptocephalus corsicus* PIC, 1909

*Cryptocephalus hummleri* PIC, 1909

*Cryptocephalus ocellatus* ab. *montanellus* G. MÜLLER, 1948

Distribution: Europe except for the North, Caucasus, Transcaucasia, Kazakhstan, Asian Russia (S Siberia).

***Cryptocephalus (Burlinius) ochroleucus* STEPHENS, 1839**

*Cryptocephalus fallax* SUFFRIAN, 1868

*Cryptocephalus ochroleucus* ab. *laghouatensis* PIC, 1903

*Cryptocephalus ochroleucus orientalis* RAPILLY, 1980

Distribution: Europe, Steppe zone of Ukraine, Transcaucasia, Middle East, NW Africa.

***Cryptocephalus (Burlinius) pallifrons* GYLLENHAL, 1813**

*Cryptocephalus insignis* PAYKULL, 1799 [nomen nudum]

*Cryptocephalus furcifrons* MANNERHEIM, 1844

*Cryptocephalus irkutensis* CLAVAREAU, 1913

*Cryptocephalus pallidifrons* CLAVAREAU, 1913

Distribution: Europe except for the South, Urals, Kazakhstan (except for the South), Asian Russia (W Siberia, Sayan, Yakutia, Maritime Prov.).

***Cryptocephalus (Burlinius) planifrons* J. WEISE, 1882**

Distribution: Southern part of Central and East Europe, Balkans, Caucasus, Kazakhstan, Asian Russia (southern part of W Siberia).

***Cryptocephalus (Burlinius) polymorphus polymorphus* SOLSKY, 1882**

*Cryptocephalus laevifrons* J. WEISE, 1894

Distribution: Mountain Areas of Central Asia and S Kazakhstan, north to Transili Alatau, Afghanistan.

***Cryptocephalus (Burlinius) polymorphus parallelus* JACOBSON, 1895**

*Cryptocephalus norensis* PIC, 1911

Distribution: Tajikistan (Vakhsh and Khozratishoh Mountain Ranges).

***Cryptocephalus (Burlinius) polymorphus schmidti* JACOBSON, 1894**

Distribution: Kirghizstan (Issyk-Kul Depression).

***Cryptocephalus (Burlinius) populi* SUFFRIAN, 1848**

*Cryptocephalus brachialis* MULSANT et REY, 1859

Distribution: W, Central and southern E Europe, Caucasus, Transcaucasia, E Kazakhstan.

***Cryptocephalus (Burlinius) pusillus* FABRICIUS, 1777**

*Cryptocephalus minutus* HERBST, 1783

*Cryptocephalus gracilis* STEPHENS, 1834

*Cryptocephalus verticalis* BOHEMAN, 1851

*Cryptocephalus raphaelensis* GAUTIER DE COTTES, 1861

*Cryptocephalus pusillus* var. *immaculatus* WESTHOFF, 1882



- Cryptocephalus pusillus* var. *marshami* J. WEISE, 1882  
*Cryptocephalus pusillus* var. *viduus* J. WEISE, 1889  
*Cryptocephalus pusillus* ab. *flavissimus* ROUBAL, 1915  
*Cryptocephalus pusillus* ab. *ernestii* PIC, 1926  
*Cryptocephalus pusillus* ab. *planchezi* PIC, 1926  
*Cryptocephalus pusillus* ab. *baudi* PIC, 1945  
*Cryptocephalus pusillus* ab. *multinigronotatus* PIC, 1945

Distribution: Europe except for the North, Caucasus, Turkey, Algeria.

***Cryptocephalus (Burlinius) pygmaeus* FABRICIUS, 1792**

- Cryptocephalus amaenus* DRAPIEZ, 1819  
*Cryptocephalus pygmaeus* var. *orientalis* J. WEISE, 1882  
*Cryptocephalus pygmaeus* ab. *lugubris* DEMAISON, 1904  
*Cryptocephalus pygmaeus mediterraneus* G. MÜLLER, 1948  
*Cryptocephalus pygmaeus picticollis* G. MÜLLER, 1948  
*Cryptocephalus pygmaeus atlanticus* BURLINI, 1970  
*Cryptocephalus gracilis* FABRICIUS, 1792  
*Cryptocephalus rufipes* ab. *hispanus* ZEIDLITZ, 1891  
*Cryptocephalus rufipes* ab. *praescutellaris* PIC, 1913  
*Cryptocephalus rufipes* ab. *dormeyeri* WAGNER, 1919  
*Cryptocephalus rufipes* ab. *bimaculatus* ČEPELAK, 1927

Distribution: W, SW, Central and southerwest parts of E Europe, N Africa.

***Cryptocephalus (Burlinius) saliceti* G. ZEBE, 1855**

- Cryptocephalus saliceti* ab. *resinaceus* PIC, 1903  
*Cryptocephalus saliceti* ab. *schaefferi* BURLINI, 1967

Distribution: N Europe, northern parts and Mountains of Central and East Europe.

***Cryptocephalus (Burlinius) shabalinae* LOPATIN, 1968**

Distribution: Kirghizstan.

***Cryptocephalus (Burlinius) tschimganensis tschimganensis* J. WEISE, 1894**

Distribution: SW Kazakhstan, Uzbekistan, Kirghizstan, S Tajikistan.

***Cryptocephalus (Burlinius) tschimganensis narzykulovi* LOPATIN, 1958**

- Cryptocephalus narzykulovi* LOPATIN, 1958  
*Cryptocephalus puzanovi* LOPATIN, 1958

Distribution: Tajikistan.

***Cryptocephalus (Burlinius) variceps* J. WEISE, 1884**

Distribution: SE Balkans, Transcaucasia, Turkey, Iran.

***Cryptocephalus (Burlinius) vittula* SUFFRIAN, 1848**

Distribution: Central, S and SE Europe, Caucasus.

***Cryptocephalus (Burlinius) xanthus* IABLOKOFF-KHNZORIAN, 1968**

- Cryptocephalus rastegari* RAPILLY, 1980

Distribution: Armenia, Azerbaijan, Iran.

**Genus: *Melixanthus* SUFFRIAN, 1854**

***Melixanthus granularis* (SUFFRIAN, 1857)**

- Cryptocephalus granularis* SUFFRIAN, 1857

*Cryptocephalus brunnicollis* SUFFRIAN, 1857

*Melixanthus jordanicus* LOPATIN, 1979

Distribution: S Transcaucasia, Turkmenistan (Kopetdagh Mountains), Jordan, Arabia.

***Melixanthus pumilio* (SUFFRIAN, 1854)**

*Cryptocephalus pumilio* SUFFRIAN, 1854

*Melixanthus babai* CHŪJŌ, 1959

Distribution: Asian Russia (Cisbaikalia, Buryatia, Sakhalin, Kurile Is.), Mongolia, Tibet, Japan.

**Genus: *Jaxartiolus* JACOBSON, 1922**

***Jaxartiolus baeckmannianus* JACOBSON, 1922**

Distribution: S Kazakhstan, Uzbekistan, Tajikistan (Rivers Amu-Darya and Vakhsh).

***Jaxartiolus reductosignatus* LOPATIN, 1963**

Distribution: Kirghizstan (Central Tian-Shan).

Tribe: *Pachybrachini* CHAPUIS, 1874

**Genus: *Pachybrachis* CHEVROLAT in DEJEAN, 1837**

*Pachybrachys* MANNERHEIM, 1843

***Pachybrachis abditus* LOPATIN, 1991**

Distribution: Kirghizstan (N Coast of Issyk-Kul Lake).

***Pachybrachis absinthii* LOPATIN, 1990**

Distribution: SE Kazakhstan.

***Pachybrachis albicans* (J. WEISE, 1882)**

*Pachybrachys albicans* J. WEISE, 1882

Distribution: Transcaucasia, Turkey.

***Pachybrachis altimontanus* (LOPATIN, 1968)**

*Pachybrachys altimontanus* LOPATIN, 1968

Distribution: Kirghizstan (Fergan and Alai Mountain Ranges).

***Pachybrachis amurensis* L. MEDVEDEV, 1973**

*Pachybrachis mohri* LOPATIN, 1974

Distribution: Asian Russia (Tuva, Amurland, Maritime Prov.), Mongolia.

***Pachybrachis arnoldii* (LOPATIN, 1967)**

*Pachybrachys arnoldii* LOPATIN, 1967

*Pachybrachis parvissimus* LOPATIN, 1975

Distribution: N and NW Kazakhstan, Mongolia.

***Pachybrachis atraphaxidis* (LOPATIN, 1968)**

*Pachybrachys atraphaxidis* LOPATIN, 1968

Distribution: S and SE Kazakhstan.

***Pachybrachis auliensis* (BREIT, 1921)***Pachybrachys auliensis* BREIT, 1921

Distribution: S and E Kazakhstan.

***Pachybrachis caprea caprea* (J. WEISE, 1887)***Pachybrachys caprea caprea* J. WEISE, 1887

Distribution: Kazakhstan and Central Asia (except for Turkmenistan).

***Pachybrachis caprea dzhungaricus* L. MEDVEDEV et RYBAKOVA, 1980**

Distribution: Mongolia.

***Pachybrachis caraganae* LOPATIN, 1977**

Distribution: Mongolia (Mongol Mountain Range and Ghobi Altai).

***Pachybrachis carpathicus* (REY, 1883)***Pachybrachys carpathicus* REY, 1883

Distribution: Ukrainian Transcarpathia, Romania.

***Pachybrachis cribricollis* (PIC, 1907)***Pachybrachys cribricollis* PIC, 1907

Distribution: SE of European Russia, Kazakhstan, Turkmenistan, Uzbekistan, Kirghizstan.

***Pachybrachis curtipennis* (BREIT, 1921)***Pachybrachys curtipennis* BREIT, 1921

Distribution: Uzbekistan.

***Pachybrachis distictopygus* (JACOBSON, 1901)***Pachybrachys distictopygus* JACOBSON, 1901*Pachybrachys kaszabi* LOPATIN, 1966

Distribution: Asian Russia (S Siberia, Maritime Prov.), Mongolia, NE China.

***Pachybrachis fimbriolatus* (SUFFRIAN, 1848)***Cryptocephalus tristis* OLIVIER, 1808, non LAICHARTING, 1781*Pachybrachys fimbriolatus* SUFFRIAN, 1848*Pachybrachys mulsanti* PERRIS, 1852*Pachybrachys fimbriolatus* ab. *kislovodskensis* ROUBAL, 1916*Pachybrachys fimbriolatus* ab. *varanensis* PIC, 1913*Pachybrachis fimbriolatus* ab. *weisei* BURLINI, 1968*Pachybrachis danieli* BURLINI, 1968

Distribution: S, Central and SE Europe, Ukraine, S European Russia, NW Caucasus, Daghestan, Turkey.

***Pachybrachis fraudator* LOPATIN, 1991**

Distribution: SE Kazakhstan, Kirghizstan (Kirghiz Alatau).

***Pachybrachis ghilarovi* LOPATIN, 1974**

Distribution: SE Kazakhstan.

***Pachybrachis glycyrrhizae* (OLIVIER, 1808)***Cryptocephalus glycyrrhizae* OLIVIER, 1808

Distribution: Transcaucasia, S Central Asia, Near and Middle East.

***Pachybrachis heptapotamicus* LOPATIN, 1997**

Distribution: Kazakhstan, Kirghizstan, E Turkmenistan.

***Pachybrachis hieroglyphicus* (LAICHARTING, 1781)**

*Cryptocephalus hieroglyphicus* LAICHARTING, 1781

*Cryptocephalus tristis* LAICHARTING, 1781

*Cryptocephalus histrio* FABRICIUS, 1781

*Pachybrachys hieroglyphicus* var. *ictericus* J. WEISE, 1882

*Pachybrachys hieroglyphicus* ab. *italicus* J. WEISE, 1882

*Pachybrachys hieroglyphicus* ab. *disconotatus* PIC, 1908

*Pachybrachys hieroglyphicus* ab. *doctoris* PIC, 1910

*Pachybrachys pseudoscriptus* WAGNER, 1927

*Pachybrachis hieroglyphicus* ab. *pseudosuturalis* BURLINI, 1968

Distribution: Central, E, and S Europe, S Urals, E Kazakhstan, Russian Far East.

***Pachybrachis hippophaes* (SUFFRIAN, 1848)**

*Pachybrachys hippophaes* SUFFRIAN, 1848

*Pachybrachys hippophaës* J. WEISE, 1882

*Pachybrachys hippophaës* ab. *lignosus* J. WEISE, 1882

*Pachybrachys hippophaes* ab. *galeazzii* PIC, 1910

Distribution: Mountains of E and Central Europe (Alps and Carpathians).

***Pachybrachis instabilis* (J. WEISE, 1887)**

*Pachybrachys instabilis* J. WEISE, 1887

Distribution: Uzbekistan and Kirghizstan (Karatau, Karzhantau, Kendyktau Mountain Ranges).

***Pachybrachis issykensis issykensis* (JACOBSON, 1901)**

*Pachybrachys sexsignatus issykensis* JACOBSON, 1901

Distribution: Kirghizstan (Central and S Tian-Shan).

***Pachybrachis issykensis gussakovskii* (LOPATIN, 1968)**

*Pachybrachys issykensis gussakovskii* LOPATIN, 1968

Distribution: Tajikistan (Hissar Mountain Range, Darvaz).

***Pachybrachis issykensis laevigatus* (BREIT, 1921)**

*Pachybrachys laevigatus* BREIT, 1921

Distribution: S Kazakhstan, Uzbekistan, Kirghizstan, Tajikistan up to Pamirs.

***Pachybrachis jacobsoni* (LOPATIN, 1968)**

*Pachybrachys jacobsoni* LOPATIN, 1968

Distribution: Uzbekistan, Kirghizstan, Tajikistan.

***Pachybrachis kaplini* LOPATIN, 1986**

Distribution: Turkmenistan (Kara-Kum).

***Pachybrachis kazachstanicus* LOPATIN, 1974**

Distribution: NW Kazakhstan.

***Pachybrachis kirgizicus kirgizicus* LOPATIN, 1974**

Distribution: Kirghizstan (N Coast of Issyk-Kul Lake).

***Pachybrachis kirgizicus salsolae* LOPATIN, 1974**

Distribution: Kirghizstan, Central Tian-Shan.

***Pachybrachis koktumensis* LOPATIN, 1991**

Distribution: SE Kazakhstan (Dzhungar Alatau).

***Pachybrachis korotjaevi* LOPATIN, 1995**

Distribution: Asian Russia (Tuva).

***Pachybrachis kuramensis* LOPATIN, 1974**

Distribution: N Tajikistan (Kuramin Mountain Range).

***Pachybrachis latipes* LOPATIN, 1971**

Distribution: E Kazakhstan, SW Mongolia.

***Pachybrachis limbatus* (MÉNÉTRIÉS, 1836)**

*Cryptocephalus limbatus* MÉNÉTRIÉS, 1836

*Pachybrachys limbatus* ab. *maculatus* SUFFRIAN, 1848

*Pachybrachys limbatus* ab. *ater* J. WEISE, 1882

Distribution: Balkans, Transcaucasia, Turkey, Near East.

***Pachybrachis lopatini* L. MEDVEDEV et RYBAKOVA, 1980**

Distribution: Asian Russia (S Siberia, Transbaikalia, Maritime Prov.), NE Mongolia, NE China.

***Pachybrachis marki* LOPATIN, 1997**

Distribution: S Tajikistan.

***Pachybrachis mendax* (SUFFRIAN, 1860)**

*Pachybrachys mendax* SUFFRIAN, 1860

*Pachybrachys probus* J. WEISE, 1882

*Pachybrachys probus sibiricus* LOPATIN, 1962

Distribution: Moldavia, S Ukraine, Crimea, Ciscaucasia, Transcaucasia, Maritime Prov., Kazakhstan, Asian Russia (E Siberia), Mongolia, W China.

***Pachybrachis merkensis* (LOPATIN, 1968)**

*Pachybrachys instabilis merkensis* LOPATIN, 1968

Distribution: S Kirghizstan, S Kazakhstan.

***Pachybrachis mitjaevi* LOPATIN et KULENOVA, 1982**

Distribution: SE Kazakhstan.

***Pachybrachis mogol* LOPATIN, 1986**

Distribution: S Tajikistan.

***Pachybrachis nigropunctatus* (SUFFRIAN, 1854)**

*Pachybrachys nigropunctatus* SUFFRIAN, 1854

*Pachybrachys paulonotatus* PIC, 1908

Distribution: Transcaucasia, Turkey, Near and Middle East, Steppe and Semi-deserts of Kazakhstan, Central Asia, Afghanistan.

***Pachybrachis nitidicollis* (J. WEISE, 1894)**

*Pachybrachys probus nitidicollis* J. WEISE, 1894

Distribution: S Kazakhstan, Central Asia (except for the Turkmenistan), Afghanistan.

***Pachybrachis ochropygus* (SOLSKY, 1872)**

*Pachybrachys ochropygus* SOLSKY, 1872

*Pachybrachys sexsignatus* J. WEISE, 1889

Distribution: Asian Russia (Amurland, Maritime Prov.), Mongolia, Korea, NE China.

***Pachybrachys paganus* LOPATIN, 1990**

Distribution: S Kazakhstan.

***Pachybrachys pallidulus* (SUFFRIAN, 1851)**

*Pachybrachys pallidulus* SUFFRIAN, 1851

*Pachybrachys suturalis* J. WEISE, 1882

Distribution: SE and southern parts of Central Europe.

***Pachybrachys pistor* (BREIT, 1921)**

*Pachybrachys pistor* BREIT, 1921

Distribution: Kazakhstan (except for the East), Uzbekistan.

***Pachybrachys pubipennis* LOPATIN, 1977**

Distribution: Armenia.

***Pachybrachys rufescens* LOPATIN, 1974**

Distribution: S Kazakhstan.

***Pachybrachys scripticollis* (FALDERMANN, 1837)**

*Pachybrachys scripticollis* FALDERMANN, 1837

*Pachybrachys israelita* TAPPES, 1871

Distribution: Armenia, Near East and Middle East, SW Turkmenistan.

***Pachybrachys scriptidorsum* (MARSEUL, 1875)**

*Pachybrachys scripticollis* SUFFRIAN, 1848, non FALDERMANN, 1837

*Pachybrachys scriptidorsum* MARSEUL, 1875

Distribution: SE Europe, N Caucasus, Kazakhstan, Asian Russia (Siberia, Khabarovsk Prov., Magadan Distr.), Mongolia, Korea, China.

***Pachybrachys semidesertus* LOPATIN, 1994**

Distribution: W Kazakhstan.

***Pachybrachys sericans* (SUFFRIAN, 1860)**

*Pachybrachys sericans* SUFFRIAN, 1860

*Pachybrachys sericans* ab. *unicolor* LOPATIN, 1962

Distribution: Asian Russia (Middle Siberia, Transbaikalia), Mongolia.

***Pachybrachys sinuatus* (MULSANT et REY, 1859)**

*Pachybrachys sinuatus* MULSANT et REY, 1859

*Pachybrachys haliciensis* L. MILLER, 1868

*Pachybrachys sinuatus* ab. *rufimanus* J. WEISE, 1882

Distribution: Central and SE Europe, Turkey.

***Pachybrachys skopini* (LOPATIN, 1967)**

*Pachybrachys skopini* LOPATIN, 1967

Distribution: SE Kazakhstan.

***Pachybrachys tekensis* LOPATIN, 1983**

Distribution: Turkmenistan (Kopetdagh Mountain Range).

***Pachybrachys tessellatus tessellatus* (OLIVIER, 1791)**

*Cryptocephalus tessellatus* OLIVIER, 1791

*Cryptocephalus histrio* OLIVIER, 1791  
*Pachybrachys tauricus* SUFFRIAN, 1848  
*Pachybrachys bisignatus* L. REDTENBACHER, 1849  
*Pachybrachys tessellatus* HAROLD, 1873  
*Pachybrachys tessellatus* ab. *picturatus* REY, 1883  
*Pachybrachys tessellatus* ab. *syriacus* J. WEISE, 1900  
*Pachybrachys tessellatus* ab. *melanopygus* G. MÜLLER, 1953

Distribution: Ukraine, Moldavia (except for the Mountains), European Russia, NW Caucasus.

***Pachybrachis tessellatus orientalis* (J. WEISE, 1894)**

*Pachybrachys tessellatus* var. *orientalis* J. WEISE, 1894

Distribution: Armenia, Turkey.

***Pachybrachis transbaicalius* L. MEDVEDEV, 1974**

Distribution: Asian Russia (Transbaikalia), Central and E Mongolia.

***Pachybrachis tuvensis tuvensis* L. MEDVEDEV, 1974**

Distribution: Asian Russia (Tuva).

***Pachybrachis tuvensis saluki* LOPATIN, 1995**

Distribution: Asian Russia (Altai).

***Pachybrachis velarum* WARCHALOWSKI, 1998**

Distribution: Transcaucasia, Turkey.

***Pachybrachis vermicularis* (SUFFRIAN, 1854)**

*Pachybrachys vermicularis* SUFFRIAN, 1854

*Pachybrachys astragali* STIERLIN, 1863

Distribution: SE European Russia, Daghestan, NW Kazakhstan.

**Genus: *Acolastus* GERSTAECKER, 1855**

*Thelyterotarsus* J. WEISE, 1882

*Serrinotus* TAN, 1992

***Acolastus anthracinus* (LOPATIN, 1976)**

*Thelyterotarsus anthracinus* LOPATIN, 1976

Distribution: Uzbekistan, Kirghizstan (Chatkal and Talas Mountain Ranges).

***Acolastus atraphaxidis* (LOPATIN, 1960)**

*Thelyterotarsus atraphaxidis* LOPATIN, 1960

Distribution: E Turkmenistan, Uzbekistan, SW Tajikistan.

***Acolastus badakshanicus* (LOPATIN, 1967)**

*Thelyterotarsus badakshanicus* LOPATIN, 1967

Distribution: E Tajikistan (Badakhshan), Afghanistan.

***Acolastus baeckmanni* (JACOBSON, 1916)**

*Thelyterotarsus baeckmanni* JACOBSON, 1916

Distribution: Deserts of Central Asia and Kazakhstan (Kyzyl-Kum, Muyn-Kum, Tau-Kum, Saryeshkotrau).

***Acolastus balchaschensis* (LOPATIN et KULENOVA, 1982)***Thelyterotarsus balchaschensis* LOPATIN et KULENOVA, 1982

Distribution: E Kazakhstan.

***Acolastus curtus* (LOPATIN, 1994)***Thelyterotarsus curtus* LOPATIN, 1994

Distribution: Israel.

***Acolastus darvazicus* (LOPATIN, 1975)***Thelyterotarsus darvazicus* LOPATIN, 1975

Distribution: E Tajikistan (Darvaz Mountain Range).

***Acolastus dzhungarus* (LOPATIN, 1976)***Thelyterotarsus dzhungarus* LOPATIN, 1976

Distribution: S and SE Kazakhstan.

***Acolastus fausti* (J. WEISE, 1882)***Thelyterotarsus fausti* J. WEISE, 1882*Thelyterotarsus armeniacus* PIC, 1906*Thelyterotarsus nigronotatus* PIC, 1906

Distribution: S Armenia and Azerbaijan, S Turkmenistan, NW Iran.

***Acolastus georgicus* (LOPATIN, 1986)***Thelyterotarsus georgicus* LOPATIN, 1986

Distribution: E Georgia, Azerbaijan.

***Acolastus gobustanus* (LOPATIN, 1991)***Thelyterotarsus gobustanus* LOPATIN, 1991

Distribution: Azerbaijan.

***Acolastus gurjevae gurjevae* (LOPATIN, 1968)***Thelyterotarsus gurjevae* (LOPATIN, 1968)

Distribution: Uzbekistan (Aktau and Nuratau Mountains Range).

***Acolastus gurjevae unicolor* (LOPATIN, 2001)***Thelyterotarsus gurjevae unicolor* LOPATIN, 2001

Distribution: Uzbekistan (Nuratau Mountain Range).

***Acolastus hauseri* (J. WEISE, 1887)***Thelyterotarsus hauseri* J. WEISE, 1887

Distribution: S Kazakhstan, Kirghizstan (W Tian-Shan Mountain Ranges).

***Acolastus iliensis* (LOPATIN, 1967)***Thelyterotarsus iliensis* LOPATIN, 1967

Distribution: SE Kazakhstan.

***Acolastus inopinatus* (LOPATIN, 1992)***Thelyterotarsus inopinatus* LOPATIN, 1992

Distribution: SE Kazakhstan.

***Acolastus intermedius* (LOPATIN, 1968)***Thelyterotarsus intermedius* LOPATIN, 1968

Distribution: Central Tajikistan.



***Acolastus issykensis* (LOPATIN, 1992)***Thelyterotarsus issykensis* LOPATIN, 1992

Distribution: Kirghizstan (Coasts of Issyk-Kul Lake).

***Acolastus ivanovi* (JACOBSON, 1925)***Thelyterotarsus ivanovi* JACOBSON, 1925

Distribution: Uzbekistan, Tajikistan, Kirghizstan (Turkestan Mountain Range and Fergana Valley).

***Acolastus jacobsoni* (LOPATIN, 1968)***Thelyterotarsus jacobsoni* LOPATIN, 1968

Distribution: SE Tajikistan.

***Acolastus karakirgiza* (JACOBSON, 1925)***Thelyterotarsus karakirgiza* JACOBSON, 1925

Distribution: Uzbekistan, Tajikistan (Turkestan Mountain Range).

***Acolastus karatavicus* (LOPATIN, 1976)***Thelyterotarsus karatavicus* LOPATIN, 1976

Distribution: Kazakhstan (Karatau Mountain Range, Syr-Darya River).

***Acolastus karateginus* (LOPATIN, 1992)***Thelyterotarsus karateginus* LOPATIN, 1992

Distribution: Tajikistan (Karatechin Mountain Range).

***Acolastus kaszabi* (LOPATIN, 1976)***Thelyterotarsus kaszabi* LOPATIN, 1976

Distribution: Mongolia.

***Acolastus khnzoriani* (LOPATIN, 1976)***Thelyterotarsus khnzoriani* LOPATIN, 1976

Distribution: SE Turkmenistan (Kugitang Tau Mountain Range).

***Acolastus korotjaevi* (LOPATIN, 1992)***Thelyterotarsus korotjaevi* LOPATIN, 1992

Distribution: S Tajikistan.

***Acolastus kryzhanovskii* (LOPATIN, 1976)***Thelyterotarsus kryzhanovskii* LOPATIN, 1976

Distribution: SE Turkmenistan.

***Acolastus kuramensis* (LOPATIN, 1992)***Thelyterotarsus kuramensis* LOPATIN, 1992

Distribution: N Tajikistan (Kuramin Mountain Range).

***Acolastus limbatus* (LOPATIN, 1992)***Thelyterotarsus limbatus* LOPATIN, 1992

Distribution: Kirghizstan (Coasts of Issyk-Kul Lake).

***Acolastus margaritae* (LOPATIN, 1997)***Thelyterotarsus margaritae* LOPATIN, 1997

Distribution: E Turkmenistan.

- Acolastus minimus* (JACOBSON, 1916)**  
*Thelyterotarsus minimus* JACOBSON, 1916  
Distribution: SE Kazakhstan, S Tajikistan.
- Acolastus mogoltavicus* (LOPATIN, 1992)**  
*Thelyterotarsus mogoltavicus* LOPATIN, 1992  
Distribution: N Tajikistan (Mogol Tau Mountain Ranges).
- Acolastus murinus* (LOPATIN, 1961)**  
*Thelyterotarsus murinus* LOPATIN, 1961  
Distribution: Tajikistan (Coasts of Iskander Kul Lake).
- Acolastus nanus* (LOPATIN, 1976)**  
*Thelyterotarsus nanus* LOPATIN, 1976  
Distribution: SW Kazakhstan.
- Acolastus nigrifrons* (JACOBSON, 1916)**  
*Thelyterotarsus nigrifrons* JACOBSON, 1916  
Distribution: Central Tajikistan (Hissar Mountain Range).
- Acolastus nuratavicus* (LOPATIN, 1992)**  
*Thelyterotarsus nuratavicus* LOPATIN, 1992  
Distribution: Uzbekistan (Nuratau Mountain Range).
- Acolastus pallidus pallidus* (LOPATIN, 1956)**  
*Thelyterotarsus pallidus* LOPATIN, 1956  
Distribution: Tajikistan.
- Acolastus pallidus montanus* (LOPATIN, 1976)**  
*Thelyterotarsus pallidus montanus* LOPATIN, 1976  
Distribution: E Tajikistan (Badakhshan).
- Acolastus pallidus tianshanicus* (LOPATIN, 1999)**  
*Thelyterotarsus pallidus tianshanicus* LOPATIN, 1999  
Distribution: Kirghizstan.
- Acolastus postfasciatus* (LOPATIN, 1975)**  
*Thelyterotarsus postfasciatus* LOPATIN, 1975  
Distribution: S Tajikistan.
- Acolastus przewalskii* (LOPATIN, 1992)**  
*Thelyterotarsus przewalskii* LOPATIN, 1992  
Distribution: Kirghizstan (Coasts of Issyk-Kul Lake).
- Acolastus regeli* (JACOBSON, 1898)**  
*Thelyterotarsus regeli* JACOBSON, 1898  
Distribution: SE Kazakhstan, NW China.
- Acolastus shahristanus* (LOPATIN, 1987)**  
*Thelyterotarsus shahristanus* LOPATIN, 1987  
Distribution: Uzbekistan (Turkestan Mountain Range).
- Acolastus similis* (LOPATIN, 1976)**  
*Thelyterotarsus similis* LOPATIN, 1976  
Distribution: SE Kazakhstan, S Tajikistan.

***Acolastus sogdianus* (LOPATIN, 1992)***Thelyterotarsus sogdianus* LOPATIN, 1992

Distribution: S Tajikistan, S Uzbekistan, SE Turkmenistan.

***Acolastus spitameni* (LOPATIN, 1990)***Thelyterotarsus spitameni* LOPATIN, 1990

Distribution: N Tajikistan (Zeravshan Mountain Range).

***Acolastus tadzhibajevi* (LOPATIN, 1975)***Thelyterotarsus tadzhibajevi* LOPATIN, 1975

Distribution: Tajikistan (Gazimailik and Hissar Mountain Ranges).

***Acolastus tadzhicus* (LOPATIN, 1968)***Thelyterotarsus tadzhicus* LOPATIN, 1968

Distribution: S Uzbekistan, S Tajikistan.

***Acolastus tatianae* (LOPATIN, 1983)***Thelyterotarsus tatianae* LOPATIN, 1983

Distribution: N Tajikistan (Zeravshan Mountain Range).

***Acolastus transcaspicus* (PIC, 1900)***Thelyterotarsus transcaspicus* PIC, 1900*Thelyterotarsus regeli transcaspica* PIC, 1900*Thelyterotarsus bergi* JACOBSON, 1916

Distribution: Turkmenistan.

***Acolastus turkestanicus* (LOPATIN, 1965)***Thelyterotarsus turkestanicus* LOPATIN, 1965

Distribution: N Tajikistan (Turkestan Mountain Range).

***Acolastus velutinus* (LOPATIN, 1967)***Thelyterotarsus velutinus* LOPATIN, 1967

Distribution: E Tajikistan.

***Acolastus volkovitschi* (LOPATIN, 1986)***Thelyterotarsus volkovitschi* LOPATIN, 1986

Distribution: S Uzbekistan.

***Acolastus zaissanicus* (LOPATIN, 1967)***Thelyterotarsus zaissanicus* LOPATIN, 1967

Distribution: E Kazakhstan.

***Acolastus zeravschanicus* (LOPATIN, 1960)***Thelyterotarsus zeravschanicus* LOPATIN, 1960

Distribution: Uzbekistan (Zeravshan Mountain Range).

Tribe: *Stylosomini* CLAVAREAU, 1913

Genus: *Stylosomus* SUFFRIAN, 1848

Subgenus: *Stylosomus (Microsomus)* BURLINI, 1968

*Stylosomus (Microsomus) costatus* LOPATIN, 1962

Distribution: Kirghizstan.

*Stylosomus (Microsomus) major* BREIT, 1918

*Stylosomus hirsutus* LOPATIN, 1956

*Stylosomus bechynei* BURLINI, 1957

Distribution: Kazakhstan and Central Asia (from Kopetdagh Mountain Range in the west, up to E Kazakhstan in the east), Mongolia.

*Stylosomus (Microsomus) weberi* REITTER, 1905

*Stylosomus sublineatus* J. SAHLBERG, 1913

*Stylosomus amoenus* J. SAHLBERG, 1913

*Stylosomus sahlbergi* WINKLER, 1929

Distribution: Mountains of S Kazakhstan, Central Asia.

Subgenus: *Stylosomus (Microstylus)* REY, 1883

*Stylosomus (Microstylus) ater* LOPATIN, 1962

Distribution: Kazakhstan (Transili and Karatau Mountain Ranges), Kirghizstan and Uzbekistan (W Tian-Shan).

*Stylosomus (Microstylus) cylindricus* MORAWITZ, 1860

Distribution: S Ukraine and European Russia, Caucasus, Kazakhstan.

*Stylosomus (Microstylus) tadzhicus* LOPATIN, 1965

Distribution: Mountains of Tajikistan.

Subgenus: *Stylosomus (Stylosomus)* SUFFRIAN, 1848

*Stylosomus (Stylosomus) cheni* LOPATIN, 1962

Distribution: S and E Kazakhstan, Kirghizstan, Uzbekistan.

*Stylosomus (Stylosomus) submetallicus* CHEN, 1941

*Stylosomus cinensis* LOPATIN, 1956

Distribution: NW China.

*Stylosomus (Stylosomus) tamaricis tamaricis* (HERRICH-SCHÄFFER, 1838)

*Cryptocephalus tamaricis* HERRICH-SCHÄFFER, 1838

*Stylosomus tamaricis* ab. *cruciatum* J. WEISE, 1882

*Stylosomus tamaricis* ab. *pallidicolor* PIC, 1909

*Stylosomus tamaricis* ab. *andalusiaca* PIC, 1909

*Stylosomus tamaricis* ab. *oberthuri* PIC, 1909

*Stylosomus tamaricis* ab. *nigripennis* ROUBAL, 1922

*Stylosomus tamaricis* ab. *leprieuri* PIC, 1923

*Stylosomus tamaricis* ab. *rufonotata* PIC, 1923

Distribution: S of European Russia, Ukraine, Caucasus.

***Stylosomus (Stylosomus) tamaricis nigrifrons* FLEISCHER, 1909***Stylosomus nigrifrons* FLEISCHER, 1909*Stylosomus macer* J. WEISE, 1882*Stylosomus fausti* REITTER, 1891

Distribution: Kazakhstan, Central Asia.

Subfamily: *Chlamisinae* GRESSITT, 1946Genus: *Chlamisus* RAFINESQUE, 1815***Chlamisus pubiceps* (CHŪJŌ, 1940)***Exema pubiceps* CHŪJŌ, 1940

Distribution: Asian Russia (S Maritime Prov.), Korea.

Subfamily: *Lamprosomatinae* GEMMINGER et HAROLD, 1874Tribe: *Lamprosomatini* GEMMINGER et HAROLD, 1874Genus: *Oomorphus* CURTIS, 1831*Lamprosoma* LACORDAIRE, 1848, non KIRBY, 1818***Oomorphus concolor* (STURM, 1807)***Byrrhus concolor* STURM, 1807*Phalacrus maritimus* STEPHENS, 1829*Oomorphus unicolor* BRULLÉ, 1834*Lamprosoma kolbei* SCHOLZ, 1926

Distribution: Central, SE, and S Europe, Caucasus.

Genus: *Oomorphoides* MONRÓS, 1956***Oomorphoides nigrocoeruleus* (BALY, 1873)***Lamprosoma nigrocoeruleus* BALY, 1873*Lamprosoma confucii* J. WEISE, 1889

Distribution: Asian Russia (Maritime Prov.), China, Japan.

Subfamily: *Eumolpinae* HOPE, 1840Tribe: *Nodinini* CHEN, 1940Genus: *Basilepta* BALY, 1860*Nodostoma* MOTSCHULSKY, 1860***Basilepta balyi* (HAROLD, 1877)***Nodostoma balyi* HAROLD, 1877

Distribution: Asian Russia (S Sakhalin, Kurile Is.), Japan.

***Basilepta fulvipes* (MOTSCHULSKY, 1860)***Nodostoma fulvipes* MOTSCHULSKY, 1860

Distribution: Asian Russia (Transbaikalia, Amurland, Maritime Prov., Kurile Is.), Mongolia, Korea, China, Japan.

***Basilepta kaszabi* LOPATIN, 1962**

Distribution: China (Timbu).

**Genus: *Weiselina* REITTER, 1890*****Weiselina lenkorana* REITTER, 1890**

Distribution: E Azerbaijan.

**Genus: *Pagria* LEFÈVRE, 1884*****Pagria signata* (MOTSCHULSKY, 1858)***Metachroma signata* MOTSCHULSKY, 1858*Nodostoma consimilis* BALY, 1874*Nodostoma flavopustulata* BALY, 1874

Distribution: Asian Russia (Maritime Prov.), Korea, China, S Asia.

**Genus: *Bedelia* LEFÈVRE, 1875*****Bedelia angustata* LEFÈVRE, 1875***Bedelia insignis* LEFÈVRE, 1875

Distribution: Transcaucasia, Iran.

***Bedelia kaschgarica* LOPATIN, 1962**

Distribution: NW China (Kaschgar).

***Bedelia kokanica* (SOLSKY, 1882)***Nodostoma kokanica* SOLSKY, 1882

Distribution: S Kazakhstan, Uzbekistan, Tajikistan.

***Bedelia viridicoerulea* REITTER, 1901**

Distribution: Tajikistan, Kirghizstan.

**Genus: *Chloropterus* MORAWITZ, 1861***Heterocnensis* MORAWITZ, 1860*Heterocnemis* MORAWITZ, 1861***Chloropterus grandis* J. WEISE, 1889**

Distribution: Kazakhstan, Turkmenistan, Uzbekistan, Tajikistan, Mongolia.

***Chloropterus lefevrei* REITTER, 1890***Chloropterus nigrofasciatus* REITTER, 1897

Distribution: Transcaucasia, Valleys of Kazakhstan and Central Asia, Near East.

***Chloropterus unguiculatus* LOPATIN, 1965**

Distribution: S Tajikistan.

***Chloropterus versicolor* (MORAWITZ, 1860)***Heterocnemis versicolor* MORAWITZ, 1860*Chloropterus moldaviensis* PIC, 1909

Distribution: Southern parts of Steppe zone of Ukraine and European Russia, Caucasus, Kazakhstan, SW Turkmenistan.

**Genus: *Cleoporus* LEFÈVRE, 1884*****Cleoporus variabilis* (BALY, 1874)***Paria variabilis* BALY, 1874*Paria robustus* BALY, 1874*Cleoporus tibialis* LEFÈVRE, 1885*Cleoporus niger* J. WEISE, 1922*Cleoporus pygmaeus* J. WEISE, 1922

Distribution: Asian Russia (Maritime Prov.), China, Korea, Japan.

Tribe: *Colasposomini* ŠPRINGLOVA, 1960**Genus: *Colasposoma* LAPORTE, 1833*****Colasposoma dauricum* MANNERHEIM, 1849***Colasposoma mongolicum* MOTSCHULSKY, 1860

Distribution: Asian Russia (Transbaikalia, Amurland, Maritime Prov., Sakhalin), Mongolia, N China, Korea.

Tribe: *Eumolpini* JACOBY, 1908**Genus: *Eumolpus* ILLIGER, 1798***Eumolpus* DUFTSCHMIDT, 1825*Chrysochus* CHEVROLAT in DEJEAN, 1836***Eumolpus asclepiadeus* (PALLAS, 1773)***Chrysomela asclepiadea* PALLAS, 1773*Chrysomela cyanea* OLIVIER, 1791*Cryptocephalus praetiosus* D. H. SCHNEIDER, 1791

Distribution: W, S, Central, and southern parts of E Europe, Turkey, Kazakhstan, Asian Russia (southern part of W Siberia).

***Eumolpus chinensis* BALY, 1859***Eumolpus singularis* LEFÈVRE, 1884*Eumolpus cyclostoma* J. WEISE, 1889

Distribution: Mongolia, Central and E China.

***Eumolpus globicollis* LEFÈVRE, 1888***Eumolpus curtus* PIC, 1927

Distribution: Asian Russia (S Amurland, Maritime Prov.), Korea, N China.

***Eumolpus goniostoma* J. WEISE, 1889**

Distribution: E Kazakhstan, Asian Russia (S Siberia), Mongolia, N China.

**Genus: *Abiromorphus* PIC, 1924*****Abiromorphus anceyi* PIC, 1924**

Distribution: Asian Russia (S Maritime Prov.), NE China, Korea.

**Genus: *Chrysochares* MORAWITZ, 1861*****Chrysochares asiaticus asiaticus* (PALLAS, 1771)***Chrysomela asiatica* PALLAS, 1771

Distribution: S of European Russia, Transcaucasia.

***Chrysochares asiaticus orientalis* LOPATIN, 1963**

Distribution: Kazakhstan, Turkmenistan, Uzbekistan, S Tajikistan.

***Chrysochares constricticollis* LOPATIN, 1963**

Distribution: E Georgia, Daghestan.

***Chrysochares punctatus* (GEBLER, 1845)***Chrysochus punctatus* GEBLER, 1845*Chrysochus aeneus* BALLION, 1878*Chrysochares aeneocupreus* CHEN, 1935

Distribution: Kazakhstan, Turkmenistan, Uzbekistan, S nad N Tajikistan, Mongolia, NW China.

**Tribe: *Adoxini* JACOBY, 1908****Genus: *Malegia* LEFÈVRE, 1883***Microlypesthes* PIC, 1936***Malegia aenea* (CHEN, 1940)***Microlypesthes aenea* CHEN, 1940*Malegia flavipes* CHŪJŪ, 1942

Distribution: Asian Russia (Amurland, Maritime Prov.), Korea, N China.

***Malegia colchica* REITTER, 1912**

Distribution: Caucasus (Georgia).

***Malegia jacobsoni* SUMAKOV, 1901***Malegia wisei* REITTER, 1912

Distribution: S Turkmenistan.

***Malegia turkestanica turkestanica* REITTER, 1890***Malegia hirsuta* JACOBSON, 1894

Distribution: Kazakhstan, Kirghizstan, Tajikistan, Uzbekistan.

***Malegia turkestanica uralensis* REITTER, 1912**

Distribution: NW Kazakhstan.



**Genus: *Lypesthes* BALY, 1863***Leprotos* BALY, 1863***Lypesthes ater* (MOTSCHULSKY, 1860)***Fidia atra* MOTSCHULSKY, 1860*Leprotos pulverulentus* JACOBY, 1885

Distribution: Asian Russia (Maritime Prov.), Korea, China.

**Genus: *Andosia* J. WEISE, 1896*****Andosia reitteri* J. WEISE, 1896**

Distribution: S Kazakhstan, Uzbekistan, S Tajikistan.

**Genus: *Caspiana* LOPATIN, 1977*****Caspiana armata* LOPATIN, 1977**

Distribution: SW Turkmenistan.

**Genus: *Parnops* JACOBSON, 1894*****Parnops glasunovi ferghanus* LOPATIN, 1976**

Distribution: Kirghizstan (Foothill Plains of Chatkal Mountain Range).

***Parnops glasunovi glasunovi* JACOBSON, 1894**

Distribution: S Turkmenistan, Iran, Uzbekistan, S Tajikistan.

**Genus: *Trichochrysea* BALY, 1861***Lefèvrella* JACOBSON, 1894*Heteraspis* CHAPUIS, 1874*Heteraspibrachys* PIC, 1907***Trichochrysea amygdali amygdali* (OGLOBLIN, 1941)***Lefèvrella amygdali* OGLOBLIN, 1941

Distribution: Tajikistan (Hissar, Babatagh, Gazimailik Mountain Ranges).

***Trichochrysea amygdali nuratavica* LOPATIN, 1976**

Distribution: Uzbekistan (Nuratau Mountain Range).

***Trichochrysea arnoldii* (L. MEDVEDEV, 1957)***Lefèvrella arnoldii* L. MEDVEDEV, 1957

Distribution: Kirghizstan (Fergan, Baubash-Ata, Bozbutau Mountain Ranges).

***Trichochrysea occidentalis* (J. WEISE, 1887)***Heteraspis occidentalis* J. WEISE, 1887*Heteraspis hauseri* J. WEISE, 1890*Trichochrysea bipubescens* PIC, 1907

Distribution: S Kazakhstan, Uzbekistan (W Tian-Shan).

**Genus: *Dermestops* JACOBSON, 1898***Allecumolpus* L. MEDVEDEV, 1956***Dermestops ahngeri* JACOBSON, 1898***Allecumolpus smirnovi* L. MEDVEDEV, 1956

Distribution: Turkmenistan.

**Genus: *Callipta* LEFÈVRE, 1885***Calliope* J. WEISE 1882*Arachnipta* L. MEDVEDEV, 1957***Callipta borealis* LOPATIN, 1976**

Distribution: Uzbekistan (Kara-Kum Desert, Bukantau, Tamdytau, Aktau Mountains).

***Callipta fausti fausti* (J. WEISE, 1882)***Calliope fausti* J. WEISE, 1882

Distribution: E Transcaucasia (Azerbaijan, S Daghestan).

***Callipta fausti badghyza* LOPATIN, 1997**

Distribution: Turkmenistan (Badkhyz).

***Callipta fausti balchana* LOPATIN, 1997**

Distribution: W Turkmenistan (Balkhan Mountains).

***Callipta fausti murgabica* LOPATIN, 1997**

Distribution: Turkmenistan (Iolatan').

***Callipta fausti palvanica* LOPATIN, 1997**

Distribution: Turkmenistan (Kaplankhyr Plateau).

***Callipta fausti turcomana* L. MEDVEDEV, 1957***Callipta smirnovi* L. MEDVEDEV, 1957

Distribution: S Turkmenistan (Kopetdagh Mountain Range).

**Genus: *Aphilenia* J. WEISE, 1889****Subgenus: *Aphilenia (Pseudoaphilenia)* LOPATIN, 1976*****Aphilenia (Pseudoaphilenia) hauseri* J. WEISE, 1894**

Distribution: Kazakhstan (Sandy Deserts), S Central Asia (Sandy Deserts).

**Subgenus: *Aphilenia (Aphilenia)* J. WEISE, 1889*****Aphilenia (Aphilenia) interrupta* J. WEISE, 1889**

Distribution: Turkmenistan, Uzbekistan, SW Tajikistan (Sandy Deserts).

***Aphilenia (Aphilenia) ornata* REITTER, 1888**

Distribution: Kazakhstan, Turkmenistan (Deserts).

***Aphilenia (Aphilenia) parvula* J. WEISE, 1894**

Distribution: Turkmenistan, Uzbekistan (Deserts).

**Genus: *Atomyria* JACOBSON, 1894*****Atomyria sarafschanica* (SOLSKY, 1882)***Nodostoma sarafschanica* SOLSKY, 1882

Distribution: SW Kazakhstan, Uzbekistan, Tajikistan (Bank of the Rivers Amu-Darya, Piandzh, Vakhsh).

**Genus: *Macrocoma* CHAPUIS, 1874***Pseudocolaspis* J. THOMSON, 1858, non LAPORTE, 1833*Eubraxis* BALY, 1878*Ischyromus* JACOBSON, 1893***Macrocoma rubripes turcmena* LOPATIN, 1976**

Distribution: S Turkmenistan (Kopetdagh Mountain Range).

***Macrocoma sarvadensis* (SOLSKY, 1882)***Pseudocolaspis sarvadensis* SOLSKY, 1882*Ischyromus sarvadensis* JACOBSON, 1893

Distribution: Kazakhstan and Central Asia.

**Genus: *Bromius* CHEVROLAT in DEJEAN, 1837***Adoxus* KIRBY, 1837***Bromius obscurus* (LINNAEUS, 1758)***Chrysomela obscura* LINNAEUS, 1758*Chrysomela villosula* SCHRANK, 1781*Bromius obscurus* var. *epilobii* J. WEISE, 1882*Bromius obscurus* var. *weisei* HEYDEN, 1883*Bromius obscurus* ab. *lewisi* J. WEISE, 1898*Bromius obscurus* ab. *concinus* J. WEISE, 1898

Distribution: Europe except for the extreme North, Asian Russia (Siberia, Russian Far East), Japan. Introduced to N America.

**Genus: *Anidania* REITTER, 1889*****Anidania luctuosa* (SOLSKY, 1882)***Pseudocolaspis luctuosa* SOLSKY, 1882*Pseudocolaspis rubripes* REITTER, 1888

Distribution: S Kazakhstan, Uzbekistan, Tajikistan, N Afghanistan.

**Genus: *Adoxinia* REITTER, 1888*****Adoxinia spinipes* REITTER, 1888***Adoxinia jacobsoni* REITTER, 1894

Distribution: Turkmenistan, Uzbekistan, W Tajikistan (Sandy Deserts).

**Genus: *Pachnephopterus* REITTER, 1892*****Pachnephopterus weisei* REITTER, 1892**

Distribution: Transcaucasia, Turkey, Turkmenistan, Uzbekistan, Afghanistan.

**Genus: *Pachnephorus* CHEVROLAT in DEJEAN, 1837****Subgenus: *Pachnephorus (Pachnephoriscus)* LOPATIN, 1976*****Pachnephorus (Pachnephoriscus) jacobsoni* LOPATIN, 1976**

Distribution: Uzbekistan.

**Subgenus: *Pachnephorus (Pachnephorus)* CHEVROLAT in DEJEAN, 1837*****Pachnephorus (Pachnephorus) brunneus* L. MEDVEDEV, 1957**

Distribution: Turkmenistan.

***Pachnephorus (Pachnephorus) canus* J. WEISE, 1882**

Distribution: S Ukraine, Daghestan, E Mediterranean.

***Pachnephorus (Pachnephorus) fulvus* LOPATIN, 1976**

Distribution: Uzbekistan.

***Pachnephorus (Pachnephorus) lewisi* BALY, 1878***Pachnephorus squamosus* CHŪJŌ, 1936

Distribution: Asian Russia (Maritime Prov.), China.

***Pachnephorus (Pachnephorus) pilosus* (ROSSI, 1790)***Cryptocephalus pilosus* ROSSI, 1790*Cryptocephalus arenarius* PANZER, 1797

Distribution: Europe except for the extreme North, Caucasus, Turkey, Siberia.

***Pachnephorus (Pachnephorus) tessellatus* (DUFTSCHMIDT, 1825)***Eumolpus tessellatus* DUFTSCHMIDT, 1825*Pachnephorus villosus* FAIRMAIRE, 1861, non DUFTSCHMIDT, 1825*Pachnephorus tessellatus* ab. *sabulosus* GEBLER, 1848

Distribution: All over the Palaeartic.

***Pachnephorus (Pachnephorus) turcomanicus* L. MEDVEDEV, 1957**

Distribution: Turkmenistan, Uzbekistan.

***Pachnephorus (Pachnephorus) villosus* (DUFTSCHMIDT, 1825)***Eumolpus villosus* DUFTSCHMIDT, 1825*Pachnephorus?* *aspericollis* FAIRMAIRE, 1861*Pachnephorus?* *rugaticollis* L. MILLER, 1862*Pachnephorus silvanae* DACCORDI, 1977*Pachnephorus villosus* ab. *chalybaea* J. WEISE, 1882

Distribution: S and SE Europe, Crimea, Turkey, Caucasus.

**Subfamily: *Chrysomelinae* LATREILLE, 1802****Tribe: *Timarchini* MOTSCHULSKY, 1860****Genus: *Timarcha* DEJEAN, 1821***Timarchistoma* MOTSCHULSKY, 1860

**Subgenus: *Timarcha (MetalloTimarcha)* MOTSCHULSKY, 1860**

***Timarcha (MetalloTimarcha) hummeli hummeli* FALDERMANN, 1837**

Distribution: N Caucasus, W Transcaucasia.

***Timarcha (MetalloTimarcha) hummeli armeniaca* FALDERMANN, 1837**

Distribution: Armenia.

***Timarcha (MetalloTimarcha) metallica* (LAICHARTING, 1781)**

*Chrysomela metallica* LAICHARTING, 1781

*Timarcha splendens* MATZEK, 1843

*Timarcha metallica* ab. *wendleri* FLEISCHER, 1909

Distribution: Mountains of SW and Central Europe, Ukrainian Carpathians.

**Subgenus: *Timarcha (Timarcha)* DEJEAN, 1821**

***Timarcha (Timarcha) goettingensis* (LINNAEUS, 1758)**

*Chrysomela göttingensis* LINNAEUS, 1758

*Chrysomela violaceo-inigra* DEGEER, 1775

*Chrysomela coriaria* LAICHARTING, 1781

*Timarcha goettingensis* ab. *purpurascens* MARSEUL, 1883

*Timarcha goettingensis* ab. *parisiensis* BECHYNÉ, 1946

Distribution: Europe except for the extreme North.

***Timarcha (Timarcha) rugulosa* HERRICH-SCHÄFFER, 1838**

*Timarcha lomnickii* L. MILLER, 1867

*Timarcha globata* FAIRMAIRE, 1873

Distribution: SE Europe, Moldavia, Ukraine, Crimea.

***Timarcha (Timarcha) tenebricosa tenebricosa* (FABRICIUS, 1775)**

*Chrysomela tenebricosa* FABRICIUS, 1775

*Timarcha tenebricosa* ab. *fraterna* BECHYNÉ, 1945

*Timarcha tenebricosa* ab. *bulgarica* BECHYNÉ, 1945

Distribution: W and S Europe, Steppe zone of Ukraine and European Russia.

***Timarcha (Timarcha) tenebricosa iberica* MOTSCHULSKY, 1849**

Distribution: Caucasus.

***Timarcha (Timarcha) tenebricosa motschulskyi* BECHYNÉ, 1945**

Distribution: Crimea.

Tribe: *Chrysomelini* LATREILLE, 1802

Subtribus: *Chrysolinina* CHEN, 1936

**Genus: *Ambrostoma* MOTSCHULSKY, 1860**

***Ambrostoma quadriimpresum* MOTSCHULSKY, 1845**

Distribution: Asian Russia (Transbaikalia, Amurland, Maritime Prov.), Mongolia, N China, Korea.

**Genus: *Crosita* MOTSCHULSKY, 1860*****Crosita afghanica* LOPATIN, 1962**

Distribution: E Tajikistan (Badakhshan), NE Afghanistan.

***Crosita altaica altaica* (GEBLER, 1823)***Chrysomela altaica* GEBLER, 1823*Chrysomela insignis* FISCHER-WALDHEIM, 1842

Distribution: Central and E Kazakhstan, Tarbagatay, Central Tian-Shan, S Asian Russia (Altai), Mongolia.

***Crosita altaica faldermanni* (KRYNICKI, 1832)***Chrysomela faldermanni* KRYNICKI, 1832

Distribution: SE European Russia, Central and E Kazakhstan (Dry Steppe), Mongolia (Mongol Altai Mountain Range).

***Crosita altaica maximowitschi* (ZUBKOV, 1833)***Chrysomela maximowitschi* ZUBKOV, 1833

Distribution: NE and Central Kazakhstan (Saur Mountain Range), SW Mongolia.

***Crosita bogutensis* LOPATIN, 1996**

Distribution: Kazakhstan (Boghuty Mountains).

***Crosita elegans* LOPATIN, 1968**

Distribution: Mongolia (E Mongol Altai, High-Mountain).

***Crosita heptapotamica* JACOBSON, 1895**

Distribution: Central Tian-Shan (Transili and Kungej-Alatau Mountain Ranges).

***Crosita kowalewskii* (GEBLER, 1836)***Chrysomela kowalewskii* GEBLER, 1836*Crosita kowalewski matronula* J. WEISE, 1894*Crosita kowalewski gobica* LOPATIN, 1968*Crosita kowalewski gobica* ab. *cyaneoviridula* LOPATIN, 1968*Crosita kowalewski gobica* ab. *transitoria* LOPATIN, 1968*Crosita kowalewski gobica* ab. *unicolor* LOPATIN, 1968

Distribution: Asian Russia (S Altai), Mongolia, NW China.

***Crosita longipes* JACOBSON, 1898**

Distribution: Asian Russia (S Altai, Tuva), Mongolia (Mountain's Steppe).

***Crosita pigra* J. WEISE, 1894***Crosita gemellata* LOPATIN, 1971

Distribution: Asian Russia (Tuva), Mongolia (High-mountain's Steppe).

**Genus: *Gnathomela* JACOBSON, 1895*****Gnathomela glasunovi* BECHYNÉ, 1962**

Distribution: Uzbekistan (Nuratau Mountain Range).

***Gnathomela laevigata laevigata* LOPATIN, 1972**

Distribution: Uzbekistan (Kuramin Mountain Range).

***Gnathomela laevigata kirghizica* LOPATIN, 1990**

Distribution: Kirghizstan (Kirghiz Alatau Mountain Range).

***Gnathomela meridionalis* LOPATIN, 1972**

Distribution: S Uzbekistan (Baisuntau Mountain Range).

***Gnathomela mutica* LOPATIN, 1972**

Distribution: W Tian-Shan (Ugam, Pskem and Karzhantau Mountain Ranges).

***Gnathomela ovczinnikovi* LOPATIN, 1990**

Distribution: Kirghizstan (Fergan Mountain Range).

***Gnathomela praestans* LOPATIN, 1972**

Distribution: W Tian-Shan (Chatkal, Little Chimghan and Mohol Tau Mountain Ranges).

***Gnathomela valida valida* LOPATIN, 1972**

Distribution: W Tian-Shan (Karatau, Karzhantau, Talas and Ugam Mountain Ranges).

***Gnathomela valida nuratavica* LOPATIN, 1972**

Distribution: Uzbekistan (Nuratau Mountain Range).

**Genus: *Chrysolina* MOTSCHULSKY, 1860***Chrysomela* auct. non LINNAEUS, 1758*Atechna* CHEVROLAT in DEJEAN, 1836**Subgenus: *Chrysolina (Bittotaenia)* MOTSCHULSKY, 1860*****Chrysolina (Bittotaenia) grata grata* (FALDERMANN, 1837)***Chrysomela grata* FALDERMANN, 1837

Distribution: Transcaucasia, SW Turkmenistan, N Iran.

***Chrysolina (Bittotaenia) grata turanica* (REITTER, 1888)***Chrysomela turanica* REITTER, 1888

Distribution: SW Turkmenistan, SW Tajikistan.

***Chrysolina (Bittotaenia) salviae salviae* (GERMAR, 1824)***Chrysomela salviae* GERMAR, 1824

Distribution: Europe, Caucasus, Turkey, Syria.

***Chrysolina (Bittotaenia) salviae sculptipennis* (FALDERMANN, 1837)***Chrysomela sculptipennis* FALDERMANN, 1837

Distribution: Caucasus, Turkey.

**Subgenus: *Chrysolina (Pseudocrosita)* LOPATIN, 1999*****Chrysolina (Pseudocrosita) bactriana* LOPATIN, 1961***Chrysolina globicollis* LOPATIN, 1977

Distribution: S Tajikistan (low mountains), S Kirghizstan (Alai Mountain Range).

**Subgenus: *Chrysolina (Lithopteroides)* E. STRAND, 1935***Lithoptera* MOTSCHULSKY, 1860***Chrysolina (Lithopteroides) exanthematica exanthematica* (WIEDEMANN, 1821)***Chrysomela exanthematica* WIEDEMANN, 1821*Chrysomela guttata* GEBLER, 1817, non FABRICIUS, 1792*Chrysomela musiva* GEBLER, 1830

Distribution: Asian Russia (Siberia, Russian Far East), China, N India, Vietnam.

***Chrysolina (Lithopteroides) exanthematica gemmifera* MOTSCHULSKY, 1860**? *Chrysomela guttifera* MOTSCHULSKY, 1860

Distribution: S Siberia, Mongolia.

**Subgenus: *Chrysolina (Crositops)* MARSEUL, 1883*****Chrysolina (Crositops) kabaki* LOPATIN, 1988**

Distribution: E Kazakhstan (High Mountains).

***Chrysolina (Crositops) pedestris* (GEBLER, 1823)***Chrysomela pedestris* GEBLER, 1823

Distribution: E Kazakhstan, Asian Russia (S Siberia, Altais).

**Subgenus: *Chrysolina (Timarchoptera)* MOTSCHULSKY, 1860*****Chrysolina (Timarchoptera) haemochlora* (GEBLER, 1823)***Chrysomela haemochlora* GEBLER, 1823*Chrysomela rubra* MOTSCHULSKY, 1845

Distribution: W Siberia, Mongolia.

**Subgenus: *Chrysolina (Chrysocrosita)* BECHYNÉ, 1950*****Chrysolina (Chrysocrosita) jakovlevi* (J. WEISE, 1894)***Chrysomela jakovlevi* J. WEISE, 1894

Distribution: Asian Russia (Tuva, Sayan).

***Chrysolina (Chrysocrosita) spectabilis spectabilis* MOTSCHULSKY, 1860**

Distribution: NE Asian Russia (Kamchatka).

***Chrysolina (Chrysocrosita) spectabilis polychroma* L. MEDVEDEV, 1975***Chrysolina cyaneovinosa* L. MEDVEDEV, 1978

Distribution: Asian Russia (Sayan), N Mongolia.

**Subgenus: *Chrysolina (Lithocrosita)* L. MEDVEDEV, 1982*****Chrysolina (Lithocrosita) rugulosa* (GEBLER, 1841)***Chrysomela rugulosa* GEBLER, 1841

Distribution: Asian Russia (Altais).

**Subgenus: *Chrysolina (Heliostola)* MOTSCHULSKY, 1860*****Chrysolina (Heliostola) carpathica* (FUSS, 1856)***Chrysomela carpathica* FUSS, 1856



*Chrysomela carpatica* ab. *hopffgarteni* J. WEISE, 1890  
*Chrysomela lichenis* var. *gabrieli* J. WEISE, 1903  
*Chrysolina carpathica* ab. *adulterata* BECHYNÉ, 1947  
*Chrysolina carpathica* ab. *jaspidea* BECHYNÉ, 1947  
*Chrysolina carpathica* ab. *lucifuga* BECHYNÉ, 1947  
*Chrysolina carpathica* ab. *internitens* BECHYNÉ, 1947  
 Distribution: SE Europe, W Ukraine (Carpathians).

***Chrysolina (Heliostola) katonica* LOPATIN, 1988**

Distribution: E Kazakhstan.

***Chrysolina (Heliostola) lichenis* (RICHTER, 1820)**

*Chrysomela lichenis* RICHTER, 1820  
*Chrysomela islandica* GERMAR, 1824  
*Chrysomela athena* GERMAR, 1824  
*Chrysomela lichenis* var. *moraviaca* J. WEISE, 1882  
*Chrysomela lichenis* var. *nigerrima* J. WEISE, 1882  
*Chrysomela lichenis* var. *quadricollis* J. WEISE, 1882  
*Chrysomela lichenis* var. *schneideri* SEIDLITZ, 1891  
*Chrysomela lichenis* var. *rhipaea* J. WEISE, 1898  
*Chrysolina lichenis* ab. *conducta* BECHYNÉ, 1947  
*Chrysolina lichenis* ab. *cunctabunda* BECHYNÉ, 1947  
*Chrysolina lichenis* ab. *februa* BECHYNÉ, 1947  
*Chrysolina lichenis* ab. *hiemata* BECHYNÉ, 1947  
*Chrysolina lichenis* ab. *jugosa* BECHYNÉ, 1947  
*Chrysolina lichenis* ab. *mendosa* BECHYNÉ, 1947  
*Chrysolina lichenis* ab. *neta* BECHYNÉ, 1947  
*Chrysolina lichenis* ab. *pervilis* BECHYNÉ, 1947  
*Chrysolina lichenis* ab. *pleziensis* BECHYNÉ, 1947  
*Chrysolina lichenis* ab. *prunicea* BECHYNÉ, 1947  
*Chrysolina lichenis* ab. *tatrica* BECHYNÉ, 1947  
*Chrysolina lichenis* ab. *urgensis* BECHYNÉ, 1947  
 Distribution: Central and SE Europe, W Ukraine.

***Chrysolina (Heliostola) montana* (GEBLER, 1847)**

*Chrysomela montana* GEBLER, 1847  
*Chrysolina despecta* KONTKANEN, 1957  
 Distribution: Asian Russia (Altai).

***Chrysolina (Heliostola) schewyrevi* (JACOBSON, 1895)**

*Chrysomela schewyrevi* JACOBSON, 1895  
 Distribution: Asian Russia (W Siberia, Altai, Transbaikalia).

**Subgenus: *Chrysolina (Paraheliostola)* L. MEDVEDEV, 1992**

***Chrysolina (Paraheliostola) lomakini* MIKHAILOV, 2002**

Distribution: Asian Russia (Khakassia).

***Chrysolina (Paraheliostola) soiota soiota* (JACOBSON, 1924)**

*Chrysomela soiota* JACOBSON, 1924  
 Distribution: Asian Russia (Sayan).

***Chrysolina (Paraheliostola) soiota khakassa* MIKHAILOV, 2002**

Distribution: Asian Russia (Khakassia).

**Subgenus: *Chrysolina (Pleurosticha)* MOTSCHULSKY, 1860*****Chrysolina (Pleurosticha) cavigera* (J. SAHLBERG, 1885)***Chrysomela cavigera* J. SAHLBERG, 1885

Distribution: Arctic, N Siberia.

***Chrysolina (Pleurosticha) gebleri gebleri* L. MEDVEDEV, 1992**

Distribution: Asian Russia (Altai).

***Chrysolina (Pleurosticha) gebleri sajanensis* L. MEDVEDEV, 1979***Chrysomela gebleri baicalica* L. MEDVEDEV, 1974, non MOHR, 1966

Distribution: Asian Russia (Cisbaikalia, Transbaikalia).

***Chrysolina (Pleurosticha) latimargo* (J. WEISE, 1896)***Chrysomela latimargo* J. WEISE, 1896*Chrysomela changaica* LOPATIN, 1968

Distribution: Asian Russia (S Siberia), Mongolia.

***Chrysolina (Pleurosticha) subcostata* (GEBLER, 1848)***Chrysomela subcostata* GEBLER, 1848*Chrysolina pirka* TAKIZAWA, 1970

Distribution: Asian Russia (Siberia, Russian Far East), N Japan.

***Chrysolina (Pleurosticha) sylvatica* (GEBLER, 1823)***Chrysomela sylvatica* GEBLER, 1823

Distribution: Asian Russia (Altai).

***Chrysolina (Pleurosticha) tolli* (JACOBSON, 1910)***Chrysomela tolli* JACOBSON, 1910*Chrysomela rufipes* MÉNÉTRIÉS, 1851, non LINNAEUS, 1758

Distribution: Arctic Asia.

***Chrysolina (Pleurosticha) uraltuensis* MIKHAILOV, 2000**

Distribution: Asian Russia (W Sayan, S of Krasnoyarsk).

**Subgenus: *Chrysolina (Arctolina)* KONTKANEN, 1959*****Chrysolina (Arctolina) ballioni* (LOPATIN, 1968)***Chrysomela ballioni* LOPATIN, 1968

Distribution: SE Kazakhstan (Tarbagatay and N Slopes of Dzhungar Alatau Mountain Ranges).

***Chrysolina (Arctolina) boeberi* (HAROLD, 1874)***Chrysomela boeberi* HAROLD, 1874*Chrysomela sulcata* GERMAR, 1824, non GEBLER, 1823*Chrysomela magniceps* J. SAHLBERG, 1885

Distribution: Asian Russia (Chukotka, Magadan Distr., Kamchatka), Alaska.

***Chrysolina (Arctolina) bungei* (JACOBSON, 1910)***Chrysomela bungei* JACOBSON, 1910

Distribution: NE Siberia (Novosiberian Is., Central part of NE Siberia)

***Chrysolina (Arctolina) cyanella* (GEBLER, 1830)***Chrysomela cyanella* GEBLER, 1830

Distribution: Asian Russia (Altai).

***Chrysolina (Arctolina) dolini* LOPATIN, 1999**

Distribution: SE Kazakhstan (Dzhungar Alatau).

***Chrysolina (Arctolina) dubeshkoe* (L. MEDVEDEV, 1974)***Chrysomela dubeshkoe* L. MEDVEDEV, 1974*Chrysomela bannikovae* L. MEDVEDEV, 1978

Distribution: Mongolia (Khanghay Mountain Range).

***Chrysolina (Arctolina) kaikana* LOPATIN, 1992**

Distribution: SE Kazakhstan (Kaikan Mountain Range).

***Chrysolina (Arctolina) kryzhanovskiy* (LOPATIN, 1968)***Chrysomela kryzhanovskiy* LOPATIN, 1968

Distribution: SE Kazakhstan (Zhabyn and Tastau Mountain Ranges).

***Chrysolina (Arctolina) octocosta* (JACOBSON, 1924)***Chrysomela octocosta* JACOBSON, 1924

Distribution: SE Kazakhstan (Dzhungar Alatau).

***Chrysolina (Arctolina) oirota* LOPATIN, 1990**

Distribution: Asian Russia (Altai).

***Chrysolina (Arctolina) saurica* (JACOBSON, 1924)***Chrysomela saurica* JACOBSON, 1924

Distribution: E Kazakhstan.

***Chrysolina (Arctolina) septentrionalis* (MÉNÉTRIÉS 1851)***Chrysomela septentrionalis* MÉNÉTRIÉS, 1851*Chrysomela kuznetzowi* JACOBSON, 1894*Chrysomela sculpturata* JACOBSON, 1895*Chrysomela tundralis* JACOBSON, 1895

Distribution: N part of European Russia, Asian Russia (Krasnoyarsk Prov., Transbaikalia).

***Chrysolina (Arctolina) subsulcata* (MANNERHEIM, 1853)***Chrysomela subsulcata* MANNERHEIM, 1853*Chrysomela birulai* JACOBSON, 1910*Chrysomela glacialis* JACOBSON, 1910

Distribution: Arctic Asia (from Taimyr up to Chukotka, Wrangel Is., New Siberian Is.), Alaska.

***Chrysolina (Arctolina) tastavica* LOPATIN, 1992**

Distribution: SE Kazakhstan (Tastau Mountain Range).

***Chrysolina (Arctolina) teleutica* (JACOBSON, 1924)***Chrysomela teleutica* JACOBSON, 1924

Distribution: Asian Russia (Altai).

***Chrysolina (Arctolina) valichanovi* LOPATIN, 1990**

Distribution: SE Kazakhstan (Dzhungar Alatau).

***Chrysolina (Arctolina) wolosowiczi* (JACOBSON, 1910)***Chrysomela wolosowiczi* JACOBSON, 1910*Chrysomela novosibirica* JACOBSON, 1910

Distribution: NE Asian Russia (Novosiberian Is.).

**Subgenus: *Chrysolina (Sibiriella)* L. MEDVEDEV, 1999*****Chrysolina (Sibiriella) paradoxa* L. MEDVEDEV, 1999**

Distribution: Asian Russia (? Altai, ? Sayan).

**Subgenus: *Chrysolina (Altailina)* MIKHAILOV, 2000*****Chrysolina (Altailina) capricornis* MIKHAILOV, 2000**

Distribution: Asian Russia (W Sayan).

***Chrysolina (Altailina) dudkoi dudkoi* MIKHAILOV, 2000**

Distribution: Asian Russia (W Sayan).

***Chrysolina (Altailina) dudkoi ivanovskiana* MIKHAILOV, 2000**

Distribution: Asian Russia (W Sayan).

***Chrysolina (Altailina) kholsunica* MIKHAILOV, 2000***Chrysolina ogloblini* MIKHAILOV, 2000

Distribution: Asian Russia (W Sayan).

**Subgenus: *Chrysolina (Pezocrosita)* JACOBSON, 1901*****Chrysolina (Pezocrosita) almaatica* (LOPATIN, 1962)***Chrysomela almaatica* LOPATIN, 1962

Distribution: S Kazakhstan (Transili Alatau).

***Chrysolina (Pezocrosita) brevilata* (HEYDEN, 1886)***Chrysomela brevilata* HEYDEN, 1886*Chrysomela cyrtanastes* J. WEISE, 1892

Distribution: Mountains of Kirghizstan and S Kazakhstan.

***Chrysolina (Pezocrosita) brunnicornis brunnicornis* (J. WEISE, 1887)***Chrysomela brunnicornis* J. WEISE, 1887*Chrysomela kozhantschikovi* JACOBSON, 1925

Distribution: Asian Russia (Tuva), W Mongolia.

***Chrysolina (Pezocrosita) brunnicornis bermani* (L. MEDVEDEV, 1978)***Chrysomela brunnicornis bermani* L. MEDVEDEV, 1978

Distribution: Asian Russia (Transbaikalia, Yakutia).

***Chrysolina (Pezocrosita) brunnicornis vrangeliana* VORONOVA, 1982**

Distribution: NE Asian Russia (Wrangel Is.).

***Chrysolina (Pezocrosita) convexicollis* (JACOBSON, 1901)***Chrysomela convexicollis* JACOBSON, 1901

Distribution: Asian Russia (Tuva), Mongolia.

***Chrysolina (Pezocrosita) discriminata* (JACOBSON, 1901)***Chrysomela discriminata* JACOBSON, 1901*Chrysomela fuscipes* J. WEISE, 1890*Chrysomela infuscipes* J. WEISE, 1916

Distribution: Asian Russia (E Siberia), Mongolia.

***Chrysolina (Pezocrosita) erzinica* MIKHAILOV, 2002**

Distribution: Asian Russia (Tuva).

***Chrysolina (Pezocrosita) glebi* LOPATIN, 1988**

Distribution: Kirghizstan (Pskem Mountain Range).

***Chrysolina (Pezocrosita) helenae* (LOPATIN, 1968)***Chrysomela helenae* LOPATIN, 1968

Distribution: Kirghizstan (Talas and Ugham Mountain Ranges).

***Chrysolina (Pezocrosita) hyperboreica* MIKHAILOV, 2002**

Distribution: Arctic Asia (Polar Urals).

***Chrysolina (Pezocrosita) juldusana* (LOPATIN, 1962)***Chrysomela juldusana* LOPATIN, 1962

Distribution: SE Kazakhstan, Kirghizstan (Kunghey and Terskey Alatau), NW China.

***Chrysolina (Pezocrosita) ketmenica* (LOPATIN, 1970)***Chrysomela ketmenica* LOPATIN, 1970

Distribution: E Kazakhstan (Ketmen Mountain Range).

***Chrysolina (Pezocrosita) kiritschenkoi* (LOPATIN, 1970)***Chrysomela kiritschenkoi* LOPATIN, 1970

Distribution: Kirghizstan (Kirghiz Alatau)

***Chrysolina (Pezocrosita) koenigi* (JACOBSON, 1895)***Chrysomela koenigi* JACOBSON, 1895*Chrysomela katarinae* LOPATIN, 1965

Distribution: Mountains of Tajikistan and Kirghizstan.

***Chrysolina (Pezocrosita) lehri* (LOPATIN, 1970)***Chrysomela lehri* LOPATIN, 1970

Distribution: SE Kazakhstan.

***Chrysolina (Pezocrosita) lopatini* (MOHR, 1966)***Chrysomela lopatini* MOHR, 1966

Distribution: Asian Russia (Cisbaikalia, Transbaikalia), N Mongolia.

***Chrysolina (Pezocrosita) medvedevi* (LOPATIN, 1970)***Chrysomela medvedevi* LOPATIN, 1970

Distribution: E Kazakhstan.

***Chrysolina (Pezocrosita) mohri* (LOPATIN, 1970)***Chrysomela mohri* LOPATIN, 1970

Distribution: SE Kazakhstan.

***Chrysolina (Pezocrosita) mongolensis* (LOPATIN, 1966)***Chrysomela mongolensis* LOPATIN, 1966

Distribution: Mongolia.

***Chrysolina (Pezocrosita) obovata* (JACOBSON, 1895)***Chrysomela obovata* JACOBSON, 1895

Distribution: Kirghizstan (Central Tian-Shan).

***Chrysolina (Pezocrosita) ordinata* (GEBLER, 1823)***Chrysomela ordinata* GEBLER, 1823

Distribution: E Kazakhstan, Asian Russia (SW Siberia, Altais).

***Chrysolina (Pezocrosita) oshanini* (LOPATIN, 1965)***Chrysomela oshanini* LOPATIN, 1965

Distribution: Kirghizstan and Kazakhstan (from Turkestan Mountain Range up to Transili Alatau Mountain Range).

***Chrysolina (Pezocrosita) petrenkoi* LOPATIN, 1992**

Distribution: SE Kazakhstan (Dzhungar Alatau).

***Chrysolina (Pezocrosita) pusa* (LOPATIN, 1962)***Chrysomela pusa* LOPATIN, 1962*Chrysomela pusa purkynei* MOHR, 1966

Distribution: Asian Russia (Buryatia), Mongolia.

***Chrysolina (Pezocrosita) rufilabris* (FALDERMANN, 1835)***Chrysomela rufilabris* FALDERMANN, 1835

Distribution: Asian Russia (E Siberia), Mongolia.

***Chrysolina (Pezocrosita) sahlbergiana* (JACOBSON, 1901)***Chrysomela sahlbergiana* JACOBSON, 1901*Chrysomela kuznetzovi* JACOBSON, 1902, non JACOBSON, 1896*Chrysomela jacobsoni* LOPATIN, 1970

Distribution: Asian Russia (Sayan), N Mongolia.

***Chrysolina (Pezocrosita) sajanica* (JACOBSON, 1925)***Chrysomela sajanica* JACOBSON, 1925

Distribution: Asian Russia (Tuva).

***Chrysolina (Pezocrosita) sarcandica* LOPATIN, 1990**

Distribution: SE Kazakhstan.

***Chrysolina (Pezocrosita) sogdiorum* (J. WEISE, 1892)***Chrysomela sogdiorum* J. WEISE, 1892

Distribution: Kirghizstan (Chatkal, Talas and Kirghiz Mountain Ranges).

***Chrysolina (Pezocrosita) tibialis* (JACOBSON, 1895)***Chrysomela tibialis* JACOBSON, 1895

Distribution: Asian Russia (Sayan, Tuva).

***Chrysolina (Pezocrosita) tschatkalica* (LOPATIN, 1970)***Chrysomela tschatkalica* LOPATIN, 1970

Distribution: Kirghizstan (Chatkal Mountain Range).

***Chrysolina (Pezocrosita) tuvensis* L. MEDVEDEV, 1976**

Distribution: Asian Russia (Tuva).

***Chrysolina (Pezocrosita) ulugkhemica* MIKHAILOV, 2002**

Distribution: Asian Russia (Tuva).

***Chrysolina (Pezocrosita) undulata undulata* (GEBLER, 1834)***Chrysomela undulata* GEBLER, 1834*Chrysomela baicalica* MOHR, 1966

Distribution: S Siberia.

***Chrysolina (Pezocrosita) undulata asperata* LOPATIN, 1990**

Distribution: E Kazakhstan.

***Chrysolina (Pezocrosita) urjanchaica* (JACOBSON, 1925)***Chrysomela urjanchaica* JACOBSON, 1925

Distribution: Asian Russia (Tuva).

**Subgenus: *Chrysolina (Stichoptera)* MOTSCHULSKY, 1860*****Chrysolina (Stichoptera) gypsophilae* (KÜSTER, 1845)***Chrysomela gypsophilae* KÜSTER, 1845*Chrysomela lucidicollis* KÜSTER, 1845*Chrysomela rufomarginata* SUFFRIAN, 1851*Chrysomela gypsophilae* ab. *gaubili* H. LUCAS, 1849

Distribution: Europe except for the extreme North, Caucasus, Turkey, NW and Central Kazakhstan.

***Chrysolina (Stichoptera) kuesteri* (HELLIESEN, 1912)***Chrysomela küsteri* HELLIESEN, 1912

Distribution: S and Central Europe.

***Chrysolina (Stichoptera) pavlenkoi* (JACOBSON, 1924)***Chrysomela pavlenkoi* JACOBSON, 1924

Distribution: Asian Russia (Maritime Prov.).

***Chrysolina (Stichoptera) sanguinolenta* (LINNAEUS, 1758)***Chrysomela sanguinolenta* LINNAEUS, 1758*Chrysomela marginalis* DUFTSCHMIDT, 1825*Chrysomela porosa* GEBLER, 1830*Chrysolina sanguinolenta* ab. *richteri* ROUBAL, 1934*Chrysomela latecincta breiti* FRANZ, 1938

Distribution: S, Central, and E Europe, Kazakhstan (Steppe zone), Asian Russia (S Siberia, Russian Far East), Mongolia.

**Subgenus: *Chrysolina (Craspeda)* MOTSCHULSKY, 1860*****Chrysolina (Craspeda) jennisjensis* (BREIT, 1920)***Chrysomela jennisjensis* BREIT, 1920

Distribution: European Russia (Tambov Distr.), Caucasus, Kazakhstan, Asian Russia (S Siberia, Yakutia), Mongolia.

***Chrysolina (Craspeda) limbata limbata* (FABRICIUS, 1775)***Chrysomela limbata* FABRICIUS, 1775*Chrysomela besseri* KRYNICKI, 1832

*Chrysomela limbata* ab. *limbifera* KÜSTER, 1848

Distribution: Europe except for the North, Caucasus, Kazakhstan, Asian Russia (S Siberia east to Baikal).

***Chrysolina (Craspeda) limbata discipennis* (FALDERMANN, 1835)**

*Chrysomela discipennis* FALDERMANN, 1835

Distribution: SE Europe, Kazakhstan, Asian Russia (S Siberia, Yakutia), Mongolia.

**Subgenus: *Chrysolina (Taeniossticha)* MOTSCHULSKY, 1860**

***Chrysolina (Taeniossticha) alatavica* (JACOBSON, 1910)**

*Chrysomela alatavica* JACOBSON, 1910

*Chrysomela dzhungarica* JACOBSON, 1910

Distribution: SE Kazakhstan (Dzhungar Alatau, Tarbagatay Mountain Range), NW China (Borokhoro Mountain Range).

***Chrysolina (Taeniossticha) changaica* (LOPATIN, 1968)**

*Chrysomela changaica* LOPATIN, 1968

Distribution: Mongolia.

***Chrysolina (Taeniossticha) imperfecta imperfecta* BREIT, 1920**

*Chrysomela kaltenbachi* JAKOB, 1954

Distribution: S Turkmenistan (Kopetdagh Mountain Range), Iran, N Afghanistan.

***Chrysolina (Taeniossticha) koktumensis* LOPATIN et KULENOVA, 1987**

Distribution: SE Kazakhstan (Dzhungar Alatau).

***Chrysolina (Taeniossticha) kuldzhensis* LOPATIN, 1976**

Distribution: E Tian-Shan (Julduz and Kuldzha Mountain Ranges).

***Chrysolina (Taeniossticha) reitteri reitteri* (J. WEISE, 1884)**

*Chrysomela lurida* LINNAEUS, 1767, non SCOPOLI, 1763

*Chrysomela reitteri* J. WEISE, 1884

Distribution: Steppe of European Russia and Ukraine, Caucasus, Kazakhstan, Asian Russia (Sayan).

***Chrysolina (Taeniossticha) reitteri bakuensis* BECHYNÉ, 1952**

Distribution: Azerbaijan.

***Chrysolina (Taeniossticha) reitteri jailensis* BECHYNÉ, 1952**

Distribution: S Crimea.

***Chrysolina (Taeniossticha) reitteri pseudolurida* (ROUBAL, 1936)**

*Chrysomela pseudolurida* ROUBAL, 1936

Distribution: W Caucasus, Turkey.

***Chrysolina (Taeniossticha) reitteri saxonica* SILFVERBERG, 1977**

Distribution: Central and western parts of E Europe.

***Chrysolina (Taeniossticha) tianshanica* (JACOBSON, 1910)**

*Chrysomela tianshanica* JACOBSON, 1910

Distribution: SE Kazakhstan, Kirghizstan (Tian-Shan).



**Subgenus: *Chrysolina (Ovosoma)* MOTSCHULSKY, 1860**

***Chrysolina (Ovosoma) susterai* BECHYNÉ, 1950**

*Chrysomela morio* KRYNICKI, 1832, non FABRICIUS, 1792

Distribution: SE and southern part of E Europe, Kazakhstan, Asian Russia (Altai).

***Chrysolina (Ovosoma) sahlbergi* (MÉNÉTRIÉS, 1832)**

*Chrysomela sahlbergi* MÉNÉTRIÉS, 1832

*Chrysomela cupreopunctata* REICHE, 1858

*Chrysolina halysa* BECHYNÉ, 1950

Distribution: Caucasus, Turkey, Syria, Israel.

**Subgenus: *Chrysolina (Chrysomorpha)* MOTSCHULSKY, 1860**

***Chrysolina (Chrysomorpha) cerealis cerealis* (LINNAEUS, 1767)**

*Chrysomela cerealis* LINNAEUS, 1767

*Chrysomela violacea* SCHALLER, 1783

*Chrysomela octovittata* SCHRANK, 1789

*Chrysomela alternans* PANZER, 1799

*Chrysomela megerlei* FABRICIUS, 1801

*Chrysomela ornata* AHRENS, 1812

*Chrysomela mixta* KÜSTER, 1844

*Chrysomela fulgens* DUFUR, 1851

*Chrysomela livonica* MOTSCHULSKY, 1860

*Chrysomela kiesenwetteri* MOTSCHULSKY, 1860

*Chrysomela costata* DUFUR, 1902

*Chrysomela cerealis plorans* BECHYNÉ, 1948

Distribution: Europe, Kazakhstan.

***Chrysolina (Chrysomorpha) cerealis cyaneoaurata* (MOTSCHULSKY, 1860)**

*Chrysomela cyaneoaurata* MOTSCHULSKY, 1860

Distribution: Siberia, Mongolia.

***Chrysolina (Chrysomorpha) cerealis rufolineata* (MOTSCHULSKY, 1860)**

*Chrysomela rufolineata* MOTSCHULSKY, 1860

Distribution: N Caucasus, Kazakhstan.

**Subgenus: *Chrysolina (Colaphosoma)* MOTSCHULSKY, 1860**

***Chrysolina (Colaphosoma) sturmi* (WESTHOFF, 1882)**

*Chrysomela goettingensis* LINNAEUS, 1761, non LINNAEUS, 1758

*Chrysomela nigrita* TOWNSON, 1797, non FABRICIUS, 1792

*Chrysomela goettingensis* var. *sturmi* WESTHOFF, 1882

*Chrysomela goettingensis* var. *polonica* J. WEISE, 1883

*Chrysomela diversipes* BEDEL, 1892

*Chrysomela violacea* J. WEISE, 1916

Distribution: Europe, Caucasus, Kazakhstan, Siberia.

**Subgenus: *Chrysolina (Colaphoptera)* MOTSCHULSKY, 1860**

***Chrysolina (Colaphoptera) abchastica* (J. WEISE, 1892)**

*Chrysomela abchastica* J. WEISE, 1892

Distribution: W Caucasus.

***Chrysolina (Colaphoptera) adzharica* LOPATIN, 1988**

Distribution: SW Georgia, N Turkey.

***Chrysolina (Colaphoptera) apsilaeana* SILFVERBERG, 1977***Chrysolina fuscicornis* J. WEISE, 1902, non LINNAEUS, 1763*Chrysolina rosti kubanensis* L. MEDVEDEV et OCHRIMENKO, 1991

Distribution: Caucasus.

***Chrysolina (Colaphoptera) caspica* (J. WEISE, 1892)***Chrysolina caspica* J. WEISE, 1892

Distribution: Caucasus.

***Chrysolina (Colaphoptera) marcasitica marcasitica* (GERMAR, 1824)***Chrysolina marcasitica* GERMAR, 1824*Chrysolina distincta* KÜSTER, 1844*Chrysolina marcasitica* ab. *pannonica* J. WEISE, 1882*Chrysolina marcasitica* ab. *cupreopurpurea* GERHARDT, 1909*Chrysolina marcasitica* ab. *gerhardti* J. WEISE, 1916

Distribution: Mountains of Central Europe, W Ukraine (Carpathians).

***Chrysolina (Colaphoptera) marcasitica turgida* (J. WEISE, 1882)***Chrysolina marcasitica* var. *turgida* J. WEISE, 1882

Distribution: Sudety and Carpathians.

***Chrysolina (Colaphoptera) pliginskii* (REITTER, 1913)***Chrysolina pliginskii* REITTER, 1913*Chrysolina taurica* BREIT, 1920

Distribution: Crimea.

***Chrysolina (Colaphoptera) porphyrea* (FALDERMANN, 1837)***Chrysolina porphyrea* FALDERMANN, 1837*Chrysolina differens* FRANZ, 1952*Chrysolina diga* BECHYNÉ, 1952*Chrysolina erivanicola* BECHYNÉ, 1952*Chrysolina kubanica* BECHYNÉ, 1952*Chrysolina minutior* BECHYNÉ, 1952

Distribution: Caucasus.

***Chrysolina (Colaphoptera) purpurascens* (GERMAR, 1822)***Chrysolina purpurascens* GERMAR, 1822*Chrysolina dahli* GRAVENHORST, 1836 [nomen nudum]*Chrysolina dahli* MATZEK, 1843*Chrysolina purpurascens* var. *avia* J. WEISE, 1887*Chrysolina purpurascens* ab. *carpathica* PAPP, 1946

Distribution: Central Europe, W Ukraine (Carpathians).

***Chrysolina (Colaphoptera) rosti* (J. WEISE, 1892)***Chrysolina rosti* J. WEISE, 1892

Distribution: W Caucasus.

***Chrysolina (Colaphoptera) rufa rufa* (DUFTSCHMIDT, 1825)***Chrysolina rufa* DUFTSCHMIDT, 1825*Chrysolina Menthae* DUFTSCHMIDT, 1825

- Chrysomela metallica* KÜSTER, 1847  
*Chrysomela Dahli* SUFFRIAN, 1851, non HERRICH-SCHÄFER, 1838  
*Chrysomela opulenta* SUFFRIAN, 1851  
*Chrysomela squalida* SUFFRIAN, 1851  
*Chrysomela rufa* var. *sulcicollis* GERHARDT, 1911  
*Chrysomela rufa diminuta* BECHYNÉ, 1950  
*Chrysomela rufa staphylaeoides* BECHYNÉ, 1950  
 Distribution: Mountains of Central Europe, Carpathians.

***Chrysolina (Colaphoptera) rufa lapidaria* BECHYNÉ, 1950**  
 Distribution: Central Europe, W Moldavia.

- Chrysolina (Colaphoptera) trapezicollis* BECHYNÉ, 1952**  
*Chrysolina exul* BECHYNÉ, 1952  
*Chrysolina kutaisa* BECHYNÉ, 1952  
 Distribution: Caucasus.

- Chrysolina (Colaphoptera) umbratilis* (J. WEISE, 1887)**  
*Chrysomela atra* MATZEK, 1843, non GOEZE, 1777  
*Chrysomela olivacea* SUFFRIAN, 1857, non SCHALLER, 1783  
*Chrysomela atra* var. *umbratilis* J. WEISE, 1887  
*Chrysomela umbratilis* ab. *olivicolor* J. WEISE, 1916  
*Chrysolina umbratilis* ab. *olivaceocuprea* BECHYNÉ, 1950  
 Distribution: Mountains of Central Europe, Carpathians.

**Subgenus: *Chrysolina (Ovostoma)* MOTSCHULSKY, 1860**

- Chrysolina (Ovostoma) globipennis* (SUFFRIAN, 1851)**  
*Chrysomela globipennis* SUFFRIAN, 1851  
*Chrysomela globipennis* ab. *marmarosensis* REITTER, 1912  
*Chrysolina globipennis* ab. *styriaca* BECHYNÉ, 1946  
*Chrysolina globipennis deubeli* BECHYNÉ, 1948  
*Chrysolina globipennis euminuta* BECHYNÉ, 1950  
 Distribution: Central and S Europe, Moldavia, Ukraine.

- Chrysolina (Ovostoma) olivieri* (BEDEL, 1892)**  
*Chrysomela coerulea* OLIVIER, 1807, non OLIVIER, 1790  
*Chrysomela Olivieri* BEDEL, 1892  
 Distribution: Mountains of Central and S Europe, Ukraine.

**Subgenus: *Chrysolina (Euchrysolina)* BECHYNÉ, 1950**

- Chrysolina (Euchrysolina) graminis graminis* (LINNAEUS, 1758)**  
*Chrysomela graminis* LINNAEUS, 1758  
*Chrysomela fulgida* FABRICIUS, 1801  
 Distribution: Europe except for the extreme North, Kazakhstan, Uzbekistan, Asian Russia (S Siberia, Tuva, Altai), Mongolia.
- Chrysolina (Euchrysolina) graminis artemisiae* MOTSCHULSKY, 1860**  
 Distribution: SE Europe, Kazakhstan, Central Asia, S Siberia
- Chrysolina (Euchrysolina) graminis auraria* MOTSCHULSKY, 1860**  
 Distribution: Russian Far East, Japan.

***Chrysolina (Euchrysolina) virgata* MOTSCHULSKY, 1860***Chrysomela eximia* BALY, 1862*Chrysomela obscurolfasciata* JACOBY, 1885

Distribution: Russian Far East, Japan.

**Subgenus: *Chrysolina (Erythrochysa)* BECHYNÉ, 1950*****Chrysolina (Erythrochysa) polita* (LINNAEUS, 1758)***Chrysomela polita* LINNAEUS, 1758*Chrysomela adamsi* BALY, 1879*Chrysomela polita* ab. *epipleuralis* JACOBSON, 1895*Chrysomela polita* ab. *kafkana* REITTER, 1898*Chrysomela polita* ab. *vanderwieli* EVERTS, 1921*Chrysomela polita* ab. *sirgueyi* PORTA, 1934

Distribution: All over Palaearctic.

**Subgenus: *Chrysolina (Menthastrilla)* BECHYNÉ, 1950*****Chrysolina (Menthastrilla) herbacea herbacea* (DUFTSCHMIDT, 1825)***Chrysomela herbacea* DUFTSCHMIDT, 1825*Chrysomela menthastrilla* SUFFRIAN, 1851*Chrysomela fulminans* SUFFRIAN, 1851*Chrysomela resplendes* SUFFRIAN, 1855*Chrysomela herbacea* ab. *cupreopurpurea* COSTA, 1838*Chrysomela herbacea* ab. *rugicollis* WEIDENBACH, 1859*Chrysomela herbacea* ab. *croatica* J. WEISE, 1884*Chrysomela herbacea* ab. *tolosana* BUYSSON, 1924*Chrysomela herbacea* ab. *elevarensis* BUYSSON, 1924*Chrysolina herbacea* ab. *cyaneonigra* JOLIVET: BECHYNÉ, 1950*Chrysolina herbacea* ab. *durabilis* BECHYNÉ, 1950*Chrysolina herbacea* ab. *expectata* BECHYNÉ, 1950*Chrysolina herbacea* ab. *solistima* BECHYNÉ, 1950*Chrysolina herbacea* ab. *thoracica* BECHYNÉ, 1950

Distribution: Europe except for the extreme North, Caucasus, Turkey, Transcaucasia, Middle East, W Kazakhstan, W Turkmenistan.

***Chrysolina (Menthastrilla) coeruleans coeruleans* (L. G. SCRIBA, 1791)***Chrysomela coeruleans* L. G. SCRIBA, 1791*Chrysomela violacea* PANZER, 1797*Chrysomela subfastuosa* MOTSCHULSKY, 1860

Distribution: Mountains of Central Europe, Karelia, Belarus.

***Chrysolina (Menthastrilla) coeruleans relictata* L. MEDVEDEV, 1977**

Distribution: Urals.

***Chrysolina (Menthastrilla) coeruleans splendorifera* MOTSCHULSKY, 1860**

Distribution: Caucasus, Iran, Iraq.

***Chrysolina (Menthastrilla) coeruleans uzbekorum* BECHYNÉ, 1950**

Distribution: Uzbekistan, Tajikistan.

**Subgenus: *Chrysolina (Colaphodes)* MOTSCHULSKY, 1860**

***Chrysolina (Colaphodes) haemoptera haemoptera* (LINNAEUS, 1758)**

- Chrysomela haemoptera* LINNAEUS, 1758  
*Chrysomela atra* GOEZE, 1777  
*Chrysomela gottengensis* SCHRANK, 1789, non LINNAEUS, 1758  
*Chrysomela hottentota* FABRICIUS, 1792  
*Chrysomela unicolor* SUFFRIAN, 1851, non GEBLER, 1845  
*Chrysomela acuticollis* FAIRMAIRE, 1877  
*Chrysomela camerani* PIOLTI, 1880

Distribution: Europe except for the extreme North, Turkey.

**Subgenus: *Chrysolina (Hypericia)* BEDEL, 1892**

***Chrysolina (Hypericia) cuprina cuprina* (DUFTSCHMIDT, 1825)**

- Chrysomela cuprina* DUFTSCHMIDT, 1825  
*Chrysomela cuprina* ab. *indigena* J. WEISE, 1884  
*Chrysomela cuprina* ab. *perviridis* REITTER, 1912  
*Chrysomela cuprina* ab. *nigra* REITTER, 1912  
*Chrysomela cuprina* ab. *bechynei* ŠPRINGLOVÁ, 1945  
*Chrysomela cuprina* ab. *hyperycicola* ŠPRINGLOVÁ, 1945  
*Chrysomela cuprina* ab. *vulgaris* ŠPRINGLOVÁ, 1945  
*Chrysomela cuprina* ab. *semiviridis* ŠPRINGLOVÁ, 1945  
*Chrysomela cuprina* ab. *semicoerulea* ŠPRINGLOVÁ, 1945  
*Chrysomela cuprina* ab. *semicuprea* ŠPRINGLOVÁ, 1945  
*Chrysomela cuprina* ab. *aeneola* ŠPRINGLOVÁ, 1945  
*Chrysolina cuprina* ab. *nigritula* BECHYNÉ, 1949  
*Chrysomela cuprina* ab. *lugubrina* CSIKI, 1953

Distribution: Europe except for the extreme North. Introduced to USA and Australia.

***Chrysolina (Hypericia) cuprina dilecta* BECHYNÉ, 1952**

Distribution: Asian Russia (Altai, Sayan).

***Chrysolina (Hypericia) didymata* (L. G. SCRIBA, 1791)**

- Chrysomela didymata* L. G. SCRIBA, 1791  
*Chrysolina didymata* ab. *euviridis* BECHYNÉ, 1952

Distribution: Europe, Turkey.

***Chrysolina (Hypericia) difficilis difficilis* (MOTSCHULSKY, 1869)**

- Chrysomela difficilis* MOTSCHULSKY, 1869

Distribution: Asian Russia (Altai, Sayan).

***Chrysolina (Hypericia) difficilis ussuriensis* (JACOBSON, 1901)**

- Chrysomela ussuriensis* JACOBSON, 1901

Distribution: Asian Russia (Amurland, Maritime Prov.), NE China.

***Chrysolina (Hypericia) difficilis yezoensis* (MATSUMURA, 1911)**

- Chrysomela yezoensis* MATSUMURA, 1911

Distribution: Asian Russia (Sakhalin), Korea, Japan.

***Chrysolina (Hypericia) geminata* (PAYKULL, 1799)**

- Chrysomela geminata* PAYKULL, 1799  
*Chrysomela approximata* ZENKER, 1815

*Chrysomela quadrigemina* SUFFRIAN, 1851  
*Chrysomela bifoveolata* BRANCSIK, 1910  
*Chrysolina geminata* ab. *utens* BECHYNÉ, 1949  
 Distribution: S, Central, and E Europe.

***Chrysolina (Hypericia) hyperici hyperici* (FORSTER, 1771)**

*Chrysomela hyperici* FORSTER, 1771  
*Chrysomela fucata* FABRICIUS, 1781  
*Chrysomela gemellata* FOURCROY, 1785, non ROSSI, 1792  
*Chrysomela aerea* GMELIN, 1790  
*Chrysomela hyperici* var. *ambigua* J. WEISE, 1884  
*Chrysomela hyperici* var. *privigna* J. WEISE, 1884  
 Distribution: Europe except for the extreme North, Kazakhstan, Uzbekistan.

***Chrysolina (Hypericia) hyperici daghestanica* (REITTER, 1912)**

*Chrysomela daghestanica* REITTER, 1912  
 Distribution: E Caucasus (Daghestan).

***Chrysolina (Hypericia) seriepunctata* (J. WEISE, 1887)**

*Chrysomela seriepunctata* J. WEISE, 1887  
 Distribution: Asian Russia (Amurland, Maritime Prov.).

**Subgenus: *Chrysolina (Allohypericia)* BECHYNÉ, 1950**

***Chrysolina (Allohypericia) aeruginosa aeruginosa* (FALDERMANN, 1835)**

*Chrysomela aeruginosa* FALDERMANN, 1835  
*Chrysomela distans* CSIKI, 1961  
*Chrysomela dimidiata* MÉNÉTRIÉS, 1836  
*Chrysomela instructa* MOTSCHULSKY, 1860  
*Chrysomela tarda* MOTSCHULSKY, 1860  
*Chrysomela pseudogeminata* BECHYNÉ, 1950  
 Distribution: Kazakhstan, Siberia, Mongolia.

***Chrysolina (Allohypericia) aeruginosa sibirica* (J. WEISE, 1887)**

*Chrysomela sibirica* J. WEISE, 1887  
 Distribution: Russian Far East.

***Chrysolina (Allohypericia) arctica* L. MEDVEDEV, 1980**

Distribution: Asian Russia (Wrangel Is.).

***Chrysolina (Allohypericia) koltzei koltzei* (J. WEISE, 1887)**

*Chrysomela koltzei* J. WEISE, 1887  
*Chrysomela daurica* HEYDEN, 1885, non GEBLER, 1832  
 Distribution: Asian Russia (S Siberia, Amurland).

***Chrysolina (Allohypericia) koltzei brunneipennis* (MATSUMURA, 1911)**

*Chrysomela brunneipennis* MATSUMURA, 1911  
 Distribution: Asian Russia (Sakhalin), Japan.

***Chrysolina (Allohypericia) perforata perforata* (GEBLER, 1830)**

*Chrysomela perforata* GEBLER, 1830  
*Chrysomela turczaninoffi* HAROLD, 1875  
*Chrysomela variolosa* MOTSCHULSKY, 1854  
 Distribution: Asian Russia (S Siberia, Altai, Sayan), N and NW Mongolia.

***Chrysolina (Allohypericia) perforata pallidipes* L. MEDVEDEV, 1980**

Distribution: Asian Russia (S Tuva), Mongolia (Mongolian Altai).

***Chrysolina (Allohypericia) perforata simillima* MOHR, 1966**

Distribution: Asian Russia (Transbaikalia), N Mongolia.

***Chrysolina (Allohypericia) purpurata* FALDERMANN, 1833***Chrysomela perforata* J. WEISE, 1916

Distribution: E Kazakhstan, S Siberia.

**Subgenus: *Chrysolina (Anopachys)* MOTSCHULSKY, 1860*****Chrysolina (Anopachys) aurichalcea aurichalcea* (MANNERHEIM, 1825)***Chrysomela aurichalcea* MANNERHEIM, 1825*Chrysomela praticola* DUFTSCHMIDT, 1825*Chrysomela asclepiadis* A. VILLA et J. B. VILLA, 1833*Chrysomela gibbipennis* FALDERMANN, 1835*Chrysomela recticollis* J. WEISE, 1887

Distribution: Forest and Steppe zone of European Russia, Crimea, Azerbaijan, Kazakhstan, Kirghizstan, Asian Russia (S Siberia, Russian Far East), Mongolia, China, Korea, Japan.

***Chrysolina (Anopachys) aurichalcea bohémica* G. MÜLLER, 1949***Chrysomela problematica* KASZAB, 1962

Distribution: Central Europe, Ukraine.

***Chrysolina (Anopachys) lineella* (J. WEISE, 1887)***Chrysomela lineella* J. WEISE, 1887

Distribution: Russian Far East.

***Chrysolina (Anopachys) lineigera* (JACOBSON, 1901)***Chrysomela lineigera* JACOBSON, 1901*Chrysomela watanabei* TAKIZAWA, 1970

Distribution: Asian Russia (Far East, Sakhalin), NE China, N Japan.

***Chrysolina (Anopachys) neglecta* BIENKOVSKY, 1998**

Distribution: Asian Russia (Khabarovsk Prov., Maritime Prov.).

***Chrysolina (Anopachys) pala* BIENKOVSKY, 1998**

Distribution: Asian Russia (Maritime Prov.).

***Chrysolina (Anopachys) quadrangulata* (MOTSCHULSKY, 1860)***Chrysomela quadrangulata* MOTSCHULSKY, 1860*Chrysomela linaeides* J. WEISE, 1896*Chrysomela omoka* JACOBSON, 1924

Distribution: Asian Russia (E Siberia, Russian Far East), N and Central Mongolia.

***Chrysolina (Anopachys) relucens* (ROSENHAUER, 1847)***Chrysomela relucens* ROSENHAUER, 1847

Distribution: Central Europe (Alps), Coasts of White Sea, Asian Russia (Siberia, Russian Far East).

**Subgenus: *Chrysolina (Caudatochrysa)* BECHYNÉ, 1950*****Chrysolina (Caudatochrysa) aino* TAKIZAWA, 1970**

Distribution: Asian Russia (S Kurile Is.: Kunashir).

***Chrysolina (Caudatochrysa) angusticollis* MOTSCHULSKY, 1860***Chrysomela japana* BALY, 1874

Distribution: Asian Russia (Amurland, S Kurile Is.), N China, Japan.

***Chrysolina (Caudatochrysa) cyanopurpurea* (BALLION, 1878)***Chrysomela cyanopurpurea* BALLION, 1878

Distribution: NW China, possible for SE Kazakhstan.

**Subgenus: *Chrysolina (Chrysolina)* MOTSCHULSKY, 1860*****Chrysolina (Chrysolina) staphylaea staphylaea* (LINNAEUS, 1758)***Chrysomela staphylaea* LINNAEUS, 1758*Chrysomela subferruginea* SUFFRIAN, 1851*Chrysomela lederi* J. WEISE in SCHNEIDER et LEDER, 1879

Distribution: All over the Holarctic.

***Chrysolina (Chrysolina) staphylaea daurica* (GEBLER, 1832)***Chrysomela daurica* GEBLER, 1832

Distribution: Asian Russia (S Siberia, Maritime Prov.), Kazakhstan, Kirghizstan, N China.

**Subgenus: *Chrysolina (Minckia)* E. STRAND, 1935*****Chrysolina (Minckia) chalcites* (GERMAR, 1824)***Chrysomela chalcites* GERMAR, 1824*Chrysomela festiva* MÉNÉTRIÉS, 1832*Chrysomela chalcitis* SUFFRIAN, 1851

Distribution: Europe, Central and S European Russia, S Ukraine, Transcaucasia, Turkey, Near East, Turkmenistan, Tajikistan.

***Chrysolina (Minckia) oricalcia* (O. F. MÜLLER, 1776)***Chrysomela oricalcia* O. F. MÜLLER, 1776*Chrysomela bulgarensis* SCHRANK, 1781*Chrysomela orychalca* HERBST, 1783*Chrysomela olivacea* SCHALLER, 1783*Chrysomela orichalcea* GMELIN, 1790*Chrysomela lamina* MATZEK, 1843, non FABRICIUS, 1792*Chrysomela oricalcia* ab. *nigricollis* J. WEISE, 1916

Distribution: Europe.

**Subgenus: *Chrysolina (Vittatochrysa)* LOPATIN, 1977*****Chrysolina (Vittatochrysa) nigrovittata* (BALLION, 1878)***Chrysomela nigrovittata* BALLION, 1878

Distribution: SE Kazakhstan, Kirghizstan, NW China.



**Subgenus: *Chrysolina (Sphaeromela)* BEDEL, 1892*****Chrysolina (Sphaeromela) varians* (SCHALLER, 1783)**

- Chrysomela varians* SCHALLER, 1783  
*Chrysomela Centaura* HERBST, 1783  
*Chrysomela centaurii* FABRICIUS, 1787  
*Chrysomela aethiops* FABRICIUS, 1792  
*Chrysomela varians* var. *pratensis* J. WEISE, 1884  
*Chrysomela incerta* MEISSNER, 1907

Distribution: Europe, Caucasus, Asian Russia (W Siberia, Altai).

**Subgenus: *Chrysolina (Bechynea)* L. MEDVEDEV, 1970*****Chrysolina (Bechynea) nikolskii* (JACOBSON, 1899)**

- Chrysomela nikolskii* JACOBSON, 1899  
*Chrysomela kawakami* NAKAMURA, 1911  
*Chrysolina kabakovi* L. MEDVEDEV, 1966

Distribution: Asian Russia (Amurland, Sakhalin, S Kurile Is.), Japan.

***Chrysolina (Bechynea) sulcicollis adzhalamica* L. MEDVEDEV, 1970**

Distribution: Asian Russia (S Khabarovsk Prov.)

***Chrysolina (Bechynea) sulcicollis sutschanica* L. MEDVEDEV, 1970**

Distribution: Asian Russia (S Maritime Prov.).

**Subgenus: *Chrysolina (Paracrosita)* DACCORDI, 1982*****Chrysolina (Paracrosita) armeniaca* (FALDERMANN, 1837)**

- Chrysomela armeniaca* FALDERMANN, 1837

Distribution: Caucasus, S Turkmenistan.

**Subgenus: *Chrysolina (Chalcoidea)* MOTSCHULSKY, 1860*****Chrysolina (Chalcoidea) analis* (LINNAEUS, 1767)**

- Chrysomela analis* LINNAEUS, 1767  
*Chrysomela lomata* HERBST, 1783  
*Chrysomela schach* FABRICIUS, 1792  
*Chrysomela prasina* SUFFRIAN, 1851  
*Chrysomela analis* ab. *tristis* TENENBAUM, 1927  
*Chrysomela analis* ab. *brunneomicans* ROUBAL, 1942

Distribution: Europe except for the extreme North and South.

***Chrysolina (Chalcoidea) bodemeyeri* (J. WEISE, 1910)**

- Chrysomela bodemeyeri* J. WEISE, 1910

Distribution: Turkmenistan (Kopetdagh Mountain Range), Iran.

***Chrysolina (Chalcoidea) carnifex* (FABRICIUS, 1792)**

- Chrysomela carnifex* FABRICIUS, 1792  
*Chrysomela aethiops* OLIVIER, 1807  
*Chrysomela carnifex* ab. *curvilinea* J. WEISE, 1884  
*Chrysomela carnifex* ab. *konowi* J. WEISE, 1889  
*Chrysomela carnifex* ab. *metallica* SIRGUEY, 1929

*Chrysolina carnifex* ab. *amata* BECHYNÉ, 1949

*Chrysolina carnifex* ab. *parisiensis* BECHYNÉ, 1949

*Chrysolina carnifex* ab. *toletensis* BECHYNÉ, 1949

*Chrysolina carnifex* ab. *allofossulata* BECHYNÉ, 1950

*Chrysolina carnifex* ab. *isofossulata* BECHYNÉ, 1950

*Chrysolina carnifex* ab. *lugdunensis* BECHYNÉ, 1950

*Chrysolina carnifex* ab. *pseudocruentata* BECHYNÉ, 1950

*Chrysolina carnifex* ab. *pseudofossulata* BECHYNÉ, 1950

Distribution: Europe, Ukraine, S European Russia, N and E Kazakhstan.

***Chrysolina (Chalcoidea) cinctipennis* (HAROLD, 1874)**

*Chrysomela cinctipennis* HAROLD, 1874

*Chrysomela besseri* MARSEUL, 1887, non KRYNICKI, 1832

*Chrysomela circumducta* SUFFRIAN, 1851, non MÉNÉTRIÉS, 1885

Distribution: Hungary, Ukraine, European Russia (Steppe and Semi-desert Zone), Kazakhstan (except for S and E).

***Chrysolina (Chalcoidea) circumducta* (MÉNÉTRIÉS, 1835)**

*Chrysomela circumducta* MÉNÉTRIÉS, 1835

*Chrysomela sulcata* FISCHER-WALDHEIM, 1843

Distribution: Kirghizstan, Uzbekistan, Tajikistan, NW China.

***Chrysolina (Chalcoidea) elbursica* LOPATIN, 1981**

Distribution: S Turkmenistan (Kopetdagh Mountain Range), N Iran.

***Chrysolina (Chalcoidea) hyrcana* (J. WEISE, 1884)**

*Chrysomela hyrcana* J. WEISE, 1884

Distribution: Daghestan, Armenia, Azerbaijan, Kazakhstan (except for the South).

***Chrysolina (Chalcoidea) instabilis* MÄKLIN, 1877**

Distribution: Asian Russia (N of Middle and Northeastern Siberia: from Yenisei river in the west, up to Kolyma river in the east; Yakutia, Irkutsk Distr.).

***Chrysolina (Chalcoidea) levi* OCHRIMENKO, 1990**

Distribution: N Caucasus.

***Chrysolina (Chalcoidea) lia* (JACOBSON, 1895)**

*Chrysomela lia* JACOBSON, 1895

*Chrysolina freyi* BECHYNÉ, 1950

*Chrysomela haarlovi* JAKOB, 1962

Distribution: Turkmenistan (Kopetdagh Mountain Range), Uzbekistan, Tajikistan (included Badakhshan), S Kirghizstan (Alai and Transalai Mountain Ranges), Iran, Afghanistan.

***Chrysolina (Chalcoidea) marginata marginata* (LINNAEUS, 1758)**

*Chrysomela marginata* LINNAEUS, 1758

*Chrysomela cinctella* GYLLENHAL, 1827

Distribution: All over the Holarctic.

***Chrysolina (Chalcoidea) marginata roubali* BECHYNÉ, 1950**

*Chrysomela rugosopunctata* ROUBAL, 1917

Distribution: N Caucasus.

***Chrysolina (Chalcoidea) marginata songorica* (GEBLER, 1843)***Chrysomela songorica* GEBLER, 1843

Distribution: SE and E Kazakhstan (Dzhungar Alatau and Tarbagatay Mountain Ranges).

***Chrysolina (Chalcoidea) marginata unificans* BECHYNÉ, 1950**

Distribution: Transcaucasia, E Turkey.

***Chrysolina (Chalcoidea) tesari* (ROUBAL, 1936)***Chrysomela tesari* ROUBAL, 1936*Chrysolina kulzeri* BECHYNÉ, 1950

Distribution: Caucasus.

***Chrysolina (Chalcoidea) turgaica* (JACOBSON, 1910)***Chrysomela turgaica* JACOBSON, 1910

Distribution: W Kazakhstan (Mugodzhary Chain).

***Chrysolina (Chalcoidea) unicolor unicolor* (GEBLER, 1845)***Chrysomela unicolor* GEBLER, 1845*Chrysomela immarginata* RYBAKOW, 1884

Distribution: Kazakhstan and Kirghizstan (N and Central Tian-Shan).

***Chrysolina (Chalcoidea) unicolor alaiensis* LOPATIN, 1998**

Distribution: Kirghizstan (Alai Mountain Range).

***Chrysolina (Chalcoidea) zamotajlovi* L. MEDVEDEV et OCHRIMENKO, 1991**

Distribution: Caucasus.

**Subgenus: *Chrysolina (Diachalcoidea)* BECHYNÉ, 1955*****Chrysolina (Diachalcoidea) sacarum sacarum* (J. WEISE, 1890)***Chrysomela sacarum* J. WEISE, 1890*Chrysomela badakshanica* IABLOKOFF-KHNZORIAN, 1978

Distribution: S and SE Kazakhstan, Central Asia (except for the Turkmenistan' Deserts and Pamirs), Afghanistan.

***Chrysolina (Diachalcoidea) sacarum embiense* LOPATIN, 1996**

Distribution: W Kazakhstan.

**Subgenus: *Chrysolina (Fastuolina)* WARCHALOWSKI, 1991*****Chrysolina (Fastuolina) fastuosa* (SCOPOLI, 1763)***Chrysomela fastuosa* SCOPOLI, 1763*Chrysomela speciosa* LINNAEUS, 1767*Chrysomela inexplicabilis* BRANCSIK, 1910*Chrysomela fastuosa* ab. *umbrata* GABRIEL in W. KOLBE, 1931*Chrysomela fastuosa* ab. *galeopsidis* SCHRANK, 1798*Chrysomela fastuosa* ab. *cupreonitens* MARSEUL, 1886*Chrysolina fastuosa* ab. *callichloris* BECHYNÉ, 1950*Chrysolina fastuosa* ab. *revyi* KASZAB, 1962

Distribution: Europe, Asian Russia (W Siberia, Altai).

**Genus: *Oreina* CHEVROLAT in DEJEAN, 1837***Orina* AGASSIZ, 1846**Subgenus: *Oreina* (*Allorina*) J. WEISE, 1902*****Oreina* (*Allorina*) *bidentata* BONTEMS, 1981***Orina tristis* auct., non FABRICIUS, 1792

Distribution: Mountains of S Europe, Carpathians.

***Oreina* (*Allorina*) *coerulea coerulea* (OLIVIER, 1790)***Chrysomela coerulea* OLIVIER, 1790*Chrysomela rugulosa* SUFFRIAN, 1851 non GEBLER, 1841*Orina rugulosa* var. *gaertneri* J. WEISE, 1894*Orina rugulosa* var. *nigritula* J. WEISE, 1894

Distribution: Mountains of Central and SE Europe, Carpathians, W Belarus (Bialowieza Primeval Forest), S Ukraine from Carpathians in the west up to Dnepr river in the east, European Russia (except for N).

***Oreina* (*Allorina*) *coerulea onega* BECHYNÉ, 1958**

Distribution: Karelia.

***Oreina* (*Allorina*) *coerulea rareua* BECHYNÉ, 1958**

Distribution: Bukowina.

**Subgenus: *Oreina* (*Romalorina*) J. WEISE, 1906*****Oreina* (*Romalorina*) *alpestris* (SCHUMMEL, 1843)***Chrysomela alpestris* SCHUMMEL, 1843*Chrysomela speciosa* SUFFRIAN, 1851*Chrysomela punctatissima* SUFFRIAN, 1851*Chrysomela gloriosa* fa. *polymorpha* KRAATZ, 1880*Orina alpestris* var. *bicolora* J. WEISE, 1883*Orina alpestris* var. *fontinalis* J. WEISE, 1883*Orina alpestris* var. *moesta* J. WEISE, 1883*Orina alpestris* var. *olivacea* J. WEISE, 1883*Orina alpestris* var. *rivularis* J. WEISE, 1883*Orina alpestris* var. *bannatica* J. WEISE, 1884*Orina alpestris* var. *umbrosa* J. WEISE, 1884*Orina alpestris* var. *vinariensis* J. WEISE, 1902*Chrysochloa cacaliae* var. *lissahorensis* ab. *stejskali* FLEISCHER, 1914*Chrysochloa polymorpha* ab. *bohyslavi* FLEISCHER, 1916*Chrysochloa alpestris* var. *alpestris* ab. *alutacea* HÄNEL, 1937*Chrysochloa alpestris* var. *alpestris* ab. *coeruleoviridis* HÄNEL, 1937*Chrysochloa alpestris* var. *alpestris* ab. *liturata* HÄNEL, 1937*Chrysochloa alpestris* var. *alpestris* ab. *nigripemis* HÄNEL, 1937*Chrysochloa alpestris* var. *alpestris* ab. *viridicollis* HÄNEL, 1937*Chrysochloa alpestris* var. *polymorpha* ab. *bicolor* HÄNEL, 1937*Chrysochloa alpestris* var. *polymorpha* ab. *coeruleoviridis* HÄNEL, 1937*Chrysochloa alpestris* var. *polymorpha* ab. *cyanicollis* HÄNEL, 1937*Chrysochloa alpestris* var. *polymorpha* ab. *viridis* HÄNEL, 1937

Distribution: Mountains of S, Central and E Europe, from Pyrenees to Carpathians.

***Oreina (Romalorina) basilea* (GEBLER, 1823)***Chrysomela sulcata* GEBLER, 1823

Distribution: Asian Russia (Altai, Tuva, Sayan, Transbaikalia), Mongolia.

***Oreina (Romalorina) bifrons* (FABRICIUS, 1792)***Chrysomela bifrons* FABRICIUS, 1792

Distribution: Mountains of Central and S Europe, Ukrainian Carpathians.

***Oreina (Romalorina) redikortzevi* (JACOBSON, 1925)***Chrysochloa redikortzevi* JACOBSON, 1925

Distribution: Asian Russia (Sayan).

***Oreina (Romalorina) viridis* (DUFTSCHMIDT, 1825)***Chrysomela viridis* DUFTSCHMIDT, 1825*Chrysomela nivalis* HEER, 1834*Lina ignita* COMOLLI, 1837*Orina viridis* var. *lugubris* J. WEISE, 1883*Orina viridis* var. *merkli* J. WEISE, 1884*Orina viridis* var. *transsylvanica* J. WEISE, 1884*Chrysochloa viridis* var. *merkli* ab. *cuprina* HÄNEL, 1937*Chrysochloa viridis* var. *merkli* ab. *cyanicollis* HÄNEL, 1937*Chrysochloa viridis* var. *merkli* ab. *nigripennis* HÄNEL, 1937*Chrysochloa viridis* var. *merkli* ab. *nigripennis* HÄNEL, 1937*Chrysochloa viridis* var. *merkli* ab. *viridis* HÄNEL, 1937

Distribution: Mountains of Central and S Europe, Ukrainian Carpathians.

**Subgenus: *Oreina (Intricatorina)* KUHNELT, 1984*****Oreina (Intricatorina) intricata* (GERMAR, 1824)***Chrysomela intricata* GERMAR, 1824*Chrysomela aurulenta* SUFFRIAN, 1851*Orina intricata* var. *amethystina* J. WEISE, 1883*Orina intricata* var. *seminigra* J. WEISE, 1883*Chrysochloa intricata* var. *anderschi* ab. *cyanea* HÄNEL, 1937*Chrysochloa intricata* var. *intricata* ab. *coeruleoviridis* HÄNEL, 1937

Distribution: Mountains of Central and S Europe, Ukrainian Carpathians.

***Oreina (Virgulatorina) virgulata* (GERMAR, 1824)***Chrysomela virgulata* GERMAR, 1824*Chrysomela alcyonea* SUFFRIAN, 1851*Orina virgulata* var. *candens* J. WEISE, 1884*Orina virgulata* var. *praefica* J. WEISE, 1884*Orina virgulata* var. *serena* J. WEISE, 1884*Chrysochloa virgulata* var. *virgulata* ab. *aurea* HÄNEL, 1937*Chrysochloa virgulata* var. *virgulata* ab. *cyanea* HÄNEL, 1937

Distribution: Mountains of Central, S and SE Europe, Ukrainian Carpathians and Ciscarpathia.

**Subgenus: *Oreina (Oreina)* CHEVROLAT in DEJEAN, 1837*****Oreina (Oreina) cacaliae* (Schränk, 1785)***Chrysomela cacaliae* SCHRÄNK, 1785

- Chrysomela coeruleolineata* DUFTSCHMIDT, 1825  
*Chrysomela senecionis* SCHUMMEL, 1844  
*Chrysomela tussilaginis* SUFFRIAN, 1851  
*Orina cacaliae* var. *cyanipennis* J. WEISE, 1883  
*Orina cacaliae* var. *fraudulenta* J. WEISE, 1883  
*Orina cacaliae* var. *tristicula* J. WEISE, 1883  
*Orina cacaliae* var. *lissahorensis* J. WEISE, 1896  
*Chrysochloa cacaliae* var. *senecionis* ab. *coeruleoviridis* HÄNEL, 1937  
*Chrysochloa cacaliae* var. *senecionis* ab. *cyanipennis* HÄNEL, 1937  
*Chrysochloa cacaliae* var. *senecionis* ab. *fraudulenta* HÄNEL, 1937  
*Chrysochloa cacaliae* var. *senecionis* ab. *lissahorensis* HÄNEL, 1937  
*Chrysochloa cacaliae* var. *senecionis* ab. *violacea* HÄNEL, 1937  
*Chrysochloa cacaliae* var. *senecionis* ab. *senecionis* HÄNEL, 1937  
 Distribution: Mountains of Central and S Europe, Ukranian Carpathians.

***Oreina (Oreina) speciosissima* (SCOPOLI, 1763)**

- Coccinella speciosissima* SCOPOLI, 1763  
*Chrysomela tristis* OLIVIER, 1807  
*Chrysomela fuscoaenea* SCHUMMEL, 1844  
*Chrysomela juncorum* SUFFRIAN, 1851  
*Chrysomela speciosissima* var. *coerulescens* LETZNER, 1853  
*Chrysomela speciosissima* var. *ignita* LETZNER, 1853  
*Chrysomela speciosissima* var. *nigrescens* LETZNER, 1853  
*Chrysomela speciosissima* var. *violaceus* LETZNER, 1853  
*Chrysomela speciosissima* var. *viridescens* LETZNER, 1853  
*Chrysomela speciosissima* var. *viridiana* LETZNER, 1853  
*Chrysomela punctata* G. ZEBE, 1853 [nomen nudum]  
*Chrysomela speciosissima* var. *carpathica* KRAATZ, 1859  
*Orina speciosissima* var. *coerulescens* J. WEISE, 1883  
*Orina speciosissima* var. *letzneri* J. WEISE, 1883  
*Orina speciosissima* var. *schummeli* J. WEISE, 1883  
*Chrysochloa speciosissima* var. *juncorum* ab. *coeruleoviridis* HÄNEL, 1937  
*Chrysochloa speciosissima* var. *silesiaca* ab. *coeruleoviridis* HÄNEL, 1937  
*Chrysochloa speciosissima* var. *silesiaca* ab. *nigra* HÄNEL, 1937  
*Chrysochloa speciosissima* var. *speciosissima* ab. *coeruleoviridis* HÄNEL, 1937  
*Chrysochloa speciosissima* var. *speciosissima* ab. *cupricollis* HÄNEL, 1937  
*Chrysochloa speciosissima* var. *speciosissima* ab. *cyanea* HÄNEL, 1937  
*Chrysochloa speciosissima* var. *speciosissima* ab. *olivacea* HÄNEL, 1937  
*Chrysochloa speciosissima* var. *speciosissima* ab. *opaca* HÄNEL, 1937  
*Chrysochloa speciosissima* var. *speciosissima* ab. *violaceicollis* HÄNEL, 1937  
 Distribution: Mountains of Central and S Europe, Ukranian Carpathians.

**Genus: *Cecchiniola* JACOBSON, 1908**

***Cecchiniola platyscelidina* (JACOBSON, 1898)**

- Chrysomela platyscelidina* JACOBSON, 1898  
 Distribution: Crimea.

Subtribus: *Chrysomelina* CHEN, 1936

Genus: *Colaphellus* J. WEISE, 1916

*Colaphus* DAHL, 1823

***Colaphellus alpinus alpinus* (GEBLER, 1834)**

*Chrysomela alpina* GEBLER, 1834

Distribution: Asian Russia (S and Middle Siberia), Mongolia.

***Colaphellus alpinus bowringii* (BALY, 1865)**

*Colaphellus bowringii* BALY, 1865

Distribution: Asian Russia (Far East), E Asia.

***Colaphellus apicalis* (MÉNÉTRIÉS, 1849)**

*Chrysomena apicalis* MÉNÉTRIÉS, 1849

Distribution: Middle East, Central Asia (Turkmenistan, Uzbekistan, Tajikistan, Kazakhstan).

***Colaphellus sophiae sophiae* (SCHALLER, 1783)**

*Chrysomela sophiae* SCHALLER, 1783

*Chrysomela erythropus* GMELIN, 1790

*Chrysomela sophiae* ab. *bicolor* G. REINECK, 1922

Distribution: Central Europe, W and S Ukraine, Central part of Russian Plain, Armenia.

***Colaphellus sophiae hoefti* (MÉNÉTRIÉS, 1832)**

*Chrysomela hoefti* MÉNÉTRIÉS, 1832

Distribution: S and E Ukraine, Crimea, southern parts of European Russia, Caucasus, Kazakhstan, Central Asia, W Siberia.

Genus: *Gastrophysa* CHEVROLAT in DEJEAN, 1837

Subgenus: *Gastrophysa (Gastrophysa)* CHEVROLAT in DEJEAN, 1837

*Gastroidea* AGASSIZ, 1846

***Gastrophysa (Gastrophysa) atrocyanea* MOTSCHULSKY, 1860**

Distribution: Asian Russia (Amurland, Maritime Prov., Sakhalin), Korea, China, Japan, Vietnam.

***Gastrophysa (Gastrophysa) mannerheimi* (STÅL, 1858)**

*Phytodecta mannerheimi* STÅL, 1858

*Gastrophysa suturalis* MOTSCHULSKY, 1860

*Gastrophysa popovi* MARSEUL, 1888

Distribution: Asian Russia (S Siberia, Amurland), Mongolia, N China.

***Gastrophysa (Gastrophysa) polygoni* (LINNAEUS, 1758)**

*Chrysomela polygoni* LINNAEUS, 1758

*Chrysomela obtusa* O. F. MÜLLER, 1776

*Chrysomela ruficollis* FABRICIUS, 1781

*Gastrophysa polygoni elongata* JOLIVET, 1951

Distribution: Europe, Turkey, Central Asia, Asian Russia (S Siberia, Amurland), N China, Korea, N America.

***Gastrophysa (Gastrophysa) viridula viridula* (DEGEER, 1775)***Chrysomela viridula* DEGEER, 1775*Chrysomela raphani* HERBST, 1783*Gastroidea viridula* ab. *cyanescens* J. WEISE, 1898*Gastroidea viridula* ab. *fulgurans* ACHARD, 1923*Gastroidea viridula* ab. *cyanicollis* MADER, 1931

Distribution: Europe, SE Kazakhstan, Asian Russia (W and Middle Siberia), N America.

***Gastrophysa (Gastrophysa) viridula caucasica* JOLIVET, 1951**

Distribution: Caucasus.

***Gastrophysa (Gastrophysa) viridula lenta* (J. WEISE, 1887)***Gastroidea lenta* J. WEISE, 1887

Distribution: Asian Russia (E Siberia, Russian Far East), Mongolia.

**Genus: *Phaedon* DAHL, 1823***Alitene* GISTEL, 1857**Subgenus: *Phaedon (Phaedon)* DAHL, 1823*****Phaedon (Phaedon) armoraciae* (LINNAEUS, 1758)***Chrysomela armoraciae* LINNAEUS, 1758*Chrysomela cochleariae* PANZER, 1797*Chrysomela salicina* HEER, 1845*Phaedon betulae* KÜSTER, 1846*Chrysomela veronicae* BEDEL, 1892

Distribution: N, Central, and E Europe, Mountains of S Kazakhstan and Central Asia (except for the Turkmenistan), Asian Russia (Siberia east to Yakutia).

***Phaedon (Phaedon) bogdanovikatjkovi* JACOBSON, 1921**

Distribution: W Kazakhstan (River Emba).

***Phaedon (Phaedon) cochleariae cochleariae* (FABRICIUS, 1702)***Chrysomela cochleariae* FABRICIUS, 1792*Chrysomela egena* GYLLENHAL, 1827*Chrysomela omissa* C. R. SAHLBERG, 1838*Chrysomela neglecta* C. R. SAHLBERG, 1838*Phaedon galeopsis* LETZNER, 1848*Chrysomela grammica* SUFFRIAN, 1851*Chrysomela hederiae* SUFFRIAN, 1851*Phaedon cochleariae* var. *obesus* J. WEISE, 1884*Phaedon regnianum* TOTTENHAM, 1941

Distribution: Europe, Caucasus, Turkey, Kazakhstan, Central Asia, Asian Russia (S Siberia, Altai, Sayan).

***Phaedon (Phaedon) cochleariae brassicae* (BALY, 1874)***Phaedon brassicae* BALY, 1874*Phaedon incertum* BALY, 1874

Distribution: Russian Far East, Korea, China, Japan.



***Phaedon (Phaedon) concinnus* STEPHENS, 1834***Phaedon concinna* STEPHENS, 1834*Phaedon commutatus* KONTKANEN, 1933*Phaedon contemptus* KONTKANEN, 1933

Distribution: N, NW, and E Europe (Baltic States, N European Russia), Asian Russia (N Siberia and N Russian Far East), Mongolia.

***Phaedon (Phaedon) laevigatus* (DUFTSCHMIDT, 1825)***Chrysomela laevigatus* DUFTSCHMIDT, 1825*Chrysomela gramica* DUFTSCHMIDT, 1825*Chrysomela sabulicola* SUFFRIAN, 1851*Phaedon laevigatus* var. *violaceus* J. WEISE, 1884

Distribution: S and Central Europe, Ukraine, S European Russia, Caucasus.

**Subgenus: *Phaedon (Hemiphaedon)* JACOBSON, 1901*****Phaedon (Hemiphaedon) magnificus* LOPATIN, 1985**

Distribution: Kazakhstan (Barsakeljmes Isle in Aral Sea).

***Phaedon (Hemiphaedon) subtilis subtilis* J. WEISE, 1900**

Distribution: Plains of Uzbekistan, Kirghizstan, Tajikistan.

***Phaedon (Hemiphaedon) subtilis opacus* LOPATIN, 1976**

Distribution: SE Kazakhstan.

**Genus: *Neophaedon* JACOBSON, 1901***Alitene* GISTEL, 1857***Neophaedon pyritosus alutaceus* FLEISCHER, 1909**

Distribution: W Kazakhstan, W Central Asia, E Iran, Afghanistan.

***Neophaedon pyritosus pyritosus* (ROSSI, 1792)***Chrysomela pyritosa* ROSSI, 1792*Altica graminis* PANZER, 1799*Chrysomela graminicola* DRAPIEZ, 1819

Distribution: S and E Europe, Caucasus, N Kazakhstan, N Africa.

**Genus: *Sternoplatys* MOTSCHULSKY, 1860****Subgenus: *Sternoplatys (Sternoplatys)* MOTSCHULSKY, 1860*****Sternoplatys (Sternoplatys) clementzi* JACOBSON, 1901***Sternoplatys weisei* CSIKI, 1901*Prasocuris weisei* REITTER, 1901

Distribution: S Asian Russia (S Siberia, Altai, Tuva, Yakutia).

***Sternoplatys (Sternoplatys) fausti* J. WEISE, 1884**

Distribution: SE and E Asian Russia (Magadan Distr., Ussuri Area).

***Sternoplatys (Sternoplatys) fulvipes fulvipes* MOTSCHULSKY, 1860**

Distribution: Asian Russia (E Amurland, Maritime prov.), NE China.

***Sternoplatys (Sternoplatys) fulvipes baicalicus* (J. WEISE, 1900)***Phaedon baicalicus* J. WEISE, 1900*Sternoplatys motschulskyi* JACOBSON, 1901

Distribution: S Asian Russia (Transbaikalia, W Amurland).

***Sternoplatys (Sternoplatys) segnis* J. WEISE, 1884***Chrysomela pyritosa* DUFTSCHMIDT, 1825, non ROSSI, 1792*Phaedon hederæ* KRAATZ, 1876, non SUFFRIAN, 1851*Phaedon segnis* var. *obscurus* J. WEISE, 1884*Phaedon tatricus* MARSEUL, 1889*Phaedon szmolanyi* MARSEUL, 1889*Phaedon deubeli* J. WEISE, 1900

Distribution: Mountains of Central and SE Europe, Ukrainian Carpathians.

**Subgenus: *Sternoplatys (Austrosternoplatys)* JACOBSON, 1921*****Sternoplatys (Austrosternoplatys) longulus* J. WEISE, 1890**

Distribution: High Mountains Ranges of S Kazakhstan, Kirghizstan, Tajikistan.

**Genus: *Sclerophaedon* J. WEISE, 1882*****Sclerophaedon carniolicus* (GERMAR, 1824)***Chrysomela carniolica* GERMAR, 1824*Phaedon transsylvanicus* FUSS, 1861*Sclerophaedon carniolicus* ab. *obscurus* J. WEISE, 1884

Distribution: Mountains of Central and SE Europe, Ukrainian Carpathians.

***Sclerophaedon carpathicus* (J. WEISE, 1875)***Phaedon carniolicus* var. *carpathicus* J. WEISE, 1875*Phaedon kenderesyi* KIESENWETTER, 1879

Distribution: Mountains of Central Europe, Ukrainian Carpathians.

***Sclerophaedon orbicularis* (SUFFRIAN, 1851)***Chrysomela orbicularis* SUFFRIAN, 1851

Distribution: Mountains of Central Europe, Ukrainian Carpathians.

**Genus: *Oreothassa* JACOBSON, 1900*****Oreothassa martjanovi* JACOBSON, 1900**

Distribution: E Kazakhstan, S Asian Russia (Altai).

**Genus: *Hydrothassa* J. THOMSON, 1866****Subgenus: *Hydrothassa (Agrostithassa)* JACOBSON, 1921*****Hydrothassa (Agrostithassa) glabra* (HERBST, 1783)***Chrysomela glabra* HERBST, 1783*Chrysomela aucta* FABRICIUS, 1787*Chrysomela egena* SUFFRIAN, 1851, non GYLLENHAL, 1827*Prasocuris aucta* var. *umhangi* WEHNCKE, 1866

Distribution: Europe except for extreme N, W Siberia, NW Africa.

**Subgenus: *Hydrothassa* (*Hydrothassa*) J. THOMSON, 1866*****Hydrothassa* (*Hydrothassa*) *hannoveriana* (FABRICIUS, 1775)***Chrysomela hannoveriana* FABRICIUS, 1775*Chrysomela ranunculi* HERBST, 1783*Chrysomela potentillae* HERBST, 1783*Hydrothassa hannoverana* ab. *degenerata* MÄKLIN, 1881*Hydrothassa hannoverana* var. *calthae* J. WEISE, 1883*Hydrothassa hannoverana* var. *germanica* J. WEISE, 1883

Distribution: Central Europe, Ukraine and European Russia except for Steppe-zone, Siberia.

***Hydrothassa* (*Hydrothassa*) *marginella marginella* (LINNAEUS, 1758)***Chrysomela marginella* LINNAEUS, 1758*Hydrothassa marginella* ab. *devillei* BULLOCK, 1928

Distribution: Europe except for S, Kazakhstan, S Asian Russia (S Siberia east to Yakutia).

***Hydrothassa* (*Hydrothassa*) *marginella eoa* LOPATIN, 1962**

Distribution: Asian Russia (Maritime Prov.).

**Genus: *Apterocuris* JACOBSON, 1900*****Apterocuris sibirica* (GEBLER, 1830)***Chrysomela sibirica* GEBLER, 1830

Distribution: S Asian Russia (Altai).

**Genus: *Prasocuris* LATREILLE, 1802***Hellodes* L. REDTENBACHER, 1845***Prasocuris junci* (BRAHM, 1790)***Chrysomela junci* BRAHM, 1790*Chrysomela violacea* FABRICIUS, 1792*Chrysomela beccabungae* ILLIGER, 1794*Helodes violacea* var. *atra* DUFTSCHMIDT, 1825

Distribution: S, Central, and E Europe, Caucasus, S Kazakhstan.

***Prasocuris phellandrii* (LINNAEUS, 1758)***Chrysomela phellandrii* LINNAEUS, 1758*Prasocuris phellandrii* var. *cicutae* J. WEISE, 1883*Prasocuris phellandrii* var. *sii* J. WEISE, 1883*Prasocuris weisei* REITTER, 1901

Distribution: Europe except for N, Asian Russia (S Siberia, Maritime Prov.), Mongolia, N China.

**Genus: *Plagiodera* CHEVROLAT in DEJEAN, 1837***Linamorpha* MOTSCHULSKY, 1860*Plagiomorpha* MOTSCHULSKY, 1860*Plagiosterna* MOTSCHULSKY, 1860***Plagiodera versicolora* (LAICHARTING, 1781)***Chrysomela armoraciae* FABRICIUS, 1775, non LINNAEUS, 1758

- Chrysomela versicolora* LAICHARTING, 1781  
*Chrysomela salicis* PANZER, 1795  
*Chrysomela cognata* HERBST, 1783  
*Phaedon clavicornis* STEPHENS, 1832  
*Plagiodera chinensis* J. WEISE, 1898  
*Plagiodera versicolora laevicollis* LOPATIN, 1962  
 Distribution: All over Holarctic, India, Taiwan.

**Genus: *Chrysomela* LINNAEUS, 1758**

- Melasoma* STEPHENS, 1832  
*Lina* CHEVROLAT in DEJEAN, 1837

**Subgenus: *Chrysomela (Pachylina)* L. MEDVEDEV in MEDVEDEV et CHERNOV, 1969**

- Chrysomela (Pachylina) blaisdelli blaisdelli* VAN DYKE, 1938**  
*Chrysomela engelhardti* HATCH, 1939  
 Distribution: NE Asian Russia (Chukotka), Alaska, N Canada.
- Chrysomela (Pachylina) blaisdelli taimyrensis* L. MEDVEDEV, 1969**  
 Distribution: N Asian Russia (Taimyr Peninsula).
- Chrysomela (Pachylina) blaisdelli wrangeliana* L. MEDVEDEV, 1973**  
 Distribution: NE Asian Russia (Wrangel Is.).
- Chrysomela (Pachylina) collaris* LINNAEUS, 1758**  
*Chrysomela salicis* FABRICIUS, 1792  
*Chrysomela alpina* ZETTERSTEDT, 1828  
*Lina escheri* HEER, 1836  
*Lina daurica* MOTSCHULSKY, 1860  
 Distribution: All over Palaearctic except for Central Asia.

**Subgenus: *Chrysomela (Macrolina)* MOTSCHULSKY, 1860**

- Strickerus* R. LUCAS, 1920

- Chrysomela (Macrolina) cuprea* FABRICIUS, 1775**  
*Chrysomela ruficaudis* DEGEER, 1775  
*Chrysomela cuprea* ab. *sarmatica* J. WEISE, 1887  
*Chrysomela cuprea* ab. *coeruleovirens* GABRIEL, 1928  
*Chrysomela cuprea* ab. *cupreobrunnea* CSIKI, 1953  
 Distribution: All over Palaearctic except for Central Asia.
- Chrysomela (Macrolina) lapponica* LINNAEUS, 1758**  
*Chrysomela bulgharensis* FABRICIUS, 1798  
*Chrysomela lapponica* ab. *altaica* J. WEISE, 1884  
*Chrysomela lapponica* ab. *litura* MARSEUL, 1888  
*Chrysomela lapponica* ab. *bimaculata* GRADL, 1881  
*Chrysomela lapponica* ab. *bipustulata* GRADL, 1881  
*Chrysomela lapponica* ab. *coerulea* GRADL, 1881  
*Chrysomela lapponica* ab. *marginata* GRADL, 1881  
*Chrysomela lapponica* ab. *quadrimaculata* GRADL, 1881  
*Chrysomela lapponica* ab. *quadrinotata* GRADL, 1881

- Chrysomela lapponica* ab. *rubrocoerulea* GRADL, 1881  
*Chrysomela lapponica* ab. *sexmaculata* GRADL, 1881  
*Chrysomela lapponica* ab. *unicolor* MARSEUL, 1888  
*Chrysomela lapponica* ab. *cruciata* JACOBSON, 1901  
*Chrysomela lapponica* ab. *violaceipennis* JACOBSON, 1901  
*Chrysomela lapponica* ab. *quadripustulata* JACOBSON, 1902  
*Chrysomela lapponica* ab. *quadripunctata* LENGERKEN, 1913  
 Distribution: N, Central, and E Europe, Kirghizstan, Asian Russia (S Siberia, Tuva, Yakutia, Far East, Sakhalin).

***Chrysomela (Macrolina) vigintipunctata* (SCOPOLI, 1763)**

- Coccinella 20-punctata* SCOPOLI, 1763  
*Chrysomela vigintipunctata* ab. *pustulata* J. WEISE, 1884  
*Chrysomela vigintipunctata* ab. *incontaminata* HEYDEN, 1887  
*Chrysomela vigintipunctata* ab. *degeneri* SCHULZE, 1911  
*Chrysomela vigintipunctata* ab. *extrema* SCHULZE, 1911  
*Chrysomela vigintipunctata* ab. *melaina* SCHULZE, 1911  
*Chrysomela vigintipunctata* ab. *quadripunctata* SCHULZE, 1911  
*Chrysomela vigintipunctata* ab. *alba* BRASSLER, 1926  
 Distribution: All over Palaearctic except for Middle Asia.

**Subgenus: *Chrysomela (Chrysomela)* LINNAEUS, 1758**

***Chrysomela (Chrysomela) populi* LINNAEUS, 1758**

- Chrysomela populi* LINNAEUS, 1758  
*Lina violaceicollis* MOTSCHULSKY, 1860 [nomen nudum]  
*Chrysomela populi* ab. *janaceki* REITTER, 1892  
*Chrysomela populi asiatica* JAKOB, 1952  
*Chrysomela populi kitaika* JAKOB, 1952  
*Chrysomela populi nigricollis* JAKOB, 1952  
*Chrysomela populi violaceicollis* BECHYNÉ, 1954  
 Distribution: All over Palaearctic, Mountains of S Kazakhstan and Middle Asia.

***Chrysomela (Chrysomela) saliceti* (J. WEISE, 1884)**

- Melasoma saliceti* J. WEISE, 1884  
*Chrysomela tremulae* SUFFRIAN, 1851, non FABRICIUS, 1787  
*Melasoma saliceti* ab. *nigripennis* REITTER, 1894  
 Distribution: All over Palaearctic except for Middle Asia.

***Chrysomela (Chrysomela) tremula* FABRICIUS, 1787**

- Chrysomela tremulae* PAYKULL, 1799  
*Chrysomela longicollis* SUFFRIAN, 1851  
 Distribution: All over Palaearctic except for Middle Asia.

***Chrysomela (Chrysomela) turkestanica* G. REINECK, 1937**

- Distribution: Mountains of Middle Asia, Afghanistan.

**Genus: *Linaeidea* MOTSCHULSKY, 1860**

***Linaeidea aenea* (LINNAEUS, 1758)**

- Chrysomela aenea* LINNAEUS, 1758  
*Chrysomela vitellinae* SCOPOLI, 1763

*Chrysomela coeruleo-violacea* DEGEER, 1775

*Chrysomela haemorroidalis* O. F. MÜLLER, 1776

*Chrysomela violacea* OLIVIER, 1790

*Melasoma aenea* var. *bicolor* SCHILSKY, 1908

*Melasoma aenea* ab. *discolor* GERHARDT, 1910

Distribution: Europe except for extreme N, E Kazakhstan, Siberia, N China, Japan.

**Genus: *Gastrolina* BALY, 1859**

***Gastrolina depressa* BALY, 1859**

*Gastrolina thoracica* BALY, 1864

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

***Gastrolina peltoidea* (GEBLER, 1832)**

*Chrysomela peltoidea* GEBLER, 1832

*Gastrolina japana* JACOBY, 1885

Distribution: Asian Russia (Central Siberia, Yakutia, Transbaikalia, Far East, Sakhalin, Kuriles Isles), NE China, Japan.

**Genus: *Gastrolinoides* CHÛJÔ et KIMOTO, 1960**

***Gastrolinoides japonica* (HAROLD, 1877)**

*Melasoma japonica* HAROLD, 1877

Distribution: Asian Russia (Amurland, Maritime Prov.), Japan.

Subtribus: *Doryphorina* YUASA, 1936

**Genus: *Leptinotarsa* CHEVROLAT in DEJEAN, 1837**

***Leptinotarsa decemlineata* (SAY, 1824)**

*Doryphora decemlineata* SAY, 1824

Distribution: Introduced from N America. Europe, Caucasus, European Russia east to Astrakhan' Distr. and Kalmykia.

**Genus: *Zygogramma* CHEVROLAT in DEJEAN, 1837**

***Zygogramma suturalis* (FABRICIUS, 1775)**

*Crioceris suturalis* FABRICIUS, 1775

Distribution: Introduced from N America. Ciscaucasia, W Caucasus.

Subtribus: *Entomoscelina* REITTER, 1912

**Genus: *Cystocnemis* MOTSCHULSKY, 1860**

***Cystocnemis concolor* (KRAATZ, 1879)**

*Chrysomela concolor* KRAATZ, 1879

Distribution: SE Kazakhstan (Saur and Tarbagatay Mountain Ranges).

***Cystocnemis discoidea* (GEBLER, 1830)***Chrysomela discoidea* GEBLER, 1830

Distribution: E Kazakhstan, S Asian Russia (S Altais).

**Genus: *Xenomela* J. WEISE, 1884*****Xenomela ballioni* LOPATIN, 1989**

Distribution: Kirghizstan (Kirghiz Alatau Mountain Range).

***Xenomela belousovi* LOPATIN, 1989**

Distribution: SW Kirghizstan and Uzbekistan (Chatkal, Ugam and Karzantau Mountain Ranges).

***Xenomela dohrni* (SOLSKY, 1881)***Entomoscelis dohrni* SOLSKY, 1881

Distribution: Plains of Uzbekistan and N Tajikistan.

***Xenomela karatavica* LOPATIN, 1989**

Distribution: Kazakhstan (Karatau Mountain Range).

***Xenomela konstantinovi* LOPATIN, 1995**

Distribution: Uzbekistan (Chimghan Mountain Range).

***Xenomela kraatzi* J. WEISE, 1884**

Distribution: Uzbekistan and Kirghizstan (Fergan Mountain Range).

***Xenomela laevigata* LOPATIN, 1989**

Distribution: S Kirghizstan (Chatkal, Fergan and Bozbutau Mountain Ranges).

***Xenomela marginicollis* (BALLION, 1878)***Chrysomela marginicollis* BALLION, 1878*Chrysomela fulvipes* BALLION, 1878*Chrysomela ovipennis* BALLION, 1878

Distribution: Central and E Tain-Shan in the borders of Kirghizstan and Kazakhstan, China (Narat Mountain Range).

***Xenomela minckwitzae minckwitzae* (JACOBSON, 1925)***Xenomela regeli morpha minckwitzae* JACOBSON, 1925

Distribution: Kirghizstan (Talas and Ugam Mountain Ranges).

***Xenomela minckwitzae kreutzbergi* LOPATIN, 1989**

Distribution: Uzbekistan (Karzantau Mountain Range).

***Xenomela minckwitzae pskemica* LOPATIN, 1989**

Distribution: Kirghizstan (Pskem Mountain Range).

***Xenomela ovczinnikovi* LOPATIN, 1995**

Distribution: Uzbekistan (Kuramin Mountain Range).

***Xenomela regeli* JACOBSON, 1897**

Distribution: Uzbekistan (Sailyk Mountain, Karzantau Mountain Range).

Genus: *Oreomela* JACOBSON, 1895Subgenus: *Oreomela (Olcomela)* JACOBSON, 1925*Oreomela (Olcomela) korolkovi* JACOBSON, 1925

Distribution: E Kazakhstan (Dzhungar Alatau Mountain Range).

*Oreomela (Olcomela) suvorovi* JACOBSON, 1910

Distribution: E Kazakhstan (Dzhungar Alatau Mountain Range), NW China (Borochoro Mountain Range).

Subgenus: *Oreomela (Oreomela)* JACOBSON, 1895*Oreomela (Oreomela) abramovi* JACOBSON, 1925

Distribution: Kirghizstan (Talas Alatau Mountain Range).

*Oreomela (Oreomela) andreevi* JACOBSON, 1910

Distribution: Kirghizstan (Terskey Alatau Mountain Range).

*Oreomela (Oreomela) bergi* JACOBSON, 1910

Distribution: S Kirghizstan, N Tajikistan (Turkestan Mountain Range).

*Oreomela (Oreomela) borealis* LOPATIN, 1999

Distribution: Kazakhstan (Chu-Ili Mountains).

*Oreomela (Oreomela) celyphoides* JACOBSON, 1896

Distribution: S Kazakhstan (Dzhungar Alatau Mountain Range).

*Oreomela (Oreomela) clypealis* JACOBSON, 1901

Distribution: Kirghizstan (Kirghizian Alatau Mountain Range).

*Oreomela (Oreomela) dungana* JACOBSON, 1925

Distribution: Kirghizstan (Sarydzhas Mountain Range).

*Oreomela (Oreomela) dzhungara* JACOBSON, 1925

Distribution: S Kazakhstan (Dzhungar Alatau Mountain Range).

*Oreomela (Oreomela) fuscipes* (J. WEISE, 1885)*Xenomela fuscipes* J. WEISE, 1885*Xenomela fausti* J. WEISE, 1885

Distribution: Kirghizstan (Kirghiz Alatau, Kunghey Alatau, Moldotau Mountain Range).

*Oreomela (Oreomela) gracilis* LOPATIN, 1999

Distribution: Kirghizstan (Terskey Alatau Mountain Range).

*Oreomela (Oreomela) heydeni* (J. WEISE, 1885)*Xenomela heydeni* J. WEISE, 1885

Distribution: Kirghizstan (Son-Kul Mountain Range).

*Oreomela (Oreomela) hohlbecki* JACOBSON, 1925

Distribution: Kirghizstan (Kirghiz Mountain Range).

*Oreomela (Oreomela) jacobsoni* (SEMENOV, 1894)*Xenomela jacobsoni* SEMENOV, 1894



*Xenomela trotzinai* JACOBSON, 1894

Distribution: Kirghizstan (Alai Mountain Range).

***Oreomela (Oreomela) joliveti* LOPATIN, 1976**

Distribution: SE Kazakhstan (Ketmen Mountain Range).

***Oreomela (Oreomela) juldusana* LOPATIN, 1962**

Distribution: NW China (Dzundaria).

***Oreomela (Oreomela) kabaki* LOPATIN, 1995**

Distribution: E Kazakhstan (Dzhungar Alatau Mountain Range).

***Oreomela (Oreomela) kaszabi* LOPATIN, 1962**

Distribution: Tibet.

***Oreomela (Oreomela) korotjaevi* LOPATIN, 1983**

Distribution: SE Kazakhstan (Transili Alatau Mountain Range).

***Oreomela (Oreomela) kungeica* LOPATIN, 1976**

Distribution: Kirghizstan (Kunghey Alatau Mountain Range).

***Oreomela (Oreomela) kutzenkoi* JACOBSON, 1925**

Distribution: Kirghizstan (Atbashy Mountain Range).

***Oreomela (Oreomela) lopatini* GRUJEV, 1991**

Distribution: Kirghizstan (Susamyr Mountain Range).

***Oreomela (Oreomela) medvedevi* LOPATIN, 1968**

Distribution: E Tajikistan (Badakhshan).

***Oreomela (Oreomela) mitjaevi* LOPATIN, 1996**

*Oreomela dzhungara* LOPATIN, 1990, non JACOBSON, 1925

Distribution: SE Kazakhstan (Dzhungar Alatau Mountain Range).

***Oreomela (Oreomela) montivaga* LOPATIN, 1987**

Distribution: Kirghizstan (Susamyr Mountain Range).

***Oreomela (Oreomela) pedaschenkoi* JACOBSON, 1925**

Distribution: Kirghizstan (Terskey Alatau and Atbashy Mountain Ranges).

***Oreomela (Oreomela) puncticollis* LOPATIN, 1999**

Distribution: E Kazakhstan (Dzhungar Alatau Mountain Range).

***Oreomela (Oreomela) radkewiczi* JACOBSON, 1925**

Distribution: S Kazakhstan (Kendyktau Mountain Range).

***Oreomela (Oreomela) recticollis* LOPATIN, 1995**

Distribution: Kirghizstan (Kunghey Alatau Mountain Range).

***Oreomela (Oreomela) ruckbeili* JACOBSON, 1925**

Distribution: Kirghizstan (Terskey Alatau Mountain Range).

***Oreomela (Oreomela) sapozhnikov* JACOBSON, 1925**

Distribution: Kirghizstan (Terskey Alatau, Ketmen, Sarydzhaz Mountain Ranges).

***Oreomela (Oreomela) sarydzhasea* JACOBSON, 1925**

Distribution: Kirghizstan (Sarydzhaz Mountain Ranges).

***Oreomela (Oreomela) schnitnikovi* JACOBSON, 1925**

Distribution: Kirghizstan (Kirghiz Alatau Mountain Ranges).

***Oreomela (Oreomela) scutellaris* JACOBSON, 1925**

Distribution: Kirghizstan (Sarydzhaz and Kunghey Alatau Mountain Ranges).

***Oreomela (Oreomela) semenovi* JACOBSON, 1895**

Distribution: S Kazakhstan (Transili Alatau Mountain Range).

***Oreomela (Oreomela) sussamyrica* LOPATIN, 1987**

Distribution: Kirghizstan (Susamyr Mountain Range).

***Oreomela (Oreomela) tarantscha* JACOBSON, 1925**

Distribution: SE Kazakhstan (Ketmen Mountain Range).

***Oreomela (Oreomela) tarbagataica* LOPATIN, 1968**

Distribution: E Kazakhstan (Tarbagatay Mountain Range).

***Oreomela (Oreomela) transalaica* LOPATIN, 1965**

Distribution: Kirghizstan (Transalai Mountain Range).

***Oreomela (Oreomela) transiliensis* LOPATIN, 1976**

Distribution: S Kazakhstan (Transili Alatau Mountain Range).

***Oreomela (Oreomela) tschernavini* JACOBSON, 1925**

Distribution: Kirghizstan (Susamyr Mountain Range).

***Oreomela (Oreomela) weisei* JACOBSON, 1895***Oreomela invisa* LOPATIN, 1962

Distribution: Kirghizstan (Terskey Alatau Mountain Range).

***Oreomela (Oreomela) zaslavskii* LOPATIN, 1976**

Distribution: Kirghizstan (Atbashy Mountain Range).

**Subgenus: *Oreomela (Entomomela)* JACOBSON, 1925*****Oreomela (Entomomela) arnoldii* LOPATIN, 1974**

Distribution: E Kazakhstan (Azutau Mountain Range).

**Genus: *Entomoscelis* CHEVROLAT in DEJEAN, 1837*****Entomoscelis adonidis* (PALLAS, 1771)***Chrysomela adonidis* PALLAS, 1771*Chrysomela 3lineata* FABRICIUS, 1777*Chrysomela dorsalis* FABRICIUS, 1777*Entomoscelis goliath* ABEILLE, 1897*Entomoscelis hammarstroemi* JACOBSON, 1901

Distribution: S and SE Europe, Lithuania, Ukraine, European Russia except for northern parts, Turkey, Kazakhstan, Central Asia except for Deserts of Turkmenistan, Asian Russia (Siberia, Yakutia, Far East), Mongolia, N China.

***Entomoscelis deserticola* LOPATIN, 1967**

Distribution: China (Sinkiang).

***Entomoscelis erythrocnema* JACOBSON, 1893**

Distribution: South parts of Central Asia, north to Nuratau Mountains in Uzbekistan.

***Entomoscelis orientalis* MOTSCHULSKY, 1860**

Distribution: Asian Russia (Transbaikalia, Amurland, Maritime Prov.), Mongolia, N China, Korea.

***Entomoscelis pilula* LOPATIN, 1967**

Distribution: Transcaucasia.

***Entomoscelis sacra* (LINNAEUS, 1758)***Chrysomela sacra* LINNAEUS, 1758*Chrysomela melanostoma* GMELIN, 1790*Chrysomela berytensis* REICHE et SAULCY, 1858

Distribution: SE Europe, Turkey, Kazakhstan, Central Asia.

***Entomoscelis suturalis* J. WEISE, 1882***Entomoscelis suturalis* ab. *sancta* MADER, 1943

Distribution: SE Europe, Caucasus, Transcaucasia, Turkey, Middle East.

Subtribus: *Gonioctenina* WILCOX, 1972**Genus: *Gonioctena* CHEVROLAT in DEJEAN, 1837***Phytodecta* KIRBY, 1837**Subgenus: *Gonioctena* (*Gonioctena*) CHEVROLAT in DEJEAN, 1837*****Gonioctena* (*Gonioctena*) *coreana* (BECHYNÉ, 1947)***Phytodecta coreana* BECHYNÉ, 1947

Distribution: Asian Russia (Magadan Distr., Khabarovsk Distr., Amurland, Maritime Prov.), NE China, Korea.

***Gonioctena* (*Gonioctena*) *decaspilota* (ACHARD, 1924)***Chrysomela affinis* GYLLENHAL, 1808, non FABRICIUS, 1787*Chrysomela arctica* MANNERHEIM, 1853*Phytodecta salicis* MOTSCHULSKY, 1860*Phytodecta affinis* var. *decaspilotus* ACHARD, 1924*Phytodecta affinis* ab. *hamata* ACHARD, 1924*Phytodecta norvegicus* A. STRAND, 1936*Phytodecta decaspilota* ab. *aterrima* HELLÉN, 1937*Phytodecta dinah* BECHYNÉ, 1947*Phytodecta springlovae* BECHYNÉ, 1947*Phytodecta decaspilota* ab. *fennica* BECHYNÉ, 1947*Phytodecta decaspilota* ab. *zinaida* BECHYNÉ, 1947*Phytodecta decaspilota* ab. *fennica* BECHYNÉ, 1947

Distribution: N and NE Europe, Asian Russia (Siberia, Far East), Mongolia, N China, Korea.

***Gonioctena (Gonioctena) decemnotata* (MARSHAM, 1802)***Chrysomela rufipes* DEGEER, 1775, non LINNAEUS, 1758*Chrysomela 6 punctata* FABRICIUS, 1787, non LINNAEUS, 1758*Chrysomela 10-notata* MARSHAM, 1802*Phytodecta decemnotata* ab. *conjuncta* E. REINECK, 1910*Phytodecta decemnotata* ab. *populi* BECHYNÉ, 1947*Phytodecta decemnotata* ab. *praecox* BECHYNÉ, 1947*Phytodecta decemnotata* ab. *taeniolata* BECHYNÉ, 1947*Phytodecta decemnotata* ab. *tremulae* BECHYNÉ, 1947

Distribution: All over Holarctic except for S Europe and Central Asia.

***Gonioctena (Gonioctena) flavicornis flavicornis* (SUFFRIAN, 1851)***Chrysomela flavicornis* SUFFRIAN, 1851*Phytodecta flavicornis* ab. *nigriventris* REITTER, 1912*Phytodecta flavicornis* ab. *limbratipennis* ACHARD, 1924

Distribution: N, Central, and E Europe, Asian Russia (Siberia, Far East, Maritime Prov.), Mongolia, Japan.

***Gonioctena (Gonioctena) flavicornis borealis* (L. MEDVEDEV, 1963)***Phytodecta flavicornis borealis* L. MEDVEDEV, 1963

Distribution: Asian Russia (Kamchatka).

***Gonioctena (Gonioctena) gracilicornis* (KRAATZ, 1879)***Phytodecta gracilicornis* KRAATZ, 1879*Gonioctena sunkangensis* KIMOTO, 1966

Distribution: Asian Russia (Siberia, Far East, Maritime Prov., Sakhalin), Mongolia, N China.

***Gonioctena (Gonioctena) honshuensis honshuensis* (NAKANE, 1963)***Phytodecta honshuensis honshuensis* NAKANE, 1963*Phytodecta chujoi* L. MEDVEDEV, 1966

Distribution: Asian Russia (Kuriles Isles), Japan.

***Gonioctena (Gonioctena) honshuensis ochotense* (L. MEDVEDEV, 1968)***Phytodecta honshuensis ochotense* L. MEDVEDEV, 1968

Distribution: Asian Russia (Magadan Distr., Kamchatka).

***Gonioctena (Gonioctena) honshuensis sachalinensis* (L. MEDVEDEV, 1968)***Phytodecta honshuensis sachalinensis* L. MEDVEDEV, 1968

Distribution: Asian Russia (Sakhalin).

***Gonioctena (Gonioctena) jacobsoni* (OGLOBLIN et L. MEDVEDEV, 1956)***Phytodecta jacobsoni* OGLOBLIN et L. MEDVEDEV, 1956

Distribution: Asian Russia (Siberia, Magadan Distr., Maritime Prov.), Mongolia.

***Gonioctena (Gonioctena) janovskii* L. MEDVEDEV, 1976**

Distribution: S Siberia (?), Mongolia.

***Gonioctena (Gonioctena) japonica* (CHŪJŌ et KIMOTO, 1960)***Phytodecta japonica* CHŪJŌ et KIMOTO, 1960

Distribution: Asian Russia (S Kuriles Isles), Japan.

***Gonioctena (Gonioctena) kamiyai* KIMOTO, 1963***Phytodecta kamiyai* KIMOTO, 1963

*Phytodecta ussuriensis* L. MEDVEDEV, 1964

Distribution: Asian Russia (Amurland, S Maritime Prov.), Japan.

***Gonioctena (Gonioctena) kaufmanni* (L. MILLER, 1881)**

*Phytodecta kaufmanni* L. MILLER, 1881

*Phytodecta kaufmanni* var. *seminigra* J. WEISE, 1893

*Phytodecta kaufmanni* var. *infernalis* PENECKE, 1901

Distribution: Mountains of Central and E Europe, Carpathians.

***Gonioctena (Gonioctena) laeta* L. MEDVEDEV, 1973**

Distribution: Asian Russia (S Maritime Prov.), Korea.

***Gonioctena (Gonioctena) linnaeana linnaeana* (SCHRANK, 1781)**

*Chrysomela Linnaeana* SCHRANK, 1781

*Chrysomela tibialis* var. *decasigma* DUFTSCHMIDT, 1825

*Chrysomela triandrae* SUFFRIAN, 1851

*Phytodecta tibialis* var. *kraatzi* WESTHOFF, 1882

*Phytodecta tibialis* var. *nigricollis* WESTHOFF, 1882

*Phytodecta tibialis* var. *satanas* WESTHOFF, 1882

*Phytodecta linnaeana* var. *nigrolocana* CZWALINA, 1885

*Phytodecta linnaeana* ab. *bohémica* ACHARD, 1924

*Phytodecta linnaeana* ab. *lunulata* BECHYNÉ, 1947

*Phytodecta linnaeana* ab. *flavotibialis* BECHYNÉ, 1947

Distribution: Europe except for Mediterranean, Caucasus, Kazakhstan except for southern parts, Asian Russia (W and S Siberia, Altai, Tuva), Mongolia.

***Gonioctena (Gonioctena) linnaeana bergrothi* (JACOBSON, 1901)**

*Phytodecta linnaeana bergrothi* JACOBSON, 1901

Distribution: Asian Russia (S Siberia from Altai to Pacific Coast).

***Gonioctena (Gonioctena) ogloblini* L. MEDVEDEV et DUBESHKO, 1992**

Distribution: Asian Russia (Siberia, Far East, Maritime Prov.), Mongolia.

***Gonioctena (Gonioctena) sibirica* (J. WEISE, 1893)**

*Phytodecta sibirica* J. WEISE, 1893

*Phytodecta sorbi* MOTSCHULSKY, 1860

*Phytodecta sobrinus* J. WEISE, 1916

*Phytodecta cheni* BECHYNÉ, 1947

Distribution: Asian Russia (Siberia, Far East, Kamchatka, Maritime Prov.), Japan.

***Gonioctena (Gonioctena) sundmanni* (JACOBSON, 1901)**

*Phytodecta sundmanni* JACOBSON, 1901

Distribution: N Europe, Asian Russia (Siberia, Far East, Kamchatka Peninsula), Mongolia.

***Gonioctena (Gonioctena) viminalis viminalis* (LINNAEUS, 1758)**

*Chrysomela viminalis* LINNAEUS, 1758

*Chrysomela decempunctata* LINNAEUS, 1758

*Coccinella haemorroidalis* SCOPOLI, 1763

*Chrysomela fridrichsdalensis* O. MÜLLER, 1764

*Crioceris calcarata* FABRICIUS, 1792

*Chrysomela baaderi* PANZER, 1794

*Phytodecta viminalis* var. *bicolor* KRAATZ, 1879

*Phytodecta viminalis* var. *collaris* GRADL, 1882  
*Phytodecta viminalis* var. *marginalis* GRADL, 1882  
*Phytodecta viminalis* var. *nigrescens* GRADL, 1882  
*Phytodecta viminalis* var. *gradli* HEYDEN, 1883  
*Phytodecta viminalis* var. *quadripunctata* HEYDEN, 1883  
*Phytodecta viminalis* var. *cincta* J. WEISE, 1884  
*Phytodecta viminalis* ab. *sempunctata* W. EICHLER, 1914  
*Phytodecta viminalis* ab. *trimaculatis* PIC, 1924  
*Phytodecta viminalis* ab. *decemstilata* BECHYNÉ, 1947  
*Phytodecta viminalis* ab. *marekoum* BECHYNÉ, 1947  
*Gonioctena viminalis* ab. *pudica* BECHYNÉ, 1947  
*Gonioctena viminalis* ab. *xnotata* BECHYNÉ, 1947

Distribution: All over Holarctic except for Mediterranean and Central Asia.

***Gonioctena (Gonioctena) viminalis rufa* (KRAATZ, 1879)**

*Phytodecta rufa* KRAATZ, 1879

Distribution: Asian Russia (Transbaikalia, Sayan, Far East), Mongolia, NE China, Korea.

**Subgenus: *Gonioctena (Spartophila)* CHEVROLAT in DEJEAN, 1837**

***Gonioctena (Spartophila) olivacea* (FORSTER, 1771)**

*Chrysomela olivacea* FORSTER, 1771  
*Chrysomela litura* FABRICIUS, 1775  
*Chrysomela flavicans* FABRICIUS, 1787  
*Phytodecta weisei* REITTER, 1896  
*Phytodecta olivacea* var. *nigricans* J. WEISE, 1884  
*Phytodecta olivacea* ab. *bipartita* ACHARD, 1924  
*Phytodecta olivacea* ab. *escorialensis* ACHARD, 1924  
*Phytodecta olivacea* ab. *umbriceps* ROUBAL, 1941

Distribution: S and Central Europe, Belarus, Ukraine east to Kharkov, NW Africa.

**Subgenus: *Gonioctena (Brachyphytodecta)* BECHYNÉ, 1947**

***Gonioctena (Brachyphytodecta) fulva* (MOTSCHULSKY, 1860)**

*Spartophila fulva* MOTSCHULSKY, 1860

Distribution: Asian Russia (Amurland, Maritime Prov.), E China, Vietnam.

**Subgenus: *Gonioctena (Goniomena)* MOTSCHULSKY, 1860**

***Gonioctena (Goniomena) pallida* (LINNAEUS, 1758)**

*Chrysomela pallida* LINNAEUS, 1758  
*Chrysomela dispar* PAYKULL, 1799 (part.)  
*Chrysomela frontalis* OLIVIER, 1807  
*Phytodecta pallida* var. *borealis* J. WEISE, 1884  
*Phytodecta pallida* var. *decipiens* J. WEISE, 1884  
*Phytodecta pallida* var. *frontalis* J. WEISE, 1884  
*Phytodecta pallida* ab. *nigripennis* J. WEISE, 1893  
*Phytodecta pallida* ab. *nigricolor* REITTER, 1914  
*Phytodecta pallida* ab. *brunneiventris* KASZAB, 1962

Distribution: N, Central, and E Europe, Kazakhstan, Asian Russia (Siberia, Maritime Prov.), N Mongolia.

***Gonioctena (Goniomena) quinquepunctata intermedia* (HELLIESEN, 1913)**

- Phytodecta intermedia* HELLIESEN, 1913  
*Phytodecta intermedia* ab. *kendii* CSIKI, 1953  
*Phytodecta intermedia* ab. *barsica* KASZAB, 1962  
*Phytodecta intermedia* ab. *brunnicollis* KASZAB, 1962  
*Phytodecta intermedia* ab. *deletepunctata* KASZAB, 1962  
*Phytodecta intermedia* ab. *falsopallida* KASZAB, 1962  
*Phytodecta intermedia* ab. *herculeana* KASZAB, 1962

Distribution: Mountains of Central and E Europe, N and NE Europe.

***Gonioctena (Goniomena) quinquepunctata quinquepunctata* (FABRICIUS, 1787)**

- Chrysomela quinquepunctata* FABRICIUS, 1787  
*Chrysomela decempunctata* SCHRANK, 1781, non LINNAEUS, 1758  
*Chrysomela dispar* var. *flavicollis* DUFTSCHMIDT, 1825  
*Paropsis obscura* GRIMMER, 1841  
*Phytodecta quinquepunctata* var. *sorbi* J. WEISE, 1884  
*Phytodecta quinquepunctata* var. *unicolor* J. WEISE, 1884  
*Phytodecta quinquepunctata* var. *melanoptera* PENECKE, 1898  
*Phytodecta quinquepunctata* var. *nigriventris* PENECKE, 1898  
*Phytodecta quinquepunctata* var. *padi* PENECKE, 1898  
*Phytodecta quinquepunctata* var. *aucupariae* JACOBSON, 1900  
*Phytodecta quinquepunctata* var. *nigritinctus* ACHARD, 1924  
*Phytodecta quinquepunctata* ab. *scutellatus* TRELLA, 1930  
*Phytodecta quinquepunctata* ab. *bechynei* ROUBAL, 1948  
Distribution: N, Central, and E Europe, Carpathians, European Russia, Asian Russia (W Siberia, Yakutia).

**Subgenus: *Gonioctena (Spartoxena)* MOTSCHULSKY, 1860**

*Spartomena* REITTER, 1912

***Gonioctena (Spartoxena) fornicata* BRÜGGEMANN, 1873**

- Chrysomela sexpunctata* PANZER, 1796, non LINNAEUS, 1758  
*Phytodecta fornicata* ab. *innotata* J. WEISE, 1884  
*Phytodecta fornicata* ab. *picea* J. WEISE, 1884  
*Phytodecta fornicata* ab. *sequensi* REITTER, 1916  
*Phytodecta fornicata* ab. *bohumilae* BECHYNÉ, 1945  
*Phytodecta fornicata* ab. *hexaspilota* BECHYNÉ, 1945  
*Phytodecta fornicata* ab. *septempunctata* BECHYNÉ, 1945  
*Phytodecta fornicata* ab. *tetraspilota* BECHYNÉ, 1945  
*Phytodecta fornicata* ab. *triangularis* BECHYNÉ, 1945  
*Phytodecta fornicata* ab. *nigriceps* CSIKI, 1953  
*Phytodecta fornicata* ab. *vittipennis* CSIKI, 1953  
*Phytodecta fornicata* ab. *conjuncta* ENDRÖDI, 1957  
*Phytodecta fornicata* ab. *diabolica* ENDRÖDI, 1957  
Distribution: SE Europe, SW Ukraine, Moldavia, Turkey.

**Genus: *Cercyonops* JACOBSON, 1900*****Cercyonops caraganae* (GEBLER, 1823)**

- Chrysomela caraganae* GEBLER, 1823  
Distribution: NE of European Russia, S Asian Russia (S Siberia, Altai, Tuva).

Subtribus: *Paropsina* J. WEISE, 1915

**Genus: *Paropsides* MOTSCHULSKY, 1860**

***Paropsides soriculatus* (SWARTZ, 1808)**

*Paropsis soriculata* SWARTZ, 1808

*Paropsis duodecimpustulata* GEBLER, 1825

*Paropsis hieroglyphica* GEBLER, 1825

Distribution: Asian Russia (Yakutia, Transbaikalia, Amurland, Maritime Prov.), China, Korea, Japan, Vietnam.

Subtribus: *Phratorina* LOPATIN, 1977

**Genus: *Phratora* CHEVROLAT in DEJEAN, 1837**

*Phyllodecta* KIRBY, 1837

**Subgenus: *Phratora* (*Phratora*) CHEVROLAT in DEJEAN, 1837**

***Phratora* (*Phratora*) *atrovirens* (CORNELIUS, 1857)**

*Chrysomela atrovirens* CORNELIUS, 1857

Distribution: Central and E Europe, Caucasus, Asian Russia (Siberia, Far East), Mongolia.

***Phratora* (*Phratora*) *horioni* (MOHR, 1968)**

*Phyllodecta horioni* MOHR, 1968

Distribution: Caucasus (Teberda).

***Phratora* (*Phratora*) *laticollis* (SUFFRIAN, 1851)**

*Chrysomela laticollis* SUFFRIAN, 1851

*Phyllodecta cavifrons* C. G. THOMSON, 1866

Distribution: All over Palaearctic except for extreme N and Central Asia.

***Phratora* (*Phratora*) *polaris polaris* (W. G. SCHNEIDER, 1886)**

*Phyllodecta polaris* W. G. SCHNEIDER, 1886

Distribution: N Europe, Asian Russia (Siberia, N Far East), Mongolia.

***Phratora* (*Phratora*) *polaris reitteri* (LOPATIN, 1962)**

*Phyllodecta reitteri* LOPATIN, 1962

Distribution: Asian Russia (Tuva, Cisbaikalia), Mongolia.

***Phratora* (*Phratora*) *tibialis* (SUFFRIAN, 1851)**

*Chrysomela tibialis* SUFFRIAN, 1851

*Phyllodecta viennensis* J. WEISE, 1882

*Phyllodecta viennensis* fa. *cornelii* J. WEISE, 1882

*Phyllodecta viennensis* var. *coerulea* J. WEISE, 1884

Distribution: Central and E Europe, Caucasus, Kazakhstan.

***Phratora* (*Phratora*) *vitellinae* (LINNAEUS, 1758)**

*Chrysomela vitellinae* LINNAEUS, 1758

*Phratora brevicollis* MOTSCHULSKY, 1860



*Phratora nigrica* MOTSCHULSKY, 1860

Distribution: Europe, Kazakhstan, Asian Russia (Siberia, Far East), Mongolia, Korea.

**Subgenus: *Phratora (Chaetoceroides)* E. STRAND, 1935**

*Chaetocera* J. WEISE, 1884, non AGASSIZ, 1846

*Chaeroseta* MROCKOWSKI, 1990

***Phratora (Chaetoceroides) obtusicollis* MOTSCHULSKY, 1860**

*Phyllodecta inhonesta* J. WEISE, 1884

Distribution: Asian Russia (Amurland, Maritime Prov., Sakhalin, Kuriles Isles), Mongolia, NE China, Korea, Japan.

***Phratora (Chaetoceroides) vulgatissima* (LINNAEUS, 1758)**

*Chrysomela vulgatissima* LINNAEUS, 1758

*Phratora coeruleascens* KÜSTER, 1848

*Phratora longulus* MOTSCHULSKY, 1860

*Phratora vulgatissima* fa. *nigripes* LENTZ, 1879 [nomen nudum]

*Phyllodecta vulgatissima* var. *aestiva* J. WEISE, 1884

*Phyllodecta vulgatissima* var. *obscura* J. WEISE, 1884

Distribution: Europe except for Mediterranean, Caucasus, E Kazakhstan, Asian Russia (Siberia, Far East), Mongolia, N America.

Subfamily: *Galerucinae* LATREILLE, 1802

Tribe: *Galerucini* LATREILLE, 1802

**Genus: *Clitena* BALY, 1864**

***Clitena fuscipennis* (JACOBY, 1885)**

*Galerucella fuscipennis* JACOBY, 1885

*Galerucella rosinae* PIC, 1905

Distribution: Asian Russia (Amurland, Maritime Prov., S Kuriles Isles), Korea, Japan.

**Genus: *Diorhabda* J. WEISE, 1883**

*Radymna* REITTER, 1912

***Diorhabda elongata elongata* (BRULLÉ, 1832)**

*Chrysomela elongata* BRULLÉ, 1832

*Galeruca sublineata* H. LUCAS, 1849

Distribution: Mediterranean, Central Asia.

***Diorhabda elongata carinata* (FALDERMANN, 1837)**

*Galeruca carinata* FALDERMANN, 1837

Distribution: E Kazakhstan, NW China.

***Diorhabda elongata deserticola* CHEN, 1961**

Distribution: E Kazakhstan, NW China.

***Diorhabda fischeri* (FALDERMANN, 1837)**

*Galeruca fischeri* FALDERMANN, 1837

Distribution: Transcaucasia, Turkey, W Iran, Turkmenistan.

***Diorhabda nigrifrons* LABOISSIÈRE, 1914**

Distribution: S Armenia, N Turkey.

***Diorhabda persica* (FALDERMANN, 1837)***Galeruca persica* FALDERMANN, 1837*Galeruca sareptana* STIERLIN, 1863

Distribution: SE of European Russia, Daghestan, Kazakhstan, Central Asia, Near and Middle East.

***Diorhabda rickmersi* J. WEISE, 1900***Lochmaea ornaticollis* REITTER, 1900*Plassiola pamirica* MANDL, 1968

Distribution: Tajikistan.

***Diorhabda rybakovi* J. WEISE, 1890**

Distribution: SE Kazakhstan, NW China.

***Diorhabda tarsalis* J. WEISE, 1889**

Distribution: Asian Russia (S Siberia), Mongolia, NW China.

**Genus: *Galerucella* CROTCH, 1873****Subgenus: *Galerucella* (*Galerucella*) CROTCH, 1873*****Galerucella* (*Galerucella*) *aquatica* (GEOFFROY in FOURCROY, 1785)***Chrysomela aquatica* GEOFFROY in FOURCROY, 1785

Distribution: Europe.

***Galerucella* (*Galerucella*) *grisescens* (JOANNIS, 1866)***Galeruca grisescens* JOANNIS, 1866*Galleruca distincta* BALY, 1874*Galleruca vittaticollis* BALY, 1874*Galerucella sagittariae* J. WEISE, 1886, non GYLLENHAL, 1813*Galerucella reducta* CHEN, 1942

Distribution: NW, Central and E Europe, Asian Russia (S Siberia), N China.

***Galerucella* (*Galerucella*) *nipponensis* LABOISSIÈRE, 1922***Galleruca sagittariae* BALY, 1874*Galerucella paludosa* J. WEISE, 1922

Distribution: Asian Russia (Maritime Prov.), Korea, Japan.

***Galerucella* (*Galerucella*) *nymphaeae* (LINNAEUS, 1758)***Chrysomela nymphaeae* LINNAEUS, 1758*Chrysomela polygonata* LAICHARTING, 1781*Galleruca luctuosa* MANNERHEIM, 1852

Distribution: Europe, Central Asia, Asian Russia (S Siberia), Mongolia, China, Japan, N America.

***Galerucella* (*Galerucella*) *sagittariae* (GYLLENHAL, 1813)***Galleruca sagittariae* GYLLENHAL, 1813

Distribution: Europe, Asian Russia (S Siberia).

**Subgenus: *Galerucella* (*Neogalerucella*) CHÛJÔ, 1962**

***Galerucella* (*Neogalerucella*) *calmariensis* (LINNAEUS, 1767)**

*Chrysomela calmariensis* LINNAEUS, 1767

*Galeruca aquatica* O. F. MÜLLER, 1776

*Galleruca lythri* GYLLENHAL, 1813

*Galerucella calmariensis* ab. *nigrina* DELAHON, 1913

Distribution: Europe except for N and SW, Caucasus, Turkey, Asian Russia (S Siberia), Mongolia, China, Japan, N Africa, introduced into North America.

***Galerucella* (*Neogalerucella*) *lineola* (FABRICIUS, 1781)**

*Crioceris lineola* FABRICIUS, 1781

*Adimonia verna* LAICHARTING, 1781

Distribution: Europe, Asian Russia (S Siberia), Mongolia, China, Japan, N Africa.

***Galerucella* (*Neogalerucella*) *pusilla* (DUFTSCHMIDT, 1825)**

*Galleruca pusilla* DUFTSCHMIDT, 1825

Distribution: S, Central, and E Europe, Turkey, Central Asia, China.

***Galerucella* (*Neogalerucella*) *tenella* (LINNAEUS, 1761)**

*Chrysomela tenella* LINNAEUS, 1761

*Chrysomela parva* HERBST, 1783

*Lochmaea semenovi* RYBAKOV, 1889

*Galerucella sibirica* CSIKI, 1901

*Galerucella jakovlevi* CSIKI, 1901

*Galerucella tenella* ab. *melanaria* EVERTS, 1920

Distribution: N, Central, and E Europe, Asian Russia (S Siberia), China, introduced into North America.

**Genus: *Xanthogaleruca* LABOISSIÈRE, 1934**

***Xanthogaleruca* *aenescens* (FAIRMAIRE, 1878)**

*Galerucella aenescens* FAIRMAIRE, 1878

Distribution: Asian Russia (Maritime Prov.), N China.

***Xanthogaleruca* *flavescens* J. WEISE, 1887**

*Galerucella flavescens* J. WEISE, 1887

*Xanthogaleruca konishii* KIMOTO, 1964

Distribution: Asian Russia (Amurland, Maritime Prov.).

***Xanthogaleruca* *luteola* (O. F. MÜLLER, 1776)**

*Chrysomela luteola* O. F. MÜLLER, 1776

*Crioceris calmariensis* FABRICIUS, 1775, non LINNAEUS, 1767

*Chrysomela xanthomelanea* SCHRANK, 1781

*Galerucella crataegi* JOANNIS, 1866

*Galerucella luteola* ab. *obscuridorsis* ROUBAL, 1926

*Galerucella luteola* ab. *nigra* CSIKI, 1953

Distribution: Europe except for N, Caucasus, Turkey, N Africa, Central Asia (Oases), introduced to N America.

***Xanthogaleruca* *maculicollis* (MOTSCHULSKY, 1853)**

*Galeruca maculicollis* MOTSCHULSKY, 1853

*Galerucella vagepicta* J. WEISE, 1924

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Japan.

**Genus: *Pyrrhalta* JOANNIS, 1866*****Pyrrhalta viburni annulicornis* BALY, 1874**

Distribution: Asian Russia (Maritime Prov.), N China, Japan.

***Pyrrhalta viburni viburni* (PAYKULL, 1799)**

*Galeruca viburni* PAYKULL, 1799

Distribution: Europe except for N and S, Caucasus, Kazakhstan, Asian Russia (W Siberia), N Africa, introduced into North America.

**Genus: *Tricholochmaea* LABOISSIÈRE, 1932*****Tricholochmaea semifulva* (JACOBY, 1885)**

*Galerucella modesta* JACOBY, 1885

*Galerucella signaticeps* J. WEISE, 1887

*Galerucella japonica* J. WEISE, 1922

Distribution: Asian Russia (Altai, Sayan, Far East).

**Genus: *Lochmaea* J. WEISE, 1883*****Lochmaea capreae capreae* (LINNAEUS, 1758)**

*Chrysomela capreae* LINNAEUS, 1758

*Adimonia polygonata* LAICHARTING, 1781

*Chrysomela pallescens* GMELIN, 1790

*Crioceris pallida* HERBST, 1783

*Galeruca saturata* STEPHENS, 1832

*Adimonia scutellata* CHEVROLAT, 1840

*Adimonia capreae* ab. *pallidipennis* KÜSTER, 1847

*Lochmaea capreae* ab. *luctuosa* J. WEISE, 1886

*Lochmaea capreae* ab. *reitteri* LABOISSIÈRE, 1914

*Lochmaea capreae* ab. *circumflexa* BURLINI, 1946

Distribution: Europe, Transcaucasia, Turkey, N Kazakhstan.

***Lochmaea capreae cribrata* SOLSKY, 1872**

Distribution: Asian Russia (Transbaikalia, S Yakutia, Far East), Mongolia, China, Korea.

***Lochmaea crataegi* (FORSTER, 1771)**

*Chrysomela crataegi* FORSTER, 1771

*Chrysomela sanguinea* FABRICIUS, 1775

*Chrysomela femoralis* DUFTSCHMIDT, 1825

*Chrysomela tibialis* DUFTSCHMIDT, 1825

*Chrysomela femoralis* ab. *binotata* DUFTSCHMIDT, 1825

*Lochmaea crataegi* ab. *pallida* JOANNIS, 1866, non HERBST, 1783

*Lochmaea crataegi* ab. *orientalis* J. WEISE, 1924

*Lochmaea crataegi* ab. *nigronotata* PIC, 1927

*Lochmaea crataegi* ab. *lineata* PIC, 1927

*Lochmaea crataegi* ab. *flava* DONISTHORPE, 1934

*Lochmaea crataegi* ab. *obscura* KASZAB, 1962

Distribution: All over Palaeartic except for extreme N.

***Lochmaea suturalis* (C. G. THOMSON, 1866)***Galeruca suturalis* C. G. THOMSON, 1866*Lochmaea suturalis* var. *nigrita* J. WEISE, 1886

Distribution: Europe except for N and S.

**Genus: *Galeruca* GEOFFROY, 1762***Adimonia* LAICHARTING, 1781*Galleruca* FABRICIUS, 1792*Ademonia* AGASSIZ, 1846**Subgenus: *Galeruca (Galeruca)* GEOFFROY, 1762*****Galeruca (Galeruca) circassica* REITTER, 1903**

Distribution: N Caucasus, Abkhasia.

***Galeruca (Galeruca) dahli dahli* (JOANNIS, 1866)***Adimonia dahlii* JOANNIS, 1866

Distribution: S and Central Europe, Moldavia, S Ukraine, Crimea, Asian Russia (W Siberia, Altai, Cisbaikalia).

***Galeruca (Galeruca) dahli vicina* SOLSKY, 1872**

Distribution: Asian Russia (Transbaikalia, Far East), Mongolia.

***Galeruca (Galeruca) daurica* (JOANNIS, 1866)***Adimonia daurica* JOANNIS, 1866*Adimonia sedakovi* JOANNIS, 1866*Galeruca mongolica* CSIKI, 1901

Distribution: Asian Russia (Sayan, Cisbaikalia, Transbaikalia, Yakutia, Far East), Mongolia.

***Galeruca (Galeruca) extensa* (MOTSCHULSKY, 1861)***Galleruca extensa* MOTSCHULSKY, 1861*Galeruca banghaasi* J. WEISE, 1894

Distribution: Asian Russia (S Maritime Prov.), China, Korea, Japan.

***Galeruca (Galeruca) heydeni* J. WEISE, 1887**

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea.

***Galeruca (Galeruca) hyrcana* L. MEDVEDEV, 1969**

Distribution: Azerbaijan.

***Galeruca (Galeruca) interrupta interrupta* (ILLIGER, 1802)***Galleruca interrupta* ILLIGER, 1802*Galeruca jucunda* FALDERMANN, 1837*Adimonia oelandica* BOHEMAN, 1850*Adimonia hungarica* FRIVALDSZKY, 1876

Distribution: Europe except for N, Turkey, Syria, N Africa.

***Galeruca (Galeruca) interrupta armeniaca* J. WEISE, 1886**

Distribution: Transcaucasica, Turkmenistan, Uzbekistan, Tajikistan.

***Galeruca (Galeruca) interrupta circumdata* (DUFTSCHMIDT, 1825)***Galleruca circumdata* DUFTSCHMIDT, 1825*Adimonia florentina* L. REDTENBACHER, 1849*Galeruca sequensi* REITTER, 1903

Distribution: S, SE and Central Europe, Asian Russia (S Siberia), Mongolia.

***Galeruca (Galeruca) interrupta fulvimargo* REITTER, 1908**

Distribution: Mountains of S Kazakhstan, E Central Asia, NW China, Korea.

***Galeruca (Galeruca) laticollis* C. R. SAHLBERG, 1838***Adimonia villae* COMOLLI, 1837*Adimonia fontinalis* BOHEMAN, 1851*Adimonia flava* KÜSTER, 1854

Distribution: N Europe, Mountains of Central and E Europe, NE Kazakhstan, Siberia, Mongolia.

***Galeruca (Galeruca) pomonae petschenega* JACOBSON, 1925**

Distribution: SE European Russia (S Cisvolga), E Daghestan.

***Galeruca (Galeruca) pomonae pomonae* (SCOPOLI, 1763)***Tenebrio pomonae* SCOPOLI, 1763*Chrysomela rustica* SCHALLER, 1783*Adimonia gredleri* JOANNIS, 1866*Adimonia brevis* JOANNIS, 1866*Adimonia rufescens* JOANNIS, 1866*Galeruca pomonae* var. *anthracina* J. WEISE, 1886*Galeruca pomonae* var. *alpestris* J. WEISE, 1886*Galeruca pomonae* var. *cretica* J. WEISE, 1889*Galeruca circassica* REITTER, 1889

Distribution: Europe, Kazakhstan, Central Asia, Asian Russia (W Siberia, Yakutia, Altai), Mongolia.

***Galeruca (Galeruca) reichardti* JACOBSON, 1925**

Distribution: Asian Russia (Maritime Prov.), China, Korea.

***Galeruca (Galeruca) spectabilis spectabilis* FALDERMANN, 1837**

Distribution: Transcaucasia, N Iran.

***Galeruca (Galeruca) spectabilis lacericollis* SEMENOV, 1909**

Distribution: E Azerbaijan (Talysh), N Iran, S Turkmenistan.

***Galeruca (Galeruca) spectabilis orientalis* OSCULATI, 1844**

Distribution: Turkey, Syria, W Iran, ?Transcaucasia.

***Galeruca (Galeruca) tanaceti tanaceti* (LINNAEUS, 1758)***Chrysomela tanaceti* LINNAEUS, 1758*Tenebrio tristis* SCOPOLI, 1763*Galleruca declivis* JOANNIS, 1866*Galleruca bonvouloiri* JOANNIS, 1866*Galeruca tanaceti* ab. *rufifrons* LABOISSIÈRE, 1929? *Galeruca hunyadensis* CSIKI, 1953

Distribution: Europe, Caucasus, Turkey, Kazakhstan, Central Asia, Asian Russia (W Siberia, Altai).

***Galeruca (Galeruca) tanaceti convexa* JACOBSON, 1925**

Distribution: Mountains of Ciscaucasia, Armenia.

***Galeruca (Galeruca) tanaceti incisicollis* (MOTSCHULSKY, 1860)***Galleruca incisicollis* MOTSCHULSKY, 1860

Distribution: Asian Russia (Siberia, Far East), Mongolia.

***Galeruca (Galeruca) weisei* REITTER, 1903**

Distribution: SE Kazakhstan, Asian Russia (Siberia, Altai, Far East), Mongolia.

**Subgenus: *Galeruca (Haptoscelis)* J. WEISE, 1886*****Galeruca (Haptoscelis) melanocephala melanocephala* (PONZA, 1805)***Chrysomela melanocephala* PONZA, 1805*Hispa aptera* BONELLI, 1812*Galeruca haematidia* GERMAR, 1824*Lochmaea kubanensis* FORMANEK, 1900*Haptoscelis melanocephala* var. *baltica* J. WEISE, 1904Distribution: S and Central Europe, Belarus, Ukraine, S European Russia, Ciscaucasia, ?  
W Siberia.**Subgenus: *Galeruca (Emarhopa)* J. WEISE, 1886*****Galeruca (Emarhopa) rufa* GERMAR, 1824**

Distribution: S and SE Europe, Ukraine, Crimea, S European Russia, Turkey.

**Genus: *Theone* GISTEL, 1857***Leptosonyx* J. WEISE, 1885*Leptonyx* JACOBSON, 1895***Theone costipennis* (KIRSCH, 1880)***Leptosonyx costipennis* KIRSCH, 1880*Theone filicornis* JAKOB, 1957*Theone filicornis bucharica* MANDL, 1968

Distribution: W Daghestan, Turkmenistan, Uzbekistan.

***Theone margelanica* (KRAATZ, 1882)***Galeruca margelanica* KRAATZ, 1882

Distribution: Uzbekistan, Kighizstan (Fergana Valley).

***Theone octocostata octocostata* (J. WEISE, 1912)***Leptosonyx octocostata* J. WEISE, 1912

Distribution: Pamirs, N India and Pakistan.

***Theone octocostata lopatini* MANDL, 1970**

Distribution: E Tajikistan (Badakhshan).

***Theone silphoides silphoides* (DALMAN, 1823)***Adimonia silphoides* DALMAN, 1823

Distribution: SE European Russia, Daghestan, Steppe of Kazakhstan.

***Theone silphoides artemisiae* (JACOBSON, 1895)***Leptosonyx artemisiae* JACOBSON, 1895

Distribution: Tajikistan.

***Theone silphoides kuldshensis* MANDL, 1968**

Distribution: E Kazakhstan, NW China.

***Theone silphoides scrobiculata* MANDL, 1968**

Distribution: E Kazakhstan (Tarbagatay Mountain Range).

**Genus: *Pallasiola* JACOBSON, 1925***Pallasia* J. WEISE, 1886***Pallasiola absinthii* (PALLAS, 1773)***Chrysomela absinthii* PALLAS, 1773

Distribution: N Kazakhstan, Kirghizstan, Pamirs, Asian Russia (S Siberia), N Mongolia, N China.

**Genus: *Nyctiphantus* SEMENOV, 1902***Nyctidromas* SEMENOV, 1895***Nyctiphantus bergi* SEMENOV, 1904**

Distribution: Kazakhstan (Shores of Balkhash Lake).

***Nyctiphantus custos* SEMENOV, 1902**

Distribution: S Kazakhstan.

***Nyctiphantus hirtus hirtus* (J. WEISE, 1885)***Leptosonyx hirtus* J. WEISE, 1885

Distribution: S and E Kazakhstan, NW China.

***Nyctiphantus hirtus nocturnus* (SEMENOV, 1891)***Leptosonyx nocturnus* SEMENOV, 1891

Distribution: Turkmenistan (Karakum Desert).

**Genus: *Apophylia* J. THOMSON, 1858***Malaxia* FAIRMARE, 1878*Galerucesthis* J. WEISE, 1896***Apophylia eoa* OGLOBLIN, 1936**

Distribution: Asian Russia (Maritime Prov.), Korea.

***Apophylia thalassina* (FALDERMANN, 1835)***Auchenia thalassina* FALDERMANN, 1835

Distribution: Asian Russia (Transbaikalia, Amurland), Mongolia, China.



Tribe: *Sermylassini* MROCKOWSKI, 1991

*Sermylini* CHAPUIS, 1875  
*Agelasini* LENG, 1920

Genus: *Sermylassa* REITTER, 1912

*Sermyla* CHAPUIS, 1875, non WALKER, 1854 (*Lepidoptera*)  
*Agelasa* G. HORN, 1893, non MOTSCHULSKY, 1860

***Sermylassa halensis* (LINNAEUS, 1767)**

*Chrysomela halensis* LINNAEUS, 1767  
*Crioceris nigricornis* FABRICIUS, 1775  
*Sermyla halensis* var. *cuprina* J. WEISE, 1886  
*Sermyla halensis* ab. *picea* LABOISSIÈRE, 1911

Distribution: Central and E Europe, Balkans, Kazakhstan, Asian Russia (Urals, W and Middle Siberia, Altais).

Tribe: *Agelasticini* CHAPUIS, 1875Genus: *Agelasa* MOTSCHULSKY, 1860***Agelasa nigriceps* MOTSCHULSKY, 1860**

Distribution: Asian Russia (Maritime Prov.), N China, Japan.

Genus: *Gallerucida* MOTSCHULSKY, 1860***Gallerucida bifasciata* MOTSCHULSKY, 1860**

*Melospila nigromaculata* BALY, 1861  
*Gallerucida nigrofasciata* BALY, 1879  
 Distribution: Asian Russia (S Amurland, Maritime Prov.), China, Korea, Japan.

***Gallerucida flavipennis* SOLSKY, 1872**

Distribution: Asian Russia (Maritime Prov.), E China.

***Gallerucida gloriosa* (BALY, 1861)**

*Eustetha gloriosa* BALY, 1861  
*Gallerucida jacobsoni* OGLOBLIN, 1936  
 Distribution: Asian Russia (Maritime Prov.), N and Central China.

Genus: *Agelastica* CHEVROLAT in DEJEAN, 1837***Agelastica alni alni* (LINNAEUS, 1758)**

*Chrysomela alni* LINNAEUS, 1758  
*Adimonia violacea* LAICHARTING, 1781  
*Agelastica alni* ab. *atra* TENENBAUM, 1927  
 Distribution: All over Palearctic except for Central Asia. Introduced to N America.

***Agelastica alni orientalis* BALY, 1878**

Distribution: Central Asia, SE Kazakhstan, Mongolia, W China.

***Agelastica coerulea* BALY, 1874**

Distribution: Asian Russia (Maritime Prov.), N China, Korea, Japan, N America.

**Genus: *Morphosphaera* BALY, 1861*****Morphosphaera japonica* (HORNSTEDT, 1788)***Chrysomela japonica* HORNSTEDT, 1788

Distribution: Asian Russia (S Amurland), China, Japan, India.

Tribe: *Luperini* LENG, 1920Subtribus: *Aucacophorina* CHAPUIS, 1875**Genus: *Aulacophora* CHEVROLAT in DEJEAN, 1837***Rhaphidopalpa* ROSENHAUER, 1856***Aulacophora foveicollis* (H. LUCAS, 1849)***Galeruca foveicollis* H. LUCAS, 1849

Distribution: S Europe (Mediterranean), Near and Middle East, Turkey, Transcaucasia, India, Burma, Sri Lanka.

***Aulacophora nigripennis* MOTSCHULSKY, 1857**

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

Subtribus: *Luperina* WILCOX, 1965**Genus: *Medythia* JACOBY, 1887***Paraluperodes* OGLOBLIN, 1936***Medythia suturalis nigrobilineatus* (MOTSCHULSKY, 1860)***Cnecodes nigrobilineatus* MOTSCHULSKY, 1860

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

***Medythia suturalis suturalis* (MOTSCHULSKY, 1860)***Cnecodes suturalis* MOTSCHULSKY, 1860

Distribution: S Asia.

**Genus: *Tschitscherinula* JACOBSON, 1908*****Tschitscherinula paradoxocara* JACOBSON, 1908**

Distribution: Uzbekistan, Turkmenistan, S Tajikistan.

**Genus: *Stenoluperus* OGLOBLIN, 1936*****Stenoluperus cyaneus* (BALY, 1874)***Arthrotus cyaneus* BALY, 1874

Distribution: Asian Russia (S Kuriles Isles), Japan.

***Stenoluperus nipponensis* (LABOISSIÈRE, 1913)***Luperus nipponensis* LABOISSIÈRE, 1913*Luperus longicornis* JACOBY, 1885, non FABRICIUS, 1781*Luperus jacobyi* J. WEISE, 1924

Distribution: Asian Russia (Amurland, Maritime Prov., Sakhalin, S Kuriles Isles), China, Korea, Japan.

**Genus: *Nymphius* J. WEISE, 1900*****Nymphius ogloblini* (BOGATCHEV, 1947)***Luperus ogloblini* BOGATCHEV, 1947

Distribution: Armenia, N Turkey.

***Nymphius pravei* (JACOBSON, 1899)***Luperus pravei* JACOBSON, 1899

Distribution: S Ukraine, SE of European Russia, NW Kazakhstan.

***Nymphius stylifer* (J. WEISE, 1899)***Luperus stylifer* J. WEISE, 1899

Distribution: Georgia, Armenia.

**Genus: *Phyllobrotica* CHEVROLAT in DEJEAN, 1837*****Phyllobrotica adusta* (CREUTZER, 1799)***Crioceris adusta* CREUTZER, 1799*Phyllobrotica adusta* ab. *litoralis* G. MÜLLER, 1924

Distribution: Central and SE Europe.

***Phyllobrotica elegans* KRAATZ, 1866***Phyllobrotica trimaculata* BALLION, 1890

Distribution: Ciscaucasia, S of Steppe Zone of Ukraine and European Russia, Crimea, Transcaucasia, Turkey.

***Phyllobrotica quadrimaculata* (LINNAEUS, 1758)***Chrysomela 4-maculata* LINNAEUS, 1758*Chrysomela melanogaster* GMELIN, 1790 (part.)*Phyllobrotica quadrimaculata* var. *munda* J. WEISE, 1886

Distribution: Europe except for Mediterranean, Ciscaucasia, S Siberia, E Kazakhstan.

***Phyllobrotica signata* (MANNERHEIM, 1825)***Galeruca signata* MANNERHEIM, 1825

Distribution: Asian Russia (S Siberia, Altai, Sayan, Amurland, Maritime Prov.), Mongolia, N and W China.

**Genus: *Euliroetis* OGLOBLIN, 1936*****Euliroetis obscuripes* OGLOBLIN, 1936**

Distribution: Asian Russia (S Amurland, Maritime Prov.).

***Euliroetis ornata* (BALY, 1874)***Phyllobrotica ornata* BALY, 1874

Distribution: Asian Russia (S Maritime Prov.), Korea, China, Japan.

**Genus: *Exosoma* JACOBY, 1903***Malacosoma* CHEVROLAT, 1845, non HÜBNER, 1816***Exosoma collare* (HUMMEL, 1825)***Luperus collaris* HUMMEL, 1825*Galleruca luteicollis* GEBLER, 1830*Galleruca cyanoptera* KRYNICKI, 1832*Malacosoma triumphans* FALDERMANN, 1837*Malacosoma collare* ab. *russica* PIC, 1915

Distribution: Ukraine, S and E of European Russia, Caucasus, Syria, N Kazakhstan, SW Siberia.

***Exosoma flavipes* (HEYDEN, 1878)***Malacosoma flavipes* HEYDEN, 1878

Distribution: Armenia, Turkey.

**Genus: *Euluperus* J. WEISE, 1886*****Euluperus xanthopus xanthopus* (DUFTSCHMIDT, 1825)***Crioceris xanthopa* DUFTSCHMIDT, 1825*Luperus fossulatus* JOANNIS, 1866*Luperus violaceus* HAROLD, 1876*Euluperus xanthopus* ab. *fuscipes* J. WEISE, 1886

Distribution: Southern Steppe of Ukraine and European Russia, NW Kazakhstan, W Transcaucasia.

***Euluperus xanthopus virescens* J. WEISE, 1886**

Distribution: E Transcaucasia.

**Genus: *Falsoexosoma* PIC, 1926*****Falsoexosoma cyanipennis* (REITTER, 1902)***Malacosoma cyanipennis* REITTER, 1902

Distribution: Azerbaijan, Turkmenistan (Kopetdagh), Iran.

**Genus: *Calomicrus* STEPHENS, 1832***Lyperus* BILLBERG, 1820***Calomicrus altaicus altaicus* (MANNERHEIM, 1825)***Luperus altaicus* MANNERHEIM, 1825

Distribution: Asian Russia (S and Middle Siberia, Far East, Amurland), N Mongolia.

***Calomicrus altaicus eous* OGLOBLIN, 1936**

Distribution: Asian Russia (Maritime Prov.).

***Calomicrus angorensis* PIC, 1913**

Distribution: Transcaucasia, Turkey.

***Calomicrus circumfusus* (MARSHAM, 1802)***Crioceris circumfusa* MARSHAM, 1802

*Haltica spartii* KOCH, 1803

*Luperus nigrofasciatus* J. WEISE, 1886

*Luperus circumfusus* ab. *nigripennis* BEDEL, 1891

*Luperus circumfusus* ab. *bipunctata* LABOISSIÈRE, 1912

*Luperus circumfusus* ab. *bedeli* J. WEISE, 1924

Distribution: Europe, W Ukraine, Belarus, W of European Russia, Moldavia, N Africa (Tunisia).

***Calomicrus darvasicus* LOPATIN, 1975**

Distribution: E Tajikistan.

***Calomicrus deserticola* OGLOBLIN, 1936**

Distribution: Turkmenistan.

***Calomicrus ghilarovi* LOPATIN, 1988**

Distribution: S Tajikistan (Babatagh Mountain Range).

***Calomicrus grandis* (JACOBSON, 1894)**

*Luperus grandis* JACOBSON, 1894

Distribution: Kirghiztan, SE Kazakhstan.

***Calomicrus gussakovskiy* OGLOBLIN, 1936**

Distribution: S Tajikistan.

***Calomicrus hissaricus* OGLOBLIN, 1936**

Distribution: Central Tajikistan.

***Calomicrus minutus* JOANNIS, 1866**

Distribution: Asian Russia (Southern E Siberia, Khabarovsk Distr., Maritime Prov.).

***Calomicrus nairicus* LOPATIN, 1990**

Distribution: Armenia.

***Calomicrus palii* LOPATIN, 1965**

Distribution: Kirghizstan.

***Calomicrus pinicola* (DUFTSCHMIDT, 1825)**

*Crioceris pinicola* DUFTSCHMIDT, 1825

*Luperus pinicola* var. *silvestris* J. WEISE, 1886

Distribution: Europe, W Ukraine, Northern and Central parts of European Russia.

***Calomicrus populi* LOPATIN, 1963**

Distribution: Central and SE Tajikistan.

***Calomicrus sericeus* (JACOBSON, 1894)**

*Luperus sericeus* JACOBSON, 1894

Distribution: S Kazakhstan, Kirghizstan (Tian-Shan).

***Calomicrus sugonjaevi* LOPATIN, 1983**

Distribution: Turkmenistan (Kopetdagh).

**Genus: *Luperus* GEOFFROY, 1762***Luperus* BEDEL, 1892***Luperus anthracinus* OGLOBLIN, 1936**

Distribution: S Asian Russia (Altai), Mongolia.

***Luperus armeniacus* KIESENWETTER, 1878**

Distribution: Bulgaria, S Crimea, Transcaucasia.

***Luperus caucasicus* J. WEISE, 1879***Luperus circassicus* L. MEDVEDEV, 1962

Distribution: Caucasus.

***Luperus cyanipennis* KÜSTER, 1848**

Distribution: S and SW Ukraine, Crimea, Moldavia, Balkans.

***Luperus discolor* FALDERMANN, 1837**

Distribution: Transcaucasia.

***Luperus ehnerbergi* JACOBSON, 1901**

Distribution: Kirghizstan, SE Kazakhstan.

***Luperus flavilabris* OGLOBLIN, 1941**

Distribution: Tajikistan.

***Luperus flavipes flavipes* (LINNAEUS, 1767)***Chrysomela flavipes* LINNAEUS, 1767*Chrysomela ochropus* GMELIN, 1790*Luperus carniolicus* KIESENWETTER, 1861

Distribution: All over Palaeartic except for Central Asia.

***Luperus flavipes obscuricornis* OGLOBLIN, 1936**

Distribution: E Asian Russia (Far East).

***Luperus floralis* FALDERMANN, 1837**

Distribution: Azerbaijan.

***Luperus kiesenwetteri* JOANNIS, 1866**

Distribution: SE of European Russia (Steppe zone), N Kazakhstan, Transcaucasia.

***Luperus kiritschenkoi* OGLOBLIN, 1936**

Distribution: Crimea.

***Luperus longicornis longicornis* (FABRICIUS, 1781)***Ptinus longicornis* FABRICIUS, 1781*Luperus betulinus* JOANNIS, 1866, non GEOFFROY, 1785*Luperus diniensis* BELLIER DE LA CHAVIGNERIE, 1870*Luperus longicornis* ab. *subviridis* LABOISSIÈRE, 1912

Distribution: Europe, Caucasus, Asian Russia (Far East).

***Luperus longicornis tenebricosus* OGLOBLIN, 1936**

Distribution: Asian Russia (Yakutia).

***Luperus lyperus* (SULZER, 1776)**

*Chrysomela lyperus* SULZER, 1776  
*Chrysomela erythromelas* GMELIN, 1790  
*Chrysomela nigricans* GMELIN, 1790  
*Luperus lyperus* BEDEL, 1897  
*Luperus niger* GOEZE, 1777 (male)  
*Luperus rufipes* GOEZE, 1777 (female)  
*Luperus dispar* L. REDTENBACHER, 1874

Distribution: Central and E Europe, Asian Russia (W Siberia), Mongolia.

***Luperus margaritae* LOPATIN, 1990**

Distribution: Armenia.

***Luperus orientalis* FALDERMANN, 1837**

*Luperus amoenus* FALDERMANN, 1837  
*Luperus oxypterus* FALDERMANN, 1837

Distribution: Daghestan, Transcaucasia, W Turkmenistan (Kopetdagh).

***Luperus perlucidus* KHNZORIAN, 1956**

Distribution: Armenia.

***Luperus semiflavus* OGLOBLIN, 1936**

*Luperus mandzhuricus* OGLOBLIN, 1936  
 Distribution: NE China.

***Luperus similis* IABLOKOFF-KHNZORIAN, 1970**

Distribution: Asian Russia (Altai).

***Luperus turkestanicus* J. WEISE, 1892**

Distribution: Kirghizstan.

***Luperus viridipennis viridipennis* GERMAR, 1824**

*Crioceris coerulea* DUFTSCHMIDT, 1825  
*Luperus viridipennis* ab. *varicollis* J. WEISE, 1886  
*Luperus viridipennis* ab. *rhenanus* STIERLIN, 1890  
*Luperus viridipennis* ab. *pallidicollis* GUILLEBEAU, 1891  
*Luperus viridipennis* ab. *maritimus* J. DANIEL, 1903  
*Luperus viridipennis* ab. *winklerii* G. MÜLLER, 1949  
*Luperus viridipennis* ab. *gredleri* G. MÜLLER, 1949

Distribution: Mountains of Europe, W Ukraine (Carpathians), Forest and Forest-Steppe zone of European Russia.

***Luperus viridipennis laricis* (MOTSCHULSKY, 1859)**

*Luperus laricis* MOTSCHULSKY, 1859  
*Luperus viridipennis murakamii* KIMOTO, 1965  
*Luperus viridipennis spurius* OGLOBLIN, 1936

Distribution: Asian Russia (Middle Siberia, Amurland), Mongolia, Japan.

***Luperus xanthopoda* (SCHRANK, 1781)**

*Chrysomela xanthopoda* SCHRANK, 1781  
*Luperus xanthopus* SCHRANK, 1785

Distribution: Southern parts of Central and E Europe, Ciscaucasia, Kirghizstan, SE Kazakhstan.

**Genus: *Atrachya* DEJEAN, 1837***Cnecodes* MOTSCHULSKY, 1858***Atrachya menetriesi* (FALDERMANN, 1835)***Galeruca menetriesi* FALDERMANN, 1835

Distribution: Asian Russia (Amurland, Maritime Prov., Sakhalin, S Kuriles Isles), Korea, China, Japan.

**Genus: *Monolepta* CHEVROLAT in DEJEAN, 1837***Ochralea* CLARK, 1865*Candezea* CHAPUIS, 1880***Monolepta dichroa dichroa* HAROLD, 1877**

Distribution: Asian Russia (Sakhalin), Japan.

***Monolepta dichroa kurilensis* L. MEDVEDEV, 1966**

Distribution: Asian Russia (Kunashir Isle).

***Monolepta eoa* OGLOBLIN, 1936**

Distribution: Asian Russia (Maritime Prov.).

***Monolepta hieroglyphica biarcuata* J. WEISE, 1889***Monolepta biarcuata* J. WEISE, 1889

Distribution: Asian Russia (Maritime Prov.), Korea, China, Japan.

***Monolepta nebulosa* OGLOBLIN, 1936**

Distribution: Asian Russia (Maritime Prov.).

***Monolepta pallidula* (BALY, 1874)***Luperodes pallidulus* BALY, 1874

Distribution: Asian Russia (S Maritime Prov.), China, Japan.

***Monolepta quadriguttata* (MOTSCHULSKY, 1860)***Luperodes quadriguttatus* MOTSCHULSKY, 1860

Distribution: Asian Russia (Amurland, Maritime Prov.), Mongolia, Korea, China, Japan.

***Monolepta russica* (GMELIN, 1790)***Chrysomela russica* GMELIN, 1790*Chrysomela ruficollis* FABRICIUS, 1787*Luperodes dubius* MOTSCHULSKY, 1838*Monolepta angustula* J. WEISE, 1889

Distribution: E Ciscaucasia, Kazakhstan, Plains of Central Asia.

***Monolepta semenovi* OGLOBLIN, 1936**

Distribution: Asian Russia (S Maritime Prov.), China.

***Monolepta subseriata* J. WEISE, 1887**

Distribution: Asian Russia (Transbaikalia, Amurland, Maritime Prov.).



**Genus: *Fleutiauxia* LABOISSIÈRE, 1933*****Fleutiauxia armata* (BALY, 1874)***Aenidea armata* BALY, 1874*Aenidea armata koltzei* HEYDEN, 1893

Distribution: Asian Russia (Amurland, Maritime Prov.), Korea, China, Japan.

Subfamily: *Alticinae* NEWMAN, 1834**Genus: *Phyllotreta* CHEVROLAT in DEJEAN, 1837***Orchestris* CROTCH, 1873*Tanygaster* BLATCHLEY, 1921***Phyllotreta acutecarinata* HEIKERTINGER, 1941**

Distribution: Southern parts of Central Europe, Balkans, S Ukraine, SE European Russia.

***Phyllotreta andreevae* LOPATIN, 1992**

Distribution: E Tajikistan (Badakhshan).

***Phyllotreta annae* KONSTANTINOV, 1992**

Distribution: Kazakhstan (Karatau Mts.).

***Phyllotreta araxicola* IABLOKOFF-KHNZORIAN, 1968**

Distribution: Armenia, Azerbaijan.

***Phyllotreta armoraciae* (KOCH, 1803)***Haltica armoraciae* KOCH, 1803*Altica nemorum* FABRICIUS, 1775*Altica vittata* STEPHENS, 1832*Phyllotreta armoraciae* var. *biplagiata* CHITTENDEN, 1927

Distribution: Europe, Kazakhstan, Central Asia, Siberia. Introduced to N America.

***Phyllotreta astrachanica* LOPATIN, 1977***Phyllotreta diademata astrachanica* LOPATIN, 1977*Phyllotreta astrachanica*: GRUEV, 1982*Phyllotreta diademata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to WARCHAŁOWSKI, 1978)

Distribution: Central, S and SE Europe, Transcaucasia, Turkey, NW Kazakhstan

***Phyllotreta atra* (FABRICIUS, 1775)***Altica atra* FABRICIUS, 1775*Chrysomela atra* FABRICIUS, 1781*Chrysomela pulex* SCHRANK, 1781*Altica aethiopissa* SCHRANK, 1789*Altica aterrima* SCHRANK, 1798*Crioceris atra* FABRICIUS, 1801*Chrysomela punctata* MARSHAM, 1802*Haltica melaena* ILLIGER, 1807*Haltica punctulata* STEPHENS, 1832*Haltica elongata* STEPHENS, 1834

*Phyllotreta obscurella* Foudras, 1860

*Phyllotreta atra* ab. *erdosi* Csiki, 1953

*Galeruca atra*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Fabricius, 1792 and Paykull, 1799)

*Haltica atra*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Koch, 1803)

Distribution: Europe, Caucasus, Turkey, Near and Middle East, Kazakhstan, Kirghizstan, Tajikistan, Asian Russia (S Siberia, Sakhalin), Mongolia.

***Phyllotreta austriaca* Heikertinger, 1909**

*Phyllotreta austriaca* var. *aligera* Heikertinger, 1909

*Phyllotreta austriaca aligera* Bechyne, 1956

Distribution: Southern parts of Central Europe, Ciscaucasia, Asian Russia (S Siberia, Amurland, western parts of Maritime Prov.), Korea.

***Phyllotreta bactriana* Heikertinger, 1941**

Distribution: Turkmenistan, S Tajikistan, SE Uzbekistan.

***Phyllotreta balcanica* Heikertinger, 1909**

*Phyllotreta foudrasi* Wanka, 1908

*Phyllotreta crassicornis* Bodemeyer, 1910

*Phyllotreta adanensis* Pic, 1919

*Phyllotreta anatolica* Pic, 1910

Distribution: S and southern parts of Central and E Europe, Caucasus, Daghestan, Turkey, Kazakhstan, Kirghizstan, Uzbekistan, Afghanistan.

***Phyllotreta banghaasi* Heikertinger, 1941**

Distribution: Turkmenistan, S Kazakhstan, Kirghizstan, Uzbekistan.

***Phyllotreta bartanga* Lopatin, 1966**

Distribution: Tajikistan (Badakhshan).

***Phyllotreta beschkentica* Lopatin, 1961**

Distribution: S Tajikistan.

***Phyllotreta brevistriata* Kimoto, 1966**

Distribution: S Kuriles Isles, Japan.

***Phyllotreta buchtarmensis* Lopatin, 1990**

Distribution: E Kazakhstan.

***Phyllotreta buhseae* Iablokoff-Khinzorian, 1978**

Distribution: Turkmenistan.

***Phyllotreta caucasicola* Heikertinger, 1941**

Distribution: Caucasus, Transcaucasia, Turkey, Middle East.

***Phyllotreta corrugata* Reiche, 1858**

*Phyllotreta corrugata* var. *galloprovincialis* Caillole, 1908

*Phyllotreta rufitarsis* var. *beauprei* Pic, 1909

*Phyllotreta discophora* Iablokov-Khinzorian, 1960

*Phyllotreta bella* Palij, 1970

Distribution: S Europe, Ukraine, Caucasus, Transcaucasia, Turkey, Near and Middle East, Turkmenistan, Kazakhstan, Uzbekistan.

***Phyllotreta cruciferae* (GOEZE, 1777)***Chrysomela cruciferae* GOEZE, 1777*Altica brassicae* GEOFFROY, 1785*Altica hortensis* OLIVIER, 1789*Haltica obscurella* ILLIGER, 1807*Phyllotreta nigroaenea* STEPHENS, 1832*Haltica elongata* STEPHENS, 1834*Phyllotreta poeciloceras* COMOLLI, 1837*Haltica punctipennis* WEIDENBACH, 1859*Phyllotreta poeciloceras* ALLARD, 1860*Phyllotreta colorea* FOUDRAS, 1860*Phyllotreta atra* var. *obscurella* REITTER, 1870*Phyllotreta atra* var. *cruciferae* HEIKERTINGER, 1930*Phyllotreta aerea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to DIRIMANOV et ANGELOVA, 1964)*Phyllotreta scheuchii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to WARCHALOWSKI, 1974)

Distribution: Europe except for Fennoscandia, Caucasus, Transcaucasia, Turkey, Near and Middle East, Central Asia, India, Mongolia, N Africa. Introduced to N America.

***Phyllotreta curvipes* HEIKERTINGER, 1941**

Distribution: S Kazakhstan.

***Phyllotreta dacica* HEIKERTINGER, 1941**

Distribution: Balkans, Carpathians, S Ukraine, Transcaucasia, Turkey.

***Phyllotreta diademata* FOUDRAS, 1860**

Distribution: Europe except for extreme N, Caucasus, Transcaucasia, Turkey, Middle East, Central Asia, N India.

***Phyllotreta dilatata* C. G. THOMSON, 1866***Phyllotreta tetrastigma* var. *dilatata* J. WEISE, 1888*Phyllotreta tetrastigma* ab. *dilatata* HEIKERTINGER et CSIKI, 1940*Phyllotreta tetrastigma dilatata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

Distribution: Central Europe, Asian Russia (Middle Siberia: Krasnoyarsk Distr.)

***Phyllotreta erysimi erysimi* J. WEISE, 1900***Phyllotreta schreineri*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KRÁL, 1973)

Distribution: SE Europe, Ukraine, S European Russia, Caucasus, Turkey, Middle East, Central Asia, Mongolia.

***Phyllotreta erysimi baicalica* HEIKERTINGER, 1941***Phyllotreta turkmenica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1939)*Phyllotreta erysimi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to SHAPIRO, 1974)

Distribution: Asian Russia (E Siberia), Mongolia.

***Phyllotreta erysimi iranella* LOPATIN, 1990**

Distribution: Iran.

***Phyllotreta erysimi kutscherai* HEIKERTINGER, 1941**

Distribution: Asian Russia (Siberia).

***Phyllotreta erysimi tekensis* LOPATIN, 1992**

Distribution: Turkmenistan.

***Phyllotreta exclamationis* (THUNBERG, 1784)***Chrysomela exclamationis* THUNBERG, 1784*Chrysomela brassicae* FABRICIUS, 1789*Galeruca quadripustulata* PAYKULL, 1799? *Crioceris brassicae* FABRICIUS, 1801*Chrysomela quadripustulata* MARSHAM, 1802*Haltica quadripustulata* KOCH, 1803? *Haltica quadriguttata* STEPHENS, 1832*Phyllotreta quadripustulata* KÜSTER, 1847*Phyllotreta exclamationis* var. *vibex* J. WEISE, 1888*Phyllotreta exclamationis* ab. *vibex*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1940)*Altica exclamationis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GMELIN, 1790)? *Altica brassicae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to OLIVIER, 1789 and FABRICIUS, 1792)*Haltica brassicae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ILLIGER, 1807)*Longitarsus quadripustulatus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BARGAGLI, 1878)

Distribution: Europe, Caucasus, Transcaucasia, Asian Russia (S Siberia, Far East).

***Phyllotreta ezoensis* KIMOTO, 1993***Phyllotreta atra*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KIMOTO, 1965)

Distribution: Asian Russia (S Kuriles Isle: Kunashir), Japan.

***Phyllotreta flexuosa* (ILLIGER, 1794)***Altica flexuosa* ILLIGER, 1794*Phyllotreta nemorum* PAYKULL, 1799*Haltica sinuata* STEPHENS, 1832*Phyllotreta fallax* ALLARD, 1860? *Phyllotreta tetrastigma* BARGAGLI, 1878*Phyllotreta flexuosa* var. *fenestrata* J. WEISE, 1888*Phyllotreta sinuata* HEIKERTINGER, 1910*Haltica flexuosa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ILLIGER, 1807)*Phyllotreta flexuosa* ab. *fenestrata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1940)

Distribution: All over Palaearctic except for E Mediterranean, Central Asia.

***Phyllotreta ganglbaueri* HEIKERTINGER, 1909**

Distribution: Southern parts of Central Europe, Balkans, Crimea.

***Phyllotreta gurskii* LOPATIN, 1966**

Distribution: Tajikistan (Badakhshan).

***Phyllotreta humilis* J. WEISE, 1887**

Distribution: Asian Russia (Transbaikalia, Amurland, Maritime Prov.), Mongolia, NE China, Korea.

***Phyllotreta judaea* PIC, 1901**

Distribution: E Bulgaria, Transcaucasia (Armenia), Turkey, Near and Middle East.

***Phyllotreta koltzei* J. WEISE, 1887**

Distribution: Asian Russia (Maritime Prov., Ussuri Area), Korea.

***Phyllotreta konevi* LOPATIN, 1985**

Distribution: Kazakhstan (Isle Barsakeljmes in Aral Sea).

***Phyllotreta krali* LOPATIN, 1990**

Distribution: Iran.

***Phyllotreta lativittata* KUTSCHERA, 1860**

*Phyllotreta lativittata* GEMMINGER et HAROLD, 1876

*Phyllotreta orientalis* BALY, 1877

*Phyllotreta latevittata* J. WEISE, 1888

*Phyllotreta ruficollis* J. WEISE, 1888

*Phyllotreta iris* REITTER, 1891

*Phyllotreta rubrithorax* PIC, 1900

*Phyllotreta ruficollis* var. *ininterrupta* PIC, 1909

*Phyllotreta latevittata* var. *bisbinotata* PIC, 1909

*Phyllotreta aegyptiaca* PIC, 1915

*Phyllotreta latevittata* ab. *orientalis* HEIKERTINGER et CSIKI, 1940

*Phyllotreta latevittata* ab. *bisbinotata* HEIKERTINGER et CSIKI, 1940

*Phyllotreta latevittata* ab. *viridicollis* HEIKERTINGER, 1941

*Phyllotreta latevittata* ab. *ruficollis* HEIKERTINGER, 1941

*Phyllotreta latevittata* ab. *vittipennis* HEIKERTINGER, 1941

*Phyllotreta latevittata* ab. *maculipennis* HEIKERTINGER, 1941

*Phyllotreta latevittata* ab. *nigrosuturata* HEIKERTINGER, 1941

Distribution: NE Mediterranean, Caucasus, Transcaucasia, Near and Middle East, Central Asia, China (Sinkiang).

***Phyllotreta lopatini* KONSTANTINOV, 1992**

Distribution: Azerbaijan.

***Phyllotreta lubischevi* LOPATIN, 1992**

Distribution: Kirghizstan.

***Phyllotreta maculosa* IABLOKOFF-KHNZORIAN, 1968**

Distribution: Armenia.

***Phyllotreta misella* JACOBSON, 1901**

*Phyllotreta austriaca aligera*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. MEDVEDEV et DUBESHKO, 1974)

Distribution: Asian Russia (Middle Siberia, Transbaikalia), Mongolia, N China.

***Phyllotreta mollis* KONSTANTINOV, 1992**

Distribution: Uzbekistan.

***Phyllotreta mongolica* L. MEDVEDEV, 1980**

Distribution: Mongolia.

***Phyllotreta nemorum* (LINNAEUS, 1758)**

*Chrysomela nemorum* LINNAEUS, 1758

? *Phyllotreta flexuosa* KOCH, 1803

*Phyllotreta nasturtii* PANZER, 1805

*Phyllotreta kuennemanni* REITTER, 1905  
*Phyllotreta nemorum* var. *theresae* PIC, 1909  
*Altica nemorum*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to O. F. MÜLLER, 1776)  
*Galleruca nemorum*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1794 et PAYKULL, 1799)  
*Crioceris nemorum*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1801)  
*Haltica flexuosa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ILLIGER, 1807) -  
*Haltica nemorum*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to OLIVIER, 1808)  
 Distribution: Europe, Caucasus, Transcaucasia, Turkey, Near and Middle East, Central Asia, Asian Russia (S Siberia to Amurland), Mongolia, Korea. Introduced to Australia.

***Phyllotreta nigripes* (FABRICIUS, 1775)**

*Altica nigripes* FABRICIUS, 1775  
*Chrysomela lens* THUNBERG, 1784  
 ? *Altica cruciferarum* GMELIN, 1790  
*Haltica lepidi* KOCH, 1803  
*Phyllotreta lepidii* KÜSTER, 1847  
*Phyllotreta nigripes* var. *arabidis* HOFFMANN, 1953  
*Altica lens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GMELIN, 1790)  
*Galleruca nigripes*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1792)  
*Haltica nigripes*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PANZER, 1794)  
*Chrysomela nigripes*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1781)  
*Haltica nigripes* var. *lens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GYLLENHAL, 1813)  
*Phyllotreta thalassicola*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. MEDVEDEV, 1983)  
 Distribution: Europe, Caucasus, Transcaucasia, Near and Middle East, Central Asia, N Africa.

***Phyllotreta nodicornis* (MARSHAM, 1802)**

*Chrysomela nodicornis* MARSHAM, 1802  
*Haltica antennata* KOCH, 1803  
*Phyllotreta oncera* MAULIK, 1926  
*Phyllotreta antennata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)  
 Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, N India.

***Phyllotreta ochripes* (CURTIS, 1837)**

*Haltica ochripes* CURTIS, 1837  
 ? *Haltica horticola* RICHTER, 1821  
*Phyllotreta excisa* L. REDTENBACHER, 1849  
 ? *Phyllotreta caucasica* HAROLD, 1875  
*Phyllotreta ochripes* var. *cruciata* J. WEISE, 1888  
 ? *Phyllotreta eximia* GUILLEBAEU, 1895  
*Phyllotreta ochripes* var. *burdigalensis* PIC, 1909  
*Phyllotreta ochripes* var. *comanensis* PIC, 1912  
*Phyllotreta kerkhoveni* EVERTS, 1919  
*Phyllotreta armoraciae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to to DIRIMANOV et ANGELOVA, 1964)  
 Distribution: Europe except for N, Caucasus, Transcaucasia, Iran, Asian Russia (S Siberia to Far East).

***Phyllotreta ogloblini* SHAPIRO, 1960**

Distribution: S Ukraine.

***Phyllotreta pallidipennis* REITTER, 1891***Phyllotreta dilutipennis* REITTER, 1891*Phyllotreta turcmenica* J. WEISE, 1900*Phyllotreta schreineri* JACOBSON, 1915*Phyllotreta turcmenica* ab. *pallidipennis* HEIKERTINGER et CSIKI, 1940*Phyllotreta turcmenica schreineri*: GRUEV and DÖBERL, 1997 (in synonymy erroneously attributed to SHAPIRO, 1957)*Phyllotreta erysimi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KRÁL, 1973)

Distribution: S Ukraine, SE European Russia, Ciscaucasia, Daghestan, Turkey, Central Asia, Afghanistan, Iran, Asian Russia (SW Siberia, Altai, Baikal Area, Transbaikalia), Mongolia, NE China.

***Phyllotreta paradoxa* LOPATIN, 1956**

Distribution: Tajikistan, Afghanistan.

***Phyllotreta parfentjevi* SHAPIRO, 1958**

Distribution: Turkmenistan, Kazakhstan, Uzbekistan.

***Phyllotreta praticola* J. WEISE, 1887***Phyllotreta fucata* J. WEISE, 1890

Distribution: SE European Russia, S Ural, Caucasus, Transcaucasia, Turkey, Afghanistan, Iran, Central Asia, Asian Russia (Altai), Mongolia, NW China, India.

***Phyllotreta procera* (L. REDTENBACHER, 1849)***Haltica procera* L. REDTENBACHER, 1849*Phyllotreta subtilis* WOLLASTON, 1864*Phyllotreta rufitarsis* J. WEISE, 1888? *Phyllotreta carreti* MONNOT, 1913*Phyllotreta nodicornis* APFELBECK, 1914 (part.)*Phyllotreta procera procerior* PEYERIMHOFF, 1941

Distribution: S and southern part of Central Europe, Caucasus, Transcaucasia, Turkey, Near and Middle East, Turkmenistan, Tajikistan, N and NE Africa.

***Phyllotreta pseudoexclamationis* KONSTANTINOV, 1992**

Distribution: Georgia.

***Phyllotreta punctulata* (MARSHAM, 1802)***Chrysomela punctulata* MARSHAM, 1802*Phyllotreta aerea* ALLARD, 1859

Distribution: Europe except for N, Caucasus, Near East, Turkey. Introduced to N America.

***Phyllotreta rectilineata* CHEN, 1939***Phyllotreta chinensis* HEIKERTINGER, 1941*Phyllotreta chinensis* fa. *shirahatai* MADAR, 1959*Phyllotreta shirahatai* CHŪJŌ et KIMOTO, 1961

Distribution: Asian Russia (Ussuri Area), China, Korea, Laos, N Vietnam, Japan.

***Phyllotreta reitteri* HEIKERTINGER, 1911**

Distribution: S Ukraine (Crimea), Kazakhstan, Uzbekistan.

***Phyllotreta sachalinensis* L. MEDVEDEV, 1973**

Distribution: Asian Russia (S Sakhalin).

***Phyllotreta scheuchi* HEIKERTINGER, 1941**

Distribution: England, Spain, southern parts of Central Europe, S Ukraine, Mongolia.

***Phyllotreta sisymbrii* J. WEISE, 1888**

Distribution: Caucasus, Transcaucasia, Iran.

***Phyllotreta striolata* (FABRICIUS, 1803)***Crioceris striolata* FABRICIUS, 1803*Crioceris vittata* FABRICIUS, 1801*Haltica sinuata* L. REDTENBACHER, 1849*Phyllotreta sinuata* L. REDTENBACHER, 1858*Phyllotreta vittata* var. *discedens* J. WEISE, 1888*Phyllotreta vittata* var. *monticola* J. WEISE, 1888*Phyllotreta vittata* ab. *lineolata* CHITTENDEN, 1927*Phyllotreta vittata* ab. *discedens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1940).*Phyllotreta vittata* ab. *monticola*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1940).

Distribution: Europe except for extreme N, Caucasus, Transcaucasia, Syria, Kazakhstan, Asian Russia (Siberia, Sakhalin, Kuriles Isles), Mongolia, China, Korea, Japan, Taiwan, SE Asia, India, Indonesia. Introduced to S Africa and N America.

***Phyllotreta talassicola* HEIKERTINGER, 1944**

Distribution: Central Asia, Afghanistan.

***Phyllotreta tetrastigma* (COMOLLI, 1837)***Altica tetrastigma* COMOLLI, 1837*Haltica flexuosa* var. *DUFTSCHMIDT*, 1825*Phyllotreta sinuata* SEIDLITZ, 1875*Phyllotreta tetrastigma* var. *dilatata* J. WEISE, 1888*Phyllotreta flexuosa* var. *fenestrata* BEDEL, 1900*Phyllotreta fallax* BEDEL, 1900*Phyllotreta tetrastigma* fa. *maculipennis* HEIKERTINGER, 1941*Phyllotreta tetrastigma* ab. *weiseana* CSIKI, 1953*Phyllotreta flexuosa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1874)

Distribution: Europe, Asian Russia (W and Middle Siberia).

***Phyllotreta tomboi* LOPATIN, 1967***Phyllotreta brachyptera* L. MEDVEDEV et VORONOVA, 1977

Distribution: Mongolia.

***Phyllotreta absunurica* L. MEDVEDEV, 1980**

Distribution: Mongolia.

***Phyllotreta undulata* KUTSCHERA, 1860***Phyllotreta undulata* var. *bilineata* J. WEISE, 1888*Phyllotreta australis* BLACKBURN, 1890*Phyllotreta vittigera* BROUN, 1893*Phyllotreta undulata* var. *biinterrupta* PIC, 1915



- Phyllotreta blackburni* BRYANT, 1925  
*Phyllotreta undulata* ab. *bilineata* HEIKERTINGER et CSIKI, 1940  
*Phyllotreta undulata* fa. *nigroconcolor* DÖBERL, 1994  
*Phyllotreta undulata* fa. *evanescens* DÖBERL, 1994  
*Altica nemorum*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PANZER, 1794)  
*Haltica nemorum*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KOCH, 1803)  
*Phyllotreta flexuosa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GYLLENHAL, 1813)  
 Distribution: Europe, Caucasus, Transcaucasia, Turkey, Central Asia, Asian Russia (Siberia, Far East), Mongolia. Introduced to N America.

***Phyllotreta ustulata* LOPATIN, 1961**

Distribution: Kazakhstan, Tajikistan, Uzbekistan.

***Phyllotreta vittula* (L. REDTENBACHER, 1849)**

- Haltica vittula* L. REDTENBACHER, 1849  
 ? *Haltica vittata* STEPHENS, 1832  
*Phyllotreta rivularis* MOTSCHULSKY, 1849  
 ? *Phyllotreta exigua* HEIKERTINGER, 1911  
*Phyllotreta vittula* var. *exigua* HEIKERTINGER, 1930  
*Phyllotreta vittata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)  
 Distribution: Europe, Caucasus, Transcaucasia, Turkey, Central Asia, Afghanistan, Asian Russia (Siberia, Far East), Mongolia, Korea, China (Heilungkiang).

***Phyllotreta wiseana* JACOBSON, 1901**

- Phyllotreta denticornis* J. WEISE, 1900  
*Phyllotreta wisei* LOPATIN, 1960 (attributed to Jacobson) [nomen nudum]  
 Distribution: S Ukraine, Crimea, SE European Russia, Caucasus, Transcaucasia, Turkey, N Iran, W Kazakhstan.

***Phyllotreta zimmermanni* (CROTCH, 1873)**

- Phyllotreta sibirica* J. WEISE, 1887  
*Phyllotreta sinuata* G. HORN, 1889  
 Distribution: N Europe, Asian Russia (Middle and E Siberia, Far East, N Cispaacific Land), N America.

**Genus: *Aphthona* CHEVROLAT in DEJEAN, 1837**

- Aphthona* SIEBOLD, 1847  
*Ectonia* J. WEISE, 1922  
*Asiatica* SCHERER, 1969  
*Aphthonotarsa* MEDVEDEV, 1984

***Aphthona abdominalis* (DUFTSCHMIDT, 1825)**

- Thyamis abdominalis* STEPHENS, 1832  
*Aphthona albinea* C. G. THOMSON, 1866  
*Aphthona foudrasi* JACOBY, 1885  
*Aphthona finitima* OGLOBLIN, 1926  
*Aphthona sajanica* OGLOBLIN, 1926  
*Aphthona licetana* CHEN, 1939  
 Distribution: Europe, Caucasus, Afghanistan, Iran, Kazakhstan, Asian Russia (S Siberia), China, Vietnam, Japan.

***Aphthona aeneomicans* ALLARD, 1875**

*Aphthona metallica* J. WEISE, 1889  
*Aphthona orientalis* ROUBAL, 1936  
*Aphthona pieniensis* HEIKERTINGER et CSIKI, 1939  
 Distribution: S Europe east to Slovakia.

***Aphthona atrocaerulea* (STEPHENS, 1832)**

*Altica atrocaerulea* STEPHENS, 1832  
*Haltica cyanella* L. REDTENBACHER, 1849  
*Aphthona puncticollis* ALLARD, 1866  
*Altica atrocaerulea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)  
 Distribution: Europe except for European Russia.

***Aphthona atrovirens* (FÖRSTER, 1849)**

*Haltica atrovirens* FÖRSTER, 1849  
*Aphthona tantilla* FOU DRAS, 1860  
*Aphthona atrovirens* var. *aeneipennis* J. WEISE, 1892  
*Aphthona atrovirens* var. *punctigera* J. WEISE, 1893  
 Distribution: W and Central Europe, Ukraine, N Caucasus, Turkey.

***Aphthona beckeri* JACOBSON, 1896**

*Aphthona brunneomicans* HEIKERTINGER, 1912  
*Aphthona daurica* L. MEDVEDEV, 1973  
*Aphthona nigriscutis beckeri*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1940)  
 Distribution: Southern E Europe, Caucasus, W Kazakhstan, Siberia, Mongolia.

***Aphthona chalchica* L. MEDVEDEV, 1980**

Distribution: Mongolia.

***Aphthona chekanovskii* KONSTANTINOV, 1998**

Distribution: Asian Russia (Yakutia).

***Aphthona chrysolina* HEIKERTINGER, 1944**

*Aphthona irisha* PALIJ, 1970  
 Distribution: Uzbekistan, SW Kirghizstan (Chatkal Mountain Range).

***Aphthona crassicornis* LOPATIN, 1990**

Distribution: Turkey.

***Aphthona cyparissiae* (KOCH, 1803)**

*Haltica cyparissiae* KOCH, 1803  
*Haltica tabida* DUFTSCHMIDT, 1825  
*Aphthona nigriventris* ALLARD, 1860  
*Aphthona cyparissiae* ab. *suturalis* DELAHON, 1922  
*Aphthona cyparissiae* ab. *obscuricornis* BAUER, 1929 [nomen nudum]  
 Distribution: Europe except for N, Caucasus, W Kazakhstan.

***Aphthona czwalinae* J. WEISE, 1889**

*Aphthona czwalinae* ab. *coerulescens* J. WEISE, 1889  
*Aphthona czwalinae asiatica* HEIKERTINGER, 1911  
 Distribution: Europe except for N, Kazakhstan, Kirghizstan, Asian Russia (W Siberia, Tuva), Mongolia.

***Aphthona erichsoni* (ZETTERSTEDT, 1838)***Haltica erichsoni* ZETTERSTEDT, 1838*Haltica nigrataris* MOTSCHULSKY, 1845*Chaetocnema erichsoni*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BARTUEVA, 1973)

Distribution: N Europe, Ukrainian Carpathians, Asian Russia (Siberia, Far East, Kamchatka), Mongolia.

***Aphthona euphorbiae* (SCHRANK, 1781)***Chrysomela euphorbiae* SCHRANK, 1781*Thyamis hilaris* STEPHENS, 1832*Aphthona atratula* ALLARD, 1859*Aphthona virescens* FOUDRAS, 1860*Aphthona punctigera* MULSANT et REY, 1874*Aphthona subaptera* MULSANT et REY, 1874*Aphthona euphorbiae* var. *cyanescens* J. WEISE, 1888*Aphthona euphorbiae* ab. *cyanescens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to WANKA, 1920)*Aphthona hilaris* ab. *subaptera* HEIKERTINGER et CSIKI, 1940*Aphthona hilaris*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)*Thyamis hilaris*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to MANNERHEIM, 1839)*Haltica hilaris*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BACH, 1839)*Haltica euphorbiae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KOCH, 1803)*Crioceris euphorbiae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to WIEGEL, 1806)*Aphthona aeneomicans*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to DONISTHOPPE, 1944)

Distribution: Europe except for N, Caucasus, Turkey, Kazakhstan, Central Asia, Siberia.

***Aphthona flava* GUILLEBEAU, 1894***Aphthona cyparissiae flava*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1939)

Distribution: S and SE Europe, Turkey.

***Aphthona flaviceps* ALLARD, 1859***Haltica pallida* BOIELDIEU, 1859*Aphthona straminea* FOUDRAS, 1859*Aphthona flavipes* ALLARD, 1866*Haltica flavipes* LETZNER, 1870

Distribution: S and SE Europe, Caucasus, Turkey, Middle East, Kazakhstan, Central Asia.

***Aphthona franzi* HEIKERTINGER, 1944**

Distribution: Southern parts of Central and SE Europe, Caucasus, Turkey, Asian Russia (Altai).

***Aphthona gracilipes* OGLOBLIN, 1926**

Distribution: S Kazakhstan, NW Kirghizstan.

***Aphthona gracilis* FALDERMANN, 1837***Aphthona picipes* J. WEISE, 1888*Aphthona armeniaca* J. WEISE, 1889*Aphthona flavofemorata* HEIKERTINGER, 1911

Distribution: SE Europe (Moldavia, Romania, S Ukraine, Crimea), Caucasus, Transcaucasia, Turkey, Near and Middle East, S Kazakhstan, Uzbekistan, Asian Russia (Transbaikalia), Mongolia.

***Aphthona hammarstroemi* JACOBSON, 1901**

*Aphthona fischeri* PIC, 1910

*Aphthona hammarstroemi issykkensis* OGLOBLIN, 1926

Distribution: SE Kazakhstan, Kirghizstan, Tajikistan, Asian Russia (S Siberia east to Transbaikalia).

***Aphthona hauseri* HEIKERTINGER, 1911**

*Aphthona claripes* RAPILLY, 1978

Distribution: Afghanistan, Iran, Central Asia.

***Aphthona herbigrada* (CURTIS, 1837)**

*Altica herbigradus* CURTIS, 1837

*Haltica campanulae* W. REDTENBACHER, 1842

*Aphthona fossulata* ALLARD, 1859

*Aphthona dimidiata* MULSANT et REY, 1873

*Aphthona laevicollis* MULSANT et REY, 1873

*Aphthona herbigrada* var. *gibbula* J. WEISE, 1891

Distribution: Europe except for N, N Africa.

***Aphthona hissarica* LOPATIN, 1975**

Distribution: Iran, NE Afghanistan, Tajikistan, India (Kashmir).

***Aphthona interstitialis* J. WEISE, 1887**

*Aphthona interstitialis* var. *tibetana* HEIKERTINGER, 1911

*Aphthona suvorovi* OGLOBLIN, 1921

*Aphthona suturanigra* CHEN, 1939

*Aphthona yuasai* OHNO, 1962

Distribution: Asian Russia (Transbaikalia, Amurland, Ussuri Area, Amurland, Maritime Prov.), Mongolia, W China, Tibet, ?Tajikistan.

***Aphthona jacobsoni* OGLOBLIN, 1926**

Distribution: Afghanistan, N Iran, Central Asia.

***Aphthona jakuta* OGLOBLIN, 1926**

Distribution: Asian Russia (Evenkia, Yakutia, middle and upper Lena Valley, Ciasbaikalia, Transbaikalia, Magadan Distr., Far East), Mongolia.

***Aphthona kaszabi* KRÁL, 1967**

*Longitarsus amygdali* L. MEDVEDEV, 1974

Distribution: Mongolia.

***Aphthona konstantinovi* LOPATIN in KONSTANTINOV, 1998**

Distribution: Crimea, N Caucasus, Turkey.

***Aphthona kuntzei* ROUBAL, 1931**

Distribution: E Mediterranean, Balkans, Ukraine, Turkey, Near East.

***Aphthona lacertosa* (ROSENHAUER, 1847)**

*Haltica lacertosa* ROSENHAUER, 1847

*Haltica divaricata* L. REDTENBACHER, 1849

Distribution: Central and SE Europe, Turkey. Introduced to N America.

***Aphthona lopatini* KONSTANTINOV, 1998**

Distribution: S Kazakhstan, Kirghizstan.

***Aphthona lubischevi* KONSTANTINOV, 1998**

Distribution: Crimea.

***Aphthona lutescens* (GYLLENHAL, 1813)**

*Thyamis lutescens* GYLLENHAL, 1813

*Thyamis sternalis* MULSANT et REY, 1873

*Aphthona praeclara* J. WEISE, 1906

*Aphthona lutescens* fa. *piceoconcolor* DÖBERL, 1994

Distribution: Europe, Caucasus, Turkey, Kazakhstan, Central Asia, Asian Russia (W Siberia, Transbaikalia), Mongolia, N Africa.

***Aphthona maculata* ALLARD, 1876**

*Aphthona suturella* J. WEISE, 1888

*Aphthona pallida maculata* HEIKERTINGER, 1930

Distribution: Caucasus, Turkey, Near East, Iran, Kazakhstan, S Tajikistan.

***Aphthona modesta* J. WEISE, 1887**

Distribution: Asian Russia (Amurland, Maritime Prov.), NE China, Korea.

***Aphthona mohri* WARCHALOWSKI, 1973**

Distribution: Turkmenistan, Iran.

***Aphthona nigriceps* (W. REDTENBACHER, 1842)**

*Longitarsus nigriceps* W. REDTENBACHER, 1842

*Aphthona sicula* FOUDRAS, 1860

*Aphthona decorata* KUTSCHERA, 1861

Distribution: S and SE Europe, S Urals, Caucasus, Turkey, Near East, N Africa.

***Aphthona nigriscutis* FOUDRAS, 1860**

*Aphthona nigriventris* ALLARD, 1860

*Aphthona nigroscutellata* REITTER, 1904

*Aphthona cyparissiae nigriscutis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

Distribution: Central and SE Europe, Caucasus, Transcaucasia, Turkey, Iran, W Kazakhstan.

Introduced to N America.

***Aphthona nonstriata* (GOEZE, 1777)**

*Chrysomela nonstriata* GOEZE, 1777

*Chrysomela pseudacori* MARSHAM, 1802

*Aphthona coerulea* var. *aenescens* J. WEISE, 1888

*Aphthona coerulea* var. *asthmatica* J. WEISE, 1888

*Aphthona coerulea* var. *metallescens* KOLBE, 1915

*Aphthona thoracica* OGLOBLIN, 1917

*Aphthona picipes* GABRIEL, 1918

*Altica coerulea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GEOFFROY, 1785)

*Haltica nonstriata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HELM, 1883)

Distribution: Europe, Caucasus, Turkey, Iran, Kazakhstan, Central Asia, Asian Russia (W Siberia).

***Aphthona ovata* FOUDRAS, 1860***Haltica euphorbiae* L. REDTENBACHER, 1849

Distribution: S, Central and SE Europe, Transcaucasia, Turkey.

***Aphthona pallida* (BACH, 1856)***Haltica pallida* BACH, 1856*Aphthona pallida* var. *geranii* J. WEISE, 1891*Aphthona nigriceps*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BARGAGLI, 1878)

Distribution: SE and Central Europe, Crimea, Caucasus.

***Aphthona perminuta* BALY, 1875***Aphthona pygmaea* BALY, 1874

Distribution: Asian Russia (Sakhalin, Kuriles Isles), Korea, Japan.

***Aphthona placida* (KUTSCHERA, 1861)***Haltica placida* KUTSCHERA, 1861*Haltica pallida* KUTSCHERA, 1861

Distribution: Central and SE Europe.

***Aphthona promissa* OGLOBLIN, 1926**

Distribution: Afghanistan, S Kazakhstan, Kirghizstan, Tajikistan.

***Aphthona przhevalskii* KONSTANTINOV, 1998**

Distribution: S Kazakhstan.

***Aphthona pygmaea* (KUTSCHERA, 1861)***Haltica pygmaea* KUTSCHERA, 1861*Aphthona euphorbiae* FOUDRAS, 1860*Haltica nigella* KUTSCHERA, 1861*Aphthona atrocoerulea* C. G. THOMSON, 1866*Aphthona orientalis* MULSANT et REY, 1874*Aphthona fulvipes* J. SAHLBERG, 1913*Aphthona pygmaea* var. *nigella* WANKA, 1920*Aphthona pygmaea* var. *orientalis* HEIKERTINGER, 1944*Aphthona substriata* L. MEDVEDEV, 1962*Aphthona pygmaea nigella*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1944)*Aphthona pygmaea orientalis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KRÁL, 1967)

Distribution: Europe except for N, Caucasus, Turkey, Near East, N Africa.

***Aphthona reinecki* LOPATIN, 1962**

Distribution: NE Afghanistan.

***Aphthona reitteri* ALLARD, 1884***Aphthona georgica* OGLOBLIN, 1926

Distribution: Caucasus, Transcaucasia.

***Aphthona rugipennis* OGLOBLIN, 1926***Aphthona aeneomicans orientalis* L. MEDVEDEV, 1962*Aphthona viridis* IABLOKOFF-KHINZORIAN, 1964*Aphthona praenuntiata* KRÁL, 1964*Aphthona aeneomicans nitidior* WARCHALOWSKI, 1995

Distribution: Caucasus, S Kazakhstan, Uzbekistan.

***Aphthona russica* KONSTANTINOV, VOLKOVITSH et CRISTOFARO, 2001**

Distribution: W Ciscaucasia (Taman' Peninsula).

***Aphthona sajanica* OGLOBLIN, 1926**

Distribution: Asian Russia (Sayan, Evenkia), Mongolia, NE China.

***Aphthona sarmatica* OGLOBLIN in GAVRILENKO, 1928**

Distribution: Southern E Europe, Crimea, Transcaucasia.

***Aphthona semicyanea* ALLARD, 1859**

*Aphthona viridula* ALLARD, 1876

*Aphthona punctatissima* J. WEISE, 1888

*Aphthona picipes* J. WEISE, 1892

*Aphthona peregrina* J. WEISE, 1892

*Aphthona punctatissima* var. *picipes* J. WEISE, 1892

*Aphthona melanopeza* JACOBSON, 1901

*Aphthona sundmani* JACOBSON, 1901

*Aphthona semicyanea* var. *phrygia* J. WEISE, 1901

*Aphthona heptapotamica* OGLOBLIN, 1917

*Aphthona pachypus* JACOBSON, 1924

*Aphthona semicyanea tenebrosa* OGLOBLIN, 1926

*Aphthona semicyanea punctatissima* OGLOBLIN, 1926

*Aphthona semicyanea heptapotamica* OGLOBLIN, 1926

*Aphthona semicyanea peregrina* OGLOBLIN, 1926

*Aphthona semicyanea sundmani* OGLOBLIN, 1926

*Aphthona semicyanea phrygia* OGLOBLIN, 1926

*Aphthona grossa* KHNZORIAN, 1959

Distribution: Central, S and SE Europe, N Caucasus, Transcaucasia, Turkey, Near East, Kazakhstan, Central Asia, Asian Russia (S Siberia), Mongolia, India.

***Aphthona semiviridis* JACOBY, 1885**

Distribution: Asian Russia (S Sakhalin), Japan.

***Aphthona stussineri* J. WEISE, 1888**

*Aphthona pyrenaea* HEIKERTINGER, 1944

Distribution: Central and S Europe, Ukrainian Carpathians.

***Aphthona testaceicornis* J. WEISE, 1894**

*Aphthona gracilis* var. *testaceicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

Distribution: Caucasus.

***Aphthona tolli* OGLOBLIN, 1926**

*Aphthona tolli mongolica* L. MEDVEDEV, 1976

Distribution: N Kazakhstan, Asian Russia (Transbaikalia, Yakutia, Amurland), Mongolia, China.

***Aphthona trivialis* J. WEISE, 1887**

*Aphthona merkli* GRUEV, 1994

Distribution: Asian Russia (Yakutia, Ussuri Area, Amurland), N Korea.

***Aphthona valachica* HEIKERTINGER, 1944**

Distribution: Romania, Turkey.

***Aphthona venustula* (KUTSCHERA, 1861)**

*Haltica venustula* KUTSCHERA, 1861  
*Aphthona cyanella* Foudras, 1859  
*Aphthona euphorbiae* Allard, 1860  
*Aphthona attica* J. Weise, 1890

Distribution: Europe except for N, Caucasus, Turkey, Middle East.

***Aphthona violacea* (Koch, 1803)**

*Haltica violacea* Koch, 1803  
*Aphthona pseudacori* L. Redtenbacher, 1849  
*Aphthona sublaevis* Boheman, 1851  
*Aphthona pseudacori* var. *discolor* J. Weise, 1893  
*Aphthona violacea* ab. *sculptipennis* Ogloblin, 1926  
*Aphthona ciscaucasica* L. Medvedev, 1962  
*Aphthona tbilisiensis* Král, 1969  
*Podagrica violacea*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Łomnicki, 1886)

Distribution: Europe except for N, Caucasus, Turkey, Kazakhstan, S Siberia.

***Aphthona wiseana* Konstantinov, 1998**

Distribution: Iran, Turkmenistan, Uzbekistan.

**Genus: *Longitarsus* Latreille, 1829**

*Thyamis* Stephens, 1832  
*Teinodactyla* Chevrolat in Dejean, 1836

**Subgenus: *Longitarsus* Latreille, 1829*****Longitarsus (Longitarsus) absinthii* Kutschera, 1862**

*Longitarsus niger* Bach, 1856  
*Longitarsus maracandius* Lopatin, 1960  
*Thyamis absinthii*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Leder, 1872)

Distribution: Europe except for extreme N, Transcaucasia, Turkey, Kazakhstan, Central Asia, Mongolia.

***Longitarsus (Longitarsus) acuticollis* Lopatin, 1976**

Distribution: S Kazakhstan.

***Longitarsus (Longitarsus) aeneicollis* (Faldermann, 1837)**

*Teinodactyla aeneicollis* Faldermann, 1837  
*Chrysomela suturalis* Marsham, 1802  
*Theinodactyla nigricollis* Foudras, 1860  
*Theinodactyla rubenticollis* Allard, 1860  
*Longitarsus atricillus* var. *similis* J. Weise, 1893  
*Longitarsus suturalis* var. *rubenticollis* J. Weise, 1893  
*Thyamis cribripennis* Abeille, 1909  
*Longitarsus suturalis*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Kutschera, 1863)  
*Thyamis aeneicollis*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Allard, 1866)  
*Thyamis suturalis*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Stephens, 1832)  
*Theinodactyla suturalis*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Allard, 1860)



*Longitarsus suturalis rubenticollis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LOPATIN, 1958)

Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, Near and Middle East, Kazakhstan, Central Asia.

***Longitarsus (Longitarsus) aeruginosus* (FOUDRAS, 1860)**

*Teinodactyla aeruginosa* FOU DRAS, 1860

*Thyamis aeruginosa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BEDEL, 1898)

Distribution: W and S Europe, N Mediterranean.

***Longitarsus (Longitarsus) albineus* (FOUDRAS, 1860)**

*Teinodactyla albinea* FOU DRAS, 1860

*Thyamis albinea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

Distribution: Southern Central Europe, Balkans, N Mediterranean, Caucasus, Turkey, Near and Middle East, Tajikistan, Uzbekistan, N Africa.

***Longitarsus (Longitarsus) alfierii alfierii* PIC, 1923**

*Longitarsus klapperichi* MOHR, 1962

Distribution: NE Mediterranean, Turkey, Near and Middle East, Kirghizstan, N Africa.

***Longitarsus (Longitarsus) alfierii furthi* GRUEV, 1982**

*Longitarsus klapperichi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GRUEV, 1973)

Distribution: N Mediterranean, Balkans, Crimea, Caucasus (Georgia).

***Longitarsus (Longitarsus) anatolicus* J. WEISE, 1900**

*Longitarsus anchusae anatolicus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed HEIKERTINGER, 1930)

Distribution: Transcaucasia, Turkey, Near and Middle East.

***Longitarsus (Longitarsus) aphthonoides* J. WEISE, 1887**

Distribution: Asian Russia (Middle Siberia, Cisbaikalia, Transbaikalia, E Siberia, Amurland, Maritime Prov., Sakhalin).

***Longitarsus (Longitarsus) apicalis* (BECK, 1817)**

*Haltica apicalis* BECK, 1817

*Haltica analis* DUFTSCHMIDT, 1825

*Longitarsus praticola* C. R. SAHLBERG, 1834

*Longitarsus fischeri* ZETTERSTEDT, 1840

*Longitarsus apicalis* var. *quadrimaculatus* J. WEISE, 1888

*Longitarsus apicalis* fa. *borealis* KOLBE, 1920

*Longitarsus apicalis* fa. *quadrimaculatus* HEIKERTINGER, 1930

*Longitarsus analis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)

*Teinodactyla apicalis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)

*Teinodactyla analis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)

*Thyamis apicalis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

Distribution: Europe, Asian Russia (W Siberia, Middle Siberia, Sayan, Cisbaikalia, Transbaikalia).

***Longitarsus (Longitarsus) afghanicus* LOPATIN, 1963**

*Longitarsus asperifoliarum afghanicus* LOPATIN, 1963

Distribution: Afghanistan.

***Longitarsus (Longitarsus) asperifoliarum* J. WEISE, 1887***Longitarsus asperifoliarum* var. *defectus* JACOBSON, 1893

Distribution: Caucasus, S and SE Kazakhstan, Kirgizstan, Turkmenistan, Uzbekistan, Afghanistan, Asian Russia (Altai).

***Longitarsus (Longitarsus) atricillus* (LINNAEUS, 1761)***Chrysomela atricilla* LINNAEUS, 1761*Haltica suturalis* DUFTSCHMIDT, 1825*Teinodactyla fuscicollis* FOU DRAS, 1860*Longitarsus atricillus* var. *declivis* J. WEISE, 1893*Longitarsus atricillus* fa. *similis* MOHR, 1965

Distribution: Europe except for extreme N, Transcaucasia, Middle East, Turkmenistan, Uzbekistan, N Africa.

***Longitarsus (Longitarsus) ballotae* (MARSHAM, 1802)***Chrysomela ballotae* MARSHAM, 1802*Teinodactyla canescens* FOU DRAS, 1860*Longitarsus ballotae* ab. *septorum* BUYSSON, 1908*Longitarsus ballotae* fa. *caligans* KRÁL, 1945

Distribution: Europe except for N, Ciscaucasia, Caucasus, Transcaucasia, Turkey, Near and Middle East, S Kazakhstan, Mongolia, N Africa.

***Longitarsus (Longitarsus) bertii* LEONARDI, 1973***Longitarsus pseudoferrugineus* WARCHALOWSKI, 1974*Longitarsus ferrugineus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KASZAB, 1962)

Distribution: Southern parts of Central Europe, Balkans, SW Ukraine, Transcaucasia, Turkey, Near and Middle East.

***Longitarsus (Longitarsus) brisouti* HEIKERTINGER, 1912***Teinodactyla piciceps*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)*Longitarsus piciceps*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BRISOUT DE BARNEVILLE, 1873)*Thyamis piciceps*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BEDEL, 1898)

Distribution: W and S Europe, Ukraine, Ciscaucasia, Turkey.

***Longitarsus (Longitarsus) brunneus* (DUFTSCHMIDT, 1825)***Haltica brunnea* DUFTSCHMIDT, 1825*Haltica castanea* DUFTSCHMIDT, 1825*Longitarsus brunneus* var. *laevicollis* J. WEISE, 1888*Longitarsus brunneus* var. *robustus* J. WEISE, 1888*Longitarsus brunneus* var. *picinus* J. WEISE, 1888*Longitarsus rubellus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to WARCHALOWSKI, 1974)

Distribution: Europe, Caucasus, Transcaucasia, SE Kazakhstan, Kirghizstan, Asian Russia (W and Middle Siberia, Cisbaikalia, Transbaikalia, Amurland, Far East), Korea.

***Longitarsus (Longitarsus) callidus* WARCHALOWSKI, 1967***Longitarsus membranaceus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BOROWIEC, 1978)

Distribution: W and Central Europe, Transcaucasia, Iran, Kazakhstan, N Africa.

***Longitarsus (Longitarsus) celticus* LEONARDI, 1975**

Distribution: Southern parts of Central Europe, Ukraine.

***Longitarsus (Longitarsus) cerinthes* (SCHRANK, 1798)**

*Altica cerinthes* SCHRANK, 1798

*Longitarsus nervosus* WOLLASTON, 1854

*Longitarsus pectoralis* FOUDRAS, 1860

*Longitarsus cerenthicola* HEIKERTINGER, 1912 [nomen nudum]

Distribution: Europe except for N, Mediterranean, N Africa.

***Longitarsus (Longitarsus) corpulentus* J. WEISE, 1887**

Distribution: S Kazakhstan, Central Asia.

***Longitarsus (Longitarsus) curtus* (ALLARD, 1860)**

*Teinodactyla curtus* ALLARD, 1860

*Teinodactyla papaveris* ALLARD, 1860

*Thyamis liliputana* ALLARD, 1866

*Thyamis vidua* ALLARD, 1866

? *Thyamis moscovita* ALLARD, 1866

*Longitarsus curtus* var. *ciliifer* MÉQUIGNON, 1948

*Longitarsus curtus* ab. *lichtmeckeri* KASZAB, 1962

*Longitarsus viduus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LEESBERG, 1882)

*Thyamis papaveris*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

*Longitarsus papaveris*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KUHN, 1913)

Distribution: Europe except for N, Caucasus, Transcaucasia, Turkmenistan, Tajikistan, Asian Russia (Transbaikalia), Afghanistan.

***Longitarsus (Longitarsus) desertorum* HEIKERTINGER, 1913**

Distribution: Turkmenistan, Kazakhstan, Afghanistan, Central Asia, Mongolia.

***Longitarsus (Longitarsus) dlabolai* KRÁL, 1964**

Distribution: Caucasus (Georgia).

***Longitarsus (Longitarsus) echii* (KOCH, 1803)**

*Haltica echii* KOCH, 1803

*Haltica tibialis* DUFTSCHMIDT, 1825

*Longitarsus excurvus* WOLLASTON, 1857

*Teinodactyla flavipes* ALLARD, 1860

*Longitarsus echii* var. *coerulescens* J. WEISE, 1888

*Longitarsus echii* var. *nigrescens* J. WEISE, 1888

*Longitarsus echii* var. *peregrinus* J. WEISE, 1888

*Longitarsus kulikovskii* JACOBSON, 1895

*Longitarsus echii* ab. *dimidiatus* GERHARDT, 1909

Distribution: Europe except for N, Turkey, Afghanistan, N Africa.

***Longitarsus (Longitarsus) eminus* WARCHALOWSKI, 1967**

*Longitarsus obliterated eminus* WARCHALOWSKI, 1973

Distribution: Near and Middle East, Kazakhstan, Afghanistan, Central Asia.

***Longitarsus (Longitarsus) exsoletus* (LINNAEUS, 1758)**

*Chrysomela exsoleta* LINNAEUS, 1758

*Altica atricilla* OLIVIER, 1789

*Chrysomela femoralis* MARSHAM, 1802

- Longitarsus boppardiensis* BACH, 1856  
*Teinodactyla rufula* FOU DRAS, 1860  
*Thyamis livens* MULSANT et REY, 1874  
*Thyamis lividus* FAUVEL, 1888  
*Longitarsus exsoletus* var. *arctulus* J. WEISE, 1893  
*Thyamis exsoletus* BEDEL, 1898  
*Thyamis exoletus* var. *arctulus* PORTEVIN, 1934  
 ? *Haltica atricillus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ILLIGER, 1807)  
*Thyamis femoralis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GYLLENHAL, 1827)  
*Teinodactyla femoralis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)  
*Teinodactyla pratensis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)  
 ? *Longitarsus rufulus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KUTSCHERA, 1863)  
*Longitarsus atricillus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KUTSCHERA, 1864)  
*Thyamis rufula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)  
*Longitarsus livens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. WEISE, 1893)  
*Longitarsus exoletus* fa. *rufulus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1912)  
*Longitarsus exoletus* var. *rufulus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PORTA, 1934)  
*Longitarsus exoletus* ab. *arctulus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1939).  
 Distribution: Europe except for extreme N, Caucasus, Transcaucaisa, Turkey, Near and Middle East, Turkmenistan.

***Longitarsus (Longitarsus) fallax* J. WEISE, 1888**

- Longitarsus linnaei* var. b *fallax* J. WEISE, 1888  
*Thyamis dimidiata* ALLARD, 1866. non STEPHENS, 1834  
*Longitarsus linnaei* var. *scrurator* J. WEISE, 1890  
*Longitarsus echii* var. c („*Thyamis dimidiata*”) J. WEISE, 1893  
*Longitarsus nitens* IABLOKOFF-KHNZORIAN, 1968  
*Longitarsus cynoglossi* PALLI, 1970  
*Longitarsus linnaei*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to DOGUET, 1973)  
*Longitarsus dimidiatus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to DOGUET, 1976)  
*Longitarsus linnaei scrurator*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LOPATIN, 1977)  
*Longitarsus echii fallax*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LOPATIN et KULENOVA, 1986)  
 Distribution: S and SE Europe, Caucasus, Transcaucaisa, Turkey, Near and Middle East, Kazakhstan, Central Asia, N Africa.

***Longitarsus (Longitarsus) ferrugineus* (FOU DRAS, 1860)**

- Teinodactyla ferruginea* FOU DRAS, 1860  
*Longitarsus waterhousei* KUTSCHERA, 1864  
*Thyamis menthae* BEDEL, 1898  
*Longitarsus lycopi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. WEISE, 1893)  
*Longitarsus juncicola*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. WEISE, 1893)

*Longitarsus rubiginosus* var. *ferrugineus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to TOMLIN et SHARP, 1912)

Distribution: Europe except for N, N Africa.

***Longitarsus (Longitarsus) finitimus* KONSTANTINOV, 1992**

Distribution: Kazakhstan (W Tian Shan).

***Longitarsus (Longitarsus) foudrasi* J. WEISE, 1893**

*Teinodactyla pallens* STEPHENS, 1832

*Thyamis foudrasi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BEDEL, 1898)

Distribution: Europe except for N, Ciscaucasia, W Caucasus, Transcaucasia, Turkey, Asian Russia (Amurland), N Africa.

***Longitarsus (Longitarsus) fulgens* (FOUDRAS, 1860)**

*Thyamis fulgens* FODRAS, 1860

*Longitarsus fulgens* var. *exhaustus* J. WEISE, 1888

*Longitarsus pinguis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KRÁL, 1947)

Distribution: Europe except for N and S, Caucasus, Transcaucasia, S Siberia.

***Longitarsus (Longitarsus) ganglbaueri* HEIKERTINGER, 1912**

*Thyamis lutescens* STEPHENS, 1832

*Teinodactyla atricilla*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)

*Longitarsus piciceps* KUTSCHERA, 1863

*Longitarsus foudrasi* CROTCH, 1871

*Thyamis senecionis* BRISOUT DE BARNEVILLE, 1873

Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, Asian Russia (Transbaikalia, Ussuri Area, Sakhalin), Mongolia. Introduced to N America.

***Longitarsus (Longitarsus) georgianus* (ALLARD, 1866)**

*Thyamis georgiana* ALLARD, 1866

Distribution: Caucasus (Georgia).

***Longitarsus (Longitarsus) godmani* (BALY, 1876)**

*Thyamis godmani* BALY, 1876

*Longitarsus subniger* CHEN, 1939

*Longitarsus daovantieni* WARCHALOWSKI, 1966

Distribution: Asian Russia (Amurland), Korea, China, N Vietnam.

***Longitarsus (Longitarsus) gracilis* KUTSCHERA, 1864**

? *Teinodactyla nigriceps* FODRAS, 1860

*Thyamis poweri* ALLARD, 1866

*Longitarsus gracilis* var. *nigrithorax* HEIKERTINGER, 1912

*Longitarsus sagittifer* MOHR, 1962

Distribution: Europe except for N, Caucasus (Arax Valley), Israel, N Africa.

***Longitarsus (Longitarsus) hissaricus* LOPATIN in KONSTANTINOV et LOPATIN, 2000**

Distribution: Tajikistan.

***Longitarsus (Longitarsus) hoberlandti* LOPATIN, 1990**

Distribution: Turkmenistan, N Iran, Tajikistan.

***Longitarsus (Longitarsus) holsaticus* (LINNAEUS, 1758)**

*Chrysomela holsatica* LINNAEUS, 1758

*Chrysomela pulicaria* LINNAEUS, 1766  
*Longitarsus haemorrhoidalis* JACOBY, 1855  
*Longitarsus holsaticus* var. *discoideus* J. WEISE, 1888  
*Longitarsus tsii* CHEN, 1941  
*Longitarsus holsaticus* var. *seriepunctatus* ROUBAL, 1943  
*Longitarsus apicalis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

Distribution: Europe, Caucasus, Asian Russia (All over Siberia), China, Korea, Japan.

***Longitarsus (Longitarsus) indigonaceus* LOPATIN, 1963**

*Longitarsus warchalowskii* SCHERER, 1969

Distribution: NE Afghanistan, Pakistan, India (Bengal).

***Longitarsus (Longitarsus) jacobaeae* (WATERHOUSE, 1858)**

*Thyamis jacobaeae* WATERHOUSE, 1858

? *Haltica laevis* DUFTSCHMIDT, 1825

Distribution: Europe, Caucasus, Transcaucasia, Kazakhstan, Kirghizstan, Mongolia, China (Tibet), Asian Russia (All over Siberia except for N).

***Longitarsus (Longitarsus) jailensis* HEIKERTINGER, 1913**

Distribution: Balkans, Crimea, Turkey.

***Longitarsus (Longitarsus) juncicola* (FOUDRAS, 1860)**

*Teinodactyla juncicola* FOU DRAS, 1860

? *Longitarsus lycopi* var. *fulviventris* J. WEISE, 1893

*Longitarsus lycopi* var. *juncicola* HEIKERTINGER, 1914

*Thyamis lycopi* var. *fulviventris* PORTEVIN, 1934

*Longitarsus seriatus* KUTSCHERA, 1864

*Thyamis juncicola*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1861)

*Thyamis seriatus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

Distribution: W and Central Europe, Caucasus, Transcaucasia, Turkey, N Africa.

***Longitarsus (Longitarsus) karlheinzi* WARCHALOWSKI, 1972**

Distribution: Crimea, Turkey, Near and Middle East.

***Longitarsus (Longitarsus) khzoriani* PALIJ, 1970**

*Longitarsus scrobipennis* GRUEV, 1982 (part.).

Distribution: W Caucasus (?), Kirghizstan.

***Longitarsus (Longitarsus) kutscherai* (RYE, 1872)**

*Thyamis kutscherai* RYE, 1872

*Chrysomela atricilla* MARSHAM, 1802

*Haltica atricapilla* DUFTSCHMIDT, 1825

*Longitarsus atriceps* KUTSCHERA, 1863

*Longitarsus articillus* J. WEISE, 1893

*Thyamis melanocephalus paludivaga* PEYERIMHOFF, 1915

*Longitarsus femoratus* CHEN, 1939

*Longitarsus melanocephalus paludivagus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1939)

*Longitarsus melanocephalus* fa. *kutscherae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to MOHR, 1965)

*Longitarsus melanocephalus* var. *paludivagus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

*Longitarsus melanocephalus* var. *kutscherae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1912)

*Thyamis atriceps*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

*Teinodactyla atricapilla*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)

*Longitarsus kutscherae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to WARCHAŁOWSKI, 1978)

Distribution: Europe except for N and E, Ciscaucasia, Caucasus, Transcaucasia, Turkey, Asian Russia (S Siberia), N China, Korea, N Africa.

***Longitarsus (Longitarsus) languidus* KUTSCHERA, 1863**

*Thyamis rubella* ALLARD, 1866

Distribution: Southern parts of Central Europe east to Transcarpathia, Caucasus.

***Longitarsus (Longitarsus) lateripunctatus lateripunctatus* (ROSENHAUER, 1856)**

*Longitarsus (Teinodactyla) lateripunctatus* ROSENHAUER, 1856

*Haltica signata* REICHE et SAULCY, 1858

*Teinodactyla biguttata* FOU DRAS, 1860

*Longitarsus lateripunctatus* var. *persimilis* NORMAND, 1937

*Longitarsus azoricus* ISRAELSON, 1991

*Thyamis lateripunctata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

Distribution: Mediterranean, Near East,? Caucasus.

***Longitarsus (Longitarsus) lateripunctatus personatus* J. WEISE, 1893**

*Longitarsus lateripunctatus* var. *personatus* J. WEISE, 1893

Distribution: Central Europe east to Ukraine, Ciscaucasia, Caucasus, Turkey, Near East.

***Longitarsus (Longitarsus) ledouxi* DOGUET, 1979**

*Longitarsus abchasicus* KONSTANTINOV, 1986

*Longitarsus georgianus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GRUEV, 1987)

Distribution: Ciscaucasia, W Caucasus, Turkey.

***Longitarsus (Longitarsus) lewisii* (BALY, 1874)**

*Thyamis lewisii* BALY, 1874

*Thyamis vidua* ALLARD, 1866

*Longitarsus lycopi* JACOBY, 1885

*Longitarsus stramineus* J. WEISE, 1887

*Longitarsus viduus* J. WEISE, 1893

*Longitarsus borodinensis* CHŪJŌ, 1940

Distribution: Europe except for N east to Ukraine, Transcaucasia, Turkey, Iran, Afghanistan, Kirghizstan, Uzbekistan, Asian Russia (Siberia, Sakhalin, Kuriles Isles), Mongolia, NE and E China, Korea, Japan, Taiwan, N Vietnam.

***Longitarsus (Longitarsus) linnaei* (DUFTSCHMIDT, 1825)**

*Haltica linnaei* DUFTSCHMIDT, 1825

*Longitarsus linnaei* var. *amoenus* J. WEISE, 1888

*Longitarsus amoenus* KUHNT, 1913

*Teinodactyla linnaei*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)

*Thyamis linnaei*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

*Longitarsus echii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to WARCHAŁOWSKI, 1967)

Distribution: Southern parts of Central, S and SE Europe, Caucasus, Transcaucasia, Turkey, Near East.

***Longitarsus (Longitarsus) longipennis* KUTSCHERA, 1863***Longitarsus femoralis* L. REDTENBACHER, 1849*Longitarsus nigrocillus* PORTA, 1934 (part.)

Distribution: Central and SE Europe, Caucasus, Transcaucasia, Turkey, Kazakhstan, Central Asia, Asian Russia (S Siberia east to Tuva and Yakutia).

***Longitarsus (Longitarsus) longiseta* J. WEISE, 1889***Haltica nigriceps* W. REDTENBACHER, 1842*Longitarsus longiseta* fa. *obsoleta* KOLBE, 1920*Longitarsus kwangsiensis* CHEN, 1939*Longitarsus clarus* ALLEN, 1967*Teinodactyla nigriceps*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOUDRAS, 1860)*Longitarsus nigriceps*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. WEISE, 1889)*Thyamis longiseta*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BEDEL, 1898)

Distribution: Europe except for S and extreme N, Caucasus, Transcaucasia, Asian Russia (E Siberia, Maritime Prov.), NE China, Korea, Japan.

***Longitarsus (Longitarsus) luridus* (SCOPOLI, 1763)***Chrysomela lurida* SCOPOLI, 1763*Haltica castanea* DUFTSCHMIDT, 1825*Haltica quadrisignata* DUFTSCHMIDT, 1825*Thyamis brunnea* STEPHENS, 1832*Thyamis fuscens* STEPHENS, 1832? *Teinodactyla conspicibilis* FALDERMANN, 1835*Longitarsus quadrisignatus* KUTSCHERA, 1863*Longitarsus fuscus* KUTSCHERA, 1864*Thyamis syriaca* ALLARD, 1866*Longitarsus amurensis* J. WEISE, 1887*Longitarsus luridus* var. *quadrisignatus* J. WEISE, 1888*Longitarsus luridus* var. *cognatus* J. WEISE, 1888*Longitarsus luridus* var. *nigricans* J. WEISE, 1888*Longitarsus piceonitens* GUILLEBEAU, 1895*Longitarsus luridus* fa. *alata* HEIKERTINGER, 1912*Longitarsus nigerrimus* FOWLER, 1913? *Longitarsus conspicibilis* HEIKERTINGER, 1930*Altica lurida*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to OLIVIER, 1789)*Haltica lurida*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ILLIGER, 1807)*Longitarsus luridus* ab. *nigricans*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GERHARDT, 1909)*Thyamis luridus* ab. *nigricans*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PORTEVIN, 1934)*Longitarsus luridus* ab. *cognatus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GERHARDT, 1909)*Longitarsus luridus conspicibilis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KRÁL, 1964)*Thyamis castanea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)*Thyamis fuscata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)*Longitarsus brunneus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)*Teinodactyla brunnea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOUDRAS, 1860)*Thyamis lurida*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)*Teinodactyla lurida*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOUDRAS, 1860)



*Longitarsus syriacus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

Distribution: All over Palaearctic except for Central Asia and Far East. Introduced to N America.

***Longitarsus (Longitarsus) lycopi* (FOUDRAS, 1860)**

*Teinodactyla lycopi* FOU DRAS, 1860

*Teinodactyla abdominalis* ALLARD, 1860

*Teinodactyla brunniceps* ALLARD, 1860

*Longitarsus seriatus* KUTSCHERA, 1863

? *Longitarsus limnophilus* ABELLE, 1907

*Thyamis lycopi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

*Thyamis abdominalis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

*Thyamis brunniceps*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, Near East, Iran, S Kazakhstan, Central Asia, N Africa.

***Longitarsus (Longitarsus) marguzoricus* KONSTANTINOV in KONSTANTINOV et LOPATIN, 2000**

Distribution: Central Asia (Tajikistan).

***Longitarsus (Longitarsus) medvedevi* SHAPIRO, 1956**

*Longitarsus pallidicornis* KRÁL, 1948

Distribution: Central, SE and E Europe, Uzbekistan.

***Longitarsus (Longitarsus) melanocephalus* (DEGEER, 1775)**

*Chrysomela melanocephalus* DEGEER, 1775

*Chrysomela atricilla* SCOPOLI, 1763

*Altica atricilla* FABRICIUS, 1775

*Haltica atricapilla* DUFTSCHMIDT, 1825

*Thyamis picipes* STEPHENS, 1832

*Teinodactyla crassicornis* FOU DRAS, 1860

? *Altica pallida* GEOFFROY, 1785

*Longitarsus melanocephalus* var. *nigrinus* J. WEISE, 1893

*Longitarsus melanocephalus* var. *nigrosuturalis* DELAHON, 1913

*Longitarsus picipes* PALII, 1960

*Galeruca melanocephala*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GOEZE, 1777)

? *Galeruca atricilla*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1787)

*Crioceris atricilla*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1801)

*Haltica atricilla*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KOCH, 1803)

*Haltica melanocephala*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GYLLENHAL, 1813)

*Longitarsus atricapillus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)

*Teinodactyla melanocephala*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)

*Thyamis crassicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1861)

*Longitarsus crassicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KUTSCHERA, 1864)

*Thyamis melanocephala*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

*Teinodactyla atricapilla*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ACLOQUE, 1896)

*Longitarsus picipes*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to TOMLIN et SHARP, 1912)

*Thyamis melanocephalus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PORTEVIN, 1934)

Distribution: Europe, Caucasus, Transcaucasia, Turkey, Near and Middle East, S Kazakhstan, Central Asia, Asian Russia (W and Middle Siberia, Transbaikalia), Mongolia.

***Longitarsus (Longitarsus) melanoxanthus* LOPATIN, 1963**

Distribution: S Afghanistan.

***Longitarsus (Longitarsus) membranaceus* (FOUDRAS, 1860)**

*Teinodactyla membranacea* FOU DRAS, 1860

*Teinodactyla teucriti* ALLARD, 1860

*Thyamis membranacea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

*Thyamis teucriti*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

Distribution: Europe except for N, Transcaucasia, Turkey, Near and Middle East, N Africa.

***Longitarsus (Longitarsus) minimus* KUTSCHERA, 1863**

*Longitarsus tantulus* var. *minimus* STIERLIN, 1898

*Thyamis minimus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

*Longitarsus pratensis minimus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

*Longitarsus pratensis* ab. *minimus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KASZAB, 1962)

Distribution: Southern parts of Central Europe, N Caucasus, Asian Russia (S Siberia).

***Longitarsus (Longitarsus) minusculus* (FOUDRAS, 1860)**

*Teinodactyla minuscula* FOU DRAS, 1860

Distribution: Europe except for N, E Caucasus, Turkey.

***Longitarsus (Longitarsus) monticola* KUTSCHERA, 1863**

*Thyamis monticola* ALLARD, 1866

*Longitarsus curtis* var. *monticola*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. WEISE, 1893)

*Longitarsus curtis* ab. *monticola*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1939)

*Longitarsus curtis* fa. *monticola*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KRÁL, 1945)

*Longitarsus curtis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LEONARDI, 1973)

Distribution: Mountains of W and Central Europe, Southern parts of European Russia, Caucasus, Tajikistan.

***Longitarsus (Longitarsus) muchei* MOHR, 1984**

Distribution: Uzbekistan (Alai Mountain Range).

***Longitarsus (Longitarsus) murteus* WARCHALOWSKI, 1972**

Distribution: SE Kazakhstan, N Afghanistan.

***Longitarsus (Longitarsus) nanus* (FOUDRAS, 1860)**

*Teinodactyla nanus* FOU DRAS, 1860

? *Teinodactyla cerina* FOU DRAS, 1860

- Longitarsus brevicollis* KUTSCHERA, 1863  
*Longitarsus vitreus* KUTSCHERA, 1863  
*Thyamis nana*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1861)  
 ? *Longitarsus cerinus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KUTSCHERA, 1863)  
*Thyamis vitreus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)  
*Thyamis brevicollis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)  
 ? *Thyamis cerina*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)  
 Distribution: Europe except for N, E Caucasus, Near East, N Africa.

***Longitarsus (Longitarsus) nasturtii* (FABRICIUS, 1792)**

- Galleruca nasturtii* FABRICIUS, 1792  
*Altica nemorum* FABRICIUS, 1775 (part.)  
*Galleruca atricilla* var. c PAYKULL, 1799  
*Haltica pratensis* var. b KOCH, 1803  
*Haltica atricilla* var. 1 ILLIGER, 1807  
*Longitarsus circumscriptus* BACH, 1856  
*Longitarsus nasturtii* var. *obscurus* DONISTHORPE, 1929  
 ? *Longitarsus dorsopictus* CHEN, 1939  
*Longitarsus nasturtii* var. *auctumnalis* J. WEISE, 1893  
*Galleruca nasturtii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PAYKULL, 1799)  
*Crioceris nasturtii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1801)  
*Altica nasturtii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LATREILLE, 1804)  
*Haltica nasturtii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PANZER, 1813)  
*Thyamis nasturtii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)  
*Teinodactyla nasturtii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)  
*Longitarsus nasturtii* ab. *auctumnalis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to TOMLIN et SHARP, 1912).  
*Thyamis nasturtii* var. *auctumnalis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KUHNT, 1913)  
*Longitarsus nasturtii* ab. *auctumnalis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to WANKA, 1920).  
*Longitarsus nasturtii* fa. *auctumnalis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to G. MÜLLER, 1953).  
 Distribution: Europe except for N, E Caucasus, Kazakhstan, Kirghizstan, Asian Russia (All over Siberia to Far East), Mongolia, NE China, Tibet.

***Longitarsus (Longitarsus) niger* (KOCH, 1803)**

- Haltica nigra* KOCH, 1803  
*Longitarsus elongatus* BACH, 1856  
*Longitarsus niger* ab. *peregii* KASZAB, 1962  
*Thyamis nigra*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)  
 Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey.

***Longitarsus (Longitarsus) nigerrimus* (GYLLENHAL, 1827)**

- Haltica nigerrima* GYLLENHAL, 1827  
*Thyamis atricornis* STEPHENS, 1834  
*Thyamis nigerrimus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)  
 Distribution: Europe except for N, Caucasus, Asian Russia (Amurland).

***Longitarsus (Longitarsus) nigrofasciatus* (GOEZE, 1777)**

- Chrysmela nigrofasciata* GOEZE, 1777  
*Altica marginata* GEOFFROY, 1785

*Haltica lateralis* ILLIGER, 1807  
*Longitarsus saltator* WOLLASTON, 1854  
*Longitarsus fractus* WOLLASTON, 1857  
*Longitarsus longipes* BACH, 1856  
*Teinodactyla trilineolata* FOUDRAS, 1860  
*Teinodactyla saltator* ALLARD, 1861  
*Thyamis patruelis* ALLARD, 1866  
*Thyamis distinguenda* RYE, 1872  
*Longitarsus patruelis* LEESBERG, 1882  
*Longitarsus lateralis* var. *patruelis* J. WEISE, 1893  
*Longitarsus lateralis* var. *domesticus* J. WEISE, 1893  
*Thyamis distinguendus* BEDEL, 1898  
*Thyamis nigrofasciata secutoria* PEYERIMHOFF, 1911  
*Longitarsus nigrofasciatus* var. *unionis* J. SAHLBERG, 1913  
*Longitarsus nigrofasciatus* ab. *ratruelis* HEIKERTINGER et CSIKI, 1939  
*Longitarsus nigrofasciatus* fa. *rudipennis* MOHR, 1965  
*Longitarsus nigrofasciatus* var. *distinguendus* KEVAN, 1967  
*Longitarsus lateralis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1858)  
*Teinodactyla lateralis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOUDRAS, 1860)  
*Thyamis lateralis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)  
*Thyamis saltator*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)  
*Thyamis fracta*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)  
*Thyamis nigrofasciata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BEDEL, 1898)  
*Longitarsus nigrofasciatus* var. *secutorius*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to NORDMAND, 1937)  
 Distribution: Europe except for extreme N, E Caucasus, Turkey, Near and Middle East, Afghanistan, Kazakhstan, Central Asia, Asian Russia (W Siberia).

***Longitarsus (Longitarsus) nitidamiculus* KIMOTO, 1965**

Distribution: Asian Russia (Maritime Prov., Ussuri Area, Kuriles Isles), Korea, Japan.

***Longitarsus (Longitarsus) nitidus* JACOBY, 1885**

Distribution: Asian Russia (Maritime Prov., Ussuri Area), NE China, Korea, Japan.

***Longitarsus (Longitarsus) noricus* LEONARDI, 1976**

Distribution: Southern parts of Central Europe, Caucasus, Transcaucasia.

***Longitarsus (Longitarsus) oblitteratoides* GRUEV, 1973**

Distribution: W and S Europe, Transcaucasia.

***Longitarsus (Longitarsus) oblitteratus* (ROSENHAUER, 1847)**

*Teinodactyla oblitterata* ROSENHAUER, 1847  
*Longitarsus consociatus* FÖRSTER, 1849  
*Teinodactyla pulex* FOUDRAS, 1860  
*Longitarsus oblitteratus* var. *meridionalis* J. WEISE, 1888  
*Thyamis oblitterata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)  
*Longitarsus pulex*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LEESBERG, 1882)  
*Longitarsus oblitteratus meridionalis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)  
*Longitarsus oblitteratus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to NORMAND, 1937)

Distribution: Central and SE Europe, Transcaucasia, Turkey, Near and Middle East.

***Longitarsus (Longitarsus) ochroleucus* (MARSHAM, 1802)***Chrysomela ochroleuca* MARSHAM, 1802*Altica albella* DUMERIL, 1816*Longitarsus cognatus* WOLLASTON, 1860*Longitarsus ochroleucus* ab. *subnigra* ROUBAL, 1943*Haltica ochroleuca*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GYLLENHAL, 1827)*Thyamis ochroleuca*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)*Altica ochroleuca*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to H. LUCAS, 1849)*Teinodactyla ochroleuca*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)*Teinodactyla cognata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1861)*Longitarsus nervosus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BOROWIEC, 1979)

Distribution: Europe, Caucasus, Transcaucasia, Turkey, N Africa, Canary Isles.

***Longitarsus (Longitarsus) pallidicornis* KUTSCHERA, 1863***Thyamis pallidicornis* ALLARD, 1866*Longitarsus hubenthali* WANKA, 1917*Longitarsus pallidicornis hubenthali*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to DOGUET, 1994)

Distribution: Mountains of W, Central, and E Europe.

***Longitarsus (Longitarsus) parvulus* (PAYKULL, 1799)***Galeruca parvula* PAYKULL, 1799*Haltica pumila* ILLIGER, 1807*Haltica pygmaea* ROGER, 1856*Thyamis maderensis* ALLARD, 1863*Longitarsus ater* LEESBERG, 1882*Longitarsus parvula* var. *concinus* J. WEISE, 1888*Longitarsus maderensis* HEIKERTINGER, 1930*Haltica parvula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KOCH, 1803)*Thyamis parvula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)*Teinodactyla parvula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)*Thyamis parvula* var. *concinus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PORTEVIN, 1934)*Longitarsus parvulus* ab. *concinus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to NORMAND, 1937)*Longitarsus parvulus* fa. *concinus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to G. MÜLLER, 1953)*Longitarsus absinthii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BOROWIEC, 1979)

Distribution: Europe, Caucasus, Transcaucasia, Turkey, Near East, Asian Russia (W and Middle Siberia, Sayan, Cisbaikalia), N Africa, Canary Isles, Madeira.

***Longitarsus (Longitarsus) pellucidus* (FOU DRAS, 1860)***Teinodactyla pellucida* FOU DRAS, 1860*Teinodactyla testacea* ALLARD, 1860*Thyamis paleacea* MULSANT et REY, 1874*Longitarsus pellucidus* var. *nigriventris* J. WEISE, 1893*Longitarsus pellucidus* ab. *nigriventris* NORMAND, 1937*Longitarsus ionius* MOHR, 1962*Longitarsus testaceus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STIERLIN, 1861)*Thyamis pellucida*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

*Longitarsus paleaceus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1912)

Distribution: Europe, Caucasus, Transcaucasia, Turkey, Near and Middle East, Kazakhstan, Central Asia, Mongolia, N Africa. Introduced to N America.

***Longitarsus (Longitarsus) picicollis* J. WEISE, 1900**

*Longitarsus tabidus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LOPATIN, 1963)

Distribution: Bulgaria, Rumania and Ukraine, Caucasus, Transcaucasia, Turkey, Middle East, Kazakhstan, Central Asia.

***Longitarsus (Longitarsus) pratensis* (PANZER, 1794)**

*Altica pratensis* PANZER, 1794

*Haltica pusilla* GYLLENHAL, 1813

*Thyamis collaris* STEPHENS, 1832

? *Longitarsus testaceus* MELSHEIMER, 1847

*Teinodactyla medicaginis* ALLARD, 1860

*Thyamis obsoleta* MULSANT et REY, 1874

*Thyamis funerea* MULSANT et REY, 1874

*Longitarsus pratensis* var. *collaris* J. WEISE, 1893

*Longitarsus pratensis* var. *funereus* J. WEISE, 1893

*Longitarsus pratensis* var. *ciliifer* MÉQUIGNON, 1948

*Thyamis pratensis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

*Thyamis pusilla*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

*Longitarsus pusillus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)

*Teinodactyla pusilla*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FODRAS, 1860)

*Longitarsus medicaginis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KUTSCHERA, 1863)

*Thyamis medicaginis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

*Longitarsus obsoletus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1913)

*Longitarsus pratensis* ab. *collaris*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1939).

*Longitarsus pratensis* ab. *funereus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1939).

*Longitarsus desertorum*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to WARCHALOWSKI, 1967)

Distribution: Europe, Caucasus, Transcaucasia, Turkey, Near and Middle East, Kazakhstan, Central Asia, N Africa, Carary Isles. Introduced to N America.

***Longitarsus (Longitarsus) pulmonariae* J. WEISE, 1893**

*Thyamis pratensis* ALLARD, 1860

*Teinodactyla femoralis* FODRAS, 1860

*Longitarsus pulmonariae* J. WEISE, 1889 [nomen nudum]

*Thyamis pulmonariae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BUYSSON, 1908)

Distribution: Central and SE Europe, Caucasus.

***Longitarsus (Longitarsus) quadriguttatus* (PONTOPPIDAN, 1765)**

*Chrysomela quadriguttata* PONTOPPIDAN, 1765

*Altica quadripustulata* FABRICIUS, 1775

*Altica quadrinotata* GMELIN, 1790

*Chrysomela cynoglossi* MARSHAM, 1802

- Haltica quadrimaculata* KOCH, 1803  
*Thyamis quadripunctulata* STEPHENS, 1839  
*Longitarsus quadrimaculatus* WEIDENBACH, 1859  
*Longitarsus quariguttatus* var. *binotatus* J. WEISE, 1888  
*Longitarsus quariguttatus* var. *immaculatus* J. WEISE, 1888  
*Longitarsus quariguttatus* var. *vittatus* J. WEISE, 1888  
*Thyamis quariguttata* BEDEL, 1898  
*Chrysomela quadripustulata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GOEZE, 1777)  
*Galeruca quadripustulata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1787)  
*Altica quadripunctata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GEOFFROY, 1789)  
*Altica quatuorpusculata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to OLIVIER, 1789)  
*Crioceris quadripustulata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1801)  
*Haltica quadripustulata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ILLIGER, 1807)  
*Haltica quadriguttata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to DUFTSCHMIDT, 1825)  
*Thyamis quadripustulata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)  
*Longitarsus quadripustulatus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)  
*Teinodactyla quadrimaculata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860 and ALLARD, 1860)  
*Longitarsus quariguttatus* ab. *binotatus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1939)  
*Longitarsus quariguttatus* ab. *immaculatus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1939)  
*Longitarsus quariguttatus* ab. *vittatus* C: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1939)  
 Distribution: Europe except for N, Ciscaucasia, E Caucasus, Turkey.

***Longitarsus (Longitarsus) ratshensis* KHNZORIAN, 1962**

Distribution: Caucasus (Georgia).

***Longitarsus (Longitarsus) rectilineatus* (FOUDRAS, 1860)**

- Teinodactyla rectilineata* FOU DRAS, 1860  
*Longitarsus spilotus* J. WEISE, 1900  
*Longitarsus rectilineatus* ab. *kaufmanni* KASZAB, 1962  
*Longitarsus rectilineatus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)  
*Longitarsus rectilineatus spilotus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GRUEV et KRÁL, 1975)  
 Distribution: W, S and southern parts of Central Europe, Caucasus, Transcaucasia, Turkey, Near and Middle East, Afghanistan, Kazakhstan, Turkey, N Africa.

***Longitarsus (Longitarsus) reichei* (ALLARD, 1860)**

- Teinodactyla reichei* ALLARD, 1860  
*Longitarsus fuscus* KUTSCHERA, 1864  
*Thyamis funerea* MULSANT et REY, 1874

*Thyamis reichei*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)  
*Longitarsus pratensis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

*Longitarsus scutellaris*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LEONARDI, 1972)

Distribution: Europe except for extreme N, Caucasus, Transcaucasia, Iran.

***Longitarsus (Longitarsus) rubellus* (FOUDRAS, 1860)**

*Teinodactyla rubellus* FOU DRAS, 1860

*Teinodactyla brunnea* ALLARD, 1860 (part.)

*Longitarsus gravidulus* KUTSCHERA, 1863

*Thyamis gravidula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

Distribution: Mountains of Central and E Europe.

***Longitarsus (Longitarsus) rubiginosus* (FOUDRAS, 1860)**

*Teinodactyla rubiginosa* FOU DRAS, 1860

*Teinodactyla flavicornis* ALLARD, 1860

*Longitarsus amurensis* J. WEISE, 1887

*Longitarsus rubiginosus* var. *fumigatus* J. WEISE, 1893

*Longitarsus rubiginosus* var. *profugus* J. WEISE, 1893

*Thyamis flavicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

*Longitarsus flavicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LEESBERG, 1882)

*Thyamis rubiginosa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BUYSSON, 1908)

*Thyamis rubiginosa* var. *fumigatus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PORTEVIN, 1934)

*Thyamis rubiginosa* var. *profugus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PORTEVIN, 1934)

*Longitarsus rubiginosus* var. *profugus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KASZAB, 1962)

Distribution: All over Palearctic except for SW and E Mediterranean, and N Africa. Introduced to N America.

***Longitarsus (Longitarsus) sahlbergi* PIC, 1907**

*Longitarsus weisei* JACOBSON, 1901

Distribution: S Kazakhstan.

***Longitarsus (Longitarsus) salarius* LOPATIN et KULENOVA, 1985**

Distribution: Kazakhstan.

***Longitarsus (Longitarsus) salviae* GRUEV, 1975**

*Longitarsus eminus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GRUEV, 1973)

Distribution: Europe except for N and E, Crimea, Caucasus, Transcaucasia.

***Longitarsus (Longitarsus) scutellaris* (REY, 1874)**

*Thyamis scutellaris* REY, 1874

*Longitarsus medicaginis* KUTSCHERA, 1863

Distribution: Europe except for extreme N, Caucasus, Transcaucasia, Turkey, Mongolia.

***Longitarsus (Longitarsus) silbursaicus* LOPATIN, n. nov.**

*Longitarsus (Longitarsus) seriepunctatus* LOPATIN, 1966, non Roubal, 1943

Distribution: Tajikistan (S Slopes of Hissar Mountain Range).



***Longitarsus (Longitarsus) strigicollis* WOLLASTON, 1864***Thyamis exiguus* BEDEL, 1898*Longitarsus decolor* HEIKERTINGER (in coll.) [nomen nudum]*Longitarsus curtis* var. *ciliifer* MÉQUIGNON, 1948*Longitarsus bombycinus* MOHR, 1962*Longitarsus fowleri*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GRUEV, 1971)

Distribution: S and southern parts of Central Europe, Caucasus (Georgia), N Africa, Canary Isles.

***Longitarsus (Longitarsus) substriatus* KUTSCHERA, 1863***Thyamis abdominalis* DUFTSCHMIDT, 1825*Longitarsus abdominalis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)*Thyamis substriata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to REY, 1873)

Distribution: W, S, SE and southern parts of Central Europe, N Caucasus, Kazakhstan.

***Longitarsus (Longitarsus) succineus* (FOUDRAS, 1860)***Teinodactyla succinea* FOU DRAS, 1860*Haltica pratensis* var. c GYLLENHAL, 1813*Teinodactyla laevis* ALLARD, 1860*Thyamis amacula* BALY, 1874*Longitarsus succineus* var. *perfectus* J. WEISE, 1893*Longitarsus aeruginosus* TOMLIN et SHARP, 1912*Longitarsus succineus* ab. *perfectus* HEIKERTINGER et CSIKI, 1939*Longitarsus arakii* CHŪJŌ, 1942*Longitarsus laevis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KUTSCHERA, 1863)*Thyamis laevis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)*Thyamis succinea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BEDEL, 1898)*Longitarsus amiculum*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

Distribution: All over Palaearctic. Introduced to N America.

***Longitarsus (Longitarsus) suturatus* (FOUDRAS, 1860)***Teinodactyla suturata* FOU DRAS, 1860*Thyamis fulvicollis* STEPHENS, 1832*Longitarsus sisymbrii* BARGAGLI, 1878*Longitarsus suturatus* var. *scophulariae* NORMAND, 1937*Thyamis suturata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

Distribution: W and S Mediterranean. ? Crimea, ? N Caucasus.

***Longitarsus (Longitarsus) suturellus* (DUFTSCHMIDT, 1825)***Haltica suturella* DUFTSCHMIDT, 1825*Thyamis fuscicollis* STEPHENS, 1832*Thyamis dimidiata* STEPHENS, 1834*Teinodactyla melanocephala* KIESENWETTER, 1847*Teinodactyla senecionis* MOTSCHULSKY, 1851*Thyamis thoracica* ALLARD, 1860*Longitarsus suturellus* var. *paludosus* J. WEISE, 1893*Longitarsus suturellus* var. *macer* J. WEISE, 1893*Longitarsus freminvillei* GUILLEBEAU, 1895*Longitarsus suturellus* fa. *limbatus* KOLBE, 1920*Longitarsus suturellus* fa. *testis* KOLBE, 1920*Thyamis suturellus* var. *paludosus* PORTEVIN, 1934

- Longitarsus melanocephalus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1858)  
*Longitarsus senecionis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BACH, 1856)  
*Longitarsus thoracicus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KUTSCHERA, 1863)  
*Thyamis thoracica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)  
*Thyamis suturella*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BEDEL, 1898)  
*Longitarsus fuscicollis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to TOMLIN et SHARP, 1912)  
*Longitarsus suturellus* ab. *paludosus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1939)

Distribution: All over Palaearctic except for Mediterranean, Transcaucasia, Central Asia.

***Longitarsus (Longitarsus) symphyti* HEIKERTINGER, 1912**

- ? *Longitarsus aeruginosus* KUTSCHERA, 1864  
*Longitarsus aeruginosus* var. *luctator* J. WEISE, 1893  
*Longitarsus luctator* HUBENTHAL, 1922  
*Thyamis symphyti* PORTEVIN, 1934  
*Longitarsus symphyti* fa. *obscura* DÖBERL, 1994  
 Distribution: Europe except for N and S, W Caucasus, Uzbekistan, Asian Russia (Tuva).

***Longitarsus (Longitarsus) tabidus* (FABRICIUS, 1775)**

- Altica tabida* FABRICIUS, 1775  
*Galleruca sisymbrii* FABRICIUS, 1792  
*Chrysomela dorsalis* ROSSI, 1792  
*Altica verbasci* PANZER, 1794  
*Chrysomela thapsi* MARSHAM, 1802  
 ? *Alticae jaccae* PANZER, 1809  
*Haltica borealis* ZETTERSTEDT, 1828  
*Longitarsus pallens* KUTSCHERA, 1863  
*Longitarsus tabidus* var. *dorsalis* OERTZEN, 1866  
*Longitarsus tabidus* var. *vulgaris* J. WEISE, 1893  
*Longitarsus tabidus* var. *elongatus* J. WEISE, 1893  
*Longitarsus tabidus* ab. *nigrinus* LEONI, 1907  
*Longitarsus tabidus* fa. *rugipennis* MOHR, 1965  
*Longitarsus tabidus* fa. *vulgaris* ROSKA, 1976  
*Chrysomela tabida*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1781)  
*Galleruca tabida*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1792)  
*Crioceris tabida*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1801)  
*Crioceris sisymbrii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1801)  
*Chrysomela verbasci*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to MARSHAM, 1802)  
*Haltica verbasci*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ILLIGER, 1807)  
*Thyamis verbasci*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)  
*Thyamis thapsi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)  
*Longitarsus verbasci*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)  
*Teinodactyla verbasci*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)  
*Teinodactyla thapsi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)  
*Teinodactyla sisymbrii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)  
*Longitarsus sisymbrii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KUTSCHERA, 1863)  
*Longitarsus borealis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to C. G. THOMSON, 1866)

- Thyamis tabida*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BEDEL, 1898)  
*Longitarsus tabidus* ab. *thapsi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1912)  
*Thyamis tabidus* ab. *thapsi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PORTEVIN, 1934)  
*Thyamis tabidus* var. *elongatus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PORTEVIN, 1934)  
*Longitarsus tabidus* ab. *sisymbrii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1939)  
*Longitarsus tabidus* ab. *elongatus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1939)  
*Longitarsus tabidus* var. *thapsi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KEVAN, 1967)  
 Distribution: Europe, Caucasus, Transcaucasia, Turkey, Near and Middle East, Kazakhstan, Turkmenistan, Kirghizstan, Mongolia, N Africa.

***Longitarsus (Longitarsus) tienshanicus* KONSTANTINOV, 1992**

Distribution: Uzbekistan (Ugam Mountain Range).

***Longitarsus (Longitarsus) tischechkini* KONSTANTINOV, 2000**

Distribution: Kazakhstan (Zaili Alatau).

***Longitarsus (Longitarsus) tournefortiae* L. MEDVEDEV et VORONOVA, 1979**

*Longitarsus tournefortii* L. MEDVEDEV, 1979

*Longitarsus rangoonensis* KRÁL, 1980

*Longitarsus longiseta*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. MEDVEDEV et VORONOVA, 1977)

Distribution: Mongolia.

***Longitarsus (Longitarsus) transbaicalicus* OGLOBLIN, 1917**

*Longitarsus exoletus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. MEDVEDEV et VORONOVA, 1977)

Distribution: S and Asian Russia (Middle Siberia, Cisbaikalia, Transbaikalia, Amurland, Amurland), Mongolia.

***Longitarsus (Longitarsus) trepidus* WARCHALOWSKI, 1973**

Distribution: Transcaucasia, Turkey, Middle East.

***Longitarsus (Longitarsus) tristis* J. WEISE, 1888**

*Longitarsus fulgens* ab. *tristis* HEIKERTINGER et CSIKI, 1939

*Longitarsus pannonicus* KASZAB, 1962

Distribution: Central and SE Europe, E Caucasus.

***Longitarsus (Longitarsus) tshikatunovi* LOPATIN, 1966**

Distribution: Kazakhstan, Kirghizstan.

***Longitarsus (Longitarsus) violentoides* KONSTANTINOV, 2000**

Distribution: Armenia (Tsahkadzor).

***Longitarsus (Longitarsus) violentus* J. WEISE, 1893**

Distribution: SE European Russia, Caucasus, Transcaucasia, Turkey, Middle East, Kazakhstan, Central Asia, Asian Russia (S and Middle Siberia, Transbaikalia, Tuva, Yakutia, Beringia), Mongolia.

Subgenus: *Testergus* J. WEISE, 1893***Longitarsus (Testergus) anchusae* (PAYKULL, 1799)***Caleruca anchusae* PAYKULL, 1799? *Chrysomela pulex* SCHRANK, 1781? *Altica ater* LAICHARTING, 1781*Haltica gagathina* DUFTSCHMIDT, 1825*Thyamis convexa* STEPHENS, 1834*Longitarsus pulex* OERTZEN, 1886*Longitarsus anchusae* var. *punctatissimus* J. WEISE, 1888*Longitarsus niger* BUDDEBERG, 1888*Haltica anchusae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KOCH, 1803)*Thyamis anchusae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)*Teinodactyla anchusae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)*Thyamis nigra*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BEDEL, 1900)

Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, Near and Middle East, Kazakhstan, Tajikistan, Asian Russia (Altai, Transbaikalia).

***Longitarsus (Testergus) anatolicus* J. WEISE, 1900***Longitarsus anatolicus* J. WEISE, 1900*Longitarsus anchusae anatolicus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

Distribution: Armenia, Azerbaijan, Turkey, Tajikistan, Uzbekistan.

***Longitarsus (Testergus) brachypterus* J. WEISE, 1890**

Distribution: Caucasus (Arax Valley).

***Longitarsus (Testergus) ellipticus* REITTER, 1909***Testergus ellipticus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BECHYNÉ, 1957)

Distribution: S Kazakhstan.

***Longitarsus (Testergus) excisipennis* LOPATIN, 1967**

Distribution: S Tajikistan.

***Longitarsus (Testergus) fuscoaeneus* L. REDTENBACHER, 1849***Longitarsus lopatini* KHNZORIAN, 1959*Longitarsus persicus* WARCHALOWSKI, 1967*Teinodactyla fuscoaenea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)*Thyamis fuscoaeneus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)*Testergus fuscoaeneus*: BECHYNE, 1957. GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BECHYNÉ, 1959)

Distribution: Europe except for N, Crimea, Caucasus, Transcaucasia, Turkey, Near East.

***Longitarsus (Testergus) hittita* BIONDI, 1995**

Distribution: Armenia, Turkey.

***Longitarsus (Testergus) imitator* LOPATIN, 1967**

Distribution: Kazakhstan, Central Asia.

***Longitarsus (Testergus) josiphi* KONSTANTINOV, 1994***Longitarsus philippi* KONSTANTINOV, 1992

Distribution: Kazakhstan (W Tian Shan).

***Longitarsus (Testergus) lederi* J. WEISE, 1889**

Distribution: Ciscaucasia, Caucasus, Transcaucasia.

***Longitarsus (Testergus) mohri* LOPATIN, 1963**

Distribution: S Afghanistan.

***Longitarsus (Testergus) pinguis* J. WEISE, 1888***Haltica? ater* ILLIGER, 1807*Longitarsus anchusae* var. *pinguis* HEIKERTINGER, 1930*Longitarsus alpigradus* KHNZORIAN, 1962

Distribution: SE and southern parts of Central Europe, Turkey, Syria.

***Longitarsus (Testergus) primaeveris* LOPATIN, 1967**

Distribution: S Kazakhstan, S Kirghizstan, SW Tajikistan, S Uzbekistan.

***Longitarsus (Testergus) pubescens* J. WEISE, 1889**

Distribution: Caucasus (Abkhazia, Georgia, Armenia).

***Longitarsus (Testergus) radiatus* KONSTANTINOV, 1992**

Distribution: Uzbekistan (W Tian Shan).

***Longitarsus (Testergus) sengloki* KONSTANTINOV, 1994**

Distribution: Tajikistan (Senglok Mountain Range).

***Longitarsus (Testergus) sogdianus* LOPATIN, 1956**

Distribution: Tajikistan (South of Hissar Mountain Range).

***Longitarsus (Testergus) tmetopterus* JACOBSON, 1893***Longitarsus nurataicus* PALIJ, 1970*Longitarsus zeravshanicus* PALIJ, 1970*Testergus tmetopterus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BECHYNÉ, 1957)

Distribution: S Turkmenistan, Uzbekistan, S Kazakhstan, Kirghizstan, Afghanistan.

***Longitarsus (Testergus) turcomanorum* LOPATIN, 1967***Longitarsus vestimnalis* PALIJ, 1970

Distribution: S Turkmenistan, Uzbekistan.

***Longitarsus (Testergus) weisei* GUILLEBEAU, 1895***Longitarsus fuscoaeneus* var. *maassi* HUBENTHAL, 1898*Longitarsus weisei* ab. *maassi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1939)

Distribution: W Europe, Albania, Poland, Turkey, Middle East, Turkmenistan, Tajikistan, Nepal, China.

**Genus: *Hermaeophaga* FOUDRAS, 1860***Orthocrepis* J. WEISE, 1888***Hermaeophaga mercurialis* (FABRICIUS, 1792)***Galleruca mercurialis* FABRICIUS, 1792*Haltica mercurialis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ILLIGER, 1807)*Sphaeroderma mercurialis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

*Graptodera mercurialis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1859)

*Altica mercurialis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to AUBÉ, 1859)

*Linostoma mercurialis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)

*Hermaphysalis cicatrix*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to MAZURKOWA, 1951)

Distribution: Europe except for N, Caucasus, Turkey.

***Hermaphysalis nigricornis* OGLOBLIN, 1921**

Distribution: Asian Russia (Transbaikalia: Dauria).

***Hermaphysalis ruficollis* (H. LUCAS, 1849)**

*Haltica ruficollis* H. LUCAS, 1849

*Leiometopona suturalis* BRANCSIK, 1899

*Orthocrepis ruficollis bamakoensis* BECHYNÉ, 1955

*Graptodera ruficollis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1859)

*Linostoma ruficollis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1859)

*Orthocrepis ruficollis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. WEISE, 1888)

Distribution: Mediterranean, Balkans (Macedonia), Turkey, Near and Middle East, Kazakhstan, Central Asia, India (Mysore), Sri Lanka, N and W Africa.

**Genus: *Altica* GEOFFROY, 1762**

*Graptodera* CHEVROLAT in DEJEAN, 1837

*Haltica* CHAPUIS, 1875

***Altica aenescens* (J. WEISE, 1888)**

*Haltica lythri* var. *aenescens* J. WEISE, 1888

*Haltica engstroemi weisei* KEMNER, 1919

*Haltica pseudolythri* HEIKERTINGER et CSIKI, 1939

*Haltica aenescens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ŁOMNICKI, 1913)

*Haltica aenescens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KUNTZE, 1930)

*Haltica engstroemi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to OGLOBLIN, 1948)

Distribution: Europe except for extreme N and S.

***Altica ancyrensis* (J. WEISE, 1897)**

*Haltica ancyrensis* J. WEISE, 1897

Distribution: Transcaucasia, Turkey, Near East.

***Altica balassogloi* (JACOBSON, 1892)**

*Haltica balassogloi* JACOBSON, 1892

*Haltica suworovi* OGLOBLIN, 1921

*Haltica lopatini* PALI, 1968

Distribution: Afghanistan, Pakistan, Kazakhstan, Kirghizstan, Mongolia.

***Altica bicarinata* (KUTSCHERA, 1860)**

*Haltica bicarinata* KUTSCHERA, 1860

*Haltica ciliciensis* J. WEISE, 1897

Distribution: E Mediterranean, Turkey, Iran.

***Altica bisulcata* (J. WEISE, 1887)**

*Haltica bisulcata* J. WEISE, 1887

Distribution: Mongolia, Asian Russia (Amurland, Maritime Prov.).

***Altica brevicollis* Foudras, 1860***Haltica oleracea* Ratzeburg, 1839*Graptodera corylii* Allard, 1860*Haltica quercetorum* var. *azurescens* J. Weise, 1888*Haltica quercetorum* var. *brevicollis*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to J. Weise, 1888)*Haltica erucacae* var. *brevicollis*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Seidlitz, 1891)*Haltica brevicollis*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Acloque, 1896)

Distribution: Europe except for extreme S and N, Caucasus, Transcaucasia, Turkey, Iran, Kazakhstan.

***Altica breviuscula* (J. Weise, 1900)***Haltica breviuscula* J. Weise, 1900

Distribution: Azerbaijan, Iran.

***Altica carduorum* Guérin-Ménéville, 1858***Graptodera carduorum* Allard, 1859*Graptodera basalis* Allard, 1859*Haltica cirsii* Israelson, 1956*Altica carduorum cirsii*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Strejček, 1993)

Distribution: Europe except for N, Transcaucasia, Turkey, Kazakhstan, Kirghizstan, Tajikistan. Introduced to N America.

***Altica carinthiaca* (J. Weise, 1888)***Haltica carinthiaca* J. Weise, 1888*Haltica issykensis* Ogloblin, 1921*Haltica daghestanica* Shapiro, 1970

Distribution: N, NW and Mountains of Central Europe, Transcaucasia, Kirghizstan, Uzbekistan, Asian Russia (N Baikal).

***Altica chamaenerii* (Lindberg, 1926)***Haltica chamaenerii* Lindberg, 1926

Distribution: N and Mountains of Central Europe, Transcaucasia, Kirghizstan, Asian Russia (Beringia, Kamchatka).

***Altica circaeae* Ohno, 1960**

Distribution: Asian Russia (S Kuriles Isles), Japan.

***Altica cornivorax* Král, 1969***Haltica ampelophaga*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Kaszab, 1962)*Altica viridula*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Král, 1964)

Distribution: Southern parts of Central Europe, Ukraine, Turkey.

***Altica deserticola* (J. Weise, 1889)***Haltica deserticola* J. Weise, 1889*Haltica glycyrrhizae* Ogloblin, 1925*Haltica glycyrrhizae eichleri* Ogloblin, 1925*Haltica glycyrrhizae chotanica* Ogloblin, 1925*Haltica deserticola glycyrrhizae*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Lopatin, 1958)

Distribution: SE European Russia, S Urals, SW Kazakhstan, W Caspian Plain, Cuscaucasia, Transcaucasia, Near East, Turkey, Afghanistan, Turkmenistan, Kazakhstan, Kirghizstan, N and NW China, Mongolia, Taiwan.

***Altica difficilis* (OGLOBLIN, 1921)**

*Haltica difficilis* OGLOBLIN, 1921

Distribution: Kirghizstan.

***Altica engstroemi* (J. SAHLBERG, 1894)**

*Haltica engstroemi* J. SAHLBERG, 1894

Distribution: N Europe, SE Kazakhstan, Asian Russia (W and Middle Siberia, Altai, Transbaikalia, Chukotka, Kamchatka).

***Altica fragariae* (NAKANE, 1955)**

*Haltica fragariae* NAKANE, 1955

*Altica sanguisorbae* OHNO, 1960

Distribution: E and NE China, Korea, Asian Russia (Sakhalin), Japan.

***Altica fruticola* (J. WEISE, 1888)**

*Haltica fruticola* J. WEISE, 1888

*Haltica lythri?* *fruticola* HEIKERTINGER et CSIKI, 1940

*Altica ampelophaga*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KRÁL, 1966)

Distribution: Central Europe, Ukrainian Carpathians.

***Altica globicollis* (J. WEISE, 1889)**

*Haltica globicollis* J. WEISE, 1889

Distribution: SW Caucasus, Transcaucasia, NW Iran, Turkey.

***Altica hampei* (ALLARD, 1867)**

*Graptodera hampei* ALLARD, 1867

*Altica khnzoriani* KRÁL, 1969

*Haltica khnzoriani*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to TOMOV et GRUEV, 1975)

Distribution: Crimea, Caucasus, Transcaucasia, Turkey.

***Altica helianthemi* (ALLARD, 1859)**

*Graptodera helianthemi* ALLARD, 1859

*Altica pusilla* DUFTSCHMIDT, 1825

*Graptodera potentillae* ALLARD, 1859

*Altica montana* FOU DRAS, 1860

*Haltica cognata* KUTSCHERA, 1864

*Graptodera montana*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1861)

*Haltica helianthemi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to OERTZEN, 1886)

*Haltica pusilla* ab. *montana* J. WEISE, 1888

Distribution: Europe except for N, Caucasus, Kazakhstan, Asian Russia (Tuva, Transbaikalia, Yakutia).

***Altica impressicollis* (REICHE, 1862)**

*Graptorera impressicollis* REICHE, 1862

*Haltica lacunosa* J. WEISE, 1900

*Haltica impressicollis*: GRUEV et DÖBERL, 1997 (in synonymy attributed to SAINTE-CLAIRE DEVILLE, 1910)

Distribution: Central, S and SE Europe, Caucasus, Turkey, Near and Middle East.



***Altica ivlievi* (L. MEDVEDEV, 1968)***Haltica ivlievi* L. MEDVEDEV, 1968

Distribution: Asian Russia (Magadan Distr.).

***Altica jarmilae* KRÁL, 1979**

Distribution: E Mediterranean, Turkey, Caucasus (Georgia).

***Altica kozlovi* (OGLOBLIN, 1921)***Haltica kozlovi* OGLOBLIN, 1921

Distribution: N China.

***Altica longicollis* (ALLARD, 1860)***Graptodera longicollis* ALLARD, 1860*Haltica britteni* SHARP, 1914*Haltica sandini* KEMNER, 1919*Haltica longicollis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. WEISE, 1803)

Distribution: Europe except for S.

***Altica lubischevi* (PALIJ, 1963)***Haltica lubischevi* PALIJ, 1963*Haltica aeneola* LOPATIN, 1956

Distribution: Kazakhstan, Kirghizstan, Tajikistan, China (Tibet).

***Altica lythri* AUBÉ, 1843***Haltica erucae* DUFESCHMIDT, 1825*Haltica indigacea* STEPHENS, 1832*Graptodera epilobii* ALLARD, 1859*Graptodera consobrina* ALLARD, 1860*Graptodera lythri*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to H. LUCAS, 1846)*Haltica lythri*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to C. G. THOMSON, 1866)*Haltica ampelophaga*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1912)*Haltica aenescens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to MOHR, 1965)

Distribution: Europe except for N.

***Altica mohri* LOPATIN, 1990**

Distribution: E Iran.

***Altica nuristanica* (LOPATIN, 1963)***Haltica nuristanica* LOPATIN, 1963

Distribution: E and NE Afghanistan.

***Altica oleracea* (LINNAEUS, 1758)***Chrysomela oleracea* LINNAEUS, 1758? *Haltica brevicollis* RATZEBURG, 1837? *Graptodera helianthemi* ALLARD, 1959*Graptodera sicula* ALLARD, 1859*Graptodera pusilla* ALLARD, 1866*Altica oleracea* ab. *splendens* REY, 1874*Graptodera longicollis* ALLARD, 1875*Haltica oleracea* var. *lugubria* J. WEISE, 1888*Haltica oleracea* var. *nobilis* J. WEISE, 1888*Haltica potentillae* BEDEL, 1900

*Haltica ytenensis* SHARP, 1914  
*Haltica oleracea* var. *geminata* OGLOBLIN, 1925  
*Haltica alchemillae* PALIJ, 1968  
*Galleruca oleracea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1792, PAYKULL, 1799 and KOCH, 1803)  
*Graptodera oleracea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1839)  
*Haltica helianthemii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GERHARDT, 1870)  
*Graptodera helianthemii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GERHARDT, 1874)  
*Altica pusilla*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to REY, 1874)  
*Altica helianthemii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to REY, 1874)  
*Haltica pusilla*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. WEISE, 1888)  
*Haltica oleracea* var. *nigra* FOWLER, 1900  
*Haltica helianthemii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BEDEL, 1900)  
Distribution: All over Palearctic except for extreme N.

***Altica opacifrons* (H. LINDBERG, 1938)**

*Haltica opacifrons* H. LINDBERG, 1938  
Distribution: N Europe.

***Altica palustris* (J. WEISE, 1888)**

*Haltica palustris* J. WEISE, 1888  
? *Haltica crassa* ALLARD, 1889  
*Haltica palustris* var. *lostiae* BURLINI, 1942  
Distribution: Europe, Caucasus, Transcaucasia, Turkey, Middle East, Kazakhstan, Central Asia.

***Altica pamiranica* (J. WEISE, 1889)**

*Haltica pamiranica* J. WEISE, 1889  
*Haltica tianshanica* OGLOBLIN, 1921  
Distribution: Kazakhstan, Kirghizstan, Tajikistan, NW China.

***Altica plotnicovi* (PALIJ, 1963)**

*Haltica plotnicovi* PALIJ, 1963  
Distribution: Kirghizstan.

***Altica pontica* (OGLOBLIN, 1925)**

*Haltica pontica* OGLOBLIN, 1925  
*Haltica lythri ampelophaga* var. *pontica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)  
Distribution: Turkey.

***Altica quercetorum quercetorum* FOU DRAS, 1860**

*Altica erucacae* OLIVIER, 1808  
*Graptodera erucacae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1859)  
*Haltica erucacae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LETZNER, 1870)  
*Haltica quercetorum*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. WEISE, 1888)  
*Haltica brevicollis quercetorum*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to NOVAK, 1952)  
Distribution: Central and S Europe, Caucasus, Transcaucasia, Turkey.

***Altica quercetorum saliceti* (J. WEISE, 1888)**

*Haltica saliceti* J. WEISE, 1888  
Distribution: Europe except for extreme N, Armenia.

***Altica rosicola* KRÁL, 1969**

Distribution: W Caucasus, Armenia.

***Altica sajanica* (CSIKI, 1901)**

*Haltica sajanica* CSIKI, 1901

*Haltica convexipennis* OGLOBLIN, 1921

*Haltica ampelophaga* var. *convexipennis* OGLOBLIN, 1921

*Haltica transbaicalica* OGLOBLIN, 1925

*Haltica mongolica* KRÁL, 1973

*Haltica deserticola*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1940).

*Haltica brevicollis transbaicalica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LOPATIN, 1967 and L. MEDVEDEV, 1974)

Distribution: Asian Russia (S Middle Siberia, Transbaikalia, Amurland).

***Altica semenovi* (JACOBSON, 1892)**

*Haltica semenovi* JACOBSON, 1892

*Haltica lazulina* OGLOBLIN, 1921

*Haltica alaiensis* PALIJ, 1968

*Haltica susamyrica* PALIJ, 1968

Distribution: Central Asia, NW China.

***Altica sibirica* (CSIKI, 1901)**

*Haltica sibirica* CSIKI, 1901

Distribution: Asian Russia (W Siberia).

***Altica stichai* KRÁL, 1965**

Distribution: W Caucasus, Transcaucasia.

***Altica talassicola* LOPATIN, 1976**

Distribution: S Kazakhstan, Kirghizstan.

***Altica talyshana* KONSTANTINOV, 1995**

*Altica lencorana* KONSTANTINOV, 1996

Distribution: Azerbaijan.

***Altica tamaricis* SCHRANK, 1785**

*Galleruca erucae* FABRICIUS, 1792

? *Haltica oleracea* var. *cyanea* ILLIGER, 1807

*Galeruca atriculata* BECK, 1817

*Haltica consobrina* DUFTSCHMIDT, 1825

? *Haltica indigaceae* STEPHENS, 1832

*Altica hippophaës* AUBÉ, 1843

*Altica azurea* KNOCH, 1843

*Graptodera hippophaes* ALLARD, 1859

*Haltica tamaricis* ab. *nigra* TENENBAUM, 1927

*Graptodera tamaricis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1865)

*Graptodera oleracea* var. *erucae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KELCH, 1846)

*Graptodera erucae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1839)

*Haltica erucae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

*Haltica tamaricis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to SCHÖNHERR, 1808)

*Chrysomela erucaae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to MARSHAM, 1802)  
*Altica erucaae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LATREILLE, 1804)  
 Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, Middle East, Kazakhstan, Uzbekistan, Asian Russia (S Siberia, Transbaikalia).

***Altica tristis* (OGLOBLIN, 1921)**

*Haltica tristis* OGLOBLIN, 1921  
 Distribution: Asian Russia (Transbaikalia).

***Altica tsharynensis* (OGLOBLIN, 1921)**

*Haltica tsharynensis* OGLOBLIN, 1921  
*Haltica deserticola*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to IABLOKOFF-KHNZORIAN, 1968)  
*Altica tamaricis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LOPATIN, 1977)  
 Distribution: Kazakhstan.

***Altica viridula* (J. WEISE, 1889)**

*Haltica viridula* J. WEISE, 1889  
*Haltica armeniaca* KHNZORIAN, 1954  
*Altica avellanae* KRÁL, 1964  
 Distribution: Caucasus, Transcaucasia, Iran.

***Altica weisei* (JACOBSON, 1892)**

*Haltica weisei* JACOBSON, 1892  
*Haltica laeviuscula* J. WEISE, 1889  
 Distribution: Asian Russia (NE and E Siberia), Korea, N China.

**Genus: *Batophila* FOUDRAS, 1860**

***Batophila acutangula* HEIKERTINGER, 1921**

*Batophila yangweii* CHEN, 1933  
*Batophila acutangula kamikochiana* NAKANE, 1958  
*Batophila acutangula omogo* NAKANE, 1958  
*Batophila acutangula yakuensis* NAKANE, 1958  
 Distribution: Asian Russia (Amurland, Ussuri Area, Maritime Prov., Sakhalin), Korea, N China, Taiwan.

***Batophila dogueti* DÖBERL, 1994**

Distribution: Azerbaijan, Iran.

***Batophila fallax* J. WEISE, 1888**

*Batophila aerata* var. *fallax* J. WEISE, 1888  
 Distribution: Southern Central and SE Europe, Caucasus, Transcaucasia, Turkey, Iran.

***Batophila olexai* KRÁL, 1964**

Distribution: Balkans, Transcaucasia, Turkey.

***Batophila rubi* (PAYKULL, 1799)**

*Crioceris rubi* PAYKULL, 1799  
*Chrysomela striatula* MARSHAM, 1802  
*Haltica rubi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KOCH, 1803)  
*Aphthona rubi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KELCH, 1846 and KALTENBACH, 1874)

*Podagrira rubi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)  
*Glyptina rubi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BEDEL, 1898)  
*Batophila fallax*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to TENENBAUM, 1929)  
 Distribution: Europe, Caucasus.

**Genus: *Mniophila* STEPHENS, 1832**

***Mniophila muscorum muscorum* (KOCH, 1803)**

*Haltica muscorum* KOCH, 1803  
*Mniophila wroblewski* WANKOWICZ, 1880  
*Mniophila muscorum* fa. *seriatopunctata* HEIKERTINGER, 1912  
*Mniophila muscorum* fa. *fortepunctata* HORION, 1939  
 Distribution: Europe except for SW and NE, Caucasus, Transcaucasia.

**Genus: *Lythraria* BEDEL, 1897**

***Lythraria salicariae* (PAYKULL, 1800)**

*Galeruca salicariae* PAYKULL, 1800  
*Haltica striatella* ILLIGER, 1807  
*Haltica nitidicollis* MOTSCHULSKY, 1860  
*Ochrosis nitidicollis* J. WEISE, 1886  
*Ochrosis nitidicollis* var. *picicollis* J. WEISE, 1886  
*Lythraria nitidicollis* var. *picicollis* EVERTS, 1903  
*Ochrosis salicariae* var. *unicolor* REITTER, 1906  
*Lythraria nitidicollis* OGLOBLIN, 1930  
*Lythraria salicariae* var. *unicolor* HEIKERTINGER, 1930  
*Paraphthonomorpha komiyamai* OHNO, 1960  
*Haltica salicariae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GYLLENHAL, 1813)  
*Crepidodera salicariae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1839)  
*Podagrira salicariae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)  
*Ochrosis salicariae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)  
 Distribution: All over Palaearctic except for E Mediterranean, N Africa and Central Asia.

**Genus: *Ochrosis* FOUDRAS, 1860**

***Ochrosis ventralis* (ILLIGER, 1807)**

*Haltica ventralis* ILLIGER, 1807  
 ? *Haltica salicaria* STEPHENS, 1832  
*Crepidodera abdominalis* KÜSTER, 1848  
*Haltica nigriventris* BACH, 1856  
*Crepidodera pisana* ALLARD, 1861  
*Haltica lubrica* WOLLASTON, 1864  
*Crepidodera rubicunda* PERRIS, 1864  
*Crepidodera corsica* ALLARD, 1866  
*Ochrosis kruperi* J. WEISE, 1886  
*Ochrosis ventralis* ab. *viridana* REITTER, 1906  
*Ochrosis ventralis* var. *corcyrea* PIC, 1909  
*Ochrosis ventralis* var. *obscuricollis* HEIKERTINGER, 1911  
*Ochrosis ventralis* ab. *nigriventris*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1911)

- Ochrosia ventralis* var. *krueperi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1911)
- Ochrosia ventralis* ab. *rubicunda*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1911)
- Ochrosia rubicunda*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. WEISE, 1886)
- Crepidodera lubrica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)
- Haltica ventralis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KUTSCHERA, 1863)
- Ochrosia corsica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to REY, 1874)
- Ochrosia ventralis* var. *pisana*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1911)
- Ochrosia ventralis* var. *lubrica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1911)
- Crepidodera ventralis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PAULINO DE OLIVEIRA, 1893)
- Ochrosia pisana*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. WEISE, 1886)
- Ochrosia ventralis* *krueperi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GRUEV et KAZAP, 1985)
- Distribution: Europe except for N, Caucasus, Turkey, Near and Middle East, N Africa, Canary Isles, Madeira.

#### Genus: *Neocrepidodera* HEIKERTINGER, 1911

*Asiolestia* JACOBSON, 1925

*Crepidodera* sensu FOU DRAS, 1860, non CHEVOLAT in DEJEAN, 1836

#### *Neocrepidodera brevicollis* (J. DANIEL, 1904)

*Crepidodera brevicollis* J. DANIEL, 1904

*Asiolestia brevicollis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BIONDI, 1989)

Distribution: Mountains of W, Central, and E Europe.

#### *Neocrepidodera corpulenta* (KUTSCHERA, 1860)

*Crepidodera corpulenta* KUTSCHERA, 1860

*Crepidodera corpulenta* var. *opaca* PETRI, 1912

*Asiolestia corpulenta*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BIONDI, 1989)

Distribution: Mountains of W, Central, E, and SE Europe.

#### *Neocrepidodera crassicornis crassicornis* (FALDERMANN, 1837)

*Haltica crassicornis* FALDERMANN, 1837

*Crepidodera crassicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)

*Asiolestia crassicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BIONDI, 1989)

Distribution: S and SE Europe, Ciscaucasia, Caucasus, Transcaucasia, Turkey, Kazakhstan.

#### *Neocrepidodera cyanescens cyanescens* (DUFTSCHMIDT, 1825)

*Haltica cyanescens* DUFTSCHMIDT, 1825

*Crepidodera alpicola* SCHMIDT, 1850

*Crepidodera cyanescens coeruleicollis* PIC, 1901

*Crepidodera sabauda* PIC, 1904

*Crepidodera cyanipennis* var. *sabauda* PIC, 1905

*Crepidodera cyanipennis coeruleicollis* PIC, 1905

*Crepidodera cyanipennis* var. *subelongata* PIC, 1910

*Crepidodera tatrica* REITTER, 1914

*Crepidodera cyanescens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)

*Crepidodera cyanescens* var. *tatrica* HEIKERTINGER, 1930

*Asiolestia cyanescens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BIONDI, 1989)

Distribution: Mountains of W and Central Europe.

***Neocrepidodera femorata* (GYLLENHALL, 1813)**

*Haltica femorata* GYLLENHALL, 1813

*Haltica femoralis* DUFTSCHMIDT, 1825

*Crepidodera femorata* var. *infuscipes* FOU DRAS, 1860

*Crepidodera femorata* var. *aeneipennis* J. WEISE, 1886

*Crepidodera peirolerii* J. DANIEL, 1904 (part.)

*Crepidodera femorata* var. *kossmani* GERHARDT, 1904

*Crepidodera femorata* var. *infuscipes* GERHARDT, 1909

*Crepidodera femorata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to *femorata* STEPHENS, 1832)

*Haltica femorata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)

*Asiolestia femorata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BIONDI, 1989)

Distribution: Mountains of W, Central, and E Europe, N Europe, Asian Russia (W and Middle Siberia, Altai, Baikal Area, Tuva).

***Neocrepidodera ferruginea* (SCOPOLI, 1763)**

*Chrysomela ferruginea* SCOPOLI, 1763

*Chrysomela exoleta* LINNAEUS, 1761

*Mordella flava* LINNAEUS, 1761

*Haltica flava* var. *similis* STEPHENS, 1832

*Crepidodera flava* var. *melanaria* EVERTS, 1919

*Altica exoleta*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1775)

*Altica ferruginea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to SCHRANK, 1781)

*Galleruca exoleta*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1787)

*Haltica exoleta*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to OLIVIER, 1789)

*Galeruca exoleta*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PAYKULL, 1799)

*Crioceris exoleta*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1801)

*Haltica ferruginea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ILLIGER, 1807)

*Crepidodera ferruginea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

*Haltica flava*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

*Crepidodera flava*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1839)

*Crepidodera exoleta*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KELCH, 1846)

*Crepidodera impressa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to DIRIMANOV et ANGELOVA, 1964)

*Asiolestia ferruginea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BIONDI, 1989)

Distribution: Europe, Caucasus, Transcaucasia, Turkey, Iran.

***Neocrepidodera impressa impressa* (FABRICIUS, 1801)**

*Galleruca impressa* FABRICIUS, 1801

*Crepidodera rufa* KÜSTER, 1849

*Crepidodera punctulata* ALLARD, 1859

*Haltica impressa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ILLIGER, 1807)

*Crepidodera impressa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to *impressa* FOU DRAS, 1860)

Distribution: S Europe, Near East, N Africa.

***Neocrepidodera interpunctata* (MOTSCHULSKY, 1859)***Crepidodera interpunctata* MOTSCHULSKY, 1859? *Crepidodera mitsuhashii* MATSUMURA, 1911

Distribution: N Europe, Middle Kazakhstan, Asian Russia (S Siberia, Tuva, Baikal Area, Transbaikalia, Yakutia, Amurland, Amurland, Maritime Prov., Ussuri Area, Sakhalin, Kuriles Isles), Mongolia, China, Korea, Japan.

***Neocrepidodera kozhantshikovi* (JACOBSON, 1925)***Asiorestia kozhantshikovi* JACOBSON, 1925*Crepidodera kozhantshikovi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

Distribution: Asian Russia (Yakutia).

***Neocrepidodera melanostoma* (L. REDTENBACHER, 1849)***Haltica melanostoma* L. REDTENBACHER, 1849*Crepidodera frigida* J. WEISE, 1886*Crepidodera theresae* PIC, 1903*Crepidodera melanostoma*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)

Distribution: Mountains of W, Central, and E Europe.

***Neocrepidodera motschulskii* (KONSTANTINOV, 1991)***Asiorestia motschulskii* KONSTANTINOV, 1991*Crepidodera sublaevis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. DANIEL, 1904)*Crepidodera interpunctata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1948)*Asiorestia sublaevis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LOPATIN, 1986)

Distribution: Europe except for S, Caucasus.

***Neocrepidodera nigrifula* (GYLLENHAL, 1813)***Haltica nigrifula* GYLLENHAL, 1813*Haltica ovulum* DUFTSCHMIDT, 1825*Haltica impressa* WALT, 1839*Crepidodera nigrifula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to SIEBOLD, 1847)*Hippuriphila nigrifula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to C. G. THOMSON, 1866)

Distribution: Europe except for SW.

***Neocrepidodera obscuritarsis* (MOTSCHULSKY, 1859)***Crepidodera obscuritarsis* MOTSCHULSKY, 1859*Crepidodera lewisii* JACOBY, 1885

Distribution: Asian Russia (Amurland, Amurland, Maritime Prov., Ussuri Area, NE Sakhalin), China, Japan.

***Neocrepidodera puncticollis* (REITTER, 1879)***Orestia puncticollis* REITTER, 1879*Crepidodera cyanipennis* ab. *puncticollis* J. DANIEL, 1904*Crepidodera picea* PETRI, 1907*Crepidodera alpina marani* ROUBAL, 1939*Crepidodera cyanipennis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to SHAPIRO, 1960)*Crepidodera puncticollis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KONNERTH-IONESCU, 1963)



*Asiorestia cyanipennis puncticollis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GRUEV, MERKL et VIG, 1993)

Distribution: Carpathians.

***Neocrepidodera sibirica* (PIC, 1909)**

*Ochrosis sibirica* PIC, 1909

Distribution: Asian Russia (Amurland, Ussuri Area), N Korea, Japan.

***Neocrepidodera sublaevis* (MOTSCHULSKY, 1859)**

*Crepidodera sublaevis* MOTSCHULSKY, 1859

*Crepidodera interpunctata* var. *sublaevis* J. DANIEL, 1904

Distribution: Kazakhstan, Afghanistan, N Tajikistan, Asian Russia (Middle and E Siberia to Far East), Mongolia.

***Neocrepidodera transsylvanica* (FUSS, 1864)**

*Crepidodera transsylvanica* FUSS, 1864

*Crepidodera transsylvanica* ab. *rufipennis* J. DANIEL, 1904

Distribution: Carpathians.

***Neocrepidodera transversa* (MARSHAM, 1802)**

*Chrysomela transversa* MARSHAM, 1802

*Haltica exoleta* PANZER, 1794

*Haltica ferruginea* ILLIGER, 1807

? *Crepidodera scutellaris* OLIVIER, 1808

*Crepidodera marginicollis* ALLARD, 1860

*Crepidodera transversa* ab. *melancholica* J. DANIEL, 1904

*Crepidodera transversa* ab. *nigra* SCHILSKY, 1908

? *Haltica marginicollis* KÜSTER, 1849

*Haltica transversa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GYLLENHAL, 1813)

*Haltica impressa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to DUFTSCHMIDT, 1825)

*Crepidodera ferruginea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

*Crepidodera transversa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)

Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, Iran.

**Genus: *Orestia* GERMAR, 1845**

*Pyladia* J. WEISE, 1886

***Orestia aubei* ALLARD, 1859**

*Orestia arcuata* L. MILLER, 1868

*Orestia pandellei apennina*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1924)

*Haltica aubei*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1874)

Distribution: Mountains of Central, E, and northern parts of SE Europe (Dynary, Carpathians).

***Orestia carpathica* REITTER, 1880**

*Crepidodera carpathica* KONNERTH-IONESCU, 1963

Distribution: Mountains of Central Europe (Czerna Gora, Carpathians).

***Orestia caucasica* REITTER, 1879***Orestia caucasica* var. *pseudoaubei* PIC, 1910

Distribution: Ciscaucasia, Caucasus, Transcaucasia.

***Orestia paveli* FRIVALDSKY, 1877**

Distribution: Mountains of Central, E, and northern parts of SE Europe (Dynary, Carpathians).

**Genus: *Crepidodera* CHEVROLAT in DEJEAN, 1836***Chalcooides* FOU DRAS, 1860***Crepidodera aurata* (MARSHAM, 1802)***Chrysomela aurata* MARSHAM, 1802? *Buprestis chrysicollis* SCOPOLI, 1763*Chrysomela nitidula* var. SCHRANK, 1781*Altica nitidula* GEOFFROY, 1785*Galeruca helxines* var. PAYKULL, 1799*Haltica helxines* KOCH, 1803*Haltica aurata* ab. *pulchella* STEPHENS, 1834*Haltica pulchella* STEPHENS, 1834*Crepidodera versicolor* KUTSCHERA, 1860*Chalcooides aurata* var. *pulchella* J. WEISE, 1886*Chalcooides nigra* KLENKA, 1914*Chalcooides aurata* ab. *subunicolor* PIC, 1918*Crepidodera helxines*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to MANNERHEIM, 1839)*Chalcooides aurata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)*Haltica aurata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to JACHNO, 1880)

Distribution: Europe, Caucasus, Transcaucasia, Turkey, Iran, Kazakhstan, Asian Russia (W, S, Middle and NE Siberia, Far East), Mongolia, N China, Korea, N Africa.

***Crepidodera aurea* (GEOFFROY, 1785)***Altica aurea* GEOFFROY, 1785*Chrysomela cyanea* MARSHAM, 1802*Haltica gaudens* STEPHENS, 1832*Haltica pulchella* STEPHENS, 1834*Crepidodera helxines* ALLARD, 1866*Chalcooides cyanea* J. WEISE, 1886*Chalcooides cyanea* var. *splendens* J. WEISE, 1886*Chalcooides cyanea* var. *laeta* J. WEISE, 1886*Chalcooides coerulea* J. WEISE, 1890*Chalcooides helxines* var. *cyanea* KUTHY, 1900*Chalcooides metallica* KUTHY, 1900*Chalcooides splendens* KUTHY, 1900*Chalcooides splendens* var. *gaudens* KUTHY, 1900*Chalcooides splendens* var. *laeta* KUTHY, 1900*Chalcooides aurea* var. *laeta* GERHARDT, 1909*Chalcooides aurea syriaca* PIC, 1918*Chalcooides aurea* ab. *purpurascens* PIC, 1918*Haltica cyanea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)*Haltica helxines*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)

*Crepidodera fulvicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)

*Chalcoides helxines*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)

*Crepidodera cyanea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to SEIDLITZ, 1891)

*Chalcoides aurea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BEDEL, 1900)

*Chalcoides gaudens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to EVERTS, 1903)

Distribution: Europe except for N, Caucasus, Turkey, E Kazakhstan, W Siberia.

### ***Crepidodera fulvicornis* (FABRICIUS, 1792)**

*Galleruca fulvicornis* FABRICIUS, 1792

? *Chrysomela helxines* LINNAEUS, 1758

? *Altica helxines* FABRICIUS, 1775

*Chrysomela viridiaurata* DEGEER, 1775

*Chrysomela violaceopunctata* DEGEER, 1775

*Chrysomela viridicoerulea* DEGEER, 1775

*Altica cyanopicta* GMELIN, 1790

*Altica coerulescens* GMELIN, 1790

*Haltica horticola* LETZNER, 1847

*Chalcoides smaragdina* FOU DRAS, 1860

*Haltica versicolor* var. c KUTSCHERA, 1860

*Crepidodera smaragdina* ALLARD, 1861

*Haltica smaragdina* LETZNER, 1870

*Chalcoides fulvipes* J. WEISE, 1886

*Chalcoides helxines* var. *jucunda* J. WEISE, 1886

*Chalcoides helxines* var. *picicornis* J. WEISE, 1886

*Chalcoides helxines* var. *fulvicornis* GERHARDT, 1891

*Chalcoides fulvicornis* var. *jucunda* SCHILSKY, 1909

*Chalcoides fulvicornis* ab. *picicornis* GERHARDT, 1910

? *Chalcoides libanicola* PIC, 1918

*Chalcoides fulvicornis lapponica* HEIKERTINGER, 1950

? *Galleruca helxines*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1792 and PAYKULL, 1799)

*Haltica helxines*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KOCH, 1803)

*Chrysomela fulvicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to WEIGEL, 1806)

*Crepidodera helxines*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

*Chalcoides helxines*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. WEISE, 1886)

*Chalcoides plutus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1940)

Distribution: Europe, Ciscaucasia, Caucasus, Transcaucasia, Turkey, Near East, N Iran, NW and E Kazakhstan, Asian Russia (W and Middle Siberia, Cisbaikalia, Transbaikalia, Amurland), Mongolia.

### ***Crepidodera japonica* BALY, 1877**

*Chalcoides chloris* JACOBY, 1885

*Chalcoides japonicus* CHŪJŌ et KIMOTO, 1961

*Chalcoides plutus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1940)

*Crepidodera fulvicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. MEDVEDEV, 1978)

Distribution: Asian Russia (Sakhalin, Kuriles Isles), Japan.

***Crepidodera lamina* (BEDEL, 1901)***Chalcoides lamina* BEDEL, 1901? *Haltica metallica* DUFTSCHMIDT, 1825*Haltica helxines* var. *metallica* L. REDTENBACHER, 1849*Crepidodera aurata* var. *lamiana* L. REDTENBACHER, 1858*Haltica helxines* var. c KUTSCHERA, 1860*Chalcoides metallica* var. *aeruginosa* J. WEISE, 1886*Chalcoides metallica* var. *cuprea* J. WEISE, 1886*Chalcoides metallica* var. *sapphirina* J. WEISE, 1886*Chalcoides metallica* var. *aeruginosa* STIERLIN, 1897*Chalcoides lamina* ab. *cuprea* SCHILSKY, 1909*Crepidodera helxines*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1861)*Chalcoides metallica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. WEISE, 1886)*Crepidodera metallica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to SEIDLITZ, 1891)

Distribution: Europe except for N, NW Caucasus.

***Crepidodera nigricoxis* ALLARD, 1878***Chalcoides aurata* var. *nigricoxis* J. WEISE, 1886? *Chalcoides aurata* var. *peloponnesiaca* HEIKERTINGER, 1910*Chalcoides aurata* ab. *nigricoxis* HEIKERTINGER, 1912*Crepidodera aurata* var. *peloponnesiaca* BECHYNÉ, 1956*Chalcoides nigricoxis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GRUEV, 1975)

Distribution: Southern parts of Central Europe, SE Europe east to Rumania, Caucasus,? N Korea.

***Crepidodera nitidula* (LINNAEUS, 1758)***Chrysomela nitidula* LINNAEUS, 1758*Altica nitidula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1781)*Galleruca nitidula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1787 and PAYKULL, 1799)*Haltica nitidula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to OLIVIER, 1789)*Chalcoides nitidula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)

Distribution: Europe except for SW; Asian Russia (W and Middle Siberia, Sayan).

***Crepidodera obscuripes* (HEIKERTINGER, 1912)***Chalcoides aurea* var. *obscuripes* HEIKERTINGER, 1912*Crepidodera aurea obscuripes* BECHYNÉ, 1956*Crepidodera picipes*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GRESSITT et KIMOTO, 1963)*Chalcoides aurea obscuripes*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BARTUEVA, 1973)

Distribution: Asian Russia (E Siberia, Far East), Mongolia, China, Korea.

***Crepidodera picipes* (J. WEISE, 1887)***Chalcoides picipes* J. WEISE, 1887*Chalcoides mroczkowski* WARCHAŁOWSKI, 1969

Distribution: Asian Russia (Cisbaikalia, Transbaikalia, Ussuri Area, Amurland, Maritime Prov., Sakhalin, Kuriles Isles), N China, Korea.

***Crepidodera plutus* (LATREILLE, 1804)***Altica plutus* LATREILLE, 1804*Altica helxineus* PANZER, 1794

- Chalcoides chloris* Foudras, 1860  
*Chalcoides chloris* var. *foudrasi* J. Weise, 1886  
*Crepidodera japonica* Baly, 1877  
*Chalcoides plutus* ab. *foudrasi* Gerhardt, 1910  
*Chalcoides aurata* var. *nigrofemorata* Heikertinger, 1912  
*Chalcoides plutus punctatithorax* Pic, 1918  
*Chalcoides plutus* var. *melanothorax* Everts, 1919  
? *Chalcoides aurata* var. *unicolor* Westhoff, 1882  
*Crepidodera aurata nigrofemorata* Bechyně, 1956  
*Chalcoides chloris*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to J. Weise, 1886)  
*Chalcoides plutus*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Bedel, 1900)  
*Haltica chloris*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to L. Redtenbacher, 1874)  
*Chalcoides japonicus*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Chûjô et Kimoto, 1961)  
Distribution: Europe except for N, Caucasus, Turkey, Iran, Kazakhstan, Kirghizstan, Tajikistan, Asian Russia (Siberia to Far East, Sakhalin, Kuriles Isles), Mongolia, N China, Korea, Japan.

***Crepidodera sahalinensis* Konstantinov, 1996**

Distribution: Asian Russia (Sakhalin, Kunashir Isle).

***Crepidodera ussuriensis* Konstantinov, 1996**

Distribution: Asian Russia (Far East).

***Crepidodera viridis* Gressitt et Kimoto, 1963**

*Crepidodera viridis*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Doguet, 1976)

Distribution: Asian Russia (Far East), NW China.

**Genus: *Derocrepis* J. Weise, 1886**

*Aeschrocnemis* J. Weise, 1888

***Derocrepis pubipennis* Reitter, 1892**

Distribution: Armenia, Georgia, Transcaucasus.

***Derocrepis rufipes* (Linnaeus, 1758)**

*Chrysomela rufipes* Linnaeus, 1758

*Galleruca ruficornis* Fabricius, 1792

*Derocrepis rufipes* var. *obscura* J. Weise, 1886

*Derocrepis rufipes* var. *latithorax* Pic, 1911

*Altica rufipes*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Fabricius, 1775)

*Crioceris ruficornis*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Paykull, 1799)

*Chrysomela ruficornis*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Marsham, 1802)

*Haltica rufipes*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Koch, 1803)

*Crepidodera rufipes*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Stephens, 1839)

Distribution: Europe except for N and Mediterranean, Transcaucasia, Turkey, Asian Russia (W and Middle Siberia, Sayan).

***Derocrepis serbica caucasica* J. Weise, 1886**

*Derocrepis serbica* var. *caucasica* J. Weise, 1886

*Derocrepis serbica*: Gruev et Döberl, 1997 (in synonymy erroneously attributed to Allard, 1878)

Distribution: Caucasus.

***Derocrepis serbica jajlensis* HEIKERTINGER, 1922**

Distribution: Crimea.

***Derocrepis serbica laterufa* PIC, 1909***Derocrepis serbica* race *laterufa* PIC, 1909*Derocrepis serbica laterufa* var. *obscuripes* PIC, 1911

Distribution: N Caucasus, Georgia.

***Derocrepis serbica ossetica* HEIKERTINGER, 1922***Derocrepis serbica* var. *ossetica* HEIKERTINGER, 1922

Distribution: Caucasus (N Ossetia).

***Derocrepis serbica serbica* (KUTSCHERA, 1860)***Haltica* (*Crepidodera*) *serbica* KUTSCHERA, 1860*Crepidodera strangulata* ALLARD, 1860*Derocrepis serbica* var. *bicolor* J. WEISE, 1886*Crepidodera serbica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1878)

Distribution: Balkans, Crimea.

**Genus: *Novofoudrasia* JACOBSON, 1901*****Novofoudrasia rufiventris* (J. WEISE, 1900)***Foudrasia rufiventris* J. WEISE, 1900

Distribution: Uzbekistan, S Kazakhstan, Afghanistan, Tajikistan, Nepal.

**Genus: *Hippuriphila* FOUDRAS, 1860***Pseudocrepidostoma* L. MEDVEDEV, 1966***Hippuriphila babai* (CHÛJÔ, 1959)***Parachalcoides babai* CHÛJÔ, 1959*Pseudocrepidostoma ohnoi* L. MEDVEDEV, 1966

Distribution: Asian Russia (Sakhalin, Kuriles Isles), Japan.

***Hippuriphila modeeri* (LINNAEUS, 1761)***Chrysomela modeeri* LINNAEUS, 1761*Altica modeeri* OLIVIER, 1789*Galeruca modeeri* FABRICIUS, 1792*Haltica modeeri* KOCH, 1803*Haltica chrysopygis* BECK, 1817*Crepidodera modeeri* STEPHENS, 1832*Hippuriphila modeeri* var. *bimaculata* J. WEISE, 1886*Hippuriphila modeeri* var. *chalybaea* J. WEISE, 1886*Hippuriphila modeeri* var. *praescutellaris* J. WEISE, 1886*Hippuriphila modeeri* var. *praescutellaris* KOLBE, 1915*Hippuriphila laevicollis* HELLÉN, 1933*Hippuriphila modeeri* ab. *violaceovittata* CSIKI, 1953

Distribution: Europe, Caucasus, Turkey, Asian Russia (Siberia), Mongolia.

**Genus: *Epitrix* FOUDRAS, 1860*****Epitrix abeillei* (BAUDUER, 1874)***Crepidodera abeillei* BAUDUER, 1874

*Crepidodera judaea* ALLARD, 1876

*Epithrix abeillei* var. *testaceipes* PIC, 1909

*Epithrix abeillei*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

Distribution: Turkey, Middle and Near East, Kazkhstan, Turkmenistan, Uzbekistan, NE China, Mongolia, N Africa.

***Epithrix atropae* FOU DRAS, 1860**

*Haltica pubescens* var. a KOCH, 1803

*Haltica pubescens* var. b KOCH, 1803

*Crepidodera pubescens* var. KÜSTER, 1853

*Epithrix atropae* var. *nigritula* J. WEISE, 1886

*Epithrix atropae* var. *quadrifaculata* J. WEISE, 1886

*Epithrix atropae* ab. *suturata* HEIKERTINGER, 1912

*Epithrix atropae* ab. *nigritula* GERHARDT, 1911

*Epithrix atropae* ab. *quadrifaculata* GERHARDT, 1911

*Epithrix atropae quadrifaculata* TRELLA, 1930

*Epithrix atropae* ab. *suturata* TENENBAUM, 1931

*Epithrix pubescens* ab. *nigritula* MAZUROWA, 1951

*Epithrix pubescens* ab. *quadrifaculata* MAZUROWA, 1951

*Epithrix atropae* ab. *nigritula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER et CSIKI, 1940)

*Haltica atropae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KUTSCHERA, 1860)

*Crepidodera atropae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LEESBERG, 1882)

*Epithrix atropae* ab. *quadrifaculata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to NOVAK, 1952)

Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, N Africa.

***Epithrix caucasica* (HEIKERTINGER, 1950)**

*Epithrix caucasica* HEIKERTINGER, 1950

Distribution: S European Russia, Caucasus, Transcaucasia, Turkey, Iran, Kazakhstan, Central Asia, India (Himalaya).

***Epithrix dieckmanni* (MOHR, 1968)**

*Epithrix dieckmanni* MOHR, 1968

*Epithrix paliji* GRUEV, 1975

Distribution: Turkmenistan, Near and Middle East, Saudi Arabia.

***Epithrix ermischii* (MOHR, 1968)**

*Epithrix ermischii* MOHR, 1968

Distribution: Turkmenistan, Uzbekistan, Iran.

***Epithrix intermedia* FOU DRAS, 1860**

*Crepidodera intermedia* ALLARD, 1861

Distribution: Southern parts of Central Europe, Balkans, Ukraine, Caucasus, Turkey.

***Epithrix ogloblini* (IABLOKOFF-KHNZORIAN, 1960)**

*Epithrix ogloblini* IABLOKOFF-KHNZORIAN, 1960

Distribution: Armenia.

***Epithrix pubescens* (KOCH, 1803)**

*Haltica pubescens* KOCH, 1803

*Epithrix pubescens* var. *ferruginea* J. WEISE, 1886

*Epithrix suturalis* BEDEL, 1897

*Epitrix lencorana* PIC, 1903  
*Epitrix pubescens* var. *ferruginea* TENENBAUM, 1931  
*Epitrix pubescens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BEDEL, 1897)  
 Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, Iran, Kazakhstan, Asian Russia (W Siberia), Azores Isles.

***Epitrix wuorentausi* (HEIKERTINGER, 1950)**

*Epitrix wuorentausi* HEIKERTINGER, 1950  
 Distribution: Asian Russia (Amurland, Ussuri Area, Maritime Prov.).

**Genus: *Minota* KUTSCHERA, 1859**

*Hypnophila* Foudras, 1860

***Minota carpathica* HEIKERTINGER, 1911**

*Minota obesa carpathica* HEIKERTINGER, 1911  
*Minota obesa* var. *carpathica* HEIKERTINGER et CSIKI, 1940  
*Hypnophila obesa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KUTHY, 1900)  
 Distribution: Mountains of W, Central, E, and SE Europe.

**Genus: *Apteropeda* CHEVOLAT in DEJEAN, 1837**

***Apteropeda globosa* (ILLIGER, 1794)**

*Altica globosa* ILLIGER, 1794  
*Haltica conglomerata* ILLIGER, 1807  
*Haltica globus* DUFTSCHMIDT, 1825  
*Apteropeda conglomerata* KÜSTER, 1853  
*Apteropeda nigroaenea* WEIDENBACH, 1859  
*Apteropeda majuscula* Foudras, 1860  
*Sphaeroderma globosa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)  
 Distribution: Europe except for extreme N.

***Apteropeda orbiculata* (MARSHAM, 1802)**

*Chrysomela orbiculata* MARSHAM, 1802  
*Haltica raminis* KOCH, 1803  
*Haltica hederæ* ILLIGER, 1807  
*Altica ciliata* OLIVIER, 1807  
*Apteropeda ciliata* KELCH, 1852  
*Apteropeda graminis* ROTTENBERG, 1868  
*Apteropeda orbiculata* var. *aurichalcea* J. WEISE, 1893  
*Apteropeda orbiculata* var. *coerulans* J. WEISE, 1893  
*Sphaeroderma orbiculata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)  
 Distribution: Europe except for extreme N and E.

***Apteropeda splendida* ALLARD, 1860**

*Apteropeda ciliata* L. REDTENBACHER, 1849  
*Apteropeda globosa* Foudras, 1860  
*Apteropeda splendidula* ROUBAL, 1937–1941  
 Distribution: Europe except for extreme N and E.



**Genus: *Podagrica* CHEVROLAT in DEJEAN, 1837*****Podagrica fuscicornis* (LINNAEUS, 1766)***Chrysomela fuscicornis* LINNAEUS, 1766*Altica rufipes* LAICHTING, 1781*Crioceris fulvipes* FABRICIUS, 1801*Haltica chrysomelina* WATTL, 1835*Podagrica fuscicornis* var. *meridionalis* J. WEISE, 1886*Podagrica meridionalis* NETOLITZKY, 1912*Podagrica fusciformis chrysomelina* HEIKERTINGER, 1930*Altica fuscicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GMELIN, 1790)*Galleruca fuscicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1792)*Galleruca rufipes*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PAYKULL, 1799)*Haltica fuscicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KOCH, 1803)*Podagrica chrysomelina*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GEMMINGER et HAROLD, 1876)*Podagrica fusciformis* var. *chrysomelina*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FURTH, 1976)

Distribution: Europe except for extreme N, Transcaucasia, Turkey, Near East, N Africa.

***Podagrica malvae* (ILLIGER, 1807)***Haltica malvae* ILLIGER, 1807*Haltica saracena* REICHE, 1858*Haltica cyanea* REICHE, 1858*Podagrica tristicula* ALLARD, 1860*Podagrica unicolor* MARSEUL, 1869*Podagrica malvae* var. *aenescens* J. WEISE, 1886*Podagrica malvae* ab. *nigricans* DEMAISON, 1903*Podagrica malvae* ab. *picicolor* HEIKERTINGER, 1912*Podagrica angusticollis* J. SAHLBERG, 1913*Podagrica cedricola* J. SAHLBERG, 1913*Podagrica convexicollis* J. SAHLBERG, 1913*Podagrica subaptera* J. SAHLBERG, 1913*Podagrica malvae* ab. (var.) *cyanella* HEIKERTINGER et CSIKI, 1940*Podagrica malvae* ab. *aenescens* HEIKERTINGER et CSIKI, 1940*Podagrica malvae* var. *flaviceps* CSIKI, 1953*Podagrica malvae* var. *nigricornis* G. MÜLLER, 1953*Podagrica malvae semirufa* MOHR, 1965 (part.)*Podagrica malvae* var. *tristicula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)*Podagrica malvae* var. *saracena*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

Distribution: S and southern parts of Central and E Europe, Caucasus, Transcaucasia, Turkey, Near and Middle East.

***Podagrica menetriesi* (FALDERMANN, 1837)***Triplax menetriesi* FALDERMANN, 1837*Podagrica punctatostrata* MOTSCHULSKY, 1845*Haltica discedens* BOIELDIEU, 1859*Podagrica rudicollis* FOUDRAS, 1860*Podagrica discedens* J. WEISE, 1886*Podagrica robusta* BALLION, 1878

- Podagrica desertorum* BRANCSIK, 1899  
*Podagrica fuscipes* KUTHY, 1900  
*Podagrica menetriesi* var. *luctuosa* DEMAISON, 1902  
*Podagrica minor* J. SAHLBERG, 1913  
*Podagrica menetriesi* ab. *scepelensis* CSIKI, 1953  
*Podagrica menetriesi* var. *desertorum*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)  
*Podagrica menetriesi* var. *discedens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)  
*Podagrica menetriesi discedens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to NOVAK, 1952)  
*Podagrica fuscipes* var. *discedens*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KONNERTH-IONESCU, 1963)  
 Distribution: S and SE Europe, Caucasus, Transcaucasia, Turkey, Iraq, Kazakhstan, Central Asia.

**Genus: *Chaetocnema* STEPHENS, 1832**

*Plectroscelis* CHEVROLAT in DEJEAN, 1837

***Chaetocnema aerosa* (LETZNER, 1846)**

- Plectroscelis aerosa* LETZNER, 1846  
*Plectroscelis punctatissima* GRAELLS, 1858  
*Chaetocnema laevilinea* J. SAHLBERG, 1903  
 Distribution: Europe except for S and SE, Caucasus, Daghestan, Turkey, Israel.

***Chaetocnema arenacea* (ALLARD, 1860)**

- Plectroscelis arenacea* ALLARD, 1860  
 Distribution: S and SE Europe, Caucasus, Daghestan, Turkey, N Africa.

***Chaetocnema arida* (FOUDRAS, 1860)**

- Plectroscelis arida* FODRAS, 1860  
*Plectroscelis confusa* ALLARD, 1866  
*Chaetocnema arida* var. *aestive* J. WEISE, 1886  
*Chaetocnema surcoufi* PIC, 1915  
 Distribution: Europe except for N, Caucasus, N Africa.

***Chaetocnema aridula* (GYLLENHAL, 1827)**

- Haltica aridula* GYLLENHAL, 1827  
*Chaetocnema aridula* var. *weisei* GERHARDT, 1906  
*Plectroscelis aridula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)  
 Distribution: Europe, Caucasus, Transcaucasia, Turkey, Kazakhstan, Kirghizstan, Asian Russia (Siberia), N Africa.

***Chaetocnema arisi* PIC, 1915**

- Chaetocnema sahlbergi* var. *arisi* PIC, 1915  
 Distribution: S Kazakhstan.

***Chaetocnema breviuscula* (FALDERMANN, 1837)**

- Haltica breviuscula* FALDERMANN, 1837  
*Plectroscelis breviuscula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

*Chaetocnema tibialis breviscula*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

Distribution: S and SE Europe, Latvia, Caucasus, Transcaucasia, Middle East, Kazakhstan, Tajikistan, Asian Russia (southern parts of W Siberia), N China, Korea.

***Chaetocnema chlorophana* (DUFTSCHMIDT, 1825)**

*Haltica chlorophana* DUFTSCHMIDT, 1825

*Plectroscelis dentipes* OLIVIER, 1808

*Chaetocnema chlorophana* var. *amoena* J. WEISE, 1886

*Chaetocnema chlorophana* var. *kolbei* J. WEISE, 1886

*Chaetocnema chlorophana* var. *laeta* J. WEISE, 1886

*Chaetocnema syriaca* DEMAISON, 1896

*Chaetocnema chlorophana pelagica* CAILLOL, 1914

*Chaetocnema chlorophana*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)

Distribution: S, SE and southern parts of Central Europe, Caucasus, Turkey, Iraq, N Africa.

***Chaetocnema compressa* (LETZNER, 1864)**

*Plectroscelis compressa* LETZNER, 1864

*Plectroscelis tarda* BACH, 1856

*Chaetocnema compressa* ab. *aneicollis* FLEISCHER, 1917

Distribution: Central and SE Europe, Caucasus, Kazakhstan.

***Chaetocnema concinna* (MARSHAM, 1802)**

*Chrysomela concinna* MARSHAM, 1802

*Haltica dentipes* KOCH, 1803

? *Chaetocnema picipes* STEPHENS, 1832

*Chaetocnema lewisii* CHŪJŌ, 1942

*Plectroscelis dentipes*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to UECHTRITZ, 1844)

*Haltica concinna*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to WALKER, 1846)

*Plectroscelis concinna*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KÜSTER, 1847)

*Plectroscelis dentipes*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)

*Phyllotreta concinna*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ORMEROD, 1882)

*Chaetocnema tibialis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to RUSZKOWSKI, 1933)

Distribution: Europe, Caucasus, Transcaucasia, Turkey, Israel, Turkmenistan, Kazakhstan, Asian Russia (Siberia, Far East, Sakhalin, Kuriles Isles), Mongolia, China, Korea, Japan, Taiwan, N Africa. Introduced to N America.

***Chaetocnema concinnicollis* (BALY, 1874)**

*Plectroscelis concinnicollis* BALY, 1874

*Plectroscelis phyloxena* BALY, 1876

*Chaetocnema aerosa* HEIKERTINGER et CSIKI, 1940

*Chaetocnema coccinelliformis* CHŪJŌ, 1951

*Chaetocnema coccinelliformis kaibarensis* MADAR, 1960

Distribution: Asian Russia (Amurland, Ussuri Area, Maritime Prov.), China, Korea, Japan, SE Asia.

***Chaetocnema conducta* (MOTSCHULSKY, 1838)**

*Plectroscelis conducta* MOTSCHULSKY, 1838

*Chaetocnema foudrasi* BAUDUER, 1874

*Chaetocnema ahngeri* JACOBSON, 1901

*Chaetocnema conducta* var. *ahngeri* HEIKERTINGER, 1930

Distribution: S, SE and southern parts of Central Europe, Caucasus, Transcaucasia, Turkey, Near East, Iran, Kazakhstan, Central Asia, N, W and S Africa.

***Chaetocnema confusa* (BOHEMAN, 1851)**

*Plectroscelis confusa* BOHEMAN, 1851

*Plectroscelis arida*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)

Distribution: Europe except for N, Caucasus, Turkey.

***Chaetocnema costulata* (MOTSCHULSKY, 1860)**

*Plectroscelis costulata* MOTSCHULSKY, 1860

*Chaetocnema aridula costulata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. MEDVEDEV, 1973)

Distribution: Afghanistan, Asian Russia (Cisbaikalia, Sayan, Transbaikalia, Buratia, Tuva, Yakutia, Amurland, Maritime Prov., Beringia, Kamchatka, Sakhalin), Mongolia, China.

***Chaetocnema coyei* (ALLARD, 1863)**

*Plectroscelis coyei* ALLARD, 1863

Distribution: SE Europe, Transcaucasia, Turkey, Near and Middle East.

***Chaetocnema gottwaldi* KRÁL, 1969**

Distribution: W Kazakhstan.

***Chaetocnema heptapotamica* LUBISCHEV, 1963**

Distribution: Caucasus, SE Kazakhstan, Kirghizstan.

***Chaetocnema hortensis* (GEOFFROY, 1785)**

*Altica hortensis* GEOFFROY, 1785

*Galeruca aridella* PAYKULL, 1799

*Plectroscelis scabricollis* ALLARD, 1860

*Plectroscelis convexa* MOTSCHULSKY, 1860

? *Chaetocnema granosa* MOTSCHULSKY, 1860

*Chaetocnema hortensis* var. *brenskoi* PIC, 1910

*Haltica aridella*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KOCH, 1803)

*Chaetocnema aridella*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

*Plectroscelis aridella*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KELCH, 1846)

*Chaetocnema scabricollis aridella*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to REY, 1875)

*Chaetocnema convexa*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to J. WEISE, 1887)

*Chaetocnema concinnipennis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to BRYANT, 1957)

*Chaetocnema montenegrina*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KRÁL, 1973)

Distribution: Europe, Caucasus, Transcaucasia, Turkey, Near and Middle East, Kazakhstan, Tajikistan, Uzbekistan, Asian Russia (W, Middle and S Siberia, Cisbaikalia, Sayan, Cispacific land, Amurland, Maritime Prov.), N Africa.

***Chaetocnema ingenua* (BALY, 1876)**

*Plectroscelis ingenua* BALY, 1876

*Chaetocnema japonica* JACOBY, 1885

*Chaetocnema aurifrons* JACOBY, 1885

*Chaetocnema fulvipes* JACOBY, 1885

*Chaetocnema cognata* CHEN, 1934

Distribution: Afghanistan, China, Asian Russia (Amurland, Maritime Prov., Kuriles Isles), Korea.

***Chaetocnema jelineki* LOPATIN, 1990**

Distribution: SE Iran.

***Chaetocnema kabakovi* LOPATIN, 1995**

Distribution: Afghanistan.

***Chaetocnema klapperichi* LOPATIN, 1963**

Distribution: Afghanistan.

***Chaetocnema laevicollis* (C. G. THOMSON, 1866)**

*Plectroscelis laevicollis* C. G. THOMSON, 1866

*Haltica dentipes* var. *d* GYLLENHAL, 1813

*Haltica concinna* var. *nitidicollis* JACOBSON, 1902

*Chaetocnema heikertingeri* LUBISCHEV, 1963

*Chaetocnema concinna*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1951)

*Chaetocnema concinna heikertingeri*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to IABLOKOFF-KHNZORIAN, 1968)

*Chaetocnema laevicollis* fa. *laevicollis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to DÖBERL, 1994)

Distribution: Europe, Caucasus, Transcaucasia, N Iran, Kazakhstan, Kirghizstan, Asian Russia (Transbaikalia, Tuva, Ussuri Area, Maritime Prov.), Mongolia, NE China.

***Chaetocnema major major* (JACQUELIN DU VAL, 1852)**

*Plectroscelis major major* JACQUELIN DU VAL, 1852

*Plectroscelis perrisi* BAUDUER, 1874

*Chaetocnema major* ab. *maesta* CSIKI, 1953

Distribution: S, SE and southern E Europe, Caucasus, Transcaucasia, Turkey, Near and Middle East, Kazakhstan.

***Chaetocnema major mandschurica* HEIKERTINGER, 1951**

Distribution: Asian Russia (Ussuri Area, Amurland, Maritime Prov.), NE China, Japan.

***Chaetocnema mannerheimii* (GYLLENHAL, 1827)**

*Haltica mannerheimii* GYLLENHAL, 1827

*Chaetocnema mannerheimii* var. *fallax* J. WEISE, 1888

*Mantura mannerheimii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

*Plectroscelis mannerheimii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KELCH, 1852)

*Plectroscelis mannerheimii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to MOTSCHULSKY, 1854)

*Chaetocnema mannerheimii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ROGER, 1856)

Distribution: Europe except for SW, Caucasus, Daghestan, Turkey, Kazakhstan, Kirghizstan, Asian Russia (W, Middle, S and NE Siberia), Mongolia.

***Chaetocnema medvedevi* PALIJ, 1968**

Distribution: Kirghizstan (E Tian-Shan).

***Chaetocnema montenegrina* HEIKERTINGER, 1912**

Distribution: SE Europe, Turkey, Middle East, Central Asia.

***Chaetocnema nebulosa* J. WEISE, 1886**

Distribution: S Ukraine, Crimea, Caucasus, Iran, Kazakhstan, Kirghizstan, Mongolia.

***Chaetocnema obesa* (BOIELDIEU, 1859)***Plectroscelis obesa* BOIELDIEU, 1859*Plectroscelis meridionalis* FOU DRAS, 1860*Plectroscelis subcoerulea* LETZNER, 1870*Chaetocnema insolita* REY, 1874*Chaetocnema meridionalis* var. *unicolor* J. WEISE, 1888*Chaetocnema obesa* var. *coreyrica* PIC, 1909*Chaetocnema jurassica* PIC, 1915? *Chaetocnema rufemorata* PIC, 1915

Distribution: Europe except for N, Caucasus, Turkey, Iraq, Kazakhstan, Asian Russia (Altai), China, Mongolia.

***Chaetocnema oblonga* LOPATIN, 1990**

Distribution: S Iran.

***Chaetocnema ogloblini* PALIJ, 1970***Chaetocnema micans* PALIJ, 1961

Distribution: Asian Russia (Ussuri Area, Amurland, Maritime Prov.), NE China.

***Chaetocnema orientalis* (BAUDUER, 1874)**

Distribution: SE Europe east to Rumania, Caucasus, Turkey, Near East, Iran, Turkmenistan.

***Chaetocnema procerula* (ROSENHAUER, 1856)***Plectroscelis procerula* ROSENHAUER, 1856*Chaetocnema procerula* var. *aenescens* J. WEISE, 1886*Chaetocnema compressa*: BACH, 1856 (attributed to Letzner).

Distribution: S and SE Europe, Turkey, N Africa.

***Chaetocnema psylloides* PIC, 1909**

Distribution: Middle East, S Kazakhstan, Central Asia.

***Chaetocnema sahlbergii* (GYLLENHAL, 1827)***Haltica sahlbergii* GYLLENHAL, 1827*Galleruca aridella* var. PAYKULL, 1799*Plectroscelis fairmairei* BOIELDIEU, 1852*Plectroscelis sahlbergii* var. *insolita* ALLARD, 1860*Chaetocnema sahlbergii* var. *cyanescens* J. WEISE, 1886*Chaetocnema sahlbergii* var. *cyanescens* KOLBE, 1918*Chaetocnema sahlbergii* ab. *similis* EVERTS, 1922*Haltica aridella*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KOCH, 1803)*Plectroscelis aridella*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GYLLENHAL, 1827)*Chaetocnema sahlbergii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

*Plectroscelis sahlbergii*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KELCH, 1846)  
 Distribution: Europe except for Mediterranean, Caucasus, Turkey, Asian Russia (S Siberia, Cisbaikalia, Cispafrican Land, Kamchatka, Amurland).

***Chaetocnema scheffleri* (KUTSCHERA, 1864)**

*Plectroscelis scheffleri* KUTSCHERA, 1864

*Chaetocnema longula* J. WEISE, 1890

*Chaetocnema tunisea* PIC, 1909

*Chaetocnema schaeffleri* PEYERIMHOFF, 1926

Distribution: S, Central, and SE Europe, Caucasus, Transcaucasia, Turkey, Near and Middle East, N Africa.

***Chaetocnema schlaeflii* (STIERLIN, 1866)**

*Plectroscelis schlaeflii* STIERLIN, 1866

*Cardiapus tarda* MOTSCHULSKY, 1845

*Chaetocnema schlaeflii* J. WEISE, 1886

*Chaetocnema persica* BALY, 1877

*Chaetocnema schlaeflini* JACOBSON, 1901

*Chaetocnema impunctifrons* PIC, 1909

Distribution: Caucasus, Middle East, Iran, Turkmenistan, NE China.

***Chaetocnema semicoerulea semicoerulea* (KOCH, 1803)**

*Haltica semicoerulea* KOCH, 1803

? *Chaetocnema saltitans* STEPHENS, 1832

*Plectroscelis meridionalis* ALLARD, 1859

*Chaetocnema semicoerulea* var. *femoralis* J. WEISE, 1886

*Chaetocnema semicoerulea* var. *saliceti* J. WEISE, 1886

*Chaetocnema semirufescens* PIC, 1909

Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, Kazakhstan, Asian Russia (W and Middle Siberia, Cisbaikalia).

***Chaetocnema semicoerulea transbaikalica* HEIKERTINGER, 1951**

Distribution: Asian Russia (Transbaikalia, Amurland, Ussuri Area, Beringia, Amurland, Maritime Prov.), Mongolia.

***Chaetocnema shabalini* PALIJ, 1968**

Distribution: Kirghizstan.

***Chaetocnema sinuata* J. WEISE, 1889**

*Chaetocnema hortensis* var. *grandis* PIC, 1909

Distribution: Kazakhstan, Central Asia, Asian Russia (Transbaikalia), Mongolia, China (Kansu).

***Chaetocnema sonkulica* PALIJ, 1968**

Distribution: Kirghizstan (Tian-Shan).

***Chaetocnema splendens splendens* (MOTSCHULSKY, 1845)**

*Udorpes splendens* MOTSCHULSKY, 1845

*Hydropus splendens* var. *aenea* MOTSCHULSKY, 1860

*Chaetocnema daurica* HEYDEN, 1881

Distribution: Asian Russia (Cisbaikalia, Transbaikalia, Ussuri Area, Amurland), Mongolia, N China.

***Chaetocnema splendens ljudmilae* LOPATIN, 1961**

Distribution: S Kazakhstan, Kirghizstan, S Tajikistan.

***Chaetocnema subcoerulea* (KUTSCHERA, 1864)***Plectroscelis subcoerulea* KUTSCHERA, 1864*Chaetocnema punctulata* REY, 1874*Chaetocnema subcoerulea christinae* HEIKERTINGER, 1909

Distribution: Europe except for N and SW, Caucasus.

***Chaetocnema tibialis* (ILLIGER, 1807)***Haltica tibialis* ILLIGER, 1807*Plectroscelis pumila* ALLARD, 1859*Chaetocnema caesaraugustana* FUENTE, 1909*Chaetocnema obscuripes* PIC, 1909

Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, Near and Middle East, Kazakhstan, Central Asia, Asian Russia (Transbaikalia), Mongolia, N Africa.

***Chaetocnema ussuriensis* HEIKERTINGER, 1951**

Distribution: Asian Russia (S Yakutia, Ussuri Area, Maritime Prov.), N China.

**Genus: *Mantura* STEPHENS, 1832***Balanomorpha* CHEVROLAT in DEJEAN, 1837***Mantura ambigua* (KUTSCHERA, 1862)***Haltica ambigua* KUTSCHERA, 1862*Balanomorpha ambigua*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1866)*Mantura obtusata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to JANSSON, 1947)

Distribution: W and N Europe, Ukrainian Carpathians.

***Mantura chrysanthemii* (KOCH, 1803)***Haltica chrysanthemii* KOCH, 1803*Balanomorpha crotchi* ALLARD, 1867*Mantura diluta* ABEILLE, 1895*Mantura reclairei* EVERTS, 1921

Distribution: Europe, Ukrainian Carpathians, Turkey, N Africa, Canary Isles, Madeira.

***Mantura cylindrica* L. MILLER, 1880**

Distribution: Italy, S Croatia, S Bulgaria, Greece, Turkey,? Caucasus.

***Mantura mathewsi* (STEPHENS, 1834)***Cardiapus mathewsi* STEPHENS, 1834*Balanomorpha mathewsi* CURTIS, 1833*Cardiapus helianthemii* CURTIS, 1837*Balanomorpha aeraria* FOU DRAS, 1860*Mantura mathewsi* var. *aeraria* KUTSCHERA, 1863*Mantura mathewsi* var. *bicolor* J. WEISE, 1886*Mantura mathewsi* var. *dichroa* BEDEL, 1900*Balanomorpha mathewsi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ALLARD, 1860)

Distribution: W, S, and southern parts of Central Europe, Ukrainian Carpathians, Azerbaijan.



***Mantura mesasiatica* LOPATIN, 1965**

Distribution: Kirghizstan, Tajikistan.

***Mantura obtusata* (GYLLENHAL, 1813)**

*Haltica obtusata* GYLLENHAL, 1813

*Haltica pallidicornis* WALTZ, 1839

*Mantura rustica* var. *concoloripennis* ROUBAL, 1922

*Mantura subobtusata* JANSSON, 1947

Distribution: Europe except for SW, Caucasus.

***Mantura rustica* (LINNAEUS, 1767)**

*Chrysomela rustica* LINNAEUS, 1767

*Mantura rustica* var. *suturalis* J. WEISE, 1886

*Altica rustica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GMELIN, 1790)

*Galleruca semiaenea* FABRICIUS, 1792

*Altica semiaenea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PANZER, 1795)

*Galeruca semiaenea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PAYKULL, 1799)

*Chrysomela semiaenea* FABRICIUS, 1801

*Haltica semiaenea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KOCH, 1803)

*Haltica rustica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ILLIGER, 1807)

Distribution: All over Palaearctic except for N Africa.

**Genus: *Sphaeroderma* STEPHENS, 1832*****Sphaeroderma balyi balyi* JACOBY, 1885**

Distribution: Asian Russia (S Maritime Prov.), Korea, China, Japan.

***Sphaeroderma fuscicorne* BALY, 1864**

Distribution: Asian Russia (S Maritime Prov.), Korea, China, Japan.

***Sphaeroderma rubidum* (GRAËLLS, 1858)**

*Argopus rubidus* GRAËLLS, 1858

? *Chrysomela testaceum* GOEZE, 1777

*Altica testacea* OLIVIER, 1789

*Sphaeroderma ocellaria* ALLARD, 1860

*Sphaeroderma rubidum* var. *testaceum* KUHNT, 1913

*Sphaeroderma rubidum* var. *gyllenhali* CAPRA, 1933

*Sphaeroderma nitidum* PORTEVIN, 1934

*Sphaeroderma rubidum* var. *ocularium* HEIKERTINGER et CSIKI, 1940

? *Galeruca testaceum*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PAYKULL, 1799)

*Sphaeroderma testaceum*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KRAATZ, 1861)

Distribution: Europe except for N, Caucasus, Turkey, Near East, N Africa, Madeira.

***Sphaeroderma testaceum* (FABRICIUS, 1775)**

*Altica testacea* FABRICIUS, 1775

? *Altica volutum* LAICHARTING, 1781

? *Altica fulvum* GEOFFROY, 1785

*Haltica cardui* GYLLENHAL, 1813

*Sphaeroderma centaureae* STEPHENS, 1832

*Sphaeroderma testacea* ab. *pacoviense* ROUBAL, 1913

*Galleruca testacea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1787)

*Chrysomela testacea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1801)

*Sphaeroderma cardui*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

*Argopus cardui*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KÜSTER, 1853)

*Argopus testaceus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KALTENBACH, 1874)

*Sphaeroderma rubidum*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to APFELBECK, 1914)

Distribution: Europe except for extreme N, Ciscaucasia, Caucasus, Transcaucasia.

***Sphaeroderma unicolor* KIMOTO, 1965**

Distribution: Asian Russia (Maritime Prov.), Japan.

**Genus: *Argopus* FISCHER-WALDHEIM, 1824**

*Dicherosis* FOUDRAS, 1860

***Argopus ahrensi* (GERMAR, 1817)**

*Haltica ahrensi* GERMAR, 1817

*Haltica hemisphaerica* DUFTSCHMIDT, 1825

*Argopus hemisphaericus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)

*Dicherosis hemisphaericus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOUDRAS, 1860)

Distribution: Central, E, and SE Europe, N Caucasus.

***Argopus bicolor* FISCHER-WALDHEIM, 1824**

*Argopus discolor* BIELZ, 1852

*Argopus punctatus* L. W. SCHAUFUSS, 1874

*Argopus ahrensi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GRUEV et TOMOV, 1965)

Distribution: Balkans, SE Europe, Caucasus.

***Argopus intermedius* J. WEISE, 1887**

Distribution: Asian Russia (Ussuri Area, Maritime Prov.).

***Argopus nigratarsis* (GEBLER, 1823)**

*Chrysomela nigratarsis* GEBLER, 1823

*Argopus intermedius* KRÁL, 1973

*Argopus clypeatus*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to CHŪJŌ, 1936)

*Dicherosis nigratarsis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOUDRAS, 1860)

Distribution: Eastern Central and E Europe, Turkmenistan, Kazakhstan, Asian Russia (Altai, Sayan, Transbaikalia, Amurland, Maritime Prov.), Mongolia, Korea, N and NE China, Japan.

***Argopus punctipennis punctipennis* (MOTSCHULSKY, 1866)**

*Dicherosis punctipennis* MOTSCHULSKY, 1866

*Argopus orientalis* BALY, 1874

*Sphaeroderma unicolor* JACOBY, 1885

*Argopus japonicus* CHEN, 1936

*Argopus unicolor* CHŪJŌ, 1936

*Argopus univestis* HEIKERTINGER, 1940

Distribution: Asian Russia (Sakhalin, Kuriles Isles), Japan.

***Argopus punctipennis substriatus* J. WEISE, 1887***Argopus riedeli* WARCHALOWSKI, 1969

Distribution: Asian Russia (Amurland, Maritime Prov., Ussuri Area), N Korea.

***Argopus unicolor* MOTSCHULSKY, 1860**

Distribution: Asian Russia (Amurland, Maritime Prov., Ussuri Area), Korea, Japan.

**Genus: *Pentamesa* HAROLD, 1876*****Pentamesa duodecimpunctata* (MOTSCHULSKY, 1860)***Colaphus duodecimpunctata* MOTSCHULSKY, 1860*Pentamesa tredecimpunctata* JACOBSON, 1902*Pentamesa duodecimmaculata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. MEDVEDEV, 1992)

Distribution: Asian Russia (Sayan, Altai, Baikal Area, Tuva, Yakutia, Amurland).

***Pentamesa kondarensis arnoldi* LOPATIN, 1996**

Distribution: Uzbekistan.

***Pentamesa kondarensis kondarensis* LOPATIN, 1956***Pentamesa duodecimmaculata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to SCHERER, 1969)

Distribution: Uzbekistan, Tajikistan.

**Genus: *Argopistes* MOTSCHULSKY, 1860*****Argopistes biplagiatus* MOTSCHULSKY, 1860***Argopistes biplagiatus* var. *flavitaris* MOTSCHULSKY, 1860*Argopistes biplagiatus* var. *limbatus* MOTSCHULSKY, 1860*Argopistes biplagiatus* var. *suturalis* MOTSCHULSKY, 1860*Argopistes undecimmaculatus* JACOBY, 1885*Argopistes biplagiatus* ab. *sequensi* REITTER, 1911*Argopistes biplagiatus* ab. *tenuelimbatus* REITTER, 1911*Argopistes biplagiatus* ab. *bipunctatus* REITTER, 1911*Argopistes biplagiatus* ab. *micraspoides* REITTER, 1911*Argopistes biplagiatus* ab. *rufus* CHEN, 1934

Distribution: Asian Russia (Amurland, Amurland, Ussuri Area, Khabarovsk Distr., Sakhalin), NE China, Korea, Japan.

***Argopistes coccinelliformis* CSIKI, 1940***Argopistes coccinelloides* BALY, 1874, non SUFFRIAN, 1868*Argopistes biplagiatus* CHEN, 1934, non MOTSCHULSKY, 1860

Distribution: NE China, Japan, Taiwan, Vietnam.

***Argopistes udege* KONSTANTINOV, 1994**

Distribution: Asian Russia (Far East).

***Argopistes unicolor* JACOBY, 1885**

Distribution: Asian Russia (Ussuri Area), Korea, Japan.

**Genus: *Hemipyxis* CHEVROLAT in DEJEAN, 1837***Sebaethe* BALY, 1864*Epiotis* SOLSKY, 1872***Hemipyxis plagioderoides* (MOTSCHULSKY, 1860)***Oedionychis plagioderoides* MOTSCHULSKY, 1860*Sebaethe amurensis* J. WEISE, 1887? *Sebaethe nila* MAULIK, 1926*Sebaethe yunnanica* CHEN, 1933*Epiotis plagioderoides*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to SOLSKY, 1872)

Distribution: Asian Russia (E Siberia, Amurland, Maritime Prov.), Korea, China, Japan.

**Genus: *Philopona* J. WEISE, 1903*****Philopona vibex* (ERICHSON, 1834)***Haltica vibex* ERICHSON, 1834*Oedionychis japonicus* BALY, 1874*Hyphasis signata* DUVIVIER, 1892*Oedionychis vibex* ab. *nigrodorsalis* HEIKERTINGER, 1922*Philopona vibex* ab. *limbata* J. WEISE, 1922*Philopona vibex* ab. *luzonicus* J. WEISE, 1922*Philopona signata* MAULIK, 1926*Philopona vibex japonica* CHŪJŌ, 1936*Oedionychis vibex* ab. *chūjōi* CSIKI, 1940*Oedionychis vibex* var. *nigra* CHŪJŌ, 1950*Oedionychis vibex*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1922)

Distribution: Asian Russia (Ussuri Area, S Kuriles: Kunashir Isle), Korea, China, Japan, Taiwan, N Vietnam, India.

**Genus: *Dibolia* LATREILLE, 1829***Petalopus* MOTSCHULSKY, 1845***Dibolia carpathica* J. WEISE, 1893***Dibolia transcaucasica* IABLOKOFF-KHINZORIAN, 1968

Distribution: SE Europe, S Urals, Ciscaucasia, Caucasus, Transcaucasia, Turkey, SE Kazakhstan.

***Dibolia cryptocephala* (KOCH, 1803)***Haltica cryptocephala* KOCH, 1803

Distribution: Central and SE Europe, S Urals, Ciscaucasia, Caucasus, Transcaucasia, Turkey, Kazakhstan.

***Dibolia cynoglossi* (KOCH, 1803)***Haltica cynoglossi* KOCH, 1803*Haltica cynoglossi* DUFTSCHMIDT, 1825*Dibolia paludina* FOUDRAS, 1860

Distribution: Europe except for extreme N, Armenia, Turkey.

***Dibolia depressiuscula* LETZNER, 1847***Dibolia laevicollis* FOU DRAS, 1860*Dibolia marrubii* HERING, 1924

Distribution: Europe except for N. Caucasus, Transcaucasia, Turkey, Near East, NE Kazakhstan, Asian Russia (W Siberia, Cisbaikalia, Transbaikalia), N Africa.

***Dibolia femoralis femoralis* L. REDTENBACHER, 1849***Haltica aerata* DALMAN, 1823*Dibolia aurichalcea* FÖRSTER, 1849*Dibolia femoralis* var. *aurichalcea* J. WEISE, 1893*Dibolia femoralis* var. *peregrina* J. WEISE, 1893*Dibolia theresae* PIC, 1909

Distribution: Europe except for N, Caucasus, Turkey.

***Dibolia foerstieri* BACH, 1856***Dibolia buglossi* FOU DRAS, 1860*Dibolia foudrasi* REY, 1874*Dibolia russica* J. WEISE, 1893 (new synonym)

Distribution: Europe except for N, (Belarus, Crimea, SE European Russia), Transcaucasia.

***Dibolia kraalii* MOHR, 1981**

Distribution: Transcaucasia, N Iran.

***Dibolia mesasiatica* LOPATIN, 1965**

Distribution: Uzbekistan, Tajikistan (Hissar Mountain Range).

***Dibolia occultans* (KOCH, 1803)***Haltica occultans* KOCH, 1803*Dibolia? paludina* FOU DRAS, 1860*Dibolia menthae* HERING, 1924

Distribution: Europe except for N, Caucasus, Turkey, Iran, N Africa, Canary Isles.

***Dibolia phoenicia* ALLARD, 1866***Dibolia orientalis* J. WEISE, 1893*Dibolia orientalis* var. *cyanescens* J. WEISE, 1893

Distribution: Balkans, Hungary, Caucasus (Georgia), Turkey, Near East.

***Dibolia rugulosa* L. REDTENBACHER, 1849**

Distribution: Europe except for N, Caucasus, Transcaucasia.

***Dibolia schillingii* (LETZNER, 1847)***Haltica (Dibolia) schillingii* LETZNER, 1847*Petalopus metallicus* MOTSCHULSKY, 1845*Dibolia punctillata* FOU DRAS, 1860*Dibolia schillingii* var. *ciliciensis* HEIKERTINGER, 1911*Dibolia schillingii ciliciensis* BECHYNÉ, 1956*Dibolia metallica*: GRJEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1930)

Distribution: Central and SE Europe, Caucasus, Transcaucasia, Turkey, W Kazakhstan.

***Dibolia timida* (ILLIGER, 1807)***Haltica timida* ILLIGER, 1807

*Dibolia eryngii* BACH, 1856

*Dibolia timida* var *oranensis* PIC, 1909

Distribution: Europe except for N and northern parts of Central, N Africa.

***Dibolia tricolor* REITTER, 1899**

Distribution: Armenia, Turkey.

***Dibolia tshatkatica corpulenta* MOHR, 1981**

Distribution: Kirghizstan.

***Dibolia tshatkatica tshatkatica* PALIJ, 1968**

Distribution: Turkmenistan, Kirghizstan.

***Dibolia turkmenica* IABLOKOFF-KHNZORIAN, 1978**

*Dibolia tshatkatica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to MOHR, 1981)

Distribution: Turkmenistan.

***Dibolia weisei* MOHR, 1981**

*Dibolia metallica* J. WEISE, 1893

Distribution: Caucasus (Georgia).

***Dibolia zaitzevi* L. MEDVEDEV, 1980**

Distribution: NE China, Mongolia.

***Dibolia zangezurica* IABLOKOFF-KHNZORIAN, 1968**

Distribution: Transcaucasia, Turkey, Iran.

**Genus: *Psylliodes* BERTHOLD, 1827**

***Psylliodes aeneolus* HEIKERTINGER, 1911**

Distribution: SE Kazakhstan.

***Psylliodes aereus austriacus* HEIKERTINGER, 1911**

Distribution: Mountains of Central, E, and SE Europe.

***Psylliodes affinis* (PAYKULL, 1799)**

*Galeruca affinis* PAYKULL, 1799

? *Chrysomela atricilla* LINNAEUS, 1761

*Haltica exoleta* ILLIGER, 1807

*Altica atricilla*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PANZER, 1791)

*Haltica affinis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KOCH, 1803)

*Macrocneema exoleta*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

Distribution: Europe, Caucasus, Transcaucasia, Turkey, Kazakhstan, Asian Russia (W Siberia), N Africa.

***Psylliodes angusticollis* BALY, 1874**

*Psylliodes sinensis* CHEN, 1934

*Psylliodes angusticollis rishiriensis* CHÛJÔ, 1959

*Psylliodes angusticollis loochooana* CHÛJÔ, 1961

*Psylliodes formosana* CHÛJÔ, 1963

Distribution: Asian Russia (Sakhalin, Kunashir Isle), China, Korea, Japan, Taiwan, N Vietnam.

***Psylliodes araraticus* IABLOKOFF-KHNZORIAN, 1968**

Distribution: Armenia.

***Psylliodes aristus* KHNZORIAN, 1962***Psylliodes arista* KHNZORIAN, 1962

Distribution: Armenia.

***Psylliodes attenuatus* (KOCH, 1803)***Haltica attenuata* KOCH, 1803*Macrocnema apicalis* STEPHENS, 1832*Macrocnema picicornis* STEPHENS, 1832*Psylliodes vicina* L. REDTENBACHER, 1874*Haltica oleracea* KÖPPEN, 1880*Psylliodes japonica* JACOBY, 1885*Haltica concinna* STREBEL, 1887*Psylliodes attenuata* var. *picicornis* J. WEISE, 1888*Psylliodes attenuata* var. *vicina* J. WEISE, 1888*Psylliodes attenuata* fa. *brunneotestacea* HEIKERTINGER, 1926*Psylliodes angusticollis* CHŪJŌ, 1936*Psylliodes attenuata* fa. *caerulea* KRÁL, 1947*Psylliodes attenuata japonica* SUZUKI, 1974*Psylliodes picicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)*Psylliodes hyoscyami*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KÖPPEN, 1880)*Psylliodes chrysocephala*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STREBEL, 1887)*Chaetocnema concinna*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ZIRNGIEBL, 1902)

Distribution: All over Palaearctic except for Near East and N Africa.

***Psylliodes callinotus* FALDERMANN, 1837**

Distribution: Caucasus.

***Psylliodes chalcomerus* (ILLIGER, 1807)***Haltica chalcomera* ILLIGER, 1807*Haltica brunnipes* DUFTSCHMIDT, 1825*Macrocnema unimaculata* STEPHENS, 1832*Psylliodes hyoscyami* var. *cardui* J. WEISE, 1888*Psylliodes chalcomera* var. *erythroceros* ABÉILLE, 1895*Psylliodes erythroceros* PEYERIMHOFF, 1911*Psylliodes chalcomera* var. *erythroceros* ab. *cyanescens* NORMAND, 1937

Distribution: All over Palaearctic except for southern Central Asia.

***Psylliodes chrysocephalus* (LINNAEUS, 1758)***Chrysomela chrysocephala* LINNAEUS, 1758*Altica chrysocephala* O. F. MÜLLER, 1776*Altica anglica* FABRICIUS, 1787*Altica anglicana* GMELIN, 1793*Altica napi* PANZER, 1794*Chrysomela nigricollis* MARSHAM, 1802*Haltica erythrocephala* KOCH, 1803*Haltica cyanoptera* ILLIGER, 1807 (part.)*Haltica nucea* ILLIGER, 1807? *Haltica elongata* GYLLENHAL, 1813

- Macrocnema sordida* STEPHENS, 1832  
*Psylliodes kunzei* FOU DRAS, 1860  
*Psylliodes rufilabris* ALLARD, 1860  
*Psylliodes allardi* BACH, 1866  
*Psylliodes catinensis* ROTTENBERG, 1871  
*Psylliodes chrysocephala* var. *anglica* J. WEISE, 1888  
*Psylliodes chrysocephala* var. *catinensis* J. WEISE, 1888  
*Psylliodes chrysocephala* var. *collaris* J. WEISE, 1888  
*Psylliodes chrysocephala* var. *nucea* J. WEISE, 1888  
*Psylliodes chrysocephala* var. *peregrina* J. WEISE, 1888  
*Psylliodes sicardi* J. WEISE, 1894  
*Psylliodes longipennis* PIC, 1903  
*Psylliodes castanea* ABEILLE, 1907  
*Psylliodes chrysocephala* ab. *collaris* PAGANETTI, 1910  
*Psylliodes chrysocephala* var. *cyanoptera* HEIKERTINGER, 1914  
*Psylliodes chrysocephala* var. *anglica* HEIKERTINGER, 1926  
*Psylliodes chrysocephala* fa. *collaris* HEIKERTINGER, 1926  
*Psylliodes chrysocephala* var. *sicardi* HEIKERTINGER, 1926  
*Psylliodes chrysocephala* var. *angusticollis* HEIKERTINGER, 1926  
*Psylliodes chrysocephala* var. *kunzei* HEIKERTINGER, 1926  
*Psylliodes nucea* HEIKERTINGER, 1926  
*Psylliodes chrysocephala* var. *kuntzei* HEIKERTINGER, 1930  
*Psylliodes chrysocephala* ab. *erythrocephala* HEIKERTINGER et CSIKI, 1940  
*Psylliodes chrysocephala* ab. *significata* CSIKI, 1953  
*Psylliodes chrysocephala* ab. *tristicula* CSIKI, 1953  
*Psylliodes chrysocephala angusticollis* BECHYNÉ, 1956  
*Psylliodes chrysocephala sicardi* MOHR, 1956  
*Psylliodes chrysocephala castanea* JOLIVET, 1967  
*Chrysomela anglica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1787)  
*Galleruca anglica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1792)  
*Crioceris anglica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1801)  
*Macrocnema nigricollis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)  
*Psylliodes nigricollis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FOU DRAS, 1860)  
*Psylliodes sordida*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to C. SCHAUFUSS, 1913)  
 Distribution: Europe, Caucasus, Transcaucasia, Turkey, Near East, Iran, N Africa, Azores, Madeira.

***Psylliodes circumdata* (W. REDTENBACHER, 1842)**

- Haltica circumdata* W. REDTENBACHER, 1842  
*Psylliodes lethierryi* ALLARD, 1860  
*Psylliodes sicana* REY, 1874  
*Psylliodes circumdata* ab. *integra* HEIKERTINGER et CSIKI, 1940  
 Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, Near East, Iran, N Africa.

***Psylliodes cucullatus* (ILLIGER, 1807)**

- Haltica cucullata* ILLIGER, 1807  
*Haltica spergulae* GYLLENHAL, 1813  
*Haltica spergulae* var. *angustata* WALTIL, 1835  
*Psylliodes vicina* BOIELDIEU, 1859  
 ? *Psylliodes augustalisi* PIC, 1909  
*Psylliodes cucullata gansuica* JACOBSON, 1922



? *Psylliodes cuculata ussuriensis* L. MEDVEDEV, 1973

Distribution: Europe except for E Mediterranean, N Caucasus, Kazakhstan, Asian Russia (All over Siberia), Mongolia, Korea, China (Kansu).

***Psylliodes cupreatus* (DUFTSCHMIDT, 1825)**

*Haltica cupreata* DUFTSCHMIDT, 1825

Distribution: Europe except for N and E Mediterranean, Transcaucasia, Kazakhstan, Kirghizstan, Uzbekistan, Asian Russia (Siberia east to Buratia), Mongolia.

***Psylliodes cupreus* (KOCH, 1803)**

*Haltica cuprea* KOCH, 1803

*Haltica polita* GRAVENHORST, 1807

*Haltica obscura* DUFTSCHMIDT, 1825

? *Psylliodes fusiformis* L. REDTENBACHER, 1849

*Psylliodes herbacea* FODRAS, 1860

? *Psylliodes laevata* FODRAS, 1860

*Psylliodes cupronitens* ALLARD, 1860

*Psylliodes cypricolor* ALLARD, 1860

*Psylliodes foudrasi* ALLARD, 1866

? *Psylliodes picipes* J. WEISE, 1888

*Psylliodes cuprea* ab. *herbacea* EVERTS, 1903

*Psylliodes cuprea* ab. *obscura* PORTA, 1934

*Psylliodes cuprea* var. *obscura* G. MÜLLER, 1953

*Psylliodes obscura*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEINHAUSEN, 1965)

Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, Near and Middle East, Turkmenistan, Kazakhstan, Kirghizstan, Asian Russia (W Siberia, Transbaikalia), Mongolia.

***Psylliodes cyanescens* J. WEISE, 1887**

Distribution: Asian Russia (Transbaikalia, Amurland, Ussuri Area).

***Psylliodes deplanatus* L. MEDVEDEV, 1962**

*Psylliodes deplanata* L. MEDVEDEV, 1962

Distribution: N Caucasus, Georgia.

***Psylliodes dilutellus* HEIKERTINGER, 1911**

*Psylliodes grigorievi* JACOBSON, 1922

Distribution: Kazakhstan, Central Asia.

***Psylliodes dulcamarae* (KOCH, 1803)**

*Haltica dulcamarae* KOCH, 1803

Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, Kazakhstan, Asian Russia (W Siberia), Mongolia.

***Psylliodes frivaldskyi* J. WEISE, 1888**

*Psylliodes glabra* KUTHY, 1900

Distribution: Carpathians.

***Psylliodes hyoscyami* (LINNAEUS, 1758)**

*Chrysomela hyoscyami* LINNAEUS, 1758

*Psylliodes cupronitens* FÖRSTER, 1849

*Psylliodes hyoscyami* var. *coerulescens* J. WEISE, 1888

*Altica hyoscyami*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1775)  
*Haltica hyoscyami*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to OLIVIER, 1789)  
*Galleruca hyoscyami*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1792)

Distribution: All over Palaeartic except for N Africa.

***Psylliodes illyricus* LEONARDI et GRUEV, 1993**

*Psylliodes picina obsкуроaenea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to ROUBAL, 1937-1941)

*Psylliodes picina picina*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GRUEV, 1970)

*Psylliodes obsкуроaenea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to LEONARDI, 1975)

Distribution: SE Europe, Ukrainian Carpathians, Turkey.

***Psylliodes instabilis* FOU DRAS, 1860**

*Psylliodes picipes* WATERHOUSE, 1858

Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, Near East, N Africa.

***Psylliodes isatidis* HEIKERTINGER, 1914**

*Psylliodes cuprea isatidis* HEIKERTINGER, 1914

Distribution: Europe except for N, N and E Caucasus, Turkey, Iran, Kazakhstan, Uzbekistan, Asian Russia (W Siberia), Mongolia.

***Psylliodes kasyi* LOPATIN, 1967**

Distribution: Transcaucasia, Turkey, Iran.

***Psylliodes longicollis* J. WEISE, 1900**

*Psylliodes caucasica* HEIKERTINGER, 1916

*Psylliodes oreophila* KHNZORIAN, 1957

Distribution: Ciscaucasia, Transcaucasia.

***Psylliodes luteolus* (O. F. MÜLLER, 1776)**

*Altica luteola* O. F. MÜLLER, 1776

*Haltica nana* DUFTSCHMIDT, 1825

*Macrocnema pallida* STEPHENS, 1832

*Psylliodes propinqua* L. REDTENBACHER, 1849

*Psylliodes luteola* var. *aberrans* ROUBAL, 1916

*Chrysomela luteola*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GOEZE, 1777)

Distribution: Europe except for N, Ciscaucasia, Caucasus, Transcaucasia, Turkey, Near and Middle East, N Africa.

***Psylliodes macellus agropyri* PALIJ, 1961**

*Psylliodes cucullata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to MOISEEV, 1956)

Distribution: E European Russia (Cisvolgensis: Saratov Distr.), N Kazakhstan.

***Psylliodes macellus macellus* (J. WEISE, 1900)**

*Semicnema macella* J. WEISE, 1900

*Psylliodes roddi* JACOBSON, 1922

*Psylliodes reitteri macella*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to DUBESHKO et L. MEDVEDEV, 1974)

*Psylliodes reitteri*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. MEDVEDEV et KOROTIAEV, 1975)

Distribution: E Kazakhstan, Asian Russia (S Siberia from Krasnoyarsk Distr. to Buriatia and Dauria), Mongolia.

***Psylliodes marcidus* (ILLIGER, 1807)**

*Psylliodes operosa* FOUDRAS, 1860

Distribution: Coastal parts of Atlantic, North Sea, Baltic Sea, Black Sea and Mediterranean countries.

***Psylliodes napi flavicornis* J. WEISE, 1883**

*Psylliodes napi* var. *flavicornis* J. WEISE, 1883

*Psylliodes lunariae* JAQUET, 1887

*Psylliodes napi* ab. *flavicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to PORTA, 1934)

*Psylliodes napi fulvicornis*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to GRUEV, 1992)

Distribution: Mountains of Central and E Europe.

***Psylliodes napi napi* (FABRICIUS, 1792)**

*Galleruca napi* FABRICIUS, 1792

*Haltica rapae* ILLIGER, 1807

*Psylliodes ecalcarata* L. REDTENBACHER, 1849

*Psylliodes napi* var. *asturiensis* J. WEISE, 1888

*Chrysomela napi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to FABRICIUS, 1801)

*Haltica napi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KOCH, 1803)

*Macrocnema napi*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

*Macrocnema rapae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

*Psylliodes rapae*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to KÜSTER, 1849)

Distribution: Europe, Caucasus, Transcaucasia, Turkey, S Kazakhstan, Asian Russia (Siberia east to Baikal and Yakutia), N Africa.

***Psylliodes nitidus* L. MEDVEDEV, 1973**

Distribution: Asian Russia (Ussuri Area).

***Psylliodes nivalis* KHNZORIAN, 1962**

Distribution: Caucasus (Georgia).

***Psylliodes ozisiki* LEONARDI et ARNOLD, 1995**

Distribution: Armenia, Turkey.

***Psylliodes pallidicornis* HEIKERTINGER, 1921**

Distribution: Caucasus, Turkmenistan

***Psylliodes persicus* ALLARD, 1866**

Distribution: Caucasus, Transcaucasia, Near and Middle East, Saudi Arabia, Kazakhstan, Central Asia.

***Psylliodes picinus* (MARSHAM, 1802)**

*Chrysomela picina* MARSHAM, 1802

*Haltica rufilabris* KOCH, 1803

*Haltica melanophthalma* DUFISCHMIDT, 1825

*Macrocnema apicalis* STEPHENS, 1832

*Psylliodes rufopicea* LEIZNER, 1847

*Psylliodes picea* L. REDTENBACHER, 1849

*Psylliodes picina* var. *bicolor* GERHARDT, 1908

*Macrocnema picina*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to STEPHENS, 1832)

? *Psylliodes rufilabris*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)

*Psylliodes melanophthalma*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to L. REDTENBACHER, 1849)

Distribution: Europe except for N.

***Psylliodes pubipennis* LOPATIN, 1958**

Distribution: S Tajikistan.

***Psylliodes punctifrons* BALY, 1974**

Distribution: Asian Russia (S Maritime Prov.), Korea, China, Japan, Taiwan, Indochina, Vietnam, Sumatra.

***Psylliodes reitteri parallelus* J. WEISE, 1890**

Distribution: SE European Russia, Kazakhstan, Central Asia, Asian Russia (S Siberia, Amurland, Maritime Prov.), Mongolia, China.

***Psylliodes reitteri reitteri* J. WEISE, 1888**

Distribution: Central and SE Europe.

***Psylliodes rhaicus* JACOBSON, 1922**

Distribution: S Ukraine. S European Russia, W Kazakhstan.

***Psylliodes rubroaeneus* HEIKERTINGER, 1916**

Distribution: Ciscaucasia, Georgia.

***Psylliodes saulcyi* ALLARD, 1867**

*Psylliodes atriplicis* JACOBSON, 1922

Distribution: SE Ukraine, S European Russia, N Caucasus, Near and Middle East, Kazakhstan, Mongolia, N Africa.

***Psylliodes sophiae* HEIKERTINGER, 1914**

? *Haltica elongata* GYLLENHAL, 1813

*Psylliodes cyanoptera* L. REDTENBACHER, 1849

*Psylliodes chrysocephala* FOU DRAS, 1860 (part.)

*Psylliodes cyanoptera* var. *tricolor* J. WEISE, 1888

*Psylliodes sophiae* ab. *tricolor* HEIKERTINGER et CSIKI, 1940

*Psylliodes cyanoptera* ab. *nigrivertex* JACOBSON, 1902

*Psylliodes cyanoptera* ab. *nigrifrons* HEIKERTINGER, 1941

*Psylliodes sophiae* fa. *nigricapilia* KRÁL, 1945

*Psylliodes sophiae* ab. *lichtmeckeri* KASZAB, 1962

*Psylliodes sophiae* ab. *annunialis* KASZAB, 1962

Distribution: Europe, Caucasus, Transcaucasia, Turkey, Near and Middle East, Kazakhstan, Central Asia, Asian Russia (S Siberia east to Baikal), N Africa.

***Psylliodes subaeneus* KUTSCHERA, 1867**

Distribution: Mountains of Central and E Europe (Dynars, Carpathians).

***Psylliodes subrugosus* JACOBY, 1885**

*Psylliodes intermedia* JACOBY, 1885

*Psylliodes mitchyi* CHŪJŌ, 1951

Distribution: Asian Russia (Sakhalin), China, Japan.

***Psylliodes testaceoconcolor* HEIKERTINGER, 1926**

*Psylliodes cuprea* fa. (var.) *testaceoconcolor* HEIKERTINGER, 1926

Distribution: Transcaucasia, Near East.

***Psylliodes thlaspi* FOU DRAS, 1860**

? *Psylliodes fusiformis* L. REDTENBACHER, 1858

Distribution: Europe except for N and northern parts of Central Europe, Caucasus.

***Psylliodes toelgi* HEIKERTINGER, 1914**

? *Psylliodes picipes* FOU DRAS, 1860

*Psylliodes napi* var. *brisouti* G. MÜLLER, 1953

Distribution: Mountains of Europe, except for Fennoscandia.

***Psylliodes validus* J. WEISE, 1889**

Distribution: Ciscaucasia, Caucasus, Transcaucasia.

***Psylliodes vindobonensis* HEIKERTINGER, 1914**

Distribution: Mountains of Central, E, and SE Europe.

***Psylliodes wrasei* LEONARDI et ARNOLD, 1995**

Distribution: Balkans, ?S Ukraine, Caucasus (Georgia).

**Genus: *Nonarthra* BALLY, 1862**

***Nonarthra cyaneum* BALLY, 1874**

*Nonarthra fulvum* BALLY, 1874

*Nonarthra nigricolle* J. WEISE, 1889

Distribution: Asian Russia (Amurland, Maritime Prov.), Korea, China, Japan, Taiwan, Vietnam.

**Genus: *Sangariola* JACOBSON, 1922**

*Charidea* BALLY, 1888

***Sangariola punctatostrata* (MOTSCHULSKY, 1860)**

*Galleruca punctatostrata* MOTSCHULSKY, 1860

*Galeruca multicosata* JACOBY, 1885

*Charidea regularis* PIC, 1928

*Sangariola costata* CHŪJŌ, 1935

*Sangariola punctatostrata aequicostata* CHŪJŌ, 1938

*Sangariola multicosata* CHŪJŌ, 1954

*Lochmaeina rosea* L. MEDVEDEV, 1956

Distribution: Asian Russia (Sakhalin, Kunashir Isle), N China, Korea, Japan.

**Genus: *Liprus* MOTSCHULSKY, 1860**

*Crepidomorpha* FLEISCHER, 1916

***Liprus punctatostratus* MOTSCHULSKY, 1860**

*Diabrotica rufotestacea* MOTSCHULSKY, 1860

*Crepidodera japonica* JACOBY, 1885

- Crepidodera japonensis* SCHÖNFELDT, 1887  
*Crepidodera carinulata* FLEISCHER, 1916  
*Crepidomorpha carinulata*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1923).  
*Crepidomorpha japonica*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to CHEN, 1935).  
*Crepidomorpha rufotestacea*: GRUEV et DÖBERL, 1997 (in synonymy erroneously attributed to HEIKERTINGER, 1948).  
 Distribution: Asian Russia (S Maritime Prov.), Korea, Japan, China (Kiangsu, Kirin), Taiwan.

**Genus: *Phygasia* CHEVROLAT in DEJEAN, 1837**

***Phygasia fulvipennis* (BALY, 1874)**

- Graptodera fulvipennis* BALY, 1874  
*Scaliodera fulvipennis* HAROLD, 1877  
*Aldrisma externecostata* FAIRMAIRE, 1888  
 Distribution: Asian Russia (Maritime Prov., Sikhote-Alin Mountains), Korea, N and NE China, Japan.

**Genus: *Luperomorpha* J. WEISE, 1887**

*Pushtunaltica* LOPATIN, 1962

***Luperomorpha funesta* (BALY, 1874)**

- Phyllostreta funesta* BALY, 1874  
*Luperomorpha trivialis* J. WEISE, 1887  
 Distribution: W China, Asian Russia (Maritime Prov.), Korea, Japan.

***Luperomorpha klapperichi* (LOPATIN, 1962)**

- Pushtunaltica klapperichi* LOPATIN, 1962  
 Distribution: NE Afghanistan

***Luperomorpha suturalis* CHEN, 1938**

- Luperomorpha suturalis similis* CHEN et KUNG, 1954  
*Luperomorpha trivialis* KRÁL, 1965  
 Distribution: S Asian Russia (S Siberia, S Baikal Area, Amurland), Mongolia, NE China.

**Subfamily: *Hispinae* GYLLENHAL, 1813**

**Genus: *Dactylispa* J. WEISE, 1897**

***Dactylispa angulosa* (SOLSKY, 1871)**

- Hispa angulosa* SOLSKY, 1871  
 Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

***Dactylispa excisa* (KRAATZ, 1879)**

- Hispa excisa* KRAATZ, 1879  
 Distribution: Asian Russia (S Amurland, Maritime Prov.), China, Korea.

***Dactylispa marginicollis borealis* L. MEDVEDEV, 1963**

Distribution: Asian Russia (S Maritime Prov.).

***Dactylispa masonii* GESTRO, 1923***Dactylispa ussurina* UHMANN, 1928

Distribution: Asian Russia (S Maritime Prov., Sakhalin), China, Japan.

**Genus: *Rhadinosa* J. WEISE, 1905*****Rhadinosa nigrocyanea* (MOTSCHULSKY, 1861)***Hispa nigrocyanea* MOTSCHULSKY, 1861*Rhadinosa reticulata* sensu Uhmann, 1940 nec Baly, 1888

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

**Genus: *Hispa* LINNAEUS, 1767***Hispella* CHAPUIS, 1875***Hispa atra* LINNAEUS, 1767***Hispa cariosa* REICHE et SAULCY, 1858*Hispa algira* MOTSCHULSKY, 1860*Hispa aptera* MOTSCHULSKY, 1860*Hispa minuta* BALBI, 1892*Hispella balbii* DONCKIER DE DONCEEL, 1899

Distribution: Europe, Caucasus, Kazakhstan, Central Asia, Asian Russia (S Siberia).

**Genus: *Hispellinus* J. WEISE, 1897***Monochirus* CHAPUIS, 1875 non RAFINESQUE***Hispellinus moerens* (BALY, 1874)***Hispa moerens* BALY, 1874

Distribution: Asian Russia (S Amurland), China, Japan.

**Genus: *Acmenychnus* J. WEISE, 1905*****Acmenychnus caucasicus* (HEYDEN, 1878)***Monochirus caucasicus* HEYDEN, 1878

Distribution: Transcaucasia.

***Acmenychnus inermis* (ZUBKOV, 1833)***Hispa inermis* ZUBKOV, 1833*Monochirus potanini* J. WEISE, 1889

Distribution: Kazakhstan, Central Asia, NE China, Mongolia.

**Genus: *Cassidispa* GESTRO, 1899*****Cassidispa relicta* L. MEDVEDEV, 1957**

Distribution: Asian Russia (S Maritime Prov.).

Subfamily: *Cassidinae* GYLLENHAL, 1813

Tribe: *Aspidimorphini* HINCKS, 1952

Genus: *Aspidimorpha* HOPE, 1840

*Aspidimorpha difformis* (MOTSCHULSKY, 1860)

*Deloyala difformis* MOTSCHULSKY, 1860

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

*Aspidimorpha transparipennis* (MOTSCHULSKY, 1860)

*Coptocycla transparipennis* MOTSCHULSKY, 1860

Distribution: Asian Russia (Amurland, Maritime Prov.), Mongolia, China, Korea.

Tribe: *Cassidini* HINCKS, 1952

Genus: *Thlaspida* J. WEISE, 1899

*Thlaspida lewisi* (BALY, 1874)

*Coptocycla lewisi* BALY, 1874

Distribution: Asian Russia (S Amurland, Maritime Prov.), China, Korea, Japan.

Genus: *Glyphocassis* SPAETH, 1914

*Hebdomecosta* SPAETH, 1915

*Glyphocassis spilota* (GORHAM, 1885)

*Coptocycla spilota* GORHAM, 1885

*Hebdomecosta reitteri* SPAETH, 1915

*Hebdomecosta shirahatai* CHŪJŌ, 1949

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

Genus: *Macromonycha* SPAETH, 1911

*Kari* MAULIK, 1923

*Macromonycha apicalis* (GEBLER, 1845)

*Cassida apicalis* GEBLER, 1845

*Cassida suberosa* J. WEISE, 1890

*Cassida discoidalis* REITTER, 1891

Distribution: Transcaucasia, S Kazakhstan, Central Asia.

Genus: *Chiridula* J. WEISE, 1889

*Chiridula semenovi* J. WEISE, 1889

*Cassida foyangula* REITTER, 1901

Distribution: Transcaucasia, Kazakhstan, Central Asia.

Genus: *Eremocassis* SPAETH, 1926

*Eremocassis weisei* (JACOBSON, 1894)

*Cassida weisei* JACOBSON, 1894



? *Eremocassis transcaspica* SPAETH, 1926  
*Cassida palyi* MATIS, 1970  
 Distribution: S Kazakhstan, Central Asia.

**Genus: *Ischyronota* J. WEISE, 1891**

***Ischyronota conicicollis* (J. WEISE, 1890)**

*Cassida conicicollis* J. WEISE, 1890  
 Distribution: Kazakhstan, Tajikistan, Mongolia.

***Ischyronota desertorum* (GEBLER, 1834)**

*Cassida desertorum* GEBLER, 1834  
 Distribution: Southern parts of E Europe, E Ciscaucasia, Kazakhstan, Central Asia,  
 NW China, Mongolia.

***Ischyronota elevata* (REITTER, 1890)**

*Cassida elevata* REITTER, 1890  
*Cassida nitidula* J. WEISE, 1890  
 Distribution: Transcaucasia, Kazakhstan, Turkmenistan, Uzbekistan, Iran.

***Ischyronota schusteri* SPAETH, 1914**

Distribution: S Kazakhstan, Turkmenistan, W China, Mongolia.

***Ischyronota spaethi* REITTER, 1901**

*Ischyronota deserticola* REITTER, 1901  
 Distribution: Kazakhstan.

**Genus: *Cassida* LINNAEUS, 1758**

**Subgenus: *Cassida (Alledoya)* HINCKS, 1950**

*Deloyala* L. REDTENBACHER, 1858 not CHEVROLAT, 1837

***Cassida (Alledoya) seraphina hablitziae* MOTSCHULSKY, 1838**

*Cassida testudo* SUFFRIAN, 1844  
 Distribution: Transcaucasia, Kazakhstan (Chu-Ili Mountains).

***Cassida (Alledoya) vespertina* BOHEMAN, 1862**

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

**Subgenus: *Cassida (Cassida)* Linnaeus, 1758**

***Cassida (Cassida) atrata* FABRICIUS, 1787**

Distribution: Central and SE Europe, Turkey.

***Cassida (Cassida) aurora* J. WEISE, 1907**

*Cassida rosea* BOHEMAN, 1854, non ILLIGER, 1798  
 Distribution: Southerneastern parts of Central Europe, Ciscaucasia.

***Cassida (Cassida) berolinensis* SUFFRIAN, 1844**

*Cassida obsoleta* HERBST, 1799, non ILLIGER, 1798  
*Cassida daurica* BOHEMAN, 1854  
*Cassida berolinensis* var. *pectoralis* J. WEISE, 1896

*Cassida berolinensis* ab. *pallidiventris* REITTER, 1912

Distribution: SE Europe, Kazakhstan, Asian Russia (S Siberia, Far East), Mongolia, N China.

***Cassida (Cassida) denticollis* SUFFRIAN, 1844**

*Cassida denticollis* var. *fuscicollis* J. WEISE, 1893

Distribution: Europe, Kazakhstan, Asian Russia (S Siberia, Far East).

***Cassida (Cassida) fausti* SPAETH, 1926**

*Cassida vibex* ab. *fausti* SPAETH, 1926

Distribution: S Ukraine, Ciscaucasia, Daghستان, Turkey.

***Cassida (Cassida) ferruginea* GOEZE, 1777**

*Cassida thoracica* GEOFFROY in FOURCROY, 1785

*Cassida tincta* J. WEISE, 1891

*Cassida ferruginea* var. *scutellaris* DELLA BEFFA, 1909

Distribution: Central and SE Europe, N Kazakhstan, Asian Russia (W Siberia).

***Cassida (Cassida) flaveola* THUNBERG, 1794**

*Cassida oscoleta* ILLIGER, 1798

*Cassida pallida* PAYKULL, 1799

*Cassida exsculpta* CHARPENTIER, 1825

*Cassida oscoleta* var. *dorsalis* DESBROCHERS DES LOGES, 1891, non HERBST, 1799

Distribution: Europe except for extreme N, S and Mediterranean, N Kazakhstan, Asian Russia (W Siberia).

***Cassida (Cassida) fusciorufa* MOTSCHULSKY, 1866**

*Cassida consocinta* BALY, 1874

*Cassida russata* J. WEISE, 1889, non FAIRMAIRE, 1887

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

***Cassida (Cassida) inquinata* BRULLÉ, 1832**

*Cassida plana* CHARPENTIER, 1825

*Cassida depressa* SUFFRIAN, 1844

*Cassida bohemanii* BRISOUT DE BARNEVILLE, 1860

*Cassida sordida* BRISOUT DE BARNEVILLE, 1860

*Cassida inquinata* var. *flavimana* DESBROCHERS DES LOGES, 1891

Distribution: Southern parts of E Europe, Caucasus, Turkmenistan.

***Cassida (Cassida) lineola* CREUTZER, 1799**

*Cassida russica* HERBST, 1799

*Cassida signata* HERBST, 1799

*Cassida sibirica* GEBLER, 1834

*Cassida bicostata* FISCHER-WALDHIEM, 1842

*Cassida suturalis* FISCHER-WALDHIEM, 1842

*Cassida nigroguttata* GORHAM, 1885

*Cassida nigrostrigata* FAIRMAIRE, 1888

Distribution: Central and southern parts of E Europe, Caucasus, Kazakhstan, Kirghizstan, Uzbekistan, Asian Russia (Siberia east to Maritime Prov.), Mongolia, China, Korea, Japan, Taiwan.

***Cassida (Cassida) mandli* SPAETH, 1921**

Distribution: Asian Russia (Amurland, Maritime Prov.), Korea, China.

***Cassida (Cassida) mongolica* BOHEMAN, 1854***Cassida rissata* FAIRMAIRE, 1887

Distribution: Asian Russia (Cisbaikalia, Amurland), Mongolia, NE China.

***Cassida (Cassida) nebulosa* LINNAEUS, 1758***Cassida affinis* FABRICIUS, 1775*Cassida tigrina* DEGEER, 1775*Cassida nebulosa* var. *subferruginea* ILLIGER, 1798, non SCHRANK, 1776*Cassida nigra* HERBST, 1799*Cassida atra* SCHWAEGRICHEN, 1844

Distribution: All over Palaearctic except for Mediterranean. Introduced to N America.

***Cassida (Cassida) palaestina* REICHE, 1858***Cassida incompta* J. WEISE, 1887

Distribution: Near East, Transcaucasia, Kazakhstan, Central Asia.

***Cassida (Cassida) pallidicollis* BOHEMAN, 1856***Cassida diabolica* KRAATZ, 1879*Cassida morawitzi* JACOBSON, 1894

Distribution: Asian Russia (S Siberia, Amurland, Maritime Prov.), Mongolia, China.

***Cassida (Cassida) pannonica* SUFFRIAN, 1848***Cassida elongata* J. WEISE, 1893*Cassida pannonica* var. *fraudulenta* SPAETH, 1914

Distribution: Southern parts of Central and SE Europe, Caucasus, Turkey, Near East, Kazakhstan, Kirghizstan, Asian Russia (W Siberia).

***Cassida (Cassida) panzeri* J. WEISE, 1907***Cassida thoracica* PANZER, 1796, non GEOFFROY, 1785

Distribution: Central, E, and SE Europe, Caucasus, Kazakhstan, Asian Russia (Siberia east to Far East).

***Cassida (Cassida) piperata* HOPE, 1842***Cassida biguttulata* KRAATZ, 1879*Cassida labilis* BOHEMAN, 1854*Cassida japana* BALY, 1874

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

***Cassida (Cassida) prasina* ILLIGER, 1798***Cassida viridana* HERBST, 1799*Cassida chloris* SUFFRIAN, 1844

Distribution: Europe except for N, SW and Mediterranean, Caucasus, Transcaucasia, Turkey, Kazakhstan, Asian Russia (Siberia east to Far East).

***Cassida (Cassida) reitteri* J. WEISE, 1892**

Distribution: Armenia.

***Cassida (Cassida) rubiginosa rubiginosa* O. F. MÜLLER, 1776***Cassida cardui* DEGEER, 1775*Cassida viridis* FABRICIUS, 1781*Cassida melanosceles* SCHRANK, 1798*Cassida similis* MARSHAM, 1802

*Cassida singularis* STEPHENS, 1834  
*Cassida graeca* KRAATZ, 1874  
*Cassida rubiginosa* var. *fuliginosa* J. WEISE, 1893  
*Cassida sparre-schneideri* HELLIESEN, 1898  
 Distribution: Europe, Asian Russia (W Siberia).

***Cassida (Cassida) rubiginosa rugosopunctata* MOTSCHULSKY, 1866**

Distribution: E Asian Russia (E Siberia, Far East, Sakhalin, Kuriles Isles), Mongolia.

***Cassida (Cassida) rufovirens* SUFFRIAN, 1844**

*Cassida rhilensis* J. WEISE, 1891  
 Distribution: Central and southern parts of E Europe, Ciscaucasia, Turkey.

***Cassida (Cassida) sanguinolenta* O. F. MÜLLER, 1776**

*Cassida cruentata* DONOVAN, 1793  
 Distribution: Europe except for N and SW, Turkey, S Urals, Kazakhstan, Asian Russia (W Siberia, Cisbaikalia).

***Cassida (Cassida) sanguinosa* SUFFRIAN, 1844**

*Cassida languida* CORNELIUS, 1851  
*Cassida sanguinosa* ab. *viridissima* REITTER, 1912  
 Distribution: Europe except for S and extreme N, Caucasus, Kazakhstan, Asian Russia (Siberia east to Yakutia).

***Cassida (Cassida) sareptana* KRAATZ, 1873**

Distribution: Steppe of Ukraine and European Russia, Ciscaucasia, W Kazakhstan.

***Cassida (Cassida) seladonia* GYLLENHAL, 1827**

*Cassida filaginis* PERRIS, 1855  
*Cassida rotundicollis* BRISOUT DE BARNEVILLE, 1860  
 Distribution: Europe except for extreme N and E, Ciscaucasia, N Africa.

***Cassida (Cassida) spaethi* J. WEISE, 1900**

Distribution: Asian Russia (Amurland, Maritime Prov.), Korea, China.

***Cassida (Cassida) stigmatica* SUFFRIAN, 1844**

Distribution: Europe (except for N, NW, Iberian, Appenine and Balkan Peninsula), Turkey, Kazakhstan, Asian Russia (S Siberia), N China.

***Cassida (Cassida) vibex* LINNAEUS, 1767**

*Cassida liriophora* KIRBY, 1837  
*Cassida angusticollis* HAGENBACH, 1822  
*Cassida dorsalis* HERBST, 1799  
*Cassida vibex* var. *discoidea* J. WEISE, 1893  
*Cassida vibex* ab. *significata* CSIKI, 1953  
 Distribution: All over Palaearctic except for extreme N.

**Subgenus: *Cassida (Cassidulella)* A. STRAND, 1928**

*Cassidula* J. WEISE, 1889

***Cassida (Cassidulella) nobilis* LINNAEUS, 1758**

*Cassida urticae* BRAHM, 1790

- Cassida pulchella* PANZER, 1797  
*Cassida viridula* PAYKULL, 1798  
*Cassida laevis* HERBST, 1799  
*Cassida nobilis* var. *obscura* J. WEISE, 1893, non FABRICIUS, 1792  
*Cassida nobilis* var. *obscurella* J. WEISE, 1907  
*Cassida nobilis* var. *dollmanni* FOWLER, 1913

Distribution: All over Palaearctic except for extreme N and N Africa.

***Cassida (Cassidulella) parvula* BOHEMAN, 1854**

- Cassida navicula* BOHEMAN, 1854  
*Cassida comparata* RYBAKOV, 1889

Distribution: Eastern S and southern E Europe, Crimea, Ciscaucasia, Caucasus, Turkmenistan, Kazakhstan, Central Asia, Asian Russia (S Siberia to Amurland), Mongolia.

***Cassida (Cassidulella) pusilla* WALTJ., 1839**

- Cassida puncticollis* SUFFRIAN, 1844  
*Cassida mollis* BOHEMAN, 1854  
*Cassida angusta* MARSEUL, 1878  
*Cassida pyrenaea* J. WEISE, 1893  
*Cassida pupillata* var. *olcesei* PIC, 1918

Distribution: E Mediterranean.

***Cassida (Cassidulella) velaris* J. WEISE, 1896**

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Japan.

***Cassida (Cassidulella) vittata* VILLERS, 1789**

- Cassida oblonga* ILLIGER, 1798  
*Cassida salicorniae* CURTIS, 1826

Distribution: Europe except for N, Caucasus, Kazakhstan, Uzbekistan, Asian Russia (S Siberia east to Maritime Prov.), Japan.

**Subgenus: *Cassida (Mionycha)* J. WEISE, 1891**

***Cassida (Mionycha) azurea* FABRICIUS, 1801**

- Cassida lucida* SUFFRIAN, 1844  
*Cassida ornata* CREUTZER, 1799, non HERBST, 1783  
*Cassida azurea* var. *femorata* CSIKI, 1953

Distribution: Central and E Europe, S Kazakhstan, Asian Russia (S Siberia), N Africa.

***Cassida (Mionycha) concha* SOLSKY, 1872**

Distribution: Asian Russia (Amurland, Maritime Prov.), China, Korea, Japan.

***Cassida (Mionycha) margaritacea* SCHALLER, 1783**

- Cassida superba* GMELIN, 1787  
*Cassida mutabilis* VILLIERS, 1789  
*Cassida spergulae* MARSHAM, 1802  
*Cassida margaritacea* var. *albiventris* SUFFRIAN, 1844  
*Cassida margaritacea* var. *melanocephala* SUFFRIAN, 1844  
*Cassida margaritacea* var. *nigrescens* J. WEISE, 1893

Distribution: Europe (except for N, SW, northern and central parts of E) Caucasus, Turkey, N Kazakhstan, Asian Russia (Cisbaikalia).

***Cassida (Mionycha) subreticulata* SUFFRIAN, 1844**

- Cassida splendidula* SUFFRIAN, 1844, non MARSHAM, 1802  
*Cassida subreticulata* var. *ventralis* J. WEISE, 1893  
*Cassida subreticulata helva* SPAETH, 1920  
*Cassida hincksi* L. MEDVEDEV, 1957

Distribution: Central, S and southern parts of E Europe, Caucasus, Transcaucasia, Turkey, Iran, Kazakhstan, Central Asia, Asian Russia (Siberia east to Amurland), Mongolia.

**Subgenus: *Cassida (Lordiconia)* REITTER, 1926*****Cassida (Lordiconia) canaliculata* LAICHARTING, 1781**

- Cassida speciosa* BRAHM, 1790  
*Cassida austriaca* FABRICIUS, 1792  
*Cassida austriaca* fa. *graminis* SUFFRIAN, 1844  
*Cassida austriaca* var. *illyrica* BOHEMAN, 1854  
*Cassida austriaca* var. *immutabilis* BOHEMAN, 1854

Distribution: SE Europe and southern parts of E Europe, Caucasus, Kazakhstan.

**Subgenus: *Cassida (Mionychella)* SPAETH, 1926 in HINCKS, 1952*****Cassida (Mionychella) hemisphaerica* HERBST, 1799**

- Cassida anglica* CURTIS, 1826  
*Cassida concinna* STEPHENS, 1834  
*Cassida rossi* WOLLASTON, 1857  
*Cassida nigriventris* HEYDEN, 1877, non BOHEMAN, 1854  
*Cassida corii* FAIRMAIRE, 1874  
*Cassida hemisphaerica* var. *peculiosa* J. WEISE, 1895  
*Cassida hemisphaerica* ab. *heydeni* SPAETH, 1914

Distribution: Europe except for extreme N, Caucasus, Turkey, N Africa.

**Subgenus: *Cassida (Lordicassis)* REITTER, 1926*****Cassida (Lordicassis) medvedevi* LOPATIN, 1965**

Distribution: Tajikistan (Mountains).

***Cassida (Lordicassis) moori* BOHEMAN, 1856**

Distribution: Kazakhstan, Kirghizstan, India (Mountains).

***Cassida (Lordicassis) tianshanica* BOROWIEC et SWIETOJANSKA, 2001**

Distribution: Kirghizstan (Tian-Shan Mountains).

***Cassida (Lordicassis) transcaucasica* BOROWIEC et SWIETOJANSKA, 2001**

Distribution: Transcaucasia.

***Cassida (Lordicassis) undecimnotata* GEBLER, 1834**

- Cassida undecimguttata* REITTER, 1890  
*Cassida impicticollis* J. WEISE, 1894

Distribution: Middle East, Kazakhstan, Central Asia.

**Subgenus: *Cassida (Odontionycha)* J. WEISE, 1891*****Cassida (Odontionycha) viridis* LINNAEUS, 1758***Cassida equestris* FABRICIUS, 1787*Cassida nigriceps* FAIRMAIRE, 1851*Cassida flaviceps* MARSEUL, 1878

Distribution: All over Palaearctic except for extreme N.

**Subgenus: *Cassida (Onychocassis)* SPAETH in SPAETH et REITTER, 1926*****Cassida (Onychocassis) bella* FALDERMANN, 1837**

Distribution: Caucasus, Turkey.

**Subgenus: *Cassida (Taiwania)* SPAETH, 1913*****Cassida (Taiwania) amurensis* KRAATZ, 1879**

Distribution: Asian Russia (Amurland, Maritime Prov.), Korea, China.

***Cassida (Taiwania) versicolor* BOHEMAN, 1855**

Distribution: Asian Russia (S Amurland, Maritime Prov.), China, Japan.

**Subgenus: *Cassida (Pseudocassida)* DESBROCHERS DES LOGES, 1891*****Cassida (Pseudocassida) murraea* LINNAEUS, 1767***Cassida maculata* LINNAEUS, 1767*Cassida variegata* GEOFFROY, 1785*Cassida varia* LATREILLE, 1804*Cassida murraea* var. *immaculata* DESBROCHERS DES LOGES, 1891*Cassida murraea* var. *dorsalis* J. WEISE, 1893, non HERBST, 1799*Cassida subobliterata* PIC, 1902*Cassida murraea* var. *inundata* J. WEISE, 1907*Cassida murraea ussuriensis* SPAETH, 1921*Cassida murraea halophila* SPAETH, 1921*Cassida pseudomurraea* GRUEV, 1978

Distribution: All over Palaearctic except for W Mediterranean, Central Asia and N Africa.

**Subgenus: *Cassida (Tylocentra)* REITTER, 1926*****Cassida (Tylocentra) saucia* J. WEISE, 1889**

Distribution: Caucasus, Turkey.

***Cassida (Tylocentra) sussamyrica* SPAETH, 1926**

Distribution: S Kazakhstan, Kirghizstan.

***Cassida (Tylocentra) turcmenica* J. WEISE, 1892***Cassida hauseri* J. WEISE, 1894*Cassida arata* LOPATIN, 1970

Distribution: Kazakhstan, Central Asia, Mongolia.

**Genus: *Pilemostoma* DESBROCHERS DES LOGES, 1891*****Pilemostoma bucharica* SPAETH, 1914**

Distribution: Uzbekistan.

***Pilemostoma fastuosa* (SCHALLER, 1783)***Cassida vittata* FABRICIUS, 1798*Cassida ocellata* HERBST, 1799*Cassida vittata* var. *delicatula* J. WEISE, 1893*Cassida vittata* var. *nigrina* J. WEISE, 1893

Distribution: Europe except for N and western Mediterranean, Caucasus, Kazakhstan, Central Asia, Asian Russia (S Siberia), Mongolia.

**Genus: *Hypocassida* J. WEISE, 1893*****Hypocassida subferruginea* (SCHRANK, 1776)***Cassida ferruginea* FABRICIUS, 1781*Cassida fusca* LAICHARTING, 1781*Cassida cornea* MARSEUL, 1869*Cassida costata* BOHEMAN, 1856*Cassida subferruginea* var. *sobrina* J. WEISE, 1893*Cassida subferruginea* ab. *marginata* BECHYNÉ, 1944

Distribution: Europe except for N, Caucasus, Transcaucasia, Turkey, Kazakhstan, Afghanistan, Central Asia, Asian Russia (S Siberia east to Maritime Prov.), W China, N Africa.



## References

- Abeille, P. E. 1895. Cliq nouvelles Altises du Nord de l'Afrique (*Coleoptera*). Bulletin de la Société Entomologique de France. Paris, 20: CDI-CDIV.
- Abeille, P.E. 1897. Description de deux *Entomoscelis* paléarctiques (*Col.*). Bulletin de la Société Entomologique de France. Paris, 22: 42.
- Abeille, P. E. 1907. Description de sept Halticidés réputées nouvelles. Annales de la Société Scientifique Naturelle de Provence, 1: LXXIX-LXXXII.
- Abeille, P. E. 1909. Coléoptères d'Algérie jugés nouveaux. Bulletin de la Société Entomologique de France. Paris, 1: 179-181.
- Achard, J. 1923. Espèces et variétés nouvelles. Fragmenta Entomologica. Prague-Smichov, fasc. 2: 49-126.
- Achard, J. 1924. Les *Phytodecta* et leurs variations. Espèces et variétés nouvelles. Časopis Česke Společnosti Entomologické. Praha, 21: 31-37.
- Acloque, A. 1896. Faune de France Coléoptères. Librairie J.-B. Baillière et Fils. Paris. 466 pp.
- Adams, M. 1817. Descriptio insectorum novorum Imperii Russici, inprimis Caucasi et Sibiriae. Memoire Societa Naturale. Moscou, 5: 278-314.
- Agassiz, L. 1846. Nomenclatoris Zoologici. Index Universalis continens nomina systematica Classium, Ordinum, Familiarum et Generum Animalium omnium, tam viventium quam fossilium, secundum ordinem alphabeticum unicum disposita, adjectis homonymiis plantarum, nec non variis adnotationibus et emendationibus. Soloduri. VII + 393 pp.
- Ahrens, A. 1810. Beiträge zu einer Monographie der Rohrkäfer. Neue Schriften der naturforschenden Gesellschaft zu Halle. Halle, 1: 9-48.
- Ahrens, A. 1812. Beiträge zur Kenntnis deutscher Käfer. Neue Schriften der Naturforschenden Gesellschaft zu Halle. Halle, 2(2): 2-40.
- Allard, E. 1859. Description de quelques Haltises nouvelles. Bulletin de la Société Entomologique de France. Paris, ser. 3, 7: 100-101.
- Allard, E. 1860. Essai monographique sur les Galerucites Anisopodes (Latr.) ou description des Altises d'Europe et des bords de la mer Méditerranée. Annales de la Société Entomologique de France. Paris, ser. 3(8): 370-418.
- Allard, E. 1861. Catalogue complémentaire des diverses espèces d'Altises qui ont été décrites tant dans cet ouvrage, par E. Allard, que par MM. Foudras, Wollaston, Kutschera etc., et qui proviennent d'Europe et du nord l'Afrique (1). Annales de la Société Entomologique de France. Paris, ser. 4(1): 307-348.
- Allard, E. 1863. Description de deux nouvelles espèces d'Altises. Annales de la Société Entomologique de France. Paris, (ser. 4) 3: 659-660.
- Allard, E. 1866. Monographie des Alticides tribu de la famille des Phytophages. L'Abeille, Journal d'entomologie. Paris, 3: 53, 169-417.
- Allard, E. 1867. Tribu des Alticides. In: Joanis L. Monographie des Galerucides d'Europe du Nord de l'Afrique et de l'Asie. L'Abeille, Journal d'entomologie. Paris, 3: 169-508.
- Allard, E. 1875. Répertoire des Coléoptères d'Europe Phytophages: *Alticidae*. L'Abeille, Journal d'entomologie. Paris, 6: 430-443.
- Allard, E. 1876. Descriptions d'espèces nouvelles. L'Abeille, Journal d'entomologie. Paris, 14: 21-25.
- Allard, E. 1878. *Crepidodera nigrivoxis* Allard. In: Schneider und Leder, Beiträge zur Kenntnis der Kaukasischen Käferfauna. Fortsetzung. Verhandlungen des Naturforschenden Vereins in Brünn. Brünn, 17: 1-104.
- Allard, E. 1884. Diagnoses de quatre nouveaux Coléoptères. Wiener Entomologische Zeitung. Wien, 3: 248.
- Allen, A. A. 1967. Two new species of *Longitarsus* Ltr. (*Col.*, *Chrysomelidae*) in Britain. The Entomologist's Monthly Magazine. London, 102(2): 75-82.
- Apfelbeck, V. 1912. Die Komponenten der Balkanfauna aus der Familie der *Chrysomelidae* (*Col.*). Wissenschaftliche Mitteilungen aus Bosnien und der Herzegovina, Landesmuseum Sarajevo. Wien, 13: 354-396.

- Apfelbeck, V. 1912. [Komponente Balkanske faune iz roda *Chrysomelidae* (Col.)]. Glasnik Zemskoga Muzeia Bosnia-Herzegovina. Sarajevo, 24: 235–263.
- Apfelbeck, V. 1914. [Komponente Balkanfauna iz roda *Chrysomelidae* (Col.)]. Glasnik Zemskoga Muzeia Bosnia-Herzegovina. Sarajevo, 26: 435–450.
- Askevold, I. S. 1990. Reconstructed phylogeny and reclassification of the genera *Donaciinae* (Coleoptera: *Chrysomelidae*). *Quaestiones Entomologicae*, Tallahassee, 26: 601–664.
- Aubé, C. 1843. Note sur quelques *Altica confondues* sous le nom d'*Altica oleracea*. *Annales de la Société Entomologique de France*, Paris, 2(1): 5–10.
- Aubé, C. 1859. Communication a la Société Entomologique de France. *Bulletin de la Société Entomologique de France*, Paris, 7: CCXL.
- Bach, M. 1856. Käferfauna für Nord- und Mitteldeutschland mit besonderer Rücksicht auf die Preussischen Rheinlande. III. Band, 5. Lieferung. Coblenz. 142 pp.
- Baertueva, V.A. 1973. Fauna and ecology of flea beetles (Coleoptera, *Chrysomelidae*, *Alticinae*) of Buryatskaja ASSR. In: *Fauna and Ecology of insects of Estern Siberia and Far East*, Irkutsk: 44–56. [in Russian].
- Balbi, E. 1892. Descrizione d'una nuova specie italiana del genere *Hispa* (Linneo). *Bollettino della Societa Romana di Zoologia*. Roma, 1: 195–196.
- Ballion, E. E. 1878. Verzeichnis der im Kreis von Kuldsha gesammelten Käfer. *Bulletin de la Société Impériale des Naturalistes de Moscou*, Moscou, 53(1): 253–389.
- Ballion, E. E. 1890. Einiges aus meinen Notizblättern. *Societas Entomologicae*. Stuttgart, 5 [1889–1890]: 5, 25, 33, 129–130.
- Balthasar, V. 1934. Ein Beitrag zur Kenntnis der Palaearktischen *Donaciini*. *Entomologische Nachrichten und Berichte*. Wien, 8: 128–130.
- Baly, J. S. 1859. Descriptions of new species of phytophagous beetles. *Annals of the Museum of Natural History*, London, 3(3): 195–209.
- Baly, J. S. 1859. Descriptions of new genera and species of phytophagous insects. *Annals of the Museum of Natural History*, London, 3(4): 55–61.
- Baly, J. S. 1859. Descriptions of new genera and species of phytophagous insects. *Annals of the Museum of Natural History*, London, 3(4): 124–128.
- Baly, J.S. 1859. Descriptions of new species of phytophagous insects. *Transactions of the Royal Entomological Society of London*, London, n.s., 5: 146–161.
- Baly, J. S. 1860. An attempt at a classification of the *Eumolpinae*. *Journal of Entomology*, London, 2: 143–163.
- Baly, J. S. 1861. Description of new Genera and Species of *Phytophaga*. *Journal of Entomology*, London, 1: 193–206.
- Baly, J. S. 1862. Description of new genera and species of *Phytophaga*. *Journal of Entomology*, London, 1: 450–459.
- Baly, J. S. 1863. An attempt at a classification of the *Eumolpidae*. *Journal of Entomology*, London, 2: 143–163.
- Baly, J. S. 1864. Description of new genera and species of *Phytophaga*. *The Annals and Magazine of Natural History*, London, 3(14): 433–442.
- Baly, J. S. 1864. Descriptions of uncharacterized genera and species of *Phytophaga*. *Transactions of the Royal Entomological Society of London*, London, ser. 3, 1864: 223–243.
- Baly, J. S. 1864. Descriptions of new genera and species of *Eumolpidae*. *The Journal of Entomology*, London, 2: 219–223.
- Baly, J. S. 1865. *Phytophaga* Malayana. *Transactions of the Entomological Society of London*, London, 4(1): 1–66.
- Baly, J. S. 1873. Catalogue of the Phytophagous *Coleoptera* of Japan with description of the species new to science. *Transactions of the Entomological Society of London*, London, 1: 69–99.
- Baly, J. S. 1874. Catalogue of the Phytophagous *Coleoptera* of Japan, with descriptions of the species new to science. *Transactions of the Royal Entomological Society of London*, London, 2: 161–211.
- Baly, J. S. 1875. [Geänderte Namen]. *Coleopterologische Hefte*, München, 14: 213.

- Baly, J. S. 1876. Descriptions of a new genus and of new species of *Halticinae*. Transactions of the Entomological Society of London. London, 3: 581–602.
- Baly, J. S. 1877. Descriptions of new genera and of uncharacterized species of *Halticinae*. Transactions of Entomological Society of London. London, 4: 157–184, 283–323.
- Baly, J. S. 1877. Descriptions of new genera and species of *Cryptocephalidae*. Transactions of Entomological Society of London. London, 4: 211–232.
- Baly, J. S. 1878. Description of new species and genera of *Eumolpidae*. The Journal of the Linnean Society of London, Zoology. London, 14: 246–265.
- Baly, J. S. 1879. An attempt to point out the differential characters of some closely allied species of *Chrysomela*, principally those contained in Suffriani's 11th group; also descriptions of some hitherto uncharacterized forms belonging to the same and other genera of the family. Transactions of the Royal Entomological Society of London. London, 1879: 179–197.
- Baly, J. S. 1888. Description of some genera and species of *Galerucinae*. The Journal of the Linnean Society of London, Zoology. London, 119(20): 156–188.
- Bargagli, P. 1878. La flora delle Altiche in Europe. *Bullettino della Societa Entomologica Italiana*, 10: 43–153.
- Bauduer, 1874. Séance de 26 Août 1874. *Bulletin de la Société Entomologique de France*. Paris, 22: CLXI–CLXV.
- Bauer, 1929. In: Rüschkamp, F. Zur rheinischen Käferfauna IV. *Entomologische Blätter*. Swabach, Nürnberg, Berlin, Krefeld, 25: 35–43.
- Bechyné, J. 1942. O variabilite *Plateumaris consimilis* Schrank (*Col. Donaciidae*). *Sborník entomologického oddělení Národního muzea v Praze*. Praha, 20: 232–336.
- Bechyné, J. 1944. *Cassididae* a *Hispidae* v Čechách a na Morave. *Entomologické Listy*. Brno, 7(1/4): 74–86.
- Bechyné, J. 1945. Príspevek k poznání našich mandelínek. *Entomologické Listy*. Brno, 8: 5–10.
- Bechyné, J. 1946. Nomenklatorické a faunistické poznámky k některým druhům rodu *Chrysolina* Motsch. (*Col. Phytoph. Chrysolimelidae*). *Folia Entomologica*. Brno, 9: 105–112.
- Bechyné, J. 1946. Příspěvek k poznání evropských druhů rodu *Timarcha* Latr. *Acta Entomologica Musei Nationalis Pragae*. Praha, 24: 129–139.
- Bechyné, J. 1947. Druhý příspěvek k poznání rodu *Chrysolina* Motsch. *Entomologické Listy*. Brno, 10: 58–64.
- Bechyné, J. 1947. Notulae ad cognitionem generis *Timarcha* Latr. VIII. *Acta Entomologica Musei Nationalis Pragae*. Praha, 25: 55–61.
- Bechyné, J. 1947. Poznámky ke studiu o rodu *Timarcha* Latr. *Časopis Česke Společnosti Entomologické*. Praha, 44(1/2): 3–9.
- Bechyné, J. 1947. Příspěvek k poznání rodu *Phytodecta* Kirby. *Sborník entomologického oddělení Národního muzea v Praze*. Praha, 3B: 85–158.
- Bechyné, J. 1948. Notulae ad cognitionem generis *Chrysolina* Motsch. III. *Folia Entomologica*. Brno, 11: 10–15.
- Bechyné, J. 1949. Contribution à la Connaissance du Groupe de *Chrysolina carnifex* (Col.). *Bulletin mensuel de la Société Linnéenne de Lyon*. Lyon, 18(5): 84–86.
- Bechyné, J. 1949. Čtvrtý příspěvek k poznání rodu *Chrysolina* Motsch. *Entomologické Listy*. Brno, 12: 48–55.
- Bechyné, J. 1950. 7 Contribution à la Connaissance du Genre *Chrysolina* Motsch. (*Col. Phytophaga, Chrysolimelidae*). *Entomologische Arbeiten aus dem Museum G. Frey Tutzing Bei München*. München, 1: 47–185.
- Bechyné, J. 1952. Achter Beitrag zur Kenntnis der Gattung *Chrysolina* Motsch. (*Col. Phytophaga, Chrysolimelidae*). *Entomologische Arbeiten aus dem Museum G. Frey Tutzing Bei München*. München, 3(2): 351–385.
- Bechyné, J. 1954. Beiträge zur Kenntnis der echten Chrysolimeliden (*Col. Phytophaga*). *Entomologische Arbeiten aus dem Museum G. Frey Tutzing Bei München*. München, 5(2): 581–674.
- Bechyné, J. 1955. Contributions a l'étude de la faune entomologique du Ruanda-Urundi (Mission P. Basilewsky

- 1953). LXIV. Coleoptera *Chrysomelidae* Chrysomelinae et Alticinae. Annales du Musée Royal du Congo Belge. Tervuren (ser. 8). Sci. zool., 40: 204–230.
- Bechyné, J. 1955. Notes sur les *Chrysolini* (Coleopt. Chrysom.) de la Méditerranée Orientale. Bulletin of the Research Council of Israel. Tel-Aviv, 4(4): 347–350.
- Bechyné, J. 1956. Über die Alticipiden-Sammlung Heikertinger (Col. *Phytophaga*). Entomologische Arbeiten aus dem Museum G. Frey Tutzing Bei München. München, 7: 577–598.
- Bechyné, J. 1957. Notes sur Chrysomeloidea Palerctiques recueillis par M. G. Fagel (Coleoptera *Phytophaga*). Bulletin de l'Institut royal des sciences naturelles de Belgique. Bruxelles, 33(31): 1–4.
- Bechyné, J. 1958. Über die taxonomische Valenz der Namen von *Oreina* s.str. (Col. *Phytophaga*). Mitteilungen der Schweizerischen Entomologischen Gesellschaft. Bern-Lausanne, 31(1): 79–95.
- Bechyné, J. 1962. Notes sur quelques Chrysomélides vrais (Coléoptères). Bulletin mensuel de la Société Linnéenne de Lyon. Lyon, 31(6): 182–185.
- Beck, L. 1817. Beiträge zur bayerischen Insectenfauna, oder Beschreibung und Abbildung neuentdeckter Käfer mit angehangtem Namenverzeichnis der Eleuteraten des Landgerichtbezirks Zusmenhausen. Augsburg, 45 pp.
- Becker, A. 1880. Beiträge zu meinen Verzeichnissen der um Sarepta und am Bogdo vorkommenden flanzten und Insecten, und Beschreibung einer *Mylabis*-larve. Bulletin de la Société Impériale des Naturalistes de Moscou. Moscou, 55(1): 145–156.
- Bedel, L. 1869. Note sur l'*Haemonia mosellae*. Annales de la Société Entomologique de France. Paris, ser. 4, 9: 4.
- Bedel, L. 1889–1901. Faune des Coléoptères du Bassin de la Seine. Annales de la Société Entomologique de France. Paris, 3: 423 pp.
- Bedel, L. 1891. Faune des Coléoptères du Bassin de la Seine. T. V. *Phytophaga*, [fasc. 3]. Annales de la Société Entomologique de France. Paris, 60. Hors série: 105–136.
- Bedel, L. 1892. Faune des Coléoptères du Bassin de la Seine. T. V. *Phytophaga*, [fasc. 4]. Annales de la Société Entomologique de France. Paris, 61. Hors série: 137–160.
- Bedel, L. 1899. Faune des Coléoptères du Bassin de la Seine. T. V. *Phytophaga*, [fasc. 7]. Annales de la Société Entomologique de France. Paris, 67. Hors série: 229–260.
- Bellevoys, A. 1870. Moeurs des *Haemonia*. L'Abeille, Journal d'entomologie. Paris, 7: 55–56.
- Bellier de la Chavignerie, J. B. M. 1870. Les descriptions qui suivent de Coleopteres nouveaux propres aux Basses-Alpes. Annales de la Société Entomologique de France. Paris, 4e ser. 10(27): XXVI–XXIV.
- Berthold, A. A. 1827. Latreille's naturliche familien des thierreichs. Aus dem frazosischen. Mit anmerkungen und zaszten. Weimer, 606 pp.
- Berti, N. and M. Rapilly. 1979. Contribution a la faune d'Iran 4 – *Cryptocephalinae* (Coleopteres: *Chrysomelidae*). Nouvelle Revue d'entomologie. Toulouse, 9(3): 239–269.
- Bielz, A. 1852. Entomologische Notizen. Verhandlungen und Mittheilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt. Hermannstadt, 3: 13–16.
- Bienkowski, A. O. 1997. New distributional records for several palaeartic *Chrysomelidae* species with some systematic remarks (*Insecta: Coleoptera*). Faunistische Abhandlungen, Dresden, 21(4): 91–104.
- Bienkowski, A. O. 1997. Some surprising discoveries of *Chrysolina relicens* (Coleoptera, *Chrysomelidae*) on the White Sea shore, in Siberia and in the Far East. Entomologica Fennica. Helsinki, 7: 195–199.
- Bienkowski, A. O. 1998. Morphology and systematics of larvae of some *Chrysolina* species (*Insecta: Coleoptera: Chrysomelidae*). Entomologische Abhandlungen. Dresden, 58(4): 73–82.
- Bienkowski, A. O. 1998. Revision of the subgenus *Anopachys* Motschulsky, 1860 of the genus *Chrysolina* Motschulsky, 1869 (Coleoptera: *Chrysomelidae: Chrysomelinae*). Genus. Wrocław, 9(2): 95–153.
- Bienkowski, A. O. 1999. Key to leaf beetles (Coleoptera *Chrysomelidae*) of the European part of Russia and adjacent countries. Moscow, 204 pp.
- Billberg, G. J. 1820. Enumeratio insectorum in museo Billberg. Stockholm. 138 pp.
- Biondi, M. 1989. Classification and phylogenetic of the Western Palearctic species of the genus *Asiolestia* Jacobson (Coleoptera, *Chrysomelidae, Alticinae*). Entomography. Sacramento, 6: 519–529.
- Biondi, M. 1995. The *Longitarsus anchusae* complex in Near Eastern and description of a new species (Coleoptera, *Chrysomelidae, Alticinae*). Nouvelle Revue d'entomologie. Toulouse, 12(4): 259–271.

- Blackburn, T. 1890. Further note on Australian *Coleoptera*. Transaction of Royal Society of south Australia. Adelaidae, 26: 113–169.
- Blatchley, W.S. 1921. Notes on Indiana *Halticini* with characterization of a new genus and descriptions of new species. Journal of the New York Entomological Society. New York, 29: 16–27.
- Bodor, J. 1965. Négy új *Crioceris* Fourc.változat (*Coleoptera*, *Chrysomelidae*). Rovartani Közlemenyek. Budapest, 18(1): 300–303.
- Bogatchev, A. 1947. Quelques especes nouvelles des Coleopteres de l'Iran. Entomologica et Phytopathologica applicata. Teheran, 4: 15–18.
- Boheman, C. H. 1850. Monographia Cassididarum. I. Holmiae, 452 pp.
- Boheman, C. H. 1851. Arsberättelse om Framstegen i Insekternas, Myriapodernas och Arachnidernas Naturalhistoria för 1847 och 1848. Vtensk. Akademia Stockholm, Norstedt. 333 pp.
- Boheman, C. H. 1854. Monographia Cassididarum. II. Holmiae. 506 pp.
- Boheman, C. H. 1855. Monographia Cassididarum. III. Holmiae. 543 pp.
- Boheman, C. H. 1856. Catalogue of Coleopterous Insects in the collection of the British Museum, Part IX, *Cassididae*. London. 225 pp.
- Boheman, C. H. 1862. Monographia Cassididarum. IV. Holmiae. 604 pp.
- Boheman, C.H. 1851. Arsberättelse om Framstegen i Insekternas, Myriapodernas och Arachnidernas Naturalhistoria för 1847 och 1848. Vtensk. Akademia Stockholm. Norstedt. 333 pp.
- Boieldieu, A. 1859. Description d'espèces nouvelles de Coléoptères. Annales de la Société Entomologique de France. Paris, 5: 461–482.
- Bonelli, F.A. 1812. Specimen faunae subalpiniae sistens *Insecta* Pedemontn hucusque inedita etc Fasciculus primus, *Coleoptera* plerumque comprehendens. Mem. Roy. Soc. Agrar. Torino, 9: 149–183.
- Bontems, C. 1981. Les espèces de Linné et Fabricius du genre *Oreina* Chevrolat, 1837 (*Col.*, *Chrysomelidae*, *Chrysomelinae*). Nouvelle Revue d'entomologie. Toulouse, 11(1): 93–109.
- Borowiec L. 1978. Nowe dane o susowkach (*Coleoptera*, *Chrysomelidae*, *Halticinae*) Polski ze szczególnym uwzględnieniem Śląska. Polskie Pismo Entomologiczne. Wrocław, 48: 423–427.
- Borowiec L. 1979. *Chrysomelidae* (*Coleoptera*) new to the Bulgarian fauna. Polskie Pismo Entomologiczne. Wrocław, 49: 481–483.
- Borowiec L. 1999. A world catalogue of the Cassidinae (*Coleoptera*: *Chrysomelidae*). Wrocław, 476 p.
- Borowiec, L., Świętojańska J. 2001. The Palaearctic species of the genus *Rhytidocassis* Spaeth, 1941 (*Coleoptera*: *Chrysomelidae*: *Cassidinae*). Annales Zoologici. Warszawa, 51: 325–329.
- Böving, A.G., Craighead, F.C. 1931. An illustrated synopsis of the principal larval forma of the order *Coleoptera*. Entomologica Americana. Lancaster, n. ser., 11(1930): 1–351, 125 pls.
- Brahm, N. J. 1790. Insektenkalender für Sammler und Oekonomen. Erster Teil. Mainz. 258 pp.
- Brancsik, K. 1899. Aliquot *Coleoptera* nova Russiae asiaticae. Trencsenvarmegyei Muzeum-Egysület, Ertesitoje. Trencsen, 21–22: 97–105.
- Brancsik, C. 1910. Species et varietates novae Coleopterorum exoticorum ac palaearticorum. Trencsenvarmegyei Muzeum-Egysület, Ertesitoje. Trencsen, 31–33 (1908–1910): 176–190.
- Brancsik C., 1914. *Coleoptera* nova. Trencsenvarmegyei Muzeum–Egysület, Ertesitoje. Trencsen. 61 pp.
- Brassler, K. 1926. Eine Mutation von *Melasoma vigintipunctata* L. Entomologisches Jahrbuch. Leipzig, 35: 111.
- Breit, J. 1918. Beitrag zur Kenntnis der Arten des Genus *Cryptocephalus* Geoffr. Wiener Entomologische Zeitung. Wien. 37. (1/3): 35–52.
- Breit, J. 1918. Eine neue *Stylosomus*-Art nebst Übersicht der bekannten mittelasiatischen Arten. Koleopterologische Rundschau. Wien, 7: 118–119.
- Breit, J. 1920. *Chrysomela* Studien II. Koleopterologische Rundschau. Wien, (1919–1920), 8: 81–88.
- Breit, J. 1921. Die Mittelasiatischen Pachybrachys-Arten aus der Verwandtschaft des *P. glycerrhizae* Ol. und *vermicularis* Suffr. Entomologische Blätter. Berlin, 17: 64–74.
- Brisout de Barneville, C. 1860. Description de quelques Coléoptères nouveau la Faune Française et spécialement aux environs de Paris. Annales de la Société Entomologique de France. Paris, sér. 3, 6: 339–350.
- Brisout de Barneville, C. 1873. [Über die synonymie von *Longitarsus atricillus*, *L. piciceps* und *L. senecionis*]. Bulletin de la Société Entomologique de France. Paris, 22: CCXVIII.

- Broun, T. 1893. Manual of the New Zealand *Coleoptera*. Wellington, 6. 1392 pp.
- Brovdiv, V. M. 1977. Leaf Beetles, *Chrysomelinae*. T. 19, vyp. 16. Fauna Ukrainy. Naukova Dumka, Kiev. 385 pp. [in Russian].
- Brovdiv, V. M. 1983. Leaf Beetles, *Cassidinae* and *Hispinae*. T. 19, vyp. 20. Fauna Ukrainy. Naukova Dumka, Kiev. 187 pp. [in Russian].
- Brüggemann, F. 1873. Systematisches Verzeichnis der bisher in der Gegend von Bremen gefundenen Käferarten. Abhandlungen der Naturforschenden Vereins in Bremen, Bremen, 3 441–524.
- Brullé, A. 1832. Expédition scientifique de Morée. III, 1, Paris. 400 pp.
- Brullé, A. 1834. Histoire naturelle des Insectes (Coléoptères, Orthoptères et Hémiptères), vol. 5, Paris. 436 pp.
- Bryant, G. E. 1925. New species of *Phytophaga* from the Fiji Islands. The Annals and Magazine of Natural History, London, 9(15): 591–599.
- Bryant, G. E. 1957. *Coleoptera: Chrysomelidae* of south-west Arabia. The Annals and Magazine of Natural History, London, (12)10: 353–363.
- Buddeberg, C. 1888. Beobachtungen über Lebensweise und Entwicklungsgeschichte einheimischer Käferarten. Jahrbücher des Nassauischen Vereins für Naturkunde, Wiesbaden, 41: 20–43.
- Bullock, E. 1928. *Coleoptera* from the Killarney district of County Kerry Ireland. The Entomologist's Monthly Magazine, London, 64, 5: 102–104.
- Burlini, M. 1942. Forme nuove di Coleotteri. Bollettino della Societa Entomologica Italiana, Genová, 74(1): 13.
- Burlini, M. 1946. Nuove forme di Coleotteri italiani. Bollettino della Societa Entomologica Italiana, Genová, 76(9/10): 78–79.
- Burlini, M. 1948. Nuove specie di *Cryptocephalus* italiani (*Coleoptera, Chrysomelidae*). Memorie del Museo Civico di Storia Naturale di Verona, Verona, 1: 221–229.
- Burlini, M. 1955. Revisione dei *Cryptocephalus* italiani e della maggior parte delle specie di Europa (*Col., Chrysomelidae*). Memorie della Societa Entomologica Italiana, Genová, 34: 5–287.
- Burlini, M. 1957. Un nuovo *Stylosomus* asiatico. Bollettino della Societa Entomologica Italiana, Genová, 87(3–4): 62.
- Burlini, M. 1961. Contributo alla conoscenza dei *Cryptocephalus* (*Col., Chrysomelidae*). Bollettino della Societa Entomologica Italiana, Genová, 91(5/6): 94–96.
- Burlini, M. 1962. Nuove forme di *Cryptocephalus* paleartici (*Coleoptera, Chrysomelidae*) appartenenti al Museo Nazionale di Storia Naturale di Budapest. Annales Historico-Naturales Musei Nationalis Hungarici. Pars Zoologica, Budapest, 54: 327–328.
- Burlini, M. 1967. Addenda alla revisione dei *Cryptocephalus*. XXI Contributo alla conoscenza dei *Cryptocephalini* (*Coleopt.*). Bollettino della Societa Entomologica Italiana, Genová, 97(7/8): 129–132.
- Burlini, M. 1968. Revisione delle specie italiane e della maggior parte delle specie europee del genere *Pachybrachis* Chev. (*Coleoptera, Chrysomelidae*). XXII Contributo alla conoscenza dei *Cryptocephalini*. Memorie della Societa Entomologica Italiana, Genová, 47: 11–116.
- Burlini, M. 1969. *Cryptocephalus fulvus* Goeze subsp. *schatzmayri* nova. XXIII Contributo alla conoscenza dei *Cryptocephalini* (*Coleoptera, Chrysomelidae*). Bollettino della Societa Entomologica Italiana, Genová, 99–101(5/6): 114–115.
- Burlini, M. 1970. Una nuova sottospecie nord africana del *Cryptocephalus pygmaeus* F. e note sul *Cryptocephalus* (*Cerodens* Burl.) *emiliae* Burl. XXV Contributo alla conoscenza dei *Cryptocephalini* (*Coleoptera, Chrysomelidae*). Bollettino della Societa Entomologica Italiana, Genová, 102(1/2): 38–40.
- Buysson, H. 1908. Recherches sur l'epoque d'eclosion des Altises et sur la biologie et les caracteres de quelques *Thyamis* Steph. de la faune de France (*Col.*). Miscellanea Entomologica. Revue entomologique internationale. Organe de l'Union Entomologique. Narbonne, Toulouse, Paris, 15: 1–39.
- Buysson, H. 1924. *Chrysomela menthastri* Suffr. Varietates novae. Miscellanea Entomologica. Revue entomologique internationale. Organe de l'Union Entomologique. Narbonne, Toulouse, Paris, 27(12): 90.
- Caillol, H. 1908. Catalogue des Coléoptères de Provence (1<sup>re</sup> partie). Annales de la Société des Sciences Naturelles de Provence. Marseille, 1: 73–584.

- Caillol, H. 1914. Catalogue des Coléoptères de Provence, 3<sup>e</sup> partie. Mémoires de la Société Linnéenne de Provence. Marseille. 596 pp.
- Capra, F. 1933. Note sugli *Sphaeroderma* Steph. (*Col.*, *Chrysom.*). Bollettino della Società Entomologica Italiana. Genovà, 65(7): 164–165.
- Čepelak, R. 1927. Nové aberrace *Osphya bipunctata* Fabr. Časopis Česke Společnosti Entomologické. Praha, 24(5/6): 123.
- Chapuis, F. 1874. Histoire naturelle des Insectes. Genera des Coléoptères ou exposé méthodique et critique de tous les genres proposés jusqu'ici dans cet ordre d'insectes. Par MM. Th. Lacordaire et F. Chapuis. Tome dixième. Famille des Phytophages. Paris, IV + 455 pp., pls. 111–124.
- Chapuis, F. 1875. Histoire naturelle des Insectes. Genera des Coléoptères ou exposé méthodique et critique de tous les genres proposés jusqu'ici dans cet ordre d'insectes. Par MM. Th. Lacordaire et F. Chapuis. Tome onzième. Famille des Phytophages. Paris, 420 pp., pls. 125–130.
- Chapuis, F., 1880. Phytophages Abyssiniens du Musée Civique d'Histoire Naturelle de Genes. Annali del Museo Civico di Storia Naturale di Genova. Genova, 15: 5–31.
- Charpentier, T. 1825. Horae entomologicae. (*Neuroptera*, *Orthoptera*, *Coleoptera*). Wratislaviae. XVI+255 pp.
- Chen, S. 1933. *Chrysomelidae* (*Coleoptera*) nouveaux de l'Asie tropicale, 2<sup>e</sup> note. Bulletin du Muséum National d'histoire naturelle. Paris, 2<sup>e</sup> serie(6): 449–456.
- Chen, S. H. 1933. Study of the *Halticinae* beetles with descriptions of some exotic new species. Sinensia, 3: 211–254.
- Chen, S. H. 1934. On some species of *Chrysomelidae* in the British Museum. Stylops. London, 3: 66–78.
- Chen, S. H. 1934. Revision of the *Halticinae* (*Col. Chrysomelidae*) of Yunnan and Tonkin. Sinensia. Nanking, 5(3/4): 225–416.
- Chen, S. H. 1935. Study of Chinese Eumolpid beetles. Sinensia. Nanking, 6(3): 221–387.
- Chen, S. H. 1936. Catalogue des *Chrysomelinae* de la Chine, de l'Indochine et du Japon. Notes d'entomologie Chinoise, Shanghai, 3(5): 63–102.
- Chen, S. H. 1936. Genera of Oriental *Halticinae*. Sinensia, 7(6): 625–667.
- Chen, S. H. 1938. *Chrysomelinae* et *Halticinae* recoltés par M. K. Kolthoff dans la Province du Kangsu. Arkiv för Zoologie. Stokholm, 30(4): 1–4.
- Chen, S. H. 1939. Flea beetles collected at Kwangsi. Sinensia. Nanking, 10(1–6): 21–55.
- Chen, S. H. 1939. New genera and species of Chinese *Halticinae*. Sinensia. Nanking, 10(1–6): 56–91.
- Chen, S. H. 1940. Attempt a new classification of the Leaf Beetles. Sinensia. Nanking, 11(5/6): 451–481.
- Chen, S. H. 1940. Notes on Chinese *Eumolpidae*. Sinensia, Nanking, 11(5/6): 483–528.
- Chen, S. H. 1940. On the *Coleoptera Chlamydinae* of China. Sinensia. Nanking, 11(3/4): 189–206.
- Chen, S. H. 1941. New leaf-beetles from China. Sinensia. Nanking, 12: 189–198.
- Chen, S. H. 1941. Notes on donaciine beetles. Sinensia. Nanking, 12: 1–17.
- Chen, S. H. 1942. *Galerucinae* nouveaux de la faune chinoise. Notes d'entomologie Chinoise. Shanghai, 9(3): 9–67.
- Chen, S. H. 1961. New species of Chinese *Chrysomelidae*. Acta Entomologica Sinica. Beijing, 10(4–6): 429–435.
- Chen, S. H. and Y. T. Kung, 1954. Notes on Chinese *Halticinae* of the genus *Luperomorpha* Weise. Acta Entomologica Sinica. Beijing, 44(1): 84–97.
- Chevrolat, L. A. A. 1836. See: Dejean, P. F. 1837.
- Chevrolat, A. 1840. Description de quelques Coléoptères de la Calice et du Portugal provenant d'envois de M. Deyrolles fils. Revue Zoologique, Paris, 3: 8–18.
- Chevrolat, L. A. A. 1844. In: Guérin-Ménéville, F. E. 1844. Iconographie du regne animal de G. Cuvier, ou representation d'après nature de l'une des especes les plus remarquables et souvent non encore figurees de chaque genre d'animaux, pouvant servir d'atlas à tous les traités de zoologie. 7. Insectes, avec le portrait de Latreille. Paris. 576 pp.
- Chevrolat, A. 1845. In: d'Orbigny C. etc. Dictionnaire universel d'histoire naturelle... etc., Paris, tome quatrième. 1839–1845, vol. 8.
- Chittenden, F.H. 1927. The species of *Phyllotreta* north of Mexico. Entomologica Americana. (N.S.). Lancaster, Brooklyn, N.Y., 8(1): 1–63.

- Chûjô, M. 1935. Studies on the *Chrysomelidae* in the Japanese Empire (VIII). Subfamily *Halticinae* (3). Transactions of the Natural History Society of Formosa. Taihoku, 25: 459–476.
- Chûjô, M. 1936. Studies on the *Chrysomelidae* in the Japanese Empire (VIII). Subfamily *Halticinae* (6). Transactions of the Natural History Society of Formosa. Taihoku, 25: 108–114.
- Chûjô, M. 1938. Beitrag zur Kenntnis der Chrysomeliden-Fauna Chinas (*Coleoptera*). Mushi. Fukuoka, 11(2): 158–169.
- Chûjô, M. 1940. Chrysomelid-beetles from Borodino Islands, Loo-choo (Daitojima), collected by Mr. M. Yanagiwara. Transactions of the Natural History Society of Formosa. Taihoku, 30: 363–365.
- Chûjô, M. 1940a. Chrysomelid beetles from Korea (I). Transactions of the Natural History Society of Formosa. Taihoku, 30: 349–362.
- Chûjô, M. 1940b. Chrysomelid beetles from Korea (II). Transactions of the Natural History Society of Formosa. Taihoku, 30: 383–398.
- Chûjô, M. 1941. Chrysomelid beetles from Korea (IV). Transactions of the Natural History Society of Formosa. Taihoku, 23: 155–174.
- Chûjô, M. 1942. First supplement to the fauna of Korean chrysomelid-beetles (II.). Transactions of the Natural History Society of Formosa. Taihoku, 32: 31–43.
- Chûjô, M. 1949. On the *Hebdomecosta* Spaeth with description of a new species from Japan (*Coleoptera* – *Chrysomelidae* – *Cassidinae*). Transactions of the Kansai Entomological Society. Kyoto, 14(2): 7–13.
- Chûjô, M. 1950. Description of a new Chrysomelid-beetles from Japan. Kontyu. Tokio, 18(5): 29–31.
- Chûjô, M. 1951. Chrysomelid beetles of Shikoku, Japan (II) (*Coleoptera*). Transactions of the Shikoku Entomological Society. Matsuyama, 2(3): 40–47.
- Chûjô, M. 1952. A taxonomic study on the *Chrysomelidae* (*Insecta-Coleoptera*) from Formosa. Bulletin of Kagawa Agricultural College of Technical. Takamatsu, 3 (3): 166–183.
- Chûjô, M. 1954. Chrysomelid-beetles from Shikoku, Japan. Transactions of the Shikoku Entomological Society. Matsuyama, 4(4): 51–62.
- Chûjô, M. 1959. Chrysomelid-beetles of Niigara Prefecture, Japan, chiefly collected by Dr. K. Baba (II). Journal of the Nagoaka Municipal Museum of Natural History. Nagoaka, 3: 13–22.
- Chûjô, M. 1961. 24. *Chrysomelidae*. Nature and Life in Southeast Asia, 1: 347–357
- Chûjô, M. 1962. A taxonomic study on the *Chrysomelidae* (*Insecta-Coleoptera*) from Formosa. Part XI. Subfamily *Galerucinae*. Philippine Journal of Science. Manila, 91: 1–239+I–IV.
- Chûjô, M. 1963. Chrysomelid-beetles from Formosa (Taiwan) collected by Dr. K. Baba in 1962. Niponius. Kanagawa, 2: 21–30.
- Chûjô, M., and S. Kimoto. 1960. Descriptions of three new genera and a new species of Chrysomelid-beetles from Japan, with some notes on the Japanese species. Niponius. Kanagawa, 1, 4: 1–10.
- Chûjô, M. and S. Kimoto. 1961. Systematic catalogue of Japanese *Chrysomelidae* (*Coleoptera*). Pac. Ins. Mon., Honolulu, 3(2): 117–202.
- Clark H. 1865. Description of species of *Phytophaga* received from Pulo Penang or it's neighbourhood.. The Annals and Magazine of Natural History. London, 15(1): 139–148.
- Clavareau H. 1913. *Chrysomelidae*. See: W. Junk und S. Schenkling. Coleopterorum Catalogus. Berlin, T. 24, pars 53. 278 pp.
- Clavareau H. 1914. *Chrysomelidae*. II *Eumolpinae*. See: W. Junk und S. Schenkling. Coleopterorum Catalogus. Berlin, pars 59. 214 pp.
- Comolli. A. 1837. De Coleopteris novis ac rarioribus minusve cognitis Provinciae Novocomi. Ticini Regii. 54 pp.
- Cornelius. C. 1850. Zur Ernährungs- und Entwicklungsgeschichte der Blattkäfer. Entomologische Zeitung. Stettin, 11( 1): 19–21.
- Cornelius. C. 1851. Zur Entwicklungs- und Ernährungs-Geschichte einiger Schildkäferarten. Entomologische Zeitung. Stettin, 12(1): 88–94.
- Cornelius. [C.] 1857. Ernährung und Entwicklung einiger Blattkäfer. Entomologische Zeitung. Stettin, 18(2): 162–171.
- Cornelius. [C.] 1857. Ernährung und Entwicklung einiger Blattkäfer. (Forts.). Entomologische Zeitung. Stettin, 18(4): 392–405.



- Costa, O. G. 1838. Fauna Vesuviana, ossia descrizione degli Insetti che vivono ne' Fumajoli del Cratere del Vesuvio. Atti della Reale Accademia delle Scienze, Sezione della Società Reale Borbonica, Napoli, 4: 21-53.
- Creutzer, C. 1799. Entomologische Versuche. Wien, 142 pp.
- Crotch, G. R. 1871. A catalogue of British *Coleoptera*. 2. Aufl. 8. Cambridge. 17 pp.
- Crotch, G. R. 1873. Materials for the study of *Phytophaga*. Proceedings of the Academy of Natural Sciences of Philadelphia. Philadelphia, 25: 19-83.
- Csiki, E. 1899. Hazánk *Orsodacne*-féléi. Rovartani Lapok. Budapest, 6: 92-95.
- Csiki, E. 1901. Coleopteren. In: Zichy E. Zoologische Ergebnisse der dritte Asiatische Forschungsreise. II. Budapest-Leipzig. 75-120.
- Csiki, E. 1901. Coleopterologiai jegyzetek. Rovartani Lapok. Budapest, 8(4): 99-103.
- Csiki, E. 1940. See: Heikertinger et Csiki, 1940.
- Csiki, E. 1953. Über neue und bekannte Coleopteren aus Ungarn und den angrenzenden Ländern. Annales Historico-naturales musei nationalis Hungarici (series nova). Budapest. 3: 115-135.
- Curtis, J. 1826. British entomology, being illustrations and descriptions of the genera of Insects found in Great Britain and Ireland. III. London, tt.: 99-146.
- Curtis, J. 1830. British entomology, being illustrations and descriptions of the genera of Insects found in Great Britain and Ireland. VII. London, tt.: 290-337.
- Curtis, J. 1831. British entomology, being illustrations and descriptions of the genera of Insects found in Great Britain and Ireland. VIII. London, tt.: 338-383.
- Curtis, J. 1833. The natural history of British insects. VII. London, 323. 435 pp.
- Curtis, J. 1837. British entomology; being illustrations and descriptions of the genera of insects found in Great Britain and Ireland: containing coloured figures from nature of the most rare and beautiful species, and in many instances of the plants upon which they are found. London, 14: 626-673.
- Czwalina G. 1885. Sammelbericht aus Ost- und Westpreußen. Deutsche Entomologische Zeitschrift. Berlin, 29: 251-256.
- Daccordi, M. 1977. Coleotteri Crisomelidi dell'Isola di Creta e descrizione di una nuova specie del genere *Pachnephorus* Redt. (*Coleoptera*, *Chrysomelidae*). Bollettino del Museo Civico di Storia Naturale. Verona, 4: 81-93.
- Daccordi, M. 1982. Nuovi taxi di *Chrysomelinae* Afrotropicali. Bollettino del Museo Civico di Storia Naturale. Verona, 7: 181-195.
- Dahl, G. 1823. *Coleoptera* und *Lepidoptera*; ein systematisches Verzeichniss mit beigesetzten Preisen der Vorräthe. Wien, 105 pp.
- Dalman, J. 1823. Analecta entomologica. Homiae. VII+104+4 pp.
- Daniel, J. 1903. Beiträge zur Koleopteren-Geographie. Münchener Koleopterologische Zeitschrift. München, 1(1902-1903), 2: 255-260.
- Daniel, J. 1904. Revision der palaearctischen *Crepidodera*-Arten. Münchener Koleopterologische Zeitschrift. München, 3: 237-297.
- Degeer, C. 1775. Memoires pour servir a l'histoire des insectes. V. Stockholm. 448 pp.
- Degeer, C. 1778. Mémoires pour servir a l'histoire des Insectes. VII. Stockholm. 6 + 950 pp.
- Degeer, C. 1783. Genera et Species Insectorum e generosissimi auctoris scriptis, extraxit, digessit, latine quoad partem reddidit, et terminologiam insectorum Linneanam addidit. Lipsiae. 220 pp.
- Dejean, P. F. M. A. 1821. Catalogue de la collection de Coléoptères de M. le baron Dejean. Crevot, Paris. 8. 136 pp.
- Dejean, P. F. M. A. 1837. Catalogue des Coléoptères de la Collection de M. le Comte Dejean. Troisième édition, revue, corrigée et augmentée. Mequignon-Marvis Père et Fils, Paris. XIV+503 pp.
- Delahon, P. 1913. Nachtrage zu „Schilsky's Systematischen Verzeichnis der Käfer Deutschlands" von 1909 mit besonderer Berücksichtigung der Formen der Mark Brandenburg. Deutsche Entomologische Zeitschrift. Berlin, 5: 528-539.
- Delahon, P. 1922. In: Everts, E. J. G. *Coleoptera* Neerlandica. II. s'Gravenhage. IV. 796 pp.
- Delherm de Larcenne, [E.]. 1885. Description de deux nouveaux coléoptères francias. Le Naturaliste. Paris, 7: 119-120.

- Della Beffa, G. 1909. Note Coleopterologiche. Rivista di coleotteri della italiana. Camerino, 7(8/11): 192–201.
- Demaison, C. 1896. Diagnoses de quelques espèces de Chrysomélides et note sur la synonymies de deux Clytrides [*Col.*]. Bulletin de la Société Entomologique de France. Paris : 12–13.
- Demaison, C. 1902. Description de trois Coléoptères phytophages nouveaux d'Europe et d'Égypte. Bulletin de la Société Entomologique de France. Paris, 7: 24–25.
- Demaison, C. 1903. Note sur une variété de *Podagrica* (*Col.*). Bulletin de la Société Entomologique de France. Paris. 8.
- Demaison, C. 1904. Description d'une variété et d'une espèce nouvelle d'Asie Mineure. Bulletin de la Société Entomologique de France. Paris, 9: 286–287.
- Depoli, G. 1926. Su alcuni coleotteri della Carnia. Bollettino della Societa Entomologica Italiana. Genová, 58(1): 14–15.
- Depoli, G. 1940. I Coleotteri della Liburnia. Supplemente alla parte VI. Rivista di Fiume, La Societa di Studi fiumani. Fiume, 15/16: 303–338.
- Depoli, G., Goidanich, A. 1926. I Coleotteri della Liburnia, pars VI, *Phytophaga*. Rivista di Fiume, La Societa di Studi fiumani. Fiume, [1]: 62–113.
- Des Gozis, M. P. 1886. Recherche de l'espece typique de quelques anciennes genres. Rectification synonymique et notes diverses. Montiuçon. 36 pp.
- Desbrochers des Loges, J. 1891. Monographie des *Cassida* de France suivie d observations sur le classement des especes de ce genre au Catalogus Coleopterorum Europae, Caucasi et Armeniae rossicae. Frelon, [1891–1892]. 1, 7: 1–48.
- Dirimanov, M. and R. Angelova. 1964. Über die Fauna der Erdflöhe (*Coleoptera*, *Halticinae*). Wissenschaften. Arbeiten der Landwirtsch. Hochschule „W. Kolarow“, Plovdiv, Fak. Weisn- und Gartenbau. Plovdiv, 13(2): 217–220.
- Döberl, M. 1994. 11. Unterfamilie: *Alticinae* (anstelle 10. U. F.). 92–142. In: Lohse, G. A. et W. H. Lucht (eds.) Die Käfer Mitteleuropas 3. Supplementband mit Katalogteil. „Goecke et Evers“. Krefeld. 403 pp.
- Döberl, M. 1994. Eine neue Alticine von Kaspischen Meer: *Batophila dogueti* nov. spec. (*Coleoptera*, *Chrysomelidae*, *Alticinae*). Nouvelle Revue d'Entomologie. Toulouse, 11(2): 149–150
- Doguet, S. 1973. Notes sur les Altises d'Algerie (2<sup>e</sup> note). Contribution a l'etude du genre *Longitarsus*. Bulletin de la Société Entomologique de France. Paris, 78: 110–113.
- Doguet, S. 1976. Notes systematiques et faunistiques sur quelques Altises palearctiques. Nouvelle Revue d'entomologie. Toulouse, 6(3): 283–290.
- Doguet, S. 1979. Notes systematiques et ecologiques sur divers *Chrysomelidae* palearctiques. Description de deux especes nouvelles. l'Entomologiste. Revue d'amateurs. Paris, 35(2): 49–55.
- Doguet, S. 1994. Coléoptères *Chrysomelidae*. Volume 2 *Alticinae*. Faune de France 80. Fédération Francaise des Sociétés naturelles. Paris. 697 pp.
- Donckier de Donceel H. 1899. Catalogue systématique des Hispidés. Annales de la Société Entomologique de France. Paris, 68: 540–615.
- Donisthorpe, H. 1929. *Longitarsus nasturtii* F. v. *obscurus* n. var., a variety of *Halticinae* (*Coleoptera*) new to science. Entomologist's Record and Journal of Variation. London, 41(9): 126.
- Donisthorpe, H. 1944. *Aphthona aeneomicans* Allard (*Col. Halticinae*), a species of *Coleoptera* new to Britain. The Entomologist's Monthly Magazine. London, 80(11): 259.
- Donovan, E. 1792–1813. The Natural History of British Insects explaining them in their several states, with the periods of their transformations, their food, oeconomy, etc. London, T. 1–16.
- Dorhn, C.A. 1884. Rosenberg 3. Stettiner Entomologische Zeitung. Stettin, 45(1–3): 84–85.
- Drapiez, [A.] 1819. Description de huit espèces d'insectes nouveaux. Ann. gén. Sci. phys. Bruxelles, 1: 290–298.
- Drapiez, [A.] 1819. Description de huit espèces d'insectes nouveaux. Ann. gén. Sci. phys. Bruxelles, 2: 42–50; 197–204.
- Dudich, E. 1924. A *Cryptocephalus gamma* H.-Schäff. változatai (*Coleopt.*, *Chrysomelidae*). Folia entomologica Hungarica. Budapest, 1, 2, 1934 (3): 22–24.
- Dufour, L. 1843. Excursion entomologique dans les montagnes de la vallée d'Ossan (Catalogue de 767 espèces de Coléoptères) Pau, 118 pp.

- Dufour, L. 1851. Des zones entomologiques dans nos Pyrénées occidentales. Acta de la Société Linnéenne de Bordeaux. Bordeaux, 17: 303–364.
- Dufschmidt, C. E. 1825. Fauna Austriae. Oder Beschreibung österreichischen Insecten, für angehende Freunde der Entomologie. Linz, 3: 289 pp.
- Dumeril, A. M. C. 1816. Dictionnaire des sciences naturelles. 1. 524 pp.
- Duvivier, A. 1885. Quatre Phytophages nouveaux. Annales de la Société Entomologique Belgique. Bruxelles, 29 (C.R.): 114–119.
- Duvivier, A. 1892. Les Phytophages du Chota-Nagpore. 2 Note. Annales de la Société Entomologique de Belgique. Bruxelles, 36: 396–449.
- Edwards, J. G. 1953. Species of the genus *Syneta* of the World (*Coleoptera: Chrysomeloidea*). Wasmann Journal of Biology. San Francisco, 11(1): 31–36.
- Eichler, W. 1914. Przyczynk do tegopokrywych Ojcowa. Pamiętnik Fizyograficzny. Warszawa, 22(3): 138–149.
- Endrődi, S. 1957. Über die Variabilität der Chrysomeliden-Art *Phytodecta formicata* Bruggm. (*Coleoptera*). Rovartani Közlemenyek. Budapest, 10(2): 341–348.
- Eschscholz J. F. 1818. Decades tres Eleutheratorum nevorum. Mémoires de l'Académie Impériale des Sciences de St. Pétersbourg. Saint Pétersbourg, 6: 451–484.
- Everts, E. J. G. 1903. *Coleoptera* Neerlandica. II. s'Gravenhage. IV 796 pp.
- Everts, E. J. G. 1919. Nieuve vondsten voor de Nederlandsche Coleopteren-fauna XXIII. Entomologische Berichten. Amsterdam, 5: 160–164.
- Everts, E. J. G. 1919. Nieuve vondsten voor de Nederlandsche Coleopteren-fauna XXIV. Entomologische Berichten. Amsterdam, 5: 172–175.
- Everts, E. J. G. 1920. Lijst van zeldzame en minder algemeene *Coleoptera*, op de excursies. Juni 1919, in Zuid-Limburg verzameld. Entomologische Berichten. Amsterdam, 5(113): 244–245.
- Everts, E. J. G. 1921. Nieuve vondsten voor de Nederlandsche Coleopteren-fauna XXXIV. Entomologische Berichten. Amsterdam, 5(119): 338–344.
- Everts, E. J. G. 1921. Nieuve vondsten voor de Nederlandsche Coleopterenfauna. XXXVII. Entomologische Berichten. Amsterdam, 6 (122): 21–26.
- Everts, E. J. G. 1922. *Coleoptera* Nederlandica. De Schildvleugelige Insecten van Nederland en het aangrenzend gebied. Martinus Nijhoff. S'Gravenhage. Volume III-IV, 796 pp.
- Everts, E. J. G. 1932. Nieuwe vondsten voor de Nederlandsche Coleopteren-fauna. Entomologische Berichten. Amsterdam, 8[188]: 432–434.
- Fabricius, J. C. 1775. Systema Entomologicae, sistens insectorum classes, ordines, genera, species, adjectis synonymis, locis, descriptionibus, observationibus. Korte, Flensburgi et Lipsiae. 8. [30]+832.
- Fabricius, J. C. 1777. Genera insectorum eorumque characteres naturales secundum numerum, figuram, situm et proportionem omnium partium oris adiecta mantissa specierum nuper detectarum. M. F. Bartschii, Chilonii [= Kiel]. [16+] 310 pp.
- Fabricius, J. C. 1781. Species insectorum exhibentes eorum differentias specificas, synonyma, auctorum, loca natalia, metamorphosin adiectis observationibus, descriptionibus. Hafniae [= Copenhagen], I and II, 517pp + 494 pp.
- Fabricius, J. C. 1787. Mantissa insectorum sistens species nuper detectas adiectis synonymis, observationibus, descriptionibus, emendationibus. Hafniae. I and II, 348 pp. + 382 pp.
- Fabricius, J. C. 1792. Entomologia Systematica emendata et aucta, secundum classes, ordines, genera, species adjectis synonymis, locis, observationibus. C.G. Proft, Hafniae. Tonus I. Pars I: 330 pp. Pars II: 538 pp.
- Fabricius, J. C. 1793. Entomologica systematica emendata et aucta. Secundum classes, ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus. Hafniae, 3: 487+349.
- Fabricius, J. C. 1794. Entomologica systematica emendata et aucta. Secundum classes, ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus. Hafniae, 4: 472.
- Fabricius, J. C. 1798. Supplementum entomologiae systematicae. Storch, Hafniae. 572 pp.
- Fabricius, J. C. 1801. Systema Eleutheratorum, ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus. Volume II. Bibliopol. acad., Kiliae, 687 pp.

- Fabricius, J. C. 1803. Index generum et specierum systematis eleutorum. Helmstadii, 93 pp.
- Fairmaire, L. 1851. Description de sept Coléoptères nouveaux provenant du Midi de l'Europe et du Maroc. Revue et magasin de zoologie pure et appliquée. Paris, 3: 527–528.
- Fairmaire, L. 1861. Coléoptères nouveaux recueillis pendant l'excursion en Savoie. Annales de la Société Entomologique de France. Paris, sér. 4(1): 647–650.
- Fairmaire, L. 1861. Miscellanea entomologica, quatrième partie. Annales de la Société Entomologique de France. Paris, sér. 4(1): 577–596.
- Fairmaire, L. 1874. [Indication des familles végétales sur les plantes desquelles vivent diverses espèces de Cassides]. Annales de la Société Entomologique de France. Paris, sér. 5(4): 97–98.
- Fairmaire, L. 1877. [Diagnoses de plusieurs Coléoptères nouveaux du genre *Cryptodus*]. Annales de la Société Entomologique de France. Paris, sér. 5(7): 179–180.
- Fairmaire, L. 1878. [Diagnoses de plusieurs Coléoptères nouveaux du genre *Malaxia*]. Annales de la Société Entomologique de France. Paris, sér. 5 (8): 139.
- Fairmaire, L. 1884. Descriptions de Coléoptères recueillis par le Baron Bonnaire en Algérie. Annales de la Société Entomologique Belgique. Bruxelles, 28: LIX–LXX.
- Fairmaire, L. 1885. [untitled]. Bulletin de la Société Entomologique de France. Paris, (6)5: 64–65.
- Fairmaire, L. 1885. Liste des Coléoptères recueillis à la terre de Feu par la mission de la Romanche et description des espèces nouvelles. Annales de la Société Entomologique de France. Paris, sér. 6(5): 33–62.
- Fairmaire, L. M. 1887. Notes sur les Coléoptères des environs de Pékin. Revue d'entomologie. Caen, 6: 312–335.
- Fairmaire, L. 1888. Descriptions de Coléoptères de l'intérieur de la Chine. Quatrième partie. Annales de la Société Entomologique Belgique. Bruxelles, 32: 7–46.
- Fairmaire, L. 1888. Notes sur les Coléoptères des environs de Pékin. Revue d'entomologie. Caen, 7: 111–160.
- Fairmaire, L., Allard, E. 1873. Révision du genre *Timarcha*. Annales de la Société Entomologique de France. Paris, sér. 5(3): 143–202.
- Faldermann, V. 1833. Bulletin de la Société Impériale des Naturalistes de Moscou. Moscou, 4: 70.
- Faldermann, F. 1835. Coleopterorum ab illustrissimo Bungio in China boreali, Mongolia, et montibus Altaicis collectorum, nec non ab ill. Turczaninoffio et Stchukino e provincia Irkutsk missorum illustrationes. Mémoires de l'Académie Impériale des Sciences de St. Pétersbourg. Saint Pétersbourg, 2: 337–464.
- Faldermann, V. 1835. Fauna entomologica Transcaucasica. *Coleoptera*. Nouvelle Memoire Societa Naturale. Moscow. 310 pp.
- Faldermann, F. 1837. Fauna entomologica Transcaucasica. *Coleoptera* II. Nouvelle Memoire Societa Naturale. Moscou, 5: 1–433.
- Faldermann, F. 1838. Fauna entomologica Transcaucasica. *Coleoptera* III. Nouvelle Memoire Societa Naturale. Moscou, 6: 1–338.
- Fauvel, A. 1888. Reclassifications aus Catalogus Coleopterorum Europae et Caucasi (suite). Revue d'entomologie. Caen, 7: 229–251.
- Fischer von Waldheim, G. 1824. Entomographia Imperii Rossici: Genera Insectorum systematicae exposita et analysi iconographica instructa. Volumen II. Moscou, 262 pp.
- Fischer de Waldheim, G. 1842. Catalogus Coleopterorum in Siberia Orientali à cel. Gregorio Silide Karelin collectorum. Mosquae, 28 p.
- Fischer de Waldheim, G. 1843. Spicilegium Entomographie Rossicae. Bulletin de la Société des Naturalistes de Moscou. Moscou, [1844]. 17(1): 3–144.
- Fleischer, A. 1909. Bemerkenswerte Koloritaberrationen. Wiener Entomologische Zeitung. Wien, 28(1): 32.
- Fleischer, A. 1914. Coleopterologische Notizen. Wiener Entomologische Zeitung. Wien, 33: 65.
- Fleischer, J. 1916. New Chrysomeliden aus Japan. Wiener Entomologische Zeitung. Wien, 35: 222–223.
- Fleischer, J. 1917. Bemerkenswerte Aberrationen einiger Coleopteren-Arten. Wiener Entomologische Zeitung. Wien, 36: 121–122.
- Formanek, R. 1900. Eine neue *Lochmoea* Wse. (*Col. Chrysom.*). Wiener Entomologische Zeitung. Wien, 19(5): 127.

- Forster J. R. 1771. *Novae species Insectorum*. Cenuria I. Londini, VIII + 100 pp.
- Förster, A. 1849. Uebersicht der Käferfauna der Rheinprovinz. Verhandlungen des Naturhistorischen Vereins der Preussischen Rheinland. Westfalens. Bonn, 6: 383–500. Erster Nachtrag. 1–45.
- Foudras, C. 1859. Alticides (*Halticinae*). pp. 138–384 *In*: Mulsant, E. Histoire naturelle des Coléoptères de France. Annales de la Société Linnéenne de Lyon. Lyon. (n.s.), 6.
- Foudras, C. 1860. Alticides (*Halticinae*). pp. 17–128. *In*: Mulsant, E. Histoire naturelle des Coléoptères de France. Annales de la Société Linnéenne de Lyon. Lyon, (n.s.), 7.
- Fourcroy, A. F. 1785. *Entomologia Parisiensis: sive Catalogus Insectorum quae in Agro Parisiensi reperiuntur; secundum methodum Geoffraenam in sectiones, genera et species distributus: cui addita sunt nomina trivialia et fere trecentae novae species*. Sub Privilegio Academiae. Parisiis, vol. I: 1–231, vol. II: 233–544.
- Fowler, W. W. 1900. *The Coleoptera of the British Islands*. London. 359.
- Fowler, W. W. 1913. *The Coleoptera of the British Islands*. VI (Supplement). London. 351 pp.
- Franz, H. 1938. Revision der Verwandtschaftsgruppe der *Chrysomela gypsophylae* Küst. Entomologische Blätter. Krefeld, 34(4): 190–210, (5): 249–273.
- Franz, H. 1949. Zur Kenntnis der Rassenbildung bei einigen Arten der Gattung *Cryptocephalus* (*Coleopt.*, *Chrysom.*). Portugalia Acta Biologica. Ser. B, Lisboa, 1949: 165–195.
- Franz, H. 1952. II. Beitrag zur Kenntnis der Gattung *Chrysomela* (*Coleopt.*, *Chrysomel.*). Entomologisches Nachrichtenblatt Österreichischer und Schweizer Entomologen. Wien, 4(1/3): 2–6.
- Fricken, W. 1888. *See*: Schilsky, J. 1888.
- Frivaldszky, J. 1876. Adatok Temes és Krassó megyék faunájához. Magyar Tudom. Akad. Math. Természet. Budapest, 13: 22–378.
- Frivaldszky, J. 1883. *Coleoptera nova ex Hungaria*. Természetrizsi Füzetek. Budapest, 7(1): 9–18.
- Fuente, J. M. 1909. Datos para la fauna de a provincia de Ciudad Real. Boletín de la Sociedad española de historia nateral. Madrid, 9: 306–310.
- Fuente, J. M. 1909. Consideraciones sobre una nota e Mr. Pic. Boletín de la Sociedad española de historia nateral. Madrid, 9: 436–437.
- Fuss C. 1856. Die siebenbürgischen Chrysomelen und Oreinen. Verhandlungen und Mittheilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt. Hermannstadt, 7: 25–32.
- Fuss C. 1861. Beiträge zur siebenbürgischen Käferfauna. Zur Gattung Chrysomela. Verhandlungen und Mittheilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt. Hermannstadt, 12: 151–163.
- Fuss, C. 1864. Berichtigungen und Beiträge zur siebenbürgischen Käferfauna. Verhandlungen und Mittheilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt. Hermannstadt, 15: 146–150.
- Gabriel, C. 1918. *Aphthona coerulea* ab. nov. *picipes*. p. 209. *In*: Kolbe, W. Beiträge zur schlesischen Käferfauna. Entomologische Mitteilungen, 7(10/12): 200–211.
- Gabriel, [C.] 1927. *In*: Kolbe, W. 1927. Beiträge zur schlesischen Käferfauna. Zeitschrift für Entomologie, Breslau. 15(1): 2–14.
- Gabriel, C. 1928. Kleine coleopterologische Mitteilungen. Zeitschrift für Entomologie. Breslau. [1929], 17(1): 8.
- Gabriel, C. 1931. *In*: Kolbe, W. 1931. Beiträge zur schlesischen Käferfauna. Zeitschrift für Entomologie. Breslau, 17(2): 8–17.
- Gautier de Cottés, [n.]. 1861. Description de Coléoptères nouveaux propres a la faune française. Annales de la Société Entomologique de France. Paris, sér. 4(1): 193–194.
- Gebler, F. A. 1817. *Insecta Sibiriae rariora*. Mémoires de la Société Imperiale des Naturalistes des Moscou. Moscou, 5: 315–333.
- Gebler, F. A. 1823. *Chrysomelae Sibiriae rariores*. Mémoires de la Société Imperiale des Naturalistes des Moscou. Moscou, 6: 117–126.
- Gebler, F. A. 1823. Observaciones entomologicae. Mémoires de la Société Ent. Moscou. VI: 115–131.
- Gebler, F. A. 1825. *Coleptorum Sibiriae species novae*. *In*: Hummel (ed.), Essais entomologiques, St.-Petersbourg. 4: 42–57.

- Gebler, F. [A.] 1830. Bemerkungen über die Insekten Sibiriens, vorzüglich des Altaj. *In*: Ledebur's Reise durch das Altaj-Gebirge und die songorische Kirgisen-Steppe. Bd. 2: 199, 201–218.
- Gebler, F. A. 1832. Notice sur les Coléoptères qui se trouvent dans le district de mines de Nertchinsk, dans la Sibérie orientale avec la description de quelques espèces nouvelles. *Nouveaux Mémoires de la Société Impériale des Naturalistes de Moscou*. Moscou, n. sér., 2: 23–78.
- Gebler, F. A. 1834. Notae et addidamenta ad catalogum Sibiriae occidentalis et confinis Tatariae Operis, C. F. von Ledebours Reise in das Altaigebirge und die soongorische Kirgisensteppe (Zweiter Theil, Berlin 1830). *Bulletin de la Société Impériale des Naturalistes de Moscou*. Moscou, 6 [1833]: 262–309.
- Gebler, F. A. 1841. Notice sur les Coléoptères qui se trouvent dans le district de mines de Nertchinsk. *Mémoires de la Société Impériale des Naturalistes de Moscou*. Moscou, 4: 544.
- Gebler, F. A. 1841. Notae et addidamenta ad catalogum Coleopterorum Sibiriae occidentalis et confinis Tatariae. *Bulletin de la Société Impériale des Naturalistes de Moscou*. Moscou, 14(4): 577–625.
- Gebler, F. A. 1843. Charakteristik der von H. Schrenk in den J. 1841–1843 in den Steppen der Dsüngorei gefundenen neuen Coleopterenarten. *Bulletin de la Classe physico-mathématique de l'Académie impériale des sciences de Saint-Petersbourg*. Saint Pétersbourg et Leipzig, pars 1: 36–40.
- Gebler, F. A. 1845. Charakteristik der von H. Schrenk in den J. 1841–1843 in den Steppen der Dsüngorei gefundenen neuen Coleopterenarten. *Bulletin de la Classe physico-mathématique de l'Académie impériale des sciences de Saint-Petersbourg*. Saint Pétersbourg et Leipzig, pars 3: 97–106.
- Gebler, F. A. 1847. Verzeichnis der im Kolywano-Woskresenskischen Hüttenbezirke Süd-West Sibiriens beobachteten Käfer mit Bemerkungen und Beschreibungen. *Bulletin de la Société Impériale des Naturalistes de Moscou*. Moscou, 20(1): 263–361, 20(2): 391–512.
- Gebler, F. A. 1848. Verzeichnis der im Kolywano-Woskresenskischen Hüttenbezirke Süd-West Sibiriens beobachteten Käfer mit Bemerkungen und Beschreibungen. *Bulletin de la Société Impériale des Naturalistes de Moscou*. Moscou, 21(3): 3–85.
- Gemminger, M. and E. von Harold. 1876. *Catalogus Coleopterorum hucusque descriptorum synonymicus et systematicus*. Monachii. Volume 12: 3479–3822.
- Geoffroy, E. L. 1762. *Histoire abrégée des Insectes qui se trouvent aux environs de Paris; dans laquelle ces animaux sont rangés suivant un ordre méthodique*. Durand, Paris, 1: 523 pp.
- Geoffroy, E. L. 1785. [New species]. *See*: Fourcroy, A. F. de. 1785.
- Gerhardt, J. 1891. [untitled]. *In*: Letzner, K. Verzeichnis der Käfer Schlesiens. *Zeitschrift für Entomologie*. Breslau, N. F. 16: 349–438.
- Gerhardt, J. 1904. Neuheiten der schlesischen Koleopterenfauna aus dem Jahre 1903. *Zeitschrift für Entomologie*. Breslau, N. F. 29: 77–78.
- Gerhardt, J. 1906. Neuheiten der schlesischen Koleopterenfauna. *Deutsche Entomologische Zeitschrift*. Berlin, 1906(1): 238.
- Gerhardt, J. 1909. Neuheiten der schlesischen Koleopterenfauna aus dem Jahre 1908. *Deutsche Entomologische Zeitschrift*. Berlin, 415–422.
- Gerhardt, J. 1910. Verzeichnis der Käfer Schlesiens prussischen und osterreichischen Anteils, geordnet nach dem *Catalogus coleopterorum* vom Jahre 1906. Berlin, 14: 431 pp.
- Gerhardt, J. 1911. Neuheiten der schlesischen Käferfauna aus dem Jahre 1910. (*Col.*). *Deutsche Entomologische Zeitschrift*. Berlin, 3: 337–338.
- Germar, E. F. 1810. [Untitle]. *In*: Ahrens, A. Beiträge zu einer Monographie der Rohrkäfer. *Neue Schriften der naturforschenden Gesellschaft zu Halle* 1: 9–48.
- Germar, E. F. 1811. Nachträge zu Ahrens Monographie der Rohrkäfer. *Neue Schriften der Naturforschenden Gesellschaft zu Halle* 1(6): 25–38.
- Germar, E. F. 1813–1821. *Magazin der Entomologie*. Halle. Ia [1813]: 150 pp., Ib [1815]: 194 pp., II [1817]: 4 nennun + 146 pp., III [1818]: 464 pp., IV [1821]: 462 pp.
- Germar, E. F. 1817. *Fauna insectorum Europae*. Halae, 3: 206.
- Germar, E. F. 1822. *Fauna Insectorum Europae*, fasc. VII. Halae, 24 num.
- Germar, E. F. 1824. *Insectorum species novae aut minus cognitae, descriptionibus illustratae*, vol. I Halae: XXIV+624 pp.
- Germar, E. F. 1831. *Fauna insectorum Europae*. Halae, 14–15: 206.
- Germar, E. F. 1845. *See*: Redtenbacher, F. 1845.

- Gestro, R. 1899. Nuovo forme del gruppo della *Platypria*. Annali del Museo Civico di Storia Naturale di Genova. Genova. (2) 20(40): 173–176.
- Gestro, R. 1923. Materiali per lo studio delle *Hispidae*. LIV. Contributi alla sistematica della tribu e descrizione di specie nove. Annali del Museo Civico di Storia Naturale di Genova. Genova, (3)11(51): 5–22.
- Gistel, J. 1857. Achthundert und zwanzig neue oder unbeschriebene virbellose Thiere. Vacuna. Straubing, 2: 513–606.
- Gmelin, F. 1790. Caroli a Linné Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima tertia aucta, reformata. I. 4. Lipsiae: 1517–2224.
- Gmelin, J. 1793. Caroli Linnai Systema Naturae, ed. XIII reformata. III. Lipsiae. 1693.
- Goecke, H. 1934. Revision asiatischer Donaciinen (*Col. Chrys.*). I. (3. Beitrag zur Kenntnis der Donaciinen). Koleopterologische Rundschau. Wien, 20: 215–230.
- Goecke, H. 1944. Revision asiatischer Donaciinen (*Col. Chrys.*). III. 14. Beitrag zur Kenntnis der Donaciinen. Entomologische Blätter für Biologie und Systematik der Käfer 40: 7–14.
- Goeze, J. A. E. 1777. Entomologische Beiträge zu des Ritter Linnè 12. Ausgabe des Natursystems. Erster Theil. Weidmanns Erben und Reich. Leipzig. 736 pp.
- Gorham, H. S. 1885. *Cassida chloris* Suffr. The Entomologist's Monthly Magazine. London, 22: 13–14.
- Gorham, H. S. 1885. Revision of the Phytophagous *Coleoptera* of the Japanese fauna: Subfamilies *Cassidinae* and *Hispinae*. Proceedings of the Zoological Society of London. London: 280–286.
- Gortani, M. 1906. Saggio sulla distribuzione geografica delli Coleotteri in Friuli. In: Alto, Cronaca della Soc. Alpina Friulana. Udine, 16/17: 1–93.
- Gradl, H. 1881. Aus der Fauna des Egerlandes. Entomologische Zeitung. Stettin, 7(20): 294–309.
- Gradl, H. 1882. Aus der Fauna des Egerlandes. Zweite Folge. Entomologische Zeitung. Stettin, 8: 323–332.
- Graells, M. P. 1858. Zoologie espagnole. Insectes. Memorias della comision del Mapa zoologico de Espana. 4. 111 pp.
- Gravenhorst, J. L. C. 1807. Vergleichende Uebersicht des Linneischen und einigen neuern zoologischen Systeme; nebst dem eingeschalteten Verzeichnisse der zoologischen Sammlung des Verfassers und der Beschreibung neuer Tierarten, die in derselben vorhanden sind. Göttingen. XX+476 pp.
- Gravenhorst, J.L.C. 1842. Beobachtungen über die Verwandlung von *Cassida*. Acta Acad. Leopold. Carol., 19, suppl. 2, 429–440 + 1 pl.
- Gredler, V. M. P. 1866. Die Käfer von Tirol nach ihrer horizontalen und vertikalen Verbreitung. 2. Bozen: 235–491.
- Gressitt J. L. 1946. Chinese Chrysomelid beetles of the subfamily *Chlamisinae*. Annals of the Entomological Society of America, Columbus, 39: 84–100.
- Gressitt J. L., Kimoto S. 1961. The *Chrysomelidae* (*Coleoptera*) of China and Korea. Part I. Pacific Insects Monograph. Honolulu: 1–299.
- Gressitt J. L., Kimoto S. 1963. The *Chrysomelidae* (*Coleoptera*) of China and Korea. Part II. Pacific Insects Monograph. Honolulu: 301–1026.
- Gressitt, J. L. 1942. Plant beetles from South and West China. I. Lingnan Science Journal. Kanton. 20: 271–287.
- Grimmer, K. H. B. 1841. Steiermark's Coleoptern mit einhundert sechs neu beschriebenen Species. Grätz. 50 pp.
- Gruev, B. 1970. New genera and species of subfamily *Halticinae* (*Chrysomelidae*, *Coleoptera*) of the fauna of Bulgaria. Nauchnye Trudove. Bio. Vissh. Ped. Institut. Plovdiv, 8(3): 151–154.
- Gruev, B. 1971. Für die Fauna bulgariens neue Arten der Unterfamilie *Halticinae* (*Col. Chrysomelidae*). Mitteilungen der Deutscher Entomologische Gessellschaft. Berlin, 29(5–6): 58–59.
- Gruev, B. 1973. Über die einheit der *Longitarsus lycopi*-gruppe und der *Longitarsus obliteratus*-gruppe mit notizen über verbreitung, nomenklatur und diagnostische merkmale einigen arten, sowie beschreibung von zwei neuen arten (*Coleoptera*, *Chrysomelidae*). Universite de Plovdiv „Paissi Hildendarski“. Travaux scientifiques, 11(5): 127–145.
- Gruev, B. 1975. Eine neue *Epithrix* art aus Asien und angaben über *Epithrix ermischii* Mohr. 1968 (*Col. Chrysomelidae*). Travaux scientifiques 13(4): 171–174.