

**Two new species of the subgenus *Egadroma* (genus *Stenolophus*) from South Asia, with redescription of *Stenolophus* (*Egadroma*) *ovatulus* (Bates, 1889) (Coleoptera: Carabidae)**

**Два новых вида подрода *Egadroma* (genus *Stenolophus*) из Южной Азии и переописания *Stenolophus* (*Egadroma*) *ovatulus* (Bates, 1889) (Coleoptera: Carabidae)**

B.M. КАТАЕВ\* & D.W. WRASE

Б.М. КАТАЕВ\*, Д.В. ВРАСЕ

B.M. Kataev, Zoological Institute, Russian Academy of Sciences, 1 Universitetskaya Emb., St Petersburg 199034, Russia. E-mail: harpal@zin.ru. \*Corresponding author.

D.W. Wrase, Dunckerstrasse 78, D-10437 Berlin, Germany. E-mail: carterus@gmx.de

Two new species of the subgenus *Egadroma* Motschulsky, 1855 of the genus *Stenolophus* Dejean, 1821 are described: *S. (Egadroma) melniki* sp. nov. from Thailand and *S. (Egadroma) ovchinnikovi* sp. nov. from Pakistan and India. *Stenolophus* (*E.*) *ovatulus* (Bates, 1889), **comb. nov.**, which was originally described within *Acupalpus* Latreille, 1829, is redescribed on the basis of material from China (Hainan and Fujiang provinces), Vietnam, Thailand, and Cambodia (first records from Thailand and Cambodia). The following new synonym is proposed: *Stenolophus* (*E.*) *ovatulus* (Bates, 1889) = *Egadroma fukiensis* Jedlička, 1953, **syn. nov.**

Описаны два новых вида подрода *Egadroma* Motschulsky, 1855 рода *Stenolophus* Dejean, 1821: *S. (Egadroma) melniki* sp. nov. из Таиланда и *S. (E.) ovchinnikovi* sp. nov. из Пакистана и Индии. *Stenolophus* (*E.*) *ovatulus* (Bates, 1889), **comb. nov.**, первоначально описанный в роде *Acupalpus* Latreille, 1829, переописан на основании материала из Китая (Хайнань, Фуцзянь), Вьетнама, Таиланда и Камбоджи (для Таиланда и Камбоджи этот вид указывается впервые). Установлена следующая синонимия: *Stenolophus* (*E.*) *ovatulus* (Bates, 1889) = *Egadroma fukiensis* Jedlička, 1953, **syn. nov.**


**Key words:** ground beetles, taxonomy, Cambodia, China, India, Pakistan, Thailand, Vietnam, new records, Coleoptera, Carabidae, Harpalini, *Stenolophus*, *Egadroma*, new synonymy, new species

**Ключевые слова:** жужелицы, таксономия, Вьетнам, Индия, Камбоджа, Китай, Пакистан, Таиланд, новые указания, Coleoptera, Carabidae, Harpalini, *Stenolophus*, *Egadroma*, новая синонимия, новые виды


## INTRODUCTION

*Egadroma* Motschulsky, 1855 is a rather diverse taxon of small harpalines treated by most of the recent authors as a subgenus of the genus *Stenolophus* Dejean, 1821, though some consider *Egadroma* as a separate genus as originally described. The subgenus includes more than a hundred species (Lo-

renz, 2005; Facchini, 2012) predominantly distributed in the tropical areas of the Eastern Hemisphere, with most species (76) located in the Afrotropical Region (Facchini, 2012) and about 30 species known from southern Palearctic, the Oriental Region, and Australia. Taxonomy and distribution of many species, particularly from the Ori-



**Fig. 1.** Map of South Asia, illustrating localities for *Stenolophus (Egadroma) ovatulus* (a), *S. (E.) melniki* sp. nov. (b), and *S. (E.) ovchinnikovi* sp. nov. (c).



**Figs 2–4.** *Stenolophus* (subgenus *Egadroma*), general appearance: 2, *S. (E.) ovatulus* (Vietnam); 3, *S. (E.) melniki* sp. nov. (holotype); 4, *S. (E.) ovchinnikovi* sp. nov. (holotype).

ental Region, are poorly known, a revision is needed. The present paper deals with the taxonomy of three species from South Asia (Fig. 1), somewhat similar to each other in general appearance, two of which, one from Pakistan and India and the other from Thailand, are described as new.

## MATERIAL AND METHODS

The following abbreviations are used for the depositories of the specimens examined: IOZ, Institute of Zoology, Chinese Academy of Sciences, Beijing, China; ISEN, Institute of Systematics and Ecology of Animals, Siberian Branch, Russian Academy of Sciences, Novosibirsk, Russia; MNHN, Muséum National d'Histoire Naturelle, Paris, France; NME, Naturkundemuseum Erfurt, Germany; NMP, National Museum Prague, Czech Republic; SIEE, reference collection of D.N. Fedorenko at A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow, Russia; ZIN, Zoological Institute, Russian Academy of Sciences, St Petersburg, Russia; cDST, collection of A. Dostal, Vienna, Austria; cIKAB, collection of I.I. Kabak (St Petersburg); cKM, collection of R. Kmeco, Litovel, Czech Republic; cPCH, collection of A.V. Puchkov, Kiev, Ukraine; cWR, collection of D.W. Wrase, Berlin, Germany.

Measurements were taken as follows: body length, measured from the anterior margin of the clypeus to the elytral apex; width of head, measured as the maximum linear distance across the head including the compound eyes (HWmax), and as the minimum linear distance across the neck constriction just behind the eyes (HWmin); length of pronotum (PL), measured along its median line; length of elytra (EL), measured from the basal border at scutellum to the apex of the sutural angle; maximum width of pronotum (PW) and of elytra (EW), both measured at their broadest point; length and width of metepisterna, measured along their inner and anterior margins, respectively.

## TAXONOMY

### Order COLEOPTERA

### Family CARABIDAE

### Tribe HARPALINI

Genus *Stenolophus* Dejean, 1821

Subgenus *Egadroma* Motschulsky, 1855

*Stenolophus (Egadroma) ovatulus* (Bates, 1889), **comb. nov.**

(Figs 2, 5, 6, 11, 12)

*Acupalpus ovatulus* Bates, 1889: 273. Type locality: "Saigon", Vietnam.

*Egadroma fukiensis* Jedlička, 1953: 142, **syn. nov.**  
Type locality: "Fukien [= Fujiang]: Shaowu", China.

*Type material.* Lectotype of *Acupalpus ovatulus* (present designation): male, labelled "Saigon", "*Acupalpus ovatulus* Bates" (MNHN). Paralectotypes: 1 male, labelled as lectotype (MNHN); 1 female, labelled "Saigon" (MNHN).

Paratypes of *Egadroma fukiensis*: 2 females, labelled "Fukien, Shaowu, Chine, 1937, Klapperich", "Cotype", "*Egadroma fukiensis* sp. n., det. Ing. Jedlička" (NMP).

*Additional material.* **China.** *Hainan:* 1 male, Tunchang, 11 June 1957, collector unknown (IOZ); many specimens (males and females), Hainan, March – Apr. 1980 (IOZ); 3 males, 1 female, Changjiang County, Bawang street, light trap, 19.11104°N, 109.08168°E, 145 m, 12 May 2007, night, H.B. Liang leg. (IOZ); 1 male, 1 female, Ledong, Jianfeng township, light trap, 18.69824°N, 108.78957°E, 70 m, 6 May 2007, night, H.B. Liang leg. (IOZ); 1 male, Ledong, Jianfengling township, on ground, 18.69804°N, 108.79277°E, 13 m, 19 March 2007, day, H.L. Shi & F. Yuan leg. (IOZ); 6 males, 9 females, Changiang, Shilu, reservoir, light trap, 19.25192°N, 109.08924°E, 132 m, 14 May 2007, H.B. Liang leg. (IOZ); 2 females, Danzhou, SC Tropical Agricultural University, light trap, 19.50903°N, 109.51508°E, 140 m, 15 May 2007, H.B. Liang leg. (IOZ). **Vietnam.** *Ho Chi Minh:* 1 male, "Ba-Chieu, Pres de Saigon, A. Pavie, 1996" (MNHN); *Quang Ninh:* 1 male, Fong Wong Is., 24–25 March 1987, V. Janushev leg. (ZIN); *Dongnai:* 1 female, N Dongnai Prov., Nam Cat Tien Nature Park, at light HOL450, 11 June 2005, D. Fedorenko leg. "Exped. Russ.–Vietnamese Tropical Centre" (SIEE); *Ba Ria – Vung Tau:* 5 males, 6 females,

Binh Chau–Phuoc Buu Nature Reserve, ca 50 m, 10°32'N, 107°29'E, June 2007, A.V. Abramov leg. "Exped. Russ.–Vietnamese Tropical Centre" (ZIN). **Thailand.** *Uthai Thani*: 3 males, 6 females, 25 km NW Lan Sak, 110 m, Oct. 1989, W. Thiel- en leg. (cWR, ZIN); many specimens (males and females), 25 km NW Lan Sak, 65 km NW Uthai Thani, June 1990, W. Thielen leg. (NME, ZIN); *Ta Phraya Prov.*: 1 female, La Lu Nature Park, 122 m, 14°02'31"N, 102°34'4.6"E, at light, 15 May 2010, V. Zinchenko leg. (ISEN); 2 females, same data as preceding but 127 m, 14°02'32"N, 102°34'32"E, 16 May 2010 (ISEN, ZIN). **Cambodia.** *Battambang*: 1 female, "Siam, Battam- bang, A. Pavie, 1886" (MNHN); *Koh Khong*: 1 female, 20 km SE of Koh Kong, Talai River, 11°34'N, 103°07'E, 50–300 m, 17 Oct.–15 Nov. 2011, Z. Linek leg. (cDST); 1 male, Cardamom Mts., Trapeang Rung, 5 m, 28 Jan.–4 Febr. 2010, S. Murzin leg. (cWR).

**Redescription.** Body wide, flattened (Fig. 2). Length 3.6–4.2 mm, width 1.7–2.0 mm.

Dark brown to almost black (in most specimens head slightly darker than pronotum and elytra), upperside moderately shining, not iridescent; pronotal margins along lateral furrows (in few specimens more widely) lighter, elytral interval 1 in apical half, elytral epipleurae in apical half, palpi, two basal antennomeres, and base of anten- nomere 3 brownish yellow; antennomeres 3–11 rather distinctly infusate, dark brown or brown; legs brown or brownish yellow, femora usually more or less infusate.

Microsculpture on head and pronotum fine, consisting of a mixture of isodiametric and slightly transverse meshes, on elytra meshes larger and more distinct, slightly transverse.

Head medium-sized (in male,  $HW_{max}/PW = 0.67–0.69$ ,  $HW_{min}/PW = 0.52–0.54$ ; in female, these indices 0.66–0.67 and 0.52–0.54), with large, hemispherical eyes ( $HW_{max}/HW_{min} = 1.29–1.30$  in male and 1.25–1.28 in female); the latter ventrally almost reaching buccal fissure. Tempora short, slightly convex, abruptly descending to neck. Labrum almost straight anteriorly. Fronto-clypeal suture distinct, superficial. Frontal foveae small; clypeo-ocular lines reaching eyes, slightly deepened at clyp-

us. Left mandible acute at apex. Anten- nae densely pubescent from antennomere 3, comparatively short, surpassing pronotal basal edge by approximately one apical anten- nomere (in female slightly shorter than in male); antennomeres 5–7 about 0.5–0.9 times as long as wide.

Pronotum moderately convex, de- pressed basally, wide ( $PW/PL = 1.49–1.59$ ), nearly equally narrowed apicad and basad, widest at middle, rather evenly rounded at sides, each with a lateral seti- gerous pore approximately in apical quarter. Lateral furrows narrow (slightly narrower than antennomere 2 basally), not widened basad, disappearing at basal angles; apical and basal angles widely rounded. Apical margin very shallowly concave, bordered only laterally; basal margin straight or slightly emarginate medially, occasionally widely rounded, not bordered. Areas at basal angles distinctly depressed, usually fused with the very wide and very shal- low basal foveae; area between basal foveae somewhat convex. Surface rather coarsely punctate along base, with finer punctures in median portion; impunctate in remain- ing portions. Basal edge glabrous.

Elytra moderately convex, slightly flat- tened medially, rather wide ( $EL/EW = 1.36–1.45$ ,  $EL/PL = 2.54–2.71$ ,  $EW/PW = 1.19–1.27$ ), widest at middle, very wide- ly rounded at sides. Humeri prominent, rounded at apices. Subapical sinuations distinct but shallow, sutural angles slightly less than 90°, blunted at tips. Basal border slightly sinuate, forming a very obtuse, al- most indistinct angle with lateral margin. Striae impunctate, almost superficial basal- ly and slightly deepened apically; intervals largely flat, moderately convex before apex; all intervals apically approximately equal in width. Parascutellar striole long, with a basal setigerous pore. Interval 3 with one dis- cal setigerous pore in apical fifth. Marginal elytral series consisting of 5 + 1 + 4 + 2 + 2 setigerous pores.

Ventral surface glabrous, last abdomi- nal sternite in both sexes rounded at apex,



with two pairs of setigerous pores and with few very short and fine setae along apical margin.

Protibia in both sexes with three preapical spines on outer margin. Apical spurs of pro-, meso- and metatibia slender, finely and densely denticulate at margins (apical spur denticulate only in basal half). Metatarsus shorter than HWmax and longer than HWmin, with rather long and slender tarsomere 1 (as in Fig. 19). Tarsomere 5 with one pair of lateroventral setae. Pro- and mesotarsi in male weakly enlarged (as in Figs 17, 18); mesotarsomere 1 without adhesive vestiture ventrally.

Female genitalia (Figs 11, 12). Basal stylomere narrow, apical stylomere slender, moderately arcuate, with one very short and fine spine at external margin.

Median lobe of aedeagus (Figs 5, 6) arcuate in lateral aspect, with apical portion curved dorsad and with tip bent ventrad; sides rounded; terminal lamella in dorsal aspect slightly longer than wide, triangular, narrowly rounded at tip, with sides roundly converging. Internal sac with two large and short spines in basal and median portion.

**Comparison.** *Stenolophus* (*Egadroma*) *ovatulus* is recognised from the other immaculate Oriental species of *Egadroma* by having a wide pronotum and elytra and a distinct microsculpture on head, pronotum and elytra. In colouration and in presence of microsculpture on dorsum, this species somewhat reminds *S. (E.) satoi* Habu, 1973 described from the Ryukyu Islands, Japan, but is distinctly distinguished from the latter by the smaller body size (5.0–5.8 mm in *S. satoi*), the much wider body and the aedeagal median lobe with wider terminal lamella and with two spines in the internal sac (*S. satoi* has three spines).

**Distribution.** Southern China (Hainan and Fujiang provinces), Vietnam, Thailand, and Cambodia (Fig. 1a). First records from Thailand and Cambodia.

**Remarks.** This species has originally been described within the genus *Acupalpus* Latreille, 1829 based on the males from Sai-

gon (Bates, 1889). Andrewes (1921) after examination of the type specimens noted that this taxon is not a representative of *Acupalpus* or *Stenolophus*, because “the hind tarsi have a shallow groove on the outer side, ...the fourth tarsal [segment] is only slightly emarginate, and the apex of the prosternal process is glabrous”. According to Andrewes (l. c.), the species may be a member of a new genus, but “without more substantial characters” he preferred to leave the species provisionally within *Acupalpus*. In modern catalogues (for example, Lorenz, 2005) this species is also cited as a member of *Acupalpus*. Re-examination of the two male syntypes of *Acupalpus ovatulus*, one of which is designated here as a lectotype (see Type material), revealed that this species belongs to *Egadroma* and is conspecific with *S. (Egadroma) fukiensis* described by Jedlička (1953) from “Fukien: Shaowu”, China. This species possesses all the distinctive features of *Egadroma* listed by Habu (1973) including the setae on the apex of the prosternal process (which can have broken off in some specimens). Thus we treat *S. (Egadroma) fukiensis* as a junior synonym of *S. (Egadroma) ovatulus*, **comb. nov.**

***Stenolophus (Egadroma) melniki* sp. nov.**  
(Figs 3, 7, 8, 13, 14, 17–19)


**Holotype.** Male, “N Thailand: Mae Hong Son Prov., env. Pai, 575–617 m, road on Mae Yen waterfall, 19°21′42″N, 98°27′46″E – 19°32′01″N, 98°30′29″E, 27.IV–9.V.2013, I. Melnik leg.” (ZIN).

**Paratypes.** 9 males, 4 females, same data as holotype (ZIN, cIKAB, cWR); 1 male, “NW Thailand V-92 MAE HONG SON, HUAI SUA TAO leg. Strnad” (cWR).

**Description.** Body length 3.9–4.5 mm, width 1.2–1.9 mm (in holotype, 4.1 and 1.8 mm, respectively).

Morphological characters as described for *S. (E.) ovatulus* except as follows:

Body more slender (Fig. 3). Legs more or less unicolourous, femora not infuscate. Pronotum slightly narrower, widest in or just before middle, narrowed basad usu-



**Figs 5–22.** *Stenolophus* (subgenus *Egadroma*): 5, 6, 11, 12, *S. (E.) ovatulus* (Thailand); 7, 8, 13, 14, 17–19, *S. (E.) melniki* sp. nov. (7, 8, 17–19, holotype; 13, 14, paratype); 9, 10, 15, 16, 20–22, *S. (E.) ovchinnikovi* sp. nov. (9, 10, 20–22, holotype; 15, 16, paratype). 5–10, median lobe of aedeagus; 11, 13, 15, stylus; 12, 14, 16, stylus and laterotergite; 17, 20, male protarsus; 18, 21, male mesotarsus; 19, 22, male metatarsus. Lateral view (5, 7, 9, 11, 13, 15), dorsal view (6, 8, 10, 17–22), ventral view (12, 14, 16). Scale bar: 0.5 mm.

ally more markedly than in *S. (E.) ovatulus*; disc less distinctly depressed basally, with slightly deeper and more distinctly delimited basal foveae; basal punctation slightly finer and sparser; lateral furrows prolonged to basal margin up to inner margins of basal foveae. Elytra noticeably narrower and longer. Pro-, meso- and metatarsi (Figs 17–19) similar to those in *S. (E.) ovatulus*.

Female genitalia (Figs 13, 14).

Median lobe of aedeagus (Figs 7, 8) bent ventrad behind basal bulb, in lateral aspect ventrad along dorsal margin and almost straight along middle portion of ventral margin, with apical portion slightly directed dorsad and with tip bent ventrad; terminal lamella in dorsal aspect triangular, slightly wider than in *S. (E.) ovatulus*, approximate-

ly as long as wide, narrowly rounded at tip, with sides evenly and roundly converging. Internal sac with one wide and short spine in middle portion of median lobe.

Proportions: HWmax/PW = 0.67–0.69 in male and 0.66–0.68 in female; HWmin/PW = 0.50–0.54 in male and 0.50–0.55 in female; HWmax/HWmin = 1.26–1.35 in male and 1.24–1.34 in female; PW/PL = 1.47–1.54; EL/EW = 1.41–1.46, EL/PL = 2.56–2.76, EW/PW = 1.21–1.26.

**Comparison.** *Stenolophus (Egadroma) melniki* sp. nov. is rather similar to *S. (E.) ovatulus* in general appearance and male genitalia, but distinctly differs from the latter species in the more slender body with longer elytra, the unicolourous pale legs, the lateral furrows of pronotum prolonged to basal margin and in the more robust median lobe of the aedeagus with only one spine in the internal sac. Other distinctive features of the new species are listed in the description.

**Etymology.** This new species is named after our friend and colleague, the entomologist Igor Melnik (Moscow), who collected the type series.

**Distribution.** Known only from two localities (environments of Pai and Huai Sua Tao, Mae Hong Son Province) in north-western Thailand (Fig. 1b).

***Stenolophus (Egadroma) ovchinnikovi* sp. nov.**

(Figs 4, 9, 10, 15, 16, 20–22)

**Holotype.** Male, “Pakistan, Jammu and Kashmir: Islamabad, SW from Garni, 1500 m, 5–15.VII.2003, V. Gurko & S. Ovchinnikov leg.” (ZIN).

**Paratypes.** **Pakistan.** 2 females, same data as holotype (ZIN); 1 male, “Pakistan, Islamabad, 10.VII.2003, S. Ovchinnikov leg.” (ZIN); 1 male, 1 female, “PAKISTAN, Islamabad, The National Forest Park, 12.VII.2003, S. Ovchinnikov leg.”, “ex coll. S.V. Ovchinnikov. Donation to ZIN” (ZIN); 1 male, “Pakistan, Suleiman mnts, Zhob vall., 1700 m, 08.05” (cPCH). **India. Uttarakhand:** 1 male, “N. INDIA, Himalaya, U. P., Dehra Dun, New Forest, 20–30.VI.1981, 700m, leg. C. Holzschuh”, “collection Naturkunde-museum

Erfurt” (NME); 1 female, “N. INDIA (Uttaranchal), ca 13 km NW Nainital, Khairna bridge, 900 m (river banks, light trap), 13.–17.VII.2003, Z. Kejval & M. Trýzna” (cWR); **Punjab:** 1 male, 5 females, “1 / Indien, Punjab, 308 m, Hoshiarpur, Rormazara, N 31°11'5.7''E 76°12'58.0'', 6–9.07.2004, leg. M. Uhler” (cDST, cWR); **Madhya Pradesh:** 1 male, “INDIA (Madhya Pradesh, ca 60 km SW of Indore, Mandu env., ca 500 m, 22°20'N / 75°26'E, W slope of table land, 22.–24.VI.2006, Z. Kejval” (cWR); **Goa:** 4 males, 5 females, “INDIA-GOIA state, Pallolem env., 25.V.2006, O. Šafránek leg.” (cKM; cWR; ZIN).

**Description.** Body length 3.6–4.5 mm, width 1.6–2.0 mm (in holotype, 4.2 and 1.8 mm, respectively).

Morphological characters as described for *S. (E.) ovatulus* except as follows:

Body more slender (Fig. 4). Antennae in most specimens paler, antennomeres 3–11 only slightly infuscate. Pronotum slightly narrower, widest slightly before middle, narrowed basad usually more markedly than in *S. (E.) ovatulus*; lateral furrows prolonged to basal margin up to inner margins of basal foveae. Elytra narrower and longer, with deeper striae and more convex intervals; interval 3 without discal setigerous pore. Metatarsus slightly shorter, about as wide as HWmin, with distinctly shorter tarsomere 1 (Fig. 22). Pro- and mesotarsi in male more strongly widened (Figs 20, 22).

Female genitalia (Figs 15, 16) with larger basal stylomere and slightly wider apical stylomere.

Median lobe of aedeagus (Figs 9, 10) more robust than in *S. (E.) ovatulus*, arcuate in lateral aspect, with almost straight apical portion slightly directed dorsad; terminal lamella in dorsal aspect triangular, approximately as long as wide, narrowly rounded at tip. Internal sac with two large, slender spines in middle portion of median lobe and with a row of medium-sized spines on left side in apical portion of median lobe.

Proportions: HWmax/PW = 0.67–0.71 in male and 0.65–0.72 in female; HWmin/PW = 0.52–0.56 in male and 0.52–0.57 in female; HWmax/HWmin = 1.26–1.30 in

male and 1.23–1.27 in female; PW/PL = 1.46–1.53; EL/EW = 1.38–1.54, EL/PL = 2.65–2.88, EW/PW = 1.20–1.33.

**Comparison.** In external characters, this new species is similar to the two preceding ones and in its general appearance it is particularly similar to *S. (Egadroma) melniki* **sp. nov.** *Stenolophus (E.) ovchinnikovi* **sp. nov.** is easily distinguished from both species by the deeper elytral striae, the absence of a discal setigerous pore on elytral interval 3, the shorter metatarsomere 1 and by the more markedly widened pro- and mesotarsi in male. Additionally, *S. (E.) ovchinnikovi* **sp. nov.** is distinguished from *S. (E.) melniki* **sp. nov.** in having the femora infusate as in *S. (E.) ovatulus*. The aedeagal median lobe of *S. (E.) ovchinnikovi* **sp. nov.** distinctly differs from that of *S. (E.) ovatulus* and *S. (E.) melniki* **sp. nov.** in having a straight (not bent ventrad at tip) terminal lamella and a different armature of the internal sac with two large, slender spines in middle portion of the median lobe and with a row of medium-sized spines on the left side in the apical portion of the median lobe. Additional differences of *S. (E.) ovchinnikovi* **sp. nov.** from *S. (E.) ovatulus* are listed in the description of the former.

**Etymology.** This new species is named after our friend and colleague, entomologist and arachnologist Sergey Ovchinnikov (1958–2007), who collected the holotype and some of the paratypes.

**Distribution.** Known from Pakistan and India (Uttarakhand, Punjab, Madhya Pradesh and Goa) (Fig. 21c).

#### ACKNOWLEDGEMENTS

We are very grateful to A.V. Abramov (St Petersburg), T. Deuve (Paris), A. Dostal (Vienna),

R.Yu. Dudko (Novosibirsk), D.N. Fedorenko (Moscow), J. Hájek (Prague), M. Hartmann (Erfurt), I.I. Kabak (St Petersburg), R. Kmeco (Litovel), H.-B. Liang (Beijing), I.V. Melnik (Moscow) and A.V. Puchkov (Kiev) for providing material used in this study and to K.V. Markarov (Moscow) who took the photographs of general appearance. We appreciate very much the support of Jon Cooter (Oxford, England) for reading an early draft of the manuscript. The study was supported (for BK) by the Russian Foundation for Basic Research (project No. 13–04–01002a) and Ministry of Education and Science of the Russian Federation.

#### REFERENCES

- Andrewes H.E.** 1921. Notes on synonymy and on some types of Oriental Carabidae in various foreign collections. *The Transactions of the Entomological Society of London*: 145–195.
- Bates H.W.** 1889. Contributions à la faune Indo-Chinoise. 3e Mémoire (1). Carabidae. *Annales de la Société Entomologique de France*, 6e Série, 9: 261–286.
- Facchini S.** 2012. Ten new taxa of the Subgenus *Egadroma* Motschulsky, 1855 of the Genus *Stenolophus* Dejean, 1821 from the Afrotropical Region and notes on some other species of the subgenus (*Coleoptera Carabidae, Harpalinae*). *Giornale Italiano di Entomologia*, 13 (57): 69–96.
- Habu A.** 1973. *Carabidae: Harpalini (Insecta, Coleoptera)*. *Fauna Japonica*. Tokyo: Keigaku Publishing Co. xiii+430 p.
- Jedlička A.** 1953. Neue Carabiden aus der chinesischen Provinz Fukien. *Entomologische Blätter für Biologie und Systematik der Käfer*, 49: 141–147.
- Lorenz W.** 2005. *A systematic list of extant ground beetles of the world (Coleoptera "Geadephaga": Trachypachidae and Carabidae incl. Paussinae, Cicindelinae, Rhysodinae)*. Second Edition. Tutzing: W. Lorenz. 530 p.

Received October 20, 2013 / Accepted November 25, 2013