



УДК 595.753

ON THE TAXONOMY OF SOME FULGOROIDEA (HEMIPTERA)

V.M. Gnezdilov

Zoological Institute of the Russian Academy of Sciences, Universitetskaya nab. 1, 199034, Saint Petersburg, Russia;
e-mails: vmgnezdilov@mail.ru, vgnezdilov@zin.ru

ABSTRACT

Issidae: *Ardelia* Melichar, 1907 is placed in synonymy under *Vishnuloka* Distant, 1906 and *Vishnuloka cuneata* Distant, 1906 – under *Vishnuloka prominula* Distant, 1906; *Delia deserta* Melichar, 1906 is transferred to the genus *Vishnuloka*. Nogodinidae: *Narayana buruensis* Schmidt, 1926 is transferred to the genus *Scalabis* Stål, 1870, *Hysteropterum solidum* Melichar, 1911 – to the genus *Mangola* Melichar, 1906, and *Hysteropterum ecarinatum* Synave, 1956 – to the genus *Telmosias* Fennah, 1967; *Oryxana subrecta* Jacobi, 1941 and *Buehleria rabana* Lallemand et Synave, 1953 are placed in synonymy under *Oryxana suturalis* (Melichar, 1906). Acanaloniidae: *Amphiscepa subpellucida* Fowler, 1904 is transferred to the genus *Acanalonia* Spinola, 1839. Tropiduchidae: *Nacmusius* Jacobi, 1944 is placed in synonymy under *Cixiopsis* Matsumura, 1900; *Nacmusius chelydinus* Jacobi, 1944 is transferred to the genus *Cixiopsis*. A key to the species of the genus *Vishnuloka* is provided. Lectotypes are designated for *Amphiscepa subpellucida* Fowler, 1904, *Delia deserta* Melichar, 1906, *Issina suturalis* Melichar, 1906, *Narayana buruensis* Schmidt, 1926, *Oryxana subrecta* Jacobi, 1941, *Vishnuloka cuneata* Distant, 1906, and *V. prominula* Distant, 1906.

Key words: Acanaloniidae, Issidae, Nogodinidae, Tropiduchidae, new combinations, new synonyms, lectotype designation

К СИСТЕМАТИКЕ НЕКОТОРЫХ FULGOROIDEA (HEMIPTERA)

В.М. Гнездилов

Зоологический институт Российской академии наук, Университетская наб. 1, 199034 Санкт-Петербург, Россия;
e-mails: vmgnezdilov@mail.ru, vgnezdilov@zin.ru

РЕЗЮМЕ

Issidae: *Ardelia* Melichar, 1907 сведена в синонимы к *Vishnuloka* Distant, 1906, а *Vishnuloka cuneata* Distant, 1906 – к *Vishnuloka prominula* Distant, 1906; *Delia deserta* Melichar, 1906 перенесена в род *Vishnuloka*. Nogodinidae: *Narayana buruensis* Schmidt, 1926 перенесена в род *Scalabis* Stål, 1870, *Hysteropterum solidum* Melichar, 1911 – в род *Mangola* Melichar, 1906, а *Hysteropterum ecarinatum* Synave, 1956 – в род *Telmosias* Fennah, 1967; *Oryxana subrecta* Jacobi, 1941 и *Buehleria rabana* Lallemand et Synave, 1953 сведены в синонимы к *Oryxana suturalis* (Melichar, 1906). Acanaloniidae: *Amphiscepa subpellucida* Fowler, 1904 перенесена в род *Acanalonia* Spinola, 1839. Tropiduchidae: *Nacmusius* Jacobi, 1944 сведен в синонимы к *Cixiopsis* Matsumura, 1900; *Nacmusius chelydinus* Jacobi, 1944 перенесен в род *Cixiopsis*. Данна определительная таблица к видам рода *Vishnuloka*. Установлены лектотипы для *Amphiscepa subpellucida* Fowler, 1904, *Delia deserta* Melichar, 1906, *Issina suturalis* Melichar, 1906, *Narayana buruensis* Schmidt, 1926, *Oryxana subrecta* Jacobi, 1941, *Vishnuloka cuneata* Distant, 1906 и *V. prominula* Distant, 1906.

Ключевые слова: Acanaloniidae, Issidae, Nogodinidae, Tropiduchidae, новые комбинации, новые синонимы, установление лектотипов

INTRODUCTION

In the course of studying the material, including types of planthoppers of the families Acanaloniidae Amyot et Serville, 1843, Issidae Spinola, 1839, Nogodinidae Melichar, 1898, and Tropiduchidae Stål, 1866 from several European museums some new synonyms and new combinations were discovered and are here established.

MATERIAL AND METHODS

The terminology of head follows Emeljanov (1995), hypostomal plate – Emeljanov (1971), and female genitalia – Bourgoin (1993). Dry and pinned specimens were examined using the light microscope. The material examined is deposited in the museums listed below.

Photographs of *Vishnuloka prominula* Distant and *Mangola solida* (Melichar) were made using Leica MZ8 with JVC video camera KY F7OB. Images are produced using the software Synoptics Automontage and Adobe Photoshop.

Institutional abbreviations:

SNSD – Staatliche Naturhistorische Sammlungen Dresden, Museum für Tierkunde, Germany.

MIZ – Museum and Institute of Zoology of the Polish Academy of Sciences, Warsaw, Poland.

MMBC – Moravian Museum, Brno, Czech Republic.

NMW – Naturhistorisches museum Wien, Austria.

MNB – Museum für Naturkunde, Berlin, Germany (formerly Zoologisches Museum, Humboldt Universität).

MNHN – Muséum National d' Histoire Naturelle, Paris, France.

BMNH – The Natural History Museum, London, United Kingdom.

IRSNB – Royal Belgian Institute of Natural Sciences, Brussels, Belgium.

NHMB – Naturhistorisches Museum, Basel, Switzerland.

TAXONOMY

Family Acanaloniidae Amyot et Serville, 1843

Genus *Acanalonia* Spinola, 1839

Type species: *Acanalonia servillei* Spinola, 1839.

Acanalonia subpellucida (Fowler, 1904) comb. nov.

Amphiscepa subpellucida Fowler, 1904: 119, tab. 12, figs 9, 9a.

Type material examined. Female (lectotype here designated) – GUATEMALA, City, Champion (BMNH).

Supplementary description. Coryphe very narrow. Fore wings with hypostomal plate. Hind tibiae without lateral spines.

Note. To stabilize the nomenclature in the group (ICZN 1999: Art. 74) the lectotype is here designated for the specimen listed above which corresponds to the original description and belongs to the syntype series (Fowler 1904).

The species is transferred to the genus *Acanalonia* according to wide metope and narrow coryphe and the shape and venation of fore wings. The ovipositor with large nearly triangular gonoplacs bearing marginal teeth is of characteristic acanaloniid type as described by Gnezdilov (2012).

Family Issidae Spinola, 1839

Tribe Issini Spinola, 1839

Genus *Vishnuloka* Distant, 1906

Vishnuloka Distant, 1906: 345;

Delia Melichar, 1906: 265 (Type species: *Delia deserta* Melichar, 1906), homonymy;

Ardelia Melichar, 1907: 324, nom. nov. pro *Delia* Melichar, 1906, nec *Delia* Robineau-Desvoidy, 1830, **syn. nov.**

Type species: *Vishnuloka prominula* Distant, 1906.

Vishnuloka prominula Distant, 1906

(Fig. 1A–C)

Vishnuloka prominula Distant, 1906: 345

Vishnuloka cuneata Distant, 1906: 346, **syn. nov.**

Type material examined. Female (lectotype of *Vishnuloka prominula*, here designated) – INDIA, Munghpu (BMNH). Female (lectotype of *Vishnuloka cuneata*, here designated) – MYANMAR, Tenass Vall, Myitta, Doherty (BMNH).

Supplementary description. Coryphe transverse, with anterior margin weakly marked (Fig. 1B). Metope with no intermediate carinae, protruding like a cone in its upper part (Fig. 1C). Fore wings narrowing apically; radius, median, and cubitus anterior bifurcate (R 2 M 2 CuA 2); median bifurcates




Fig. 1. Issidae and Tropiduchidae, external view: A, *Vishnuloka prominula* Distant (lectotype of *Vishnuloka cuneata*), lateral view; B, same, dorsal view; C, same (lectotype of *Vishnuloka prominula*), frontal view; D, *Cixiopsis chelydinus* (Jacobi), dorsal view. Scale bar (1 mm): a = A, B; b = C; c = D.

distally. Hind wings well developed, 3-lobed. Hind tibia with 2 lateral spines. First metatarsomere with 2 latero-apical and 2 intermediate spines.

Distribution. Northeastern India (Sikkim) and Myanmar (Distant 1906).

Note. The lectotypes are here designated for the specimens listed above which correspond to the original descriptions and belong to the syntype series (Distant 1906).

Vishnuloka deserta (Melichar, 1906) comb. nov.

Delia deserta Melichar, 1906: 265;

Ardelia deserta: Melichar, 1907: 324.

Type material examined. Female (lectotype, here designated) – INDONESIA, Sumatra, Deli, ex. coll. Fruhstorfer, “deserta det. Melichar” (handwritten), “*Delia deserta* [?] n sp” (handwritten) (NMW); 1 male (paralectotype) – INDONESIA, Sumatra, Soekaranda, Dr. H. Dohrn S., “Co=Typus.” (printed), “Ar. deserta det. Melichar” (handwritten), “*Ardelia* dbl.” (handwritten) (SNSD).

Distribution. Sumatra (Melichar 1906).

Note. The species was described after 3 specimens (Melichar 1906). The lectotype is here designated for the female specimen mentioned above which corresponds to the original description, even Melichar did not give the sex of the specimens described, but gave the length for the male only, he mentioned that one specimen from Deli is deposited in Wien (Melichar 1906). Subsequently other specimen which corresponds to the original description as well and apparently belongs to the syntype serie, originally labeled as the cotype, became the paralectotype. Very probably Melichar put the label “Ar. deserta det. Melichar” under the specimen after he replaced the name *Delia* by *Ardelia* (Melichar, 1907). Apparently “Java: Soekaranda” (Melichar 1906) in the original description is *lapsus calami*. As stated above Soekaranda is located in Sumatra. Tonkin (Northern Vietnam) mentioned by Lallemand (1942) seems to be a dubious record for the species. In Vietnam a closely related yet undescribed genus and species was discovered (Gnezdilov, unpublished).

Key to species of the genus *Vishnuloka*

1(2). Each fore wing light yellowish brown, with wide longitudinal dark brown stripe (Fig. 1A–B) *V. prominula* Distant

2(1). Fore wings brown, without stripe *V. deserta* (Melichar)

Family Nogodinidae Melichar, 1898

Tribe Tongini Fennah, 1954

Genus *Scalabis* Stål, 1870

Type species: *Tylana philippina* Stål, 1870.

Scalabis buruensis (Schmidt, 1926) comb. nov.

Narayana buruensis Schmidt, 1926: 244.

Type material examined. Male (lectotype, here designated) – INDONESIA, Buru 1921, Station 9, 21 June, L.J. Toxopeus (MIZ); 1 female (paralectotype) – INDONESIA, same place, 23 June, L.J. Toxopeus (MIZ); 1 female (paralectotype) – INDONESIA, Buru 1921, Station 13, L.J. Toxopeus (SNSD).

Note. The lectotype is here designated for the specimen listed above which corresponds to the original description and belongs to the syntype serie (Schmidt 1926). Subsequently, other specimens which belong to the syntype serie as well became paralectotypes.

The species is placed in the nogodinid genus *Scalabis* according to carination of the metope and venation of the fore wings as described by Fennah (1954). The ovipositor with narrow and long gonoplaques and anterior connective laminae of gonapophyses VIII is of characteristic nogodinid type as described by Gnezdilov (2003) for the tribe Tongini. Thus the issid genus *Narayana* Distant, 1906 is limited in its distribution to Sri Lanka and Southern India.

Genus *Oryxana* Distant, 1910

Type species: *Flata subacuta* Walker, 1870.

Oryxana suturalis (Melichar, 1906)

Issina suturalis Melichar, 1906: 210;

Issella suturalis: Metcalf, 1952: 227;

Oryxana suturalis: Gnezdilov, 2009: 86;

Oryxana subrecta Jacobi, 1941: 292, **syn. nov.**;

Buehleria rabana Lallemand et Synave, 1953: 249, **syn. nov.**;

Oryxana rabana: Gnezdilov, 2009: 86.

Type material examined. Female (lectotype of *Issina suturalis*, here designated) – INDONESIA,

Sumbawa, "Wien / Dr. Melichar", "*suturalis* det. Melichar", "Invent. č 3778 / Ent. Mor. Museum, Brno" (MMBC); 1 female (paralectotype of *Issina suturalis*) – INDONESIA, "Java / Coll. Signoret", "*suturalis* det. Melichar", "Issina" (NMW); 1 male (lectotype of *Oryxana subrecta*, here designated) – INDONESIA, W. Soembawa, Soembawa Besar, 24 April – 2 May 1927, Sunda-Exp. Rensch (SNSD); 1 male (paralectotype of *Oryxana subrecta*) – INDONESIA, W. Soembawa, Soembawa Besar, 24 April – 2 May 1927, Sunda-Exp. Rensch (MNB); 1 male and 1 female (holotype and paratype of *Buehleria rabana*) – INDONESIA, Sumbawa, Raba, 20 May 1949, Dr. Bühler & Dr. Sutter (NHMB).

Note. The lectotypes are here designated for the specimens listed above which correspond to the original descriptions and belong to the syntype series (Melichar 1906; Jacobi 1941). Subsequently, other specimens which belong to the syntype series as well became the paralectotypes.

Tribe Epaciini Fennah, 1978

Genus *Mangola* Melichar, 1906

Type species: *Mangola reticulata* Melichar, 1906.

Mangola solida (Melichar, 1911) comb. nov.

(Fig. 2A–D)

Hysteropterum solidum Melichar, 1911: 115.

Type material examined. Female (holotype) – KENYA, Afrique Orient. Angl., Mont Nyro, Maurice de Rothschild 1905, "*Hysteropt. solidum* n. sp." (handwritten) (MNHN).

Other material examined. 3 males, 3 females, KENYA, Kenya Marsabit N.P., Forêt Lac Paradis, 1400 m, 11–12 December 1972, M. Boulard leg. (MNHN).

Note. Melichar (1911) noted that only one specimen was available to him and this is here recognised as the holotype.

The species is transferred to the genus *Mangola* according to the characteristic robust shape of the body including fore wings, the transverse short coryphe, and the flat metope with a distinct median carina. The ovipositor is of characteristic nogodinid type as described by Gnezdilov (2003).

Tribe Mithymnini Fennah, 1967

Genus *Telmosias* Fennah, 1967

Type species: *Telmosias crito* Fennah, 1967.

Telmosias ecarinatus (Synave, 1956) comb. nov.

Hysteropterum ecarinatum Synave, 1956: 16, figs 14, 30, 31.

Type material examined. 1 male, 2 females (paratypes) – REPUBLIC OF SOUTH AFRICA, Lammerskraal, Prince Albert Distr., C. P. (IRSNB).

Note. The species is transferred to the genus *Telmosias* according to the carination of metope, the shape and venation of the fore wings, and the structure of male genitalia as described by Fennah (1967).

Family Tropiduchidae Stål, 1866

Tribe Cixiopsini Fennah, 1982

Fennah (1982) erected the tribe Cixiopsini for the genera *Cixiopsis* Matsumura, 1900, *Zema* Fennah, 1956, *Duriopsis* Melichar, 1902, *Padanda* Distant, 1906, and *Olontheus* Jacobi, 1944. Subsequently the genus *Caffrommatissus* Fennah, 1967 was added to the tribe (Huang and Bourgoin 1993). Matsumura (1914) placed *Padanda* Distant, 1906 in synonymy under *Cixiopsis* Matsumura, 1900 and *Padanda atkinsoni* Distant, 1906 – in synonymy under *Cixiopsis punctatus* Matsumura, 1900. He treated *P. atkinsoni* as brachypterous form of *C. punctatus*. Finally Anufriev and Emeljanov (1988) have regarded *Olontheus obscurus* Jacobi, 1944 as a junior synonym of *C. punctatus*. As *O. obscurus* is the type species of the genus *Olontheus* Jacobi, 1944 it means the synonymization of the genera *Cixiopsis* and *Olontheus* as well. Our study of the type specimen of *Nacmusius chelydinus* Jacobi, 1944, which was treated as belonging to the family Issidae (Jacobi 1944; Metcalf 1958), shows that this species belongs to the genus *Cixiopsis* and accordingly *Nacmusius* Jacobi, 1944 is a junior synonym of *Cixiopsis* Matsumura, 1900.

Taking into consideration the synonymies mentioned above the tribe Cixiopsini consist of four genera: *Cixiopsis* Matsumura, 1900, *Duriopsis* Melichar, 1902, *Zema* Fennah, 1956, and *Caffrommatissus* Fennah, 1967 (Fennah 1982; Huang and Bourgoin 1993). The first three genera are known from Asia (Matsumura 1900, 1914; Melichar 1902; Fennah




Fig. 2. Nogodinidae, *Mangola solida* (Melichar), holotype: A, lateral view; B, dorsal view; C, frontal view; D, ovipositor. Scale bar (1 mm): a = A, B; b = C; c = D.

1956; Vilbaste 1968; Anufriev and Emeljanov 1988; Wang and Liang 2007; Rahman et al. 2011) and *Caf-frommatissus* – from southern Africa (Fennah 1967). However the genus *Cixiopsis* is very peculiar within all of the mentioned genera (except the genus *Duriopsis* for which the structure of the male genitalia is not described), in the structure of penis with the character of a large laterally flattened phallobase totally covering the aedeagus. The tribe is in need of revision.

Genus *Cixiopsis* Matsumura, 1900

Cixiopsis Matsumura, 1900: 207;
Padanda Distant, 1906: 331 (Type species: *Padanda atkinsoni* Distant, 1906), syn. fide Matsumura, 1914: 268;
Olontheus Jacobi, 1944: 17 (Type species: *Olontheus obscurus* Jacobi, 1944), syn. fide Anufriev and Emeljanov, 1988: 488;
Nacmusius Jacobi, 1944: 19 (Type species: *Nacmusius chelydinus* Jacobi, 1944), **syn. nov.**

Type species: *Cixiopsis punctatus* Matsumura, 1900.

Note. We have studied the material on the genus *Cixiopsis* from Nepal (Gnezdilov, unpublished) and found it significantly different in the structure of penis from *Cixiopsis punctatus*. Thus despite the synonymy proposed by Matsumura (1914) and supported by Anufriev and Emeljanov (1988) for the species, we suggest treating all type species mentioned above as distinct species of the genus *Cixiopsis* until the males of the species from type localities are examined.

Cixiopsis punctatus Matsumura, 1900

Cixiopsis punctatus Matsumura, 1900: 208.

Distribution. Japan (Hokkaido, Honshu, Kiushu) (Matsumura 1900, 1914), Soutern Korea (Rahman et al. 2011), Russian Far East (Vilbaste 1968; Anufriev and Emeljanov 1988).

Cixiopsis atkinsoni (Distant, 1906)

Padanda atkinsoni Distant, 1906: 332, fig. 166.

Distribution. Northeastern India (Sikkim).

Cixiopsis chelydinus (Jacobi, 1944) comb. nov. (Fig. 1D)

Nacmusius chelydinus Jacobi, 1944: 19, fig. 6.

Material examined. 1 female – CHINA, Fukien, Shaowu (500 m), 7 August 1937, J. Klapperich leg. (SNSD).

Distribution. Southeastern China.

Note. The species was described after 3 males from Kwangtseh in China collected by J. Klapperich (Jacobi 1944). According to the list of types deposited in the Zoologisches Forschungsmuseum Alexander König in Bonn (Lampe et al. 2006) there is one specimen deposited there (incorrectly named as the holotype) with the labels corresponding to the original description and this may be designated later after examination as the lectotype; we have examined the photo of this specimen (Fig. 1D). Another specimen mentioned in the same list (as the paratype) does not completely correspond to Jacobi's label data (1944) as it was collected in October, but according to the abbreviation given by Jacobi all 3 syntypes were collected during July–September.

Cixiopsis obscurus (Jacobi, 1944)

Olontheus obscurus Jacobi, 1944: 17, fig. 5.

Type material examined. 1 specimen with abdomen missing (holotype) – VIETNAM, Tonkin, Chapa, 28 June 1917, Jeanvoine (SNSD).

Distribution. Northern Vietnam.

Note. The species was described after a single male (Jacobi 1944). The specimen listed above corresponds to the original description and this is recognised here as the holotype. Lampe et al. (2006) listed the holotype of the species as deposited in Bonn, however, the labels mentioned in the list is not corresponding to the original description.

ACKNOWLEDGEMENTS

The study was financially supported by the Royal Society (London, UK), the Ministry of Education and Science of the Russian Federation (project No. 16.518.11.7070), Muséum National d'Histoire Naturelle (Paris, France), and the Alexander von Humboldt Stiftung (Bonn, Germany). I give thanks to Mr. Mick Webb (London, UK), Dr. Mike Wilson (Cardiff, UK), Dr. Igor Malenovský (Brno, Czech Republic), Prof. Dr. Thierry Bourgoin and Dr. Adeline Soulier-Perkins (Paris, France), Mr. Jérôme Constant (Brussels, Belgium), Dr. Daniel Burckhardt (Basel, Switzerland), Dr. Adam Stroiński (Warsaw, Poland), Dr. Herbert Zettel (Wien, Austria), Dr. Christian Schmidt (Dresden, Germany), and Dr. Jürgen Deckert (Berlin, Germany).

for the opportunity to examine the material and their hospitality, and late Dr. Karl-Heinz Lampe (Bonn, Germany) for his photo of the type specimen of *Nacmusius chelydinus* Jacobi. I am glad to thank Dr. Lois O'Brien (Tucson, USA) for language editing of the manuscript, Prof. Dr. Hannelore Hoch (Berlin, Germany) for her valuable comments on the text, and Dr. Leonid Anisyutkin (St. Petersburg, Russia) for his help with making figure plates.

REFERENCES

- Anufriev G.A. and Emeljanov A.F. 1988.** Cicadinea (Auchenorrhyncha). In: P.A. Lehr (Ed.). Key to the insects of the Far East of the USSR, 2. Leningrad, Nauka: 12–495.
- Bourgoin T.** 1993. Female genitalia in Hemiptera Fulgoromorpha, morphological and phylogenetic data. *Annales de la Société Entomologique de France. (N.S.)*, 29(3): 225–244.
- Distant W.L. 1906.** Rhynchota (Heteroptera – Homoptera). *The fauna of British India, including Ceylon and Burma*, 3: 1–503.
- Emeljanov A.F. 1971.** New genera of planthoppers of the families Cixiidae and Issidae (Homoptera, Auchenorrhyncha) from the fauna of the USSR. *Entomologicheskoe obozrenie*, 50(3): 619–627.
- Emeljanov A.F. 1995.** On the problem of classification and phylogeny of the family Delphacidae (Homoptera, Cicadina) taking into consideration larval characters. *Entomologicheskoe obozrenie*, 74(4): 780–794.
- Fennah R.G. 1954.** The higher classification of the family Issidae (Homoptera: Fulgoroidea) with descriptions of new species. *Transactions of the Royal Entomological Society of London* 105(19): 455–474.
- Fennah R.G. 1956.** Fulgoroidea from southern China. *Proceedings of the California Academy of Sciences*, 4(28): 441–527.
- Fennah R.G. 1967.** New and little known Fulgoroidea from South Africa (Homoptera). *Annals of the Natal Museum*, 18(3): 655–714.
- Fennah R.G. 1982.** A tribal classification of the Tropiduchidae (Homoptera: Fulgoroidea), with the description of a new species on tea in Malaysia. *Bulletin of Entomological Research*, 72: 631–643.
- Fowler W.W. 1904.** Insecta. Rhynchota. Hemiptera–Homoptera. *Biologia Centrali–Americana*, 1: 45–139.
- Gnezdilov V.M. 2003.** Review of the family Issidae (Homoptera, Cicadina) of the European fauna, with notes on the structure of ovipositor in planthoppers. *Chteniya pamyati N.A. Kholodkovskogo (Meetings in memory of N.A. Cholodkovsky)*, St. Petersburg, 56(1): 1–145.
- Gnezdilov V.M. 2009.** Revisionary notes on some tropical Issidae and Nogodinidae (Hemiptera: Fulgoroidea). *Acta Entomologica Musei Nationalis Pragae*, 49(1): 75–92.
- Gnezdilov V.M. 2012.** On the composition and distribution of the family Acanaloniidae Amyot et Serville (Homoptera, Fulgoroidea). *Entomologicheskoe obozrenie*, 91(3): 157–161.
- Huang J. and Bourgoin T. 1993.** The planthopper genus Trypetimorpha: systematics and phylogenetic relationships (Hemiptera: Fulgoromorpha: Tropiduchidae). *Journal of Natural History*, 27(3): 609–629.
- International Code of Zoological Nomenclature.** 1999. Fourth Edition. The International Trust for Zoological Nomenclature c/o the Natural History Museum, London, 306 p.
- Jacobi A. 1941.** Die Zikadenfauna der Kleinen Sundainseln. Nach der Expeditionsausbeute von B. Rensch. *Zoologische Jahrbücher. Abteilung für Systematik, Ökologie und Geographie der Tiere*, 74: 277–322.
- Jacobi A. 1944.** Die Zikadenfauna der Provinz Fukien in Südchina und ihre tiergeographischen Beziehungen. *Mitteilungen der Münchener Entomologischen Gesellschaft*, 34: 5–66.
- Lallemand H. 1942.** Notes sur quelques espèces recueillies par le R. Piel (Musée Haude Shanghai) et le R.P. de Cooman (Hoa Binh, Tonkin). *Notes d'Entomologie Chinoise*, 9(4): 69–77.
- Lallemand H. and Synave H. 1953.** Homoptères de Sumba et Florès. *Verhandlungen der Naturforschenden Gesellschaft in Basel*, 64(1–2): 229–254.
- Lampe K.-H., Rohwedder D. and Rach B. 2006.** Insect types in the ZFMK collection, Bonn: "Homoptera" (<http://www.zfmk.de/web/Forschung/Sammlungen/index.de.html>), last update: April 2006.
- Matsumura S. 1900.** Uebersicht der Fulgoriden Japans. *Entomologische Nachrichten*, 13: 205–213.
- Matsumura S. 1914.** Beitrag zur Kenntnis der Fulgoriden Japans. *Annales Musei Nationalis Hungarici*, 12: 261–305.
- Melichar L. 1902.** Homopteren aus West-China, Persien und dem Süd-Ussuri-Gebiete gesammelt von Potanin, Berezovski, Zarudny und Jankovski. *Annuaire du Musée Zoologique de l'Académie Impériale des Sciences de St. Pétersbourg*, 7: 76–146.
- Melichar L. 1906.** Monographie der Issiden (Homoptera). *Abhandlungen der K.K. Zoologisch-botanischen Gesellschaft in Wien*, 3(4): 1–327.
- Melichar L. 1907.** Bemerkungen zur Monographie der Issiden (Homoptera). *Wiener Entomologische Zeitung*, 26: 323–324.
- Melichar L. 1911.** Collections recueillies par M. M. de Rothschild dans l'Afrique Orientale. Homoptères. *Bulletin du Muséum National d'Histoire Naturelle*, 17: 106–117.
- Metcalf Z.P. 1952.** New names in the Homoptera. *Journal of the Washington Academy of Sciences*, 42(7): 226–231.
- Metcal Z.P. 1958.** General catalogue of the Homoptera. Fulgoroidea. Issidae. Vol. 4(15). Waverly Press, INC, Baltimore, 561 p.

- Rahman M.A., Kwon Y.J. and Suh S.J.** 2011. Taxonomic revision of family Tropiduchidae (Hemiptera: Fulgoromorpha) from Korea. *Entomological Research*, **41**(6): 247–256.
- Robineau-Desvoidy A.J.B.** 1830. Essai sur les Myodaires. *Mémoires présentés par divers savants à l'Académie royale des Sciences de l'Institut de France (sciences mathématiques et physiques)* **2**(2): 1–813.
- Schmidt E.** 1926. Fauna Buruana. Homoptera. *Treubia*, **7**: 217–258.

- Synave H.** 1956. Contribution a l'Etude des Issidae Africains (Homoptera–Fulgoroidea). *Bulletin de l'Institut royal des Sciences naturelles de Belgique*, **32**(57): 1–22.
- Vilbaste Y.** 1968. To the fauna of Cicadina of Primorsky Territory. Valgus, Tallin, 180 p.
- Wang R.R. and Liang A.P.** 2007. Taxonomic study of the genus *Zema* Fennah (Hemiptera: Fulgoromorpha: Tropiduchidae) from China. *Zootaxa*, **1436**: 61–68.

Submitted April 23, 2012; accepted August 30, 2012.