

Records of weevils new to the Russian fauna (Coleoptera: Curculionidae)

B.A. Korotyaev

Korotyaev, B.A. 1999. Records of weevils new to the Russian fauna (Coleoptera: Curculionidae). *Zoosystematica Rossica*, 8(1): 174.

7 species of weevils are recorded from Russia for the first time.

B.A. Korotyaev, Zoological Institute, Russian Academy of Sciences, Universitetskaya nab. 1, St.Petersburg 199034, Russia.

Magdalis (Odontomagdalis) gurjevae Barrios, 1984

Material. 1 ♂, Primorsk Terr., Ussuriysk, Gorno-tayozhnoye, on elm, 6.VI.1996 (M.I. Falkovitsh).

Distribution. E and SE Mongolia.

Magdalis (Aika) margaritae Barrios, 1984

Material. 1 ♀, 1 ♂, Primorsk Terr., Spassk, 8.VI.1989 and 30.VI.-6.VII.1995 (S.A. Belokobylskij).

Distribution. E Mongolia.

Cardipennis shaowuensis Voss, 1958

Material. 1 ♀, Primorsk Terr., 20 km SW of Putsilovka, Monakino vill., clearings in the forest, 24-28.VI.1993 (S.A. Belokobylskij).

Distribution. Japan, N Korea, E China.

Ceutorhynchus simulans Korotyaev, 1980

Material. 1 ♀, S Tuva, Ovyur Distr., valley of the Irbiti River N of Lake Ubsunur, 13-16.VI.1995 (Yu.M. Marusik).

Distribution. Mongolia.

Rhynchaenus? stobieckyi Smreczynski, 1957

Material. 1 ♀, Kursk Prov., Central Chernozem Reserve, Kazatskiy Meadow, southern slope of the Barybin Gully, 29.VI.1957 (K.V. Arnoldi); 1 ♀, Tambov Prov., Michurinsk (coll. Solsky); 1 ♀, Bashkiria, 1978 (G.V. Miländer).

Distribution. West Ukraine, East Ukraine (new record); 2 ♂, 1 ♀, Lugansk prov., Provalye, 13-14.VI.-1929 (V.I. Talitzky); Turkey (new record); 1 ♂, Ispir Prov., Koprubasi, 6.VI.1980 (Ozisik).

The systematic position of this species is uncertain. All tibiae in both sexes are not mucronate, and this makes it similar to the Oriental genus *Indodinorrhopalus* Pajni & Sood with 3 species (Kojima & Morimoto, 1996). The host plant is unknown; the occurrence of the species in the steppe reserve may indicate its association with herbs rather than with trees.

Pseudorchestes ermischii Dieckmann, 1958

Material. 1 ♂, Kursk Prov., Central Black Soil Reserve, 1957 (K.V. Arnoldi); 17 spms., Stavropol Terr., 2 km E of Novoaleksandrovsk, steppe slope on right bank of the Rashevatska River upstream of the Vinogradnyi vill., on *Centaurea orientalis* L., 10-11.VI.1996 (Korotyaev).

Distribution. Western, Central, and Eastern Europe. The species was previously known east to West Ukraine. In Europe, only *Centaurea scabiosa* L. is known as host plant.

Trachyphloeus caenopsisformis Formánek, 1908

Material. 2 ♀, NW Caucasus, Krasnodar Terr., Taman' peninsula, steppe near Taman' vill., 27-28.IV.1980 (Korotyaev).

Distribution. Turkey. The holotype from the Prague Museum has been examined.

Acknowledgements

I am grateful to Prof. H. Özbek and L. Gültekin of the Atatürk University, Erzurum, Turkey, for the opportunity to examine the specimen of *Rhynchaenus stobieckyi* from their collection; Dr. J. Jelinek (Prague) for the loan of the holotype of *Trachyphloeus caenopsisformis*; M.A. Korotyaev, T.V. Nogina and V.A. Sergeyev (Novoaleksandrovsk) for their help in my field research; and to S.A. Belokobylskij, M.I. Falkovitsh (St.Petersburg), and Yu.M. Marusik (Magadan) for collecting the material reported here. The investigation was supported by grant No. 98-04-49763 from the Russian Foundation for Basic Research.

References

- Kojima, H. & Morimoto, K. 1996. Systematics of the flea weevils of the tribe Ramphini (Coleoptera, Curculionidae) from East Asia II. Phylogenetic analysis and higher classification. *Esakia*, 36: 97-134.
Received 18 January 1999