

First record of *Sejanus juglandis* Yas. from Russia and note on *S. interruptus* Reut. (Heteroptera: Miridae)

N.N. Vinokurov

Vinokurov, N.N. 2006. First record of *Sejanus juglandis* Yas. from Russia and note on *S. interruptus* Reut. (Heteroptera: Miridae). *Zoosystematica Rossica*, **15**(1): 56.

Sejanus juglandis Yas. is found in the Far East of Russia (Primorsk Terr.) for the first time. It is compared with *S. potanini* Dist. and *S. interruptus* Reut. Vesicae of *S. juglandis* and *S. interruptus* are figured.

N.N. Vinokurov, Institute for Biological Problems of Cryolithozone, Siberian Division, Russian Academy of Sciences, ul. Lenina 41, Yakutsk 677980, Russia.

Capsid bugs of the genus *Sejanus* Dist. (Phyllinae) are distributed in SE Asia, Australia, New Zealand and Pacific Islands (Schuh, 1984; Yasunaga, 2001). Hitherto, only a single species, *S. potanini* Reut., was recorded from Russia (Kerzhner, 1988). An additional species, *S. juglandis*, described from Japan (Yasunaga, 2001), is found in the collections of Zoological Institute, St.Petersburg, and Lazo Nature Reserve (Primorsk Terr.).

Sejanus juglandis Yasunaga, 2001 (Figs 1-3)

Material. **Russia, Primorsk Terr.:** 1 ♂, Kedrovaya Pad' Nature Reserve, 11.VIII.1985 (Sinev); 1 ♂, Lozovoy Range, 22 km NE Nakhodka, 21.VII.1999 (Sinev); 1 ♀, Lazo Nature Reserve, mouth of the Kievka R., Chekhunenko Lake, 4.VIII.2005 (Yu. Sundukov).

Remarks. The species differs from *S. potanini* in the presence of a white, sometimes interrupted band at the base of cuneus (Figs 1, 2). A similar band is present also in *S. interruptus* Reut. from China (Sichuan) (Fig. 4), but the two species are clearly distinguished by the shapes of the vesicae (Figs 3, 5).

Sejanus interruptus (Reuter, 1906) (Figs 4, 5)

The type series of this species has been examined. The figure of vesica is made from the lectotype.


Acknowledgements

The work was carried out with support of grant no. 05-04-49917 from the Russian Foundation for Basic Research.

References

- Kerzhner, I.M.** 1988. Fam. Miridae. In: Lehr, P.A. (Ed.). *Opredelitel' nasekomykh Dalnego Vostoka SSSR* [Keys to the insects of the Far East of the USSR], **2**: 778-857. Leningrad: Nauka.
- Schuh, R.T.** 1984. Revision of the Phyllinae (Hemiptera, Miridae) of the Indo-Pacific. *Bull. Amer. Mus. natur. Hist.*, **177**: 1-462.
- Yasunaga, T.** 2001. A review of the phylline plant bug genus *Sejanus* Distant in Japan (Heteroptera: Miridae: Phyllinae), with descriptions of three new species. *Entomol. Sci.*, **4**(1): 121-126.

Received 18 September 2006



Figs 1-5. 1-3, *Sejanus juglandis*; 4, 5, *S. interruptus*. 1, 2, 4, cuneus; 3, 5, vesica.