

A. S. Lelej
List of Palaearctic Mutilidae (Hymenoptera)

This list based on **Lelej A. S. Catalogue of the Mutilidae (Hymenoptera) of the Palaearctic region.** – Vladivostok: Dalnauka, 2002. – 171 p. with some additions and corrections.

Explication to distribution			
EUROPE (excluding Russia)			
A	– Austria	IRE	– Ireland
AL	– Albania	LA	– Latvia
B	– Belgium	LI	– Lithuania
BE	– Belarus	M	– Malta
BG	– Bulgaria	MO	– Moldova
BH	– Bosnia–Herzegovina	N	– Norway
CH	– Switzerland	NL	– Netherlands
CZ	– Czechia	P	– Portugal
D	– Germany	PL	– Poland
DK	– Denmark	R	– Romania
E	– Spain	S	– Sweden
EST	– Estonia	SF	– Finland
F	– France	SL	– Slovakia
GB	– United Kingdom	TR	– Turkey (European part)
GR	– Greece	U	– Ukraine
H	– Hungary	YU	– Yugoslavia
I	– Italy		

Other European countries are given without abbreviation: Croatia, Macedonia, Slovenia.

RUSSIA

NET	– North European territory, north of 60° North Latitude
CET	– Central European territory, limited by 60° North Latitude in the north and by 52° in the south
SET	– South European territory, south of 52° North Latitude
WS	– Western Siberia, limited by watershed of the Urals in the west and by the river Yenisei in the east
ES	– East Siberia, limited by the river Yenisei in the west and by the border of Magadanskaya oblast, Khabarovskii krai, and Amurskaya oblast in the east
FE	– Far East, limited by the border of Magadanskaya oblast, Khabarovskii krai, and Amurskaya oblast in the west

ASIA (excluding Russia)

ARM	– Armenia
AZ	– Azerbaijan
CY	– Cyprus
GE	– Georgia

KY	– Kyrgyzstan
KZ	– Kazakhstan (including European part)
TJ	– Tajikistan
TR	– Turkey (Asian part)
TU	– Turkmenistan
UZ	– Uzbekistan

Other Asian are given without abbreviation: Arabic states (North of the Tropic of Cancer), Lebanon, Syria, Jordan*, Israel*, Oman, Iraq, Iran, Afghanistan, Mongolia, Korea, China North of 30° North Latitude, Japan North of 30° North Latitude.

NORTH AFRICA (North of tropic Cancer)
Canary Is., Western Sahara, Morocco, Algeria, Tunisia, Libya, Egypt

*As type localities or distribution points, the name Palestina (Palestine) is written without any attributes. When, in the case of actual locality names, it can be ascertained that they lie in Israel or in Jordan, they are listed under the name of the respective country. The same consideration applies to Jerusalem.

Table 1
**Distribution of 968 nominal specific and infraspecific names
of Palaearctic Mutilidae among the subfamilies**

Subfamilies and tribes	Synonyms and invalid names	Valid names			Total
		Varieties and forms	Subspecies	Species	
I. Myrmosinae	13	1	1	31	46
II. Kudakrumiinae	2	0	0	13	15
III. Pseudophotopsidinae	14	0	0	24	38
IV. Ticoplinae	0	0	1	6	7
V. Myrmillinae	58	3	1	68	130
VI. Mutilinae	190	33	19	256	498
Mutilini	59	7	4	31	101
Smicromyrmini	109	24	14	191	338
Petersenidiini	4	0	0	5	9
Trogaspidiini	18	2	1	29	50
VII. Sphaeropthalminae	2	0	0	2	4
VIII. Dasylabrinae	88	10	9	103	210
IX. Ephutinae	0	0	0	3	3
X. Mutilidae incertae sedis	0	0	0	17	17
Total		367	47	31	523
					968

Family MUTILLIDAE Latreille, 1802

I. Subfamily MYRMOSINAE Fox, 1894

1. Genus *Carinomyrmosa* Lelej, 1981

gussakovskiji Lelej, 1981 (*Carinomyrmosa*). – Asia: TJ.

2. Genus *Erimyrmosa* Lelej, 1984

frater (E. Saunders, 1899) (*Mutilla*). – North Africa: Algeria, Tunisia.

3. Genus *Krombeinella* Pate, 1947

Paramyrmosa Suárez, 1963.

aramaea (Suárez, 1963) (*Myrmosa*). – Asia: Syria, Israel, TR, Iran.

atterima (Suárez, 1959) (*Myrmosa*). – Europe: F, E, P.

iberica (Suárez, 1963) (*Myrmosa*).

baetica Suárez, 1981 (*Krombeinella*). – Europe: E.

beaumonti (Invrea, 1953) (*Myrmosa*). – Europe: R, I (Sicily) E, P; North Africa: Morocco.

scutellata (Invrea, 1953) (*Myrmosa*).

hebraea (Suárez, 1963) (*Myrmosa*). – Asia: Israel, TR.

longicollis (Tournier, 1889) (*Myrmosa*). – Europe: SL, R, BG, Croatia, YU, Macedonia, GR, TR, U; Russia: SET.

maroccana (Suárez, 1960) (*Myrmosa*). – North Africa: Morocco.

naeffi (Invrea, 1952) (*Myrmosa*). – North Africa: Algeria.

nigriceps (S.S. Saunders, 1850) (*Myrmosa*). – Europe: AL, YU, GR, TR.

gaullei (Invrea, 1950) (*Myrmosa*). – Europe: TR.

obscuripes (Tournier, 1889) (*Myrmosa*). – Europe: E; North Africa: Morocco.

spinolae (Lepeletier, 1845) (*Myrmosa*). – Europe: I.

tangerina (Mercet, 1913) (*Myrmosa*). – North Africa: Morocco.

tenietensis (André, 1900) (*Myrmosa*). – North Africa: Algeria, Tunisia.

thoracica (Fabricius, 1793) (*Hyleus*). – Europe: I, F, S.

dubia (Costa, 1858) (*Myrmosa*).

thracia (Suárez, 1963) (*Myrmosa*). – Europe: BG.

wolfi (Invrea, 1963) (*Myrmosa*). – Europe: YU, BG, U; Russia: SET; Asia: TR, GE, AZ.

4. Genus *Myrmosa* Latreille, 1796

Ischioceras Provancher, 1882.

atra Panzer, 1801 (*Myrmosa*). – Europe: S, SF, GB, IRE, B, NL, D, PL, EST, BE, U, A, CZ, SL, H, CH, F, E, P, I; Slovenia, Croatia, BH, YU,

-
- R; Russia: NET, CET, SET, WS, ES (Yakutsk); Asia: TR, KZ, KY, ? Iran.
- melanocephala* (Fabricius, 1793) (*Mutilla*).
nigra Latreille, 1809 (*Myrmosa*).
atra Lepeletier, 1845 (*Myrmosa*).
nigra Lepeletier, 1845 (*Myrmosa*).
bicolor Baer, 1848 (*Myrmosa*).
melanaria (Blanchard, 1849) (*Mutilla*).
rufomaculata Hoffer, 1938 (*Myrmosa*).
erythrocephala Yarrow, 1954 (*Myrmosa*). – Europe: IRE.
eos Lelej, 1981 (*Myrmosa*). – Russia: FE; Asia: Korea.
moesica Suárez, 1981 (*Myrmosa*). – Europe: YU.
mongolica Suárez, 1974 (*Myrmosa*). – Russia: ES (Irkutskaya oblast, Transbaicalia), FE (Amurskaya oblast, Khabarovskii krai, Primorskii krai, Sakhalin); Asia: Mongolia, China: NE (Heilongjiang), Korea, Japan (Honshu).
tsunekii Haneda, 1979 (*Myrmosa*).
nigrofasciata Yasumatsu, 1931 (*Myrmosa*). – Asia: Japan (Honshu, Kyushu).

5. Genus **Paramyrmosa** Saussure, 1880

- Morysmula* Suárez, 1963.
- bischoffi** (Invrea, 1952) (*Myrmosa*). – Europe: GR.
brunnipes (Lepeletier, 1845) (*Myrmosa*). – Europe: D, PL, BE, U, CZ, SL, A, H, R, BG, GR, AL, Slovenia, Croatia, BH, YU, Macedonia, I, F, E, P; Russia: CET, SET; Asia: ARM, TR.
cognata (Costa, 1858) (*Myrmosa*).
hispanica Suárez, 1980: (*Paramyrmosa*). – Europe: E.
myrmilicephala (Skorikov, 1935) (*Myrmosa*). – Asia: TU, UZ, TJ, Mongolia.
pulla (Nylander, 1847) (*Myrmosa*). – Russia: ES (Transbaicalia), FE (Amurskaya oblast, Evreyskaya oblast, Primorskii krai); Asia: Mongolia, China: NE.
czerskii (Skorikov, 1929) (*Myrmosa*).
punica Suárez, 1979 (*Paramyrmosa*). – North Africa: Tunisia.
radoszkowskyi (Saussure, 1880) (*Myrmosa*). – Asia: TR, GE, AZ, ARM.
sbuaensis (Giner, 1945) (*Myrmosa*). – North Africa: Western Sahara.

II. Subfamily KUDAKRUMIINAE Krombein, 1979

6. Genus **Protomutilla** Bischoff, 1916

- archaeotrichosoma** Lelej, 1986 (fossil) (*Protomutilla*).

- castanea** Bischoff, 1916 (fossil) (*Protomutilla*).
dentata Bischoff, 1916 (fossil) (*Protomutilla*).
incerta Bischoff, 1916 (fossil) (*Protomutilla*).
megalophtalma Bischoff, 1916 (fossil) (*Protomutilla*).
nana Bischoff, 1916 (fossil) (*Protomutilla*).
rASNITSYNI Lelej, 1986 (fossil) (*Protomutilla*).
succinalis Bischoff, 1916 (fossil) (*Protomutilla*).
succinicola Bischoff, 1916 (fossil) (*Protomutilla*).

7. Genus *Pseudomyrmosa* Suárez, 1980

III. Subfamily PSEUDOPHOTOPSISINAE Bischoff, 1920

8. Genus *Pseudophotopsis* André, 1896

- Alloneurion** Ashmead, 1899.
Sphinctomutilla André, 1899.

aegyptiaca (Bischoff, 1920) (*Ephutomma*). – Asia: Israel; North Africa: Egypt, Libya, Sudan, Chad.

afghanica Lelej, 1980 (*Pseudophotopsis*). – Asia: Afghanistan.

armeniaca (Skorikov, 1935) (*Ephutomma*). – Asia: CY, ARM.

cypriaca (Hammer, 1950) (*Ephutomma*).

baktriana (Skorikov, 1935) (*Ephutomma*). – Asia: TU, TJ.

binghami (Bischoff, 1920) (*Pseudophotopsis*). – Asia: KZ, UZ, TJ, TU, Iran, Afghanistan, India (Rajasthan).

zarudnyi (Skorikov, 1935) (*Pseudophotopsis*).

biskrensis (André, 1900) (*Mutilla*). – North Africa: Algeria, Tunisia, Libya, Egypt

brachythorax (Skorikov, 1935) (*Ephutomma*). – Asia: TU, TJ, East Iran, Afghanistan.

brunnescens Lelej, 1971 (*Pseudophotopsis*). – Asia: KZ (south-east), KY, UZ.

caspica (Radoszkowski, 1885) (*Agama*). – Asia: KZ (south), TU, UZ, KY, TR, Iran, Afghanistan.

radoszkowskyi (Morawitz, 1890) (*Mutilla*).

selevkida (Skorikov, 1935) (*Ephutomma*).

-
- caucasica** (Radoszkowski, 1885) (*Agama*). – Asia: GE, AZ, ARM, TU, CY, Palestine, TR, Iran, Afghanistan.
caucasicola (Dalla Torre, 1897) (*Mutilla*).
schachruda (Skorikov, 1935) (*Ephutomma*).
robusta (Skorikov, 1935) (*Ephutomma*).
mavromoustakisi (Suárez, 1959) (*Ephutomma*).
fumata Bischoff, 1920 (*Pseudophotopsis*). – Asia: Israel; North Africa: Western Sahara, Tunisia, Libya, Chad.
geyri (Bischoff, 1920) (*Ephutomma*). – North Africa: Algeria, Sudan.
irania (Skorikov, 1935) (*Ephutomma*). – Asia: KZ (south), UZ, TU, TJ, Iran, Afghanistan, India (Rajasthan).
cithara (Skorikov, 1935) (*Ephutomma*).
kabakovii Lelej, 1980 (*Pseudophotopsis*). – Asia: Afghanistan
kermania Skorikov, 1935 (*Pseudophotopsis*). – Asia: Iran.
kokpetica (Radoszkowski, 1885) (*Agama*). – Asia: KZ (south), UZ, KY, TU, TJ.
komarovii (Radoszkowski, 1885) (*Agama*). – Asia: KZ (south), UZ, TU, TJ, CY, Iran; ? North Africa: Algeria, Egypt.
askhabadensis (Radoszkowski, 1885).
oculosa Skorikov, 1935 (*Pseudophotopsis*).
inductrix Skorikov, 1935 (*Pseudophotopsis*).
asiatica Suárez, 1965 (*Pseudophotopsis*).
mateui Giner, 1945 (*Pseudophotopsis*). – North Africa: Western Sahara.
maura Bischoff, 1920 (*Pseudophotopsis*). – Asia: Israel; North Africa: Algeria, Tunisia, Libya, Egypt, Chad.
obliterata (Smith, 1855) (*Mutilla*). – Europe: AL; Asia: CY.
orthophtalma (Skorikov, 1935) (*Ephutomma*). – Asia: KZ (south), UZ, TU, TJ.
pavlovskyi Skorikov, 1935 (*Pseudophotopsis*). – Asia: KY, UZ, TU, TJ, Iran (north), Afghanistan.
melanocephala Skorikov, 1935 (*Pseudophotopsis*).
rufothoracica Bischoff, 1920 (*Pseudophotopsis*). – North Africa: Egypt.
syriaca (André, 1900) (*Mutilla*). – Europe: GR Asia: Syria, Palestine.

IV. Subfamily TICOPLINAE Nagy, 1970

Nanomutillinae Suárez, 1975.

Tribe Ticoplini Nagy, 1970

9. Genus *Nanomutilla* André, 1900

Ticopla Nagy, 1970.

- nada** Argaman, 1988 (*Nanomutilla*). – Europe: E (south).
parila (Nagy, 1970) (*Ticopla*). – Asia: Jordan, Israel.

vaucheri (Tournier, 1895) (*Mutilla*). – Europe: E; Asia: Syria; North Africa: Morocco, Algeria.
yoca (Nagy, 1970) (*Ticopla*). – Asia: Jordan, Israel.

Tribe Smicromyrmillini Argaman, 1988

10. Genus *Smicromyrmilla* Suárez, 1965

ariasi (André, 1896) (*Mutilla*). – Europe: E; North Africa: Morocco.
garciapavoni Suárez, 1975 (*Smicromyrmilla*). – North Africa: Morocco.
miranda Nonveiller et Gros, 1996 (*Smicromyrmilla*). – Europe: E.

V. Subfamily MYRMILLINAE Bischoff, 1920

11. Genus *Bischoffitilla* Lelej, 2002

afghanica (Lelej, 1980) (*Myrmilla*). – Asia: Afghanistan.
ardescens (Smith, 1873) (*Mutilla*). – Asia: Japan (Honshu, Kyushu, Amami-oshima).
bidentata (Tsuneki, 1972) (*Squamulotilla*). – Asia: Japan (Honshu, Yaku-shima).
exilipunctata (Chen, 1957) (*Squamulotilla*). – Asia: China: NO (Hebei), CE (Jiangsu, Zhejiang), SE (Fujian), Korea.
pungens (Smith, 1873) (*Mutilla*). – Asia: Japan (Honshu, Kyushu).
 pungens (Smith, 1874) (*Mutilla*).
 militans (Dalla Torre, 1897) (*Mutilla*).
strangulata (Smith, 1879) (*Mutilla*). – Asia: China: CE (Jiangxi, Jiangsu, Anhui, Zhejiang), SE (Fujian, Guangdong).
unidentata (Tsuneki, 1972) (*Squamulotilla*). – Asia: Japan (Honshu).

12. Genus *Blakeius* Ashmead, 1903

Bisigilla Skorikov, 1927:

bipunctatus (Latreille, 1792) (*Mutilla*). – Europe: E, F, I, GR; Asia: CY, Syria; North Africa: Morocco, Algeria, Tunisia.
angusticollis (Spinola, 1843) (*Mutilla*).
tuberculata (Lepeletier 1845) (*Mutilla*).
bituberculata (Smith, 1855) (*Mutilla*).
erythrothorax (Sichel et Radoszkowski, 1870) (*Mutilla*).
cobosi (Suárez, 1958) (*Myrmilla*). – North Africa: Morocco.
chiesi (Spinola, 1839) (*Mutilla*). – Europe: E, P, F (Corsica), I; North Africa: Morocco, Algeria, Tunisia.
diophthalma (Costa, 1882) (*Mutilla*).
ibericus (Suárez, 1958) (*Myrmilla*). – Europe: E, P, I; North Africa:

Morocco, ? Algeria, Tunisia.

negrei (Suárez, 1958) (*Myrmilla*). – Europe: E, P, I; North Africa: Morocco, Algeria.

obscura (Suárez, 1958) (*Myrmilla*).

rufonota (Suárez, 1958) (*Myrmilla*).

nana (Suárez, 1958) (*Myrmilla*).

invreai (Suárez, 1958) (*Myrmilla*). – Europe: E; North Africa: Morocco.

leopoldinus (Invrea, 1955) (*Myrmilla*). – Europe: I (Sardinia), M.

ortizi (Suárez, 1954) (*Myrmilla*). – Europe: E; North Africa: Morocco.

otini (Giner, 1959) (*Myrmilla*).

13. Genus *Liomutilla* André, 1907

canariensis André, 1907 (*Liomutilla*). – North Africa: Canary Is.

14. Genus *Myrmilla* Wesmael, 1852

Rudia Costa, 1858.

Subgenus *Eurygnathilla* Skorikov, 1927

emiliae (Dalla Torre, 1897) (*Mutilla*). – Asia: AZ, ARM, TU, TR, Iraq, Iran.

laticeps (Radoszkowski, 1888) (*Mutilla*).

chusistana Skorikov, 1927 (*Myrmilla*).

ephutommatica Skorikov, 1927 (*Myrmilla*). – Asia: TU, UZ, TJ.

Subgenus *Myrmilla* Wesmael, 1852

anopla Skorikov, 1927 (*Myrmilla*). – Europe: YU.

astrabada Skorikov, 1927 (*Myrmilla*). – Asia: Iran.

bilobicornis Skorikov, 1927 (*Myrmilla*). – Asia: TU, Iran.

calva (Villers, 1789) (*Mutilla*). – Europe: A, CZ, SL, H, D, CH, R, AL, Slovenia, Croatia, BH, YU, Macedonia, BG, GR, I, F, E, P; Asia: TR; North Africa: Morocco, Tunisia, Libya, Egypt.

calva (Fabricius, 1798) (*Mutilla*).

triareolata (Spinola, 1841) (*Mutilla*).

distincta (Lepeletier, 1845) (*Mutilla*).

agrestis (Lepeletier, 1845) (*Mutilla*).

incompleta (Lepeletier, 1845) (*Mutilla*).

hastata (Costa, 1858) (*Rudia*).

andalusiana Skorikov, 1927 (*Myrmilla*).

seminigra Hoffer, 1938 (*Myrmilla*).

perparvula Invrea, 1958 (*Myrmilla*).

semirubra Nonveiller, 1972 (*Myrmilla*).

-
- caucasica** (Kolenati, 1846) (*Mutilla*). – Europe: U, GR, BG; Russia: SET; Asia: GE, AZ, ARM, TR, Iran.
jasonia Skorikov, 1927 (*Myrmilla*).
kiritschenkoi Skorikov, 1927 (*Myrmilla*).
corniculata (Sichel et Radoszkowski, 1869) (*Mutilla*). – Europe: ?F (south), GR;
 Asia: Syria, Palestine, TR, Iran.
corniculata (Gerstaecker, 1862) (*Mutilla*).
corniculatina Skorikov, 1927 (*Myrmilla*).
doumerguei (André, 1901) (*Mutilla*). – Europe: ?R; North Africa: Algeria.
edentata Skorikov, 1927 (*Myrmilla*). – Asia: KZ (south), UZ, TJ.
kumakana Skorikov, 1935 (*Myrmilla*).
medoides Skorikov, 1935 (*Myrmilla*).
erythrocephala (Latreille, 1792) (*Mutilla*). – Europe: A, CZ, SL, H, Croatia,
 YU, Macedonia, BG, GR, I, F (south), E, P; Asia: CY, Lebanon,
 Palestine; North Africa: Tunisia, Egypt.
latreillei (Meyer, 1794) (*Mutilla*).
cornuta (Olivier, 1811) (*Mutilla*).
megacephala (Costa, 1858) (*Rudia*).
bison (Costa, 1887) (*Mutilla*).
etzchmiadzini (Radoszkowski, 1885) (*Mutilla*). – Asia: AZ, ARM.
armeniaca Skorikov, 1927 (*Myrmilla*).
jakobsoni Skorikov, 1927 (*Myrmilla*). – Asia: KZ (south), UZ.
longicapitata Skorikov, 1935 (*Myrmilla*).
kabulensis Lelej, 1980 (*Myrmilla*). – Asia: Afghanistan.
labcuca Nagy, 1968 (*Myrmilla*). – Europe: R.
lezginica (Radoszkowski, 1885) (*Mutilla*). – Europe: GR, YU, Macedonia, BG,
 R, U; Russia: SET; Asia: GE, AZ, UZ, TU, Syria, Palestine, TR,
 Iran.
subcornuta (Morawitz, 1893) (*Mutilla*).
nigrifrons (André, 1901) (*Mutilla*). – Asia: Syria.
kokujevi Skorikov, 1927 (*Myrmilla*).
susiana Skorikov, 1927 (*Myrmilla*).
macrura Nagy, 1968 (*Myrmilla*). – Europe: R.
menozzii Invrea, 1950 (*Myrmilla*). – Asia: Palestine.
mutica (André, 1903) (*Mutilla*). – Europe: R, Croatia, BH, YU, Macedonia, GR,
 I; North Africa: Egypt.
ecornuta Skorikov, 1927 (*Myrmilla*).
hamata Skorikov, 1927 (*Myrmilla*).
propodealis Skorikov, 1935 (*Myrmilla*). – Asia: KZ (south-east), KY, China:
 NW (Xinjiang).
shnitnikovi Skorikov, 1935 (*Myrmilla*).
rostriformis Lelej, 1980 (*Myrmilla*). – Asia: Afghanistan (east).
saharae (Cockerell, 1907) (*Mutilla*). – North Africa: Egypt.

-
- fasciata** (Klug, 1829) (*Mutilla*).
sarta Skorikov, 1927 (*Myrmilla*). – Asia: KZ (south), TU, UZ, TJ, Mongolia.
schmiedeknechti (André, 1903) (*Mutilla*). – North Africa: Egypt.
tadzhikistanica Lelej, 1985 (*Myrmilla*). – Asia: KY, TJ.
variolosa Skorikov, 1935 (*Myrmilla*). – Asia: UZ, TJ.
zarudnyi Skorikov, 1927 (*Myrmilla*). – Asia: Iran, India (Rajasthan).

Subgenus **Pseudomutilla** Costa, 1885

Edrionotus Radoszkowski, 1885.

- atalanta** Nagy, 1967 (*Myrmilla*). – Europe: R; Asia: TR.
badchysiana Lelej, 1980 (*Myrmilla*). – Asia: TU, TJ, Afghanistan.
bytinskii Invrea, 1965 (*Myrmilla*). – Asia: Palestine.
capitata (Lucas, 1849) (*Mutilla*). – Europe: H, A, YU, BG, GR, I, F (south), E, P; Asia: Syria; North Africa: Morocco, Algeria, Tunisia, Libya, Egypt.
parvicollis (Costa, 1860) (*Mutilla*).
sardiniensis (Costa, 1885) (*Pseudomutilla*).
olcesei (Tournier, 1895) (*Mutilla*).
robustior (André, 1901) (*Mutilla*). – North Africa: Algeria.
cyrenaica Invrea, 1950 (*Myrmilla*). – North Africa: Libya.
filippovi Lelej, 1985 (*Myrmilla*). – Asia: TR.
glabrata (Fabricius, 1775) (*Mutilla*). – Europe: A, H, R, BG, YU, Macedonia, GR, U; Russia: SET; Asia: KZ, UZ, TR, Syria.
ciliata (Fabricius, 1793) (*Mutilla*).
fraterna (Baer, 1848) (*Mutilla*).
cephalica (Sichel et Radoszkowski, 1869) (*Mutilla*).
sarmatica Skorikov, 1927 (*Myrmilla*).
belgovskii Skorikov, 1927 (*Myrmilla*).
kochi Invrea, 1950 (*Myrmilla*). – North Africa: Libya.
lagmana Lelej, 1980 (*Myrmilla*). – Asia: Afghanistan.
mateui Giner, 1945 (*Myrmilla*). – North Africa: Western Sahara .
mauritanica Invrea, 1953 (*Myrmilla*). – North Africa: Morocco.
mavromoustakisi Hammer, 1950 (*Myrmilla*). – Asia: CY.
meda Skorikov, 1927 (*Myrmilla*). – Russia: SET (Daghestan); Asia: AZ, TU, Iran (north).
appropinqua Skorikov, 1927 (*Myrmilla*).
moralesi Giner, 1945 (*Myrmilla*). – North Africa: Western Sahara.
praetermissa Skorikov, 1927 (*Myrmilla*). – North Africa: Algeria, Tunisia.
skorikovi Lelej, 1985 (*Myrmilla*). – Russia: SET Asia: AZ, GE.
caucasica Skorikov, 1927 (*Myrmilla*).
tenuitruncata Skorikov, 1935 (*Myrmilla*). – Asia: TJ.
troodosica Hammer, 1950 (*Myrmilla*). – Asia: CY.

vutshetitschi Skorikov, 1927 (*Myrmilla*). – Europe: A, R, BG, GR, U; Asia: TR, Syria.
kusdasi Invrea, 1965 (*Myrmilla*).

15. Genus **Myrmotilla** Bischoff, 1920

lucasiella Lelej, 2002. – North Africa: Algeria, Tunisia.
lucasi (Sichel et Radoszkowski, 1870) (*Mutilla*).

16. Genus **Odontotilla** Bischoff, 1920

Subgenus **Conjunctotilla** Nonveiller, 1996

con juncta (Klug, 1829) (*Mutilla*). – Asia: Palestine, Arabia, Yemen; Africa: Egypt, Sudan, Chad, Ethiopia, Eritrea, Niger, Mauritania.

17. Genus **Platymyrmilla** André, 1900

quinquefasciata (Olivier, 1811) (*Mutilla*). – Europe: H, R, Macedonia, GR, BG, MO; Asia: GE, ARM, AZ, Palestine, Syria, TR, Iran, Iraq.

18. Genus **Sigilla** Skorikov, 1927

angelae Suárez, 1958 (*Sigilla*). – North Africa: Algeria.
dorsata (Fabricius, 1798) (*Mutilla*). – Europe: I, Monaco, F (south), E; North Africa: Morocco, Algeria, Tunisia, Libya, Egypt.
excoriata (Lepeletier, 1845) (*Mutilla*).
spinolae (Lepeletier, 1845) (*Mutilla*).
punctum (Lepeletier, 1845) (*Mutilla*).
affinis (Lucas, 1849) (*Mutilla*).
cognata (Smith, 1855) (*Mutilla*).
calcariventris (Sichel et Radoszkowski, 1870) (*Mutilla*).

19. Genus **Spilomutilla** Ashmead, 1903

perfecta (Sichel et Radoszkowski, 1870) (*Mutilla*). – Asia: Iran (south).

VI. Subfamily MUTILLINAE Latreille, 1802

Tribe Mutillini Latreille, 1802

20. Genus *Ctenotilla* Bischoff, 1920

caeca (Radoszkowski, 1879) (*Mutilla*). – Europe: U (Crimea), R, BG, GR, Croatia, YU, Macedonia; Asia: ARM, AZ, TR, Iran, Syria.
pectinifera (André, 1893) (*Mutilla*).
obscura Nonveiller, 1959 (*Ctenotilla*).

21. Genus *Macromyrme* Lelej, 1984

binotata (Radoszkowski, 1879) (*Mutilla*). Asia: GE, ARM, AZ, Iran (north-west), TR (Ararat Mt.).
araratica (Radoszkowski, 1890) (*Mutilla*).
sinuata (Olivier, 1811) (*Mutilla*). – Asia: Iran (west), Syria, Lebanon, Palestine, CY; TR (Ararat Mt.) North Africa: Morocco.
villosa (Klug, 1829) (*Mutilla*).

22. Genus *Mutilla* Linnaeus, 1758

[*caucasica* Radoszkowski, 1865 (*Mutilla*). – "Caucasus", actually Australia.]
elbana Invrea, 1963 (*Mutilla*). – North Africa: Egypt.
erschoffii Sichel et Radoszkowski, 1870 (*Mutilla*). – Asia: GE, ARM, AZ, TU, UZ, TJ, Iran (north), Afghanistan.
rugiceps Morawitz, 1890 (*Mutilla*).
bareyi Radoszkowski, 1893 (*Mutilla*).
schulthessi André, 1893 (*Mutilla*).
europaea Linnaeus, 1758 (*Mutilla*). – Europe: from Sweden and Finland to Spain; Russia: NET, CET, SET, South Ural, WS (south); Asia: KZ, TR; North Africa.
aptera Uddman, 1753 (*Apis*).
similis Harris, 1782 (*Mutilla*).
schaefferi Schrank 1802 (*Mutilla*).
coerulans Lepeletier, 1845 (*Mutilla*).
cyanea Lepeletier, 1845 (*Mutilla*).
panzeri Lepeletier, 1845 (*Mutilla*).
obscura Nylander, 1847 (*Mutilla*).
kaschiriensis Baer, 1848 (*Mutilla*).
laevigata Sichel et Radoszkowski, 1870 (*Mutilla*).
notomelas André, 1902 (*Mutilla*).
rufula Hoffer, 1938 (*Mutilla*).
nigrescens Hoffer, 1938 (*Mutilla*).

- ruficeps* Hoffer, 1938 (*Mutilla*).
 [*kuschakewitzi* Radoszkowski, 1888 (*Mutilla*). "Wiernoje" (=Alma-Ata)
 (Kazakhstan), actually Papua New Guinea.]
- marginata** Baer, 1848 (*Mutilla*). – Europe: CZ, SL, D, A, CH, H, R, U, BG, I,
 E; Russia: NET, CET, SET, South Ural; Asia: GE, TR, Syria.
- trifasciata* Radoszkowski, 1865 (*Mutilla*).
monstruosa André, 1901 (*Mutilla*). – Europe: F.
insueta Invrea, 1964 (*Mutilla*).
mikado Cameron, 1900 (*Mutilla*). – Russia: ES (south), FE (Amurskaya oblast,
 Khabarovskii krai, Primorskii krai); Asia: Korea, China (Gansu,
 Heilongjiang, Jilin, Neimenggu, Shanxi, Zhejiang), Japan
 (Hokkaido, Honshu, Kyushu).
maesta Chen, 1957 (*Mutilla*).
panfilovi Lelej, 1981 (*Mutilla*). – Russia: WS (Altai); Asia: KZ, KY, UZ (east),
 China: NW (Xinjiang).
quinquemaculata Cyrillo, 1787 (*Mutilla*). – Europe: E, F, I, M, Croatia, GR, U
 (Crimea); Asia: CY, Syria, Palestine, TR; North Africa: Algeria,
 Egypt.
quinquepunctata Olivier, 1811 (*Mutilla*).
quadrinotata Klug, 1829 (*Mutilla*).
melanaria (André, 1898) (*Mutilla*). – Asia: ? Syria; North Africa: ?
 Egypt.
richterae Lelej, 1984 (*Mutilla*). – Asia: ARM, Iraq.
saltensis Radoszkowski, 1885 (*Mutilla*). – Russia: SET (Daghestan); Asia: GE,
 ARM, AZ, NW Iran, Iraq.
scorteccii Invrea, 1939 (*Mutilla*). – North Africa: Libya.

23. Genus *Nanomyrme* Lelej, 1977

- turkmenensis** Lelej, 1977 (*Nanomyrme*). – Asia: TU.

24. Genus *Ronisia* Costa, 1858

- Pycnotilla* Bischoff, 1920.
- barbara** Linnaeus, 1758 (*Mutilla*). – Europe: M; North Africa: Morocco,
 Algeria, Tunisia.
pervittata (André, 1902) (*Mutilla*). – North Africa: Algeria.
rubrinota (Invrea, 1958) (*Mutilla*). – North Africa: Morocco.
barbarula (Petersen, 1988) (*Mutilla*). – Europe: F, E; North Africa: Morocco.
brutia (Petagna, 1787) (*Mutilla*). – Europe: A, CZ, SL, D (south), F, I, M,
 Croatia, YU, Macedonia, GR, BG, TR, U; Russia: SET, South
 Ural; Asia: KZ (west), TU (Kopet-Dag), GE, ARM, AZ, Iran
 (north), TR, CY.
brutia brutia (Petagna, 1787). – Europe: A, CZ, SL, D (south), F, I, M,

-
- Croatia, YU, Macedonia, GR, BG, TR, U; Russia: SET, South Ural; Asia: TR, KZ (west).
- sexmaculata* (Cyrillo, 1787) (*Mutilla*).
hungarica (Fabricius, 1793) (*Mutilla*).
confusa (Lepeletier, 1845) (*Mutilla*).
monspeliensis (Lepeletier, 1845) (*Mutilla*).
sexmaculata (Baer, 1848) (*Mutilla*).
robusta (Smith, 1855) (*Mutilla*).
torosa Costa, 1858 (*Ronisia*).
sericeiventris (Costa, 1864) (*Mutilla*).
taurica (Radoszkowski, 1865) (*Mutilla*).
discoidalis (Radoszkowski, 1865) (*Mutilla*).
morawitzii (Dalla Torre, 1897) (*Mutilla*).
brutia duplex (Radoszkowski, 1885) (*Mutilla*). – Asia: TR, GE, ARM, AZ, Iran (north), TU (Kopet-Dag).
brutia minoensis Nonveiller, 1972 (*Ronisia*). – Europe: GR (Crete, Karpatos).
brutia valca (Petersen, 1988) (*Mutilla*). – Asia: CY.
calva (Panzer, 1801) (*Mutilla*).
ghilianii (Spinola, 1843) (*Mutilla*). – Europe: I (Sicily, Sardinia), F (south, Corsica), E, P; North Africa: Morocco.
differens Lepeletier, 1845 (*Mutilla*).
decoratifrons (Costa, 1860) (*Mutilla*).
razzaautii Invrea, 1957 (*Ronisia*).
ambigua Invrea, 1964 (*Ronisia*).
maculosa (Olivier, 1811) (*Mutilla*). – Asia: Lebanon; North Africa: Egypt.
marocana (Olivier, 1811) (*Mutilla*). – Europe: I (Sicily), M, F, E, P; Asia: Israel, Jordan, Syria, TR; North Africa: Morocco, Algeria, Tunisia, Egypt.
novemguttata (Klug, 1835) (*Mutilla*).
octomaculata (Spinola, 1843) (*Mutilla*).
algira (Lepeletier, 1845) (*Mutilla*).
rubricollis (Lucas, 1849) (*Mutilla*).
dorsalis (Lucas, 1849) (*Mutilla*).
humeralis (Sichel et Radoszkowski, 1870) (*Mutilla*).
bodenheimeri (Invrea, 1950) (*Mutilla*).
flavocornuta Invrea, 1965 (*Ronisia*).
parasinuata Invrea, 1965 (*Ronisia*). – Asia: Israel.
wahrmani Invrea, 1965 (*Ronisia*). – Asia: Jordan.

25. Genus *Spinulotilla* Bischoff, 1920

- kruegeri** (Invrea, 1932) (*Smicromyrme*). – North Africa: Libya.

26. Genus *Tropidotilla* Bischoff, 1920

- cypriadis** Invrea, 1940 (*Tropidotilla*). – Asia: CY, Iraq.
- fimbriata** (Klug, 1829) (*Mutilla*). – Asia: Syria, Arabia; Africa: Egypt, Somali, Chad, Ethiopia, Eritrea.
- frontalis** (Klug, 1829) (*Mutilla*). – North Africa: Egypt.
- grisescens** (Lepeletier, 1845) (*Mutilla*). – Europe: U (Crimea), GR, BG, Croatia, YU, Macedonia, I, F; North Africa: Algeria.
- vulnericeps** (Costa, 1860) (*Mutilla*).
- stribligata** (Sichel et Radoszkowski, 1870) (*Mutilla*).
- litoralis** (Petagna, 1787) (*Mutilla*). – Europe: E, P, F, B, CH, A, H, CZ, R, BG, Croatia, YU, Macedonia, AL, GR, I, U, Russia: SET; Asia: GE, ARM, AZ, TU (Kopet-Dag), Iran (north), TR, Syria, Palestine; North Africa: Morocco, Algeria, Egypt.
- italica** (Fabricius, 1793) (*Mutilla*).
- indigena** (Illiger, 1807) (*Mutilla*).
- graeca** (Lepeletier, 1845) (*Mutilla*).
- iberica** (Kolenati, 1846) (*Mutilla*).
- salentina** (Costa, 1858) (*Mutilla*).
- simplica** (Radoszkowski, 1865) (*Mutilla*).
- pauperata** (Sichel et Radoszkowski, 1870) (*Mutilla*).
- nigra** (Saunders, 1890) (*Mutilla*). – Europe: E (south); North Africa: Morocco.
- cruenticeps** (André, 1901) (*Mutilla*). – Asia: Syria, Palestine.
- soconusca** (Mercet, 1905) (*Mutilla*). – Europe: E.
- bimaculata** Invrea, 1965 (*Tropidotilla*).
- verticichroma** Invrea, 1965 (*Tropidotilla*).
- pharaonis** (André, 1902) (*Mutilla*). – North Africa: Egypt.
- sareptana** (André, 1901) (*Mutilla*). – Europe: Macedonia, GR; Russia: SET; Asia: TR, ARM, AZ.
- semirufa** (André, 1893) (*Mutilla*). – Asia: ARM, AZ, Iran.

Tribe *Smicromyrmini* Bischoff, 1920

27. Genus *Andreimyrme* Lelej, 1995

- davidi** (André, 1898) (*Mutilla*). – Asia: China: CE (Jiangsu, Jiangxi), SE (Fujian, Taiwan).
- substriolata** (Chen, 1957) (*Smicromyrme*). – Asia: China: (Jiangxi, Anhui, Zhejiang, Taiwan).

28. Genus *Dentilla* Lelej, 1980

- beludzhistanica** Lelej, 1985 (*Dentilla*). – Asia: Iran.
- caucasica** Lelej, 1985 (*Dentilla*). – Asia: AZ.

-
- curtiventris** (André, 1901) (*Mutilla*). – Europe: I, GR, YU, Macedonia, AL; Asia: TR, Syria, AZ, ARM.
rubescens (André, 1901) (*Mutilla*).
errana (Nonveiller, 1958) (*Smicromyrme*).
nigrescens (Nonveiller, 1958) (*Smicromyrme*).
rutilans (Nonveiller, 1972) (*Smicromyrme*).
cretica (Nonveiller, 1972) (*Smicromyrme*).
dichroa (Sichel et Radoszkowski, 1870) (*Mutilla*). – Asia: TU, Afghanistan, Iran, Iraq, Palestine, Egypt (Sinai).
bicolor (Olivier, 1811) (*Mutilla*).
quinquedentata (Morawitz, 1890) (*Mutilla*).
erronea (André, 1902) (*Mutilla*). – Europe: AL, YU, GR, BG, I; Asia: TR; North Africa: Algeria.
iran Lelej, 1985 (*Dentilla*). – Asia: Iran.
persica (Sichel et Radoszkowski, 1870) (*Mutilla*). – Asia: AZ, ARM, Iran, TR, Palestine.
sabulicola (Skorikov, 1935) (*Smicromyrme*). – Asia: Iran.
sabulosa (Skorikov, 1935) (*Smicromyrme*). – Asia: Iran.
saharica (Giner, 1945) (*Ephutomma*). – North Africa: Western Sahara, Morocco, Algeria, Sudan.
speciosa (Lelej, 1980) (*Smicromyrme*). – Asia: Afghanistan.
testacea (Klug, 1829) (*Mutilla*). – North Africa: Egypt.
ursina (Nagy, 1972) (*Smicromyrme*). – Asia: AZ, TR.
weidneri (Nagy, 1971) (*Smicromyrme*). – Asia: Iran.
zarudnyi Lelej, 1985 (*Dentilla*). – Asia: TU (Badkhyz), Iran, Afghanistan.

29. Genus *Ephutomma* Ashmead, 1899

Eremomyrme Suárez, 1965.

- angustata** (Skorikov, 1935) (*Smicromyrme*). – Asia: KZ (south), KY, TU, UZ, TJ.
sogdiana (Lelej, 1971) (*Smicromyrme*).
coronata (Lelej, 1976) (*Eremomyrme*). – Russia: South Ural; Asia: KZ, TU.
deserta Lelej, 1985 (*Ephutomma*). – Asia: KZ, UZ, TU, Afghanistan, Iran.
ferghanica Ovchinnikov, 2002 (*Ephutomma*). – Asia: KY.
gabesiana Bischoff, 1920 (*Ephutomma*). – North Africa: Tunisia.
kazenasi (Lelej, 1976) (*Eremomyrme*). – Asia: KZ (west), UZ, Mongolia.
kerzhneri (Lelej, 1977) (*Eremomyrme*). – Asia: KZ (south), UZ, TU, TJ, China: (Xijiang).
mesopotamica Bischoff, 1920 (*Ephutomma*). – Asia: Iraq.
minuta Lelej, 1985 (*Ephutomma*). – Asia: TU, TJ, Iran (north), Mongolia.
mongolensis (Suárez, 1974) (*Smicromyrme*). – Russia: ES (Transbaicalia); Asia: Mongolia.

- montarcense** (Giner, 1944) (*Smicromyrme*). – Europe: E.
sanguinicollis (Klug, 1829) (*Mutilla*). – Asia: Arabia; North Africa: Libya.
semirubra Bischoff, 1920 (*Ephutomma*). – North Africa: Libya.
turcestanica (Dalla Torre, 1897) (*Mutilla*). – Asia: KZ (south), TU, UZ, TJ.
incerta (Radoszkowski, 1877) (*Mutilla*).
semenovi (Skorikov, 1935) (*Smicromyrme*).
rufothoracica (Skorikov, 1935) (*Ephutomma*).
turgayensis Lelej, 1985 (*Ephutomma*). – Asia: KZ.

30. Genus *Mickelomyrme* Lelej, 1995

- abnorma** (Chen, 1957) (*Smicromyrme*). – Asia: China: CE (Anhui, Jiangxi, Zhejiang), SE (Fujian), SW (Yunnan), Vietnam.
exacta (Smith, 1879) (*Mutilla*). – Asia: China: CE (Shanghai, Anhui, Jiangsu, Jiangxi), SW (Sichuan), SE (Guangdong).

31. Genus *Nemka* Lelej, 1985

Horaia Tsuneki, 1993.

- aurantiaca** (Skorikov, 1935) (*Smicromyrme*). – Asia: TU, Iran, Afghanistan.
limi (Chen, 1957) (*Smicromyrme*). – Asia: China (Jiangsu, Zhejiang, Hainan).
pondicherensis (Sichel et Radoszkowski, 1870) (*Mutilla*). – Asia: Iran (south), India (east).
taiwanensis (Mickel, 1933) (*Timulla*). – Asia: China: CE (Jiangsu, Anhwei, Zhejiang), SE (Fujian, Taiwan).
viduata (Pallas, 1773) (*Mutilla*). – Europe: E, P, F, I, M, GR, Slovenia, Croatia, BH, YU, Macedonia, AL, A, D (south), BG, H, R, CZ, SL, U; Russia: SET, South Ural, WS (Altai); Asia: CY, TR, Syria, GE, AZ, ARM, KZ, KY, UZ, TU, Iran (north), TR, Mongolia (north-west), China: NW (Xinjiang); North Africa: Morocco, Algeria, Tunisia.
viduata viduata (Pallas, 1773). – Europe: U, E, F, I, M, GR, Slovenia, Croatia, BH, YU, Macedonia, AL, A, D (south), BG, H, R, CZ, SL; Russia: SET, South Ural, WS (Altai); Asia: TR, Syria, GE, AZ, ARM, KZ, Mongolia (north-west).
stridula (Rossi, 1790) (*Tiphia*).
coronata (Rossi, 1792) (*Mutilla*).
coronata (Fabricius, 1793) (*Mutilla*).
paedemontana (Fabricius, 1794) (*Mutilla*).
togata (Ahrens, 1812) (*Mutilla*).
viduata tunensis (Fabricius, 1804) (*Mutilla*). – North Africa: Morocco, Algeria, Tunisia.
bipunctata (Lepeletier, 1845) (*Mutilla*).
dispar (Smith, 1855) (*Mutilla*). – North Africa: Algeria.
triplex (Invrea, 1953) (*Smicromyrme*).

- viduata macquarti** (Lepeletier, 1845) (*Mutilla*). – Europe: GR, F.
viduata bartholomaei (Radoszkowski, 1865) (*Mutilla*). – Asia: GE, AZ,
 ARM, KZ (south of Aral Sea and Balkhash Lake), KY, UZ, TU,
 TR, Iran (north), TR, China: NW (Xinjiang).
luctuosa (Radoszkowski, 1865) (*Mutilla*).
bareyi (Radoszkowski, 1893) (*Mutilla*).
protunensis (Skorikov, 1935) (*Smicromyrme*).
simia (Skorikov, 1935) (*Smicromyrme*).
anacta (Skorikov, 1935) (*Smicromyrme*).
intermedia (Skorikov, 1935) (*Smicromyrme*).
viduata andalusiana (Skorikov, 1935) (*Smicromyrme*). – Europe: E, P.
viduata insulae (Invrea, 1940) (*Smicromyrme*). – Asia: CY.
wotani (Zavattari, 1913) (*Mutilla*). – Asia: Japan (Honshu, Kyushu, Tanegashima), Korea, China (Liaoning, Shanxi, Taiwan), Vietnam.
yasumatsui (Mickel, 1936) (*Smicromyrme*).

32. Genus Nuristanilla Lelej, 1980

kabakovi Lelej, 1980 (*Nuristanilla*). – Asia: Afghanistan.

33. Genus *Physetopoda* Schuster, 1949

Paramyrme Lelej, 1977.

- asiatica** (Lelej, 1984) (*Paramyrme*). – Asia: ARM, AZ, KZ, KY, UZ, Iran (north).

betpakdalensis (Lelej, 1984) (*Paramyrme*). – Asia: KZ.

cingulata (Costa, 1858) (*Mutilla*). – Europe: I, GR, H; Russia: SET (south-east); Asia: KZ.

rudolfae (Lelej, 1984) (*Smicromyrme*).

compacta (Invrea, 1957) (*Smicromyrme*). – Europe: I, Croatia, F, E, P, R.

daghستانica (Radoszkowski, 1885) (*Mutilla*). – Europe: U, A, CZ, SL, F, I, Croatia, GR, CH, D, R, H; Russia: SET; WS (Altai); Asia: KZ.

moravica (Hoffer, 1936) (*Smicromyrme*).

variabilis (Hoffer, 1938) (*Smicromyrme*).

maculata (Hoffer, 1938) (*Smicromyrme*).

lineata (Hoffer, 1938) (*Smicromyrme*).

obenbergeri (Hoffer, 1938) (*Smicromyrme*).

conversa (Hoffer, 1938) (*Smicromyrme*).

deserta (Lelej, 1984) (*Paramyrme*). – Asia: TU, UZ, KZ (south-east).

fusculina (Invrea, 1955) (*Smicromyrme*). – Europe: I (Sardinia).

gorbatovskii (Lelej, 1984) (*Paramyrme*). – Asia: TJ.

halensis (Fabricius, 1787) (*Mutilla*). – Europe: BG, GR, H, D, CZ, SL, A, CH, R, Slovenia, Croatia, BH, YU, Macedonia, I, U; Russia: SET, south Ural; WS (Altai); Asia: KZ.

- ephippium* (Fabricius, 1793) (*Mutilla*).
montana (Panzer, 1805) (*Mutilla*).
schenki (Schmiedeknecht, 1907) (*Mutilla*).
pseudomontana (Hoffer, 1936) (*Smicromyrme*).
susterai (Hoffer, 1936) (*Smicromyrme*).
melaclena (Invrea, 1958) (*Smicromyrme*).
subalpina (Ressl, 1960) (*Smicromyrme*).
hissarica (Lelej, 1984) (*Paramyrmecia*). – Asia: TJ.
isis (Suárez, 1975) (*Smicromyrme*). – North Africa: Egypt.
kermanensis (Lelej, 1984) (*Paramyrmecia*). – Asia: Iran (south-east).
kozlovi (Lelej, 1976) (*Smicromyrme*). – Asia: KZ, Mongolia.
ligustica (Invrea, 1951) (*Smicromyrme*). – Europe: R, I, M, E, P, F.
borgiana (Invrea, 1951) (*Smicromyrme*). – Europe: I.
myrtacea (Invrea, 1951) (*Smicromyrme*). – Europe: I.
lucasii (Smith, 1855) (*Mutilla*). – Europe: I, F, E; North Africa: Algeria.
bicolor (Lucas, 1849) (*Mutilla*).
cerrutii (Invrea, 1956) (*Smicromyrme*).
pseudobimaculata (Suárez, 1959) (*Smicromyrme*). – Europe: E.
mendizabali (Suárez, 1956) (*Smicromyrme*). – Europe: E (south).
oratoria (Chen, 1957) (*Smicromyrme*). – Russia: ES (south, Chitinskaya oblast, Yakutia), FE (Amurskaya oblast, Khabarovskii krai, Primorskii krai); Asia: China (Jilin, Neimenggu, Shandong), Mongolia.
vaticinia (Chen, 1957) (*Smicromyrme*).
miscisis (Chen, 1957) (*Smicromyrme*).
rectangulata (Chen, 1957) (*Glossotilla*).
eoa (Lelej, 1977) (*Smicromyrme*).
pierrei (Suárez, 1958) (*Smicromyrme*). – North Africa: Algeria, Mauritania, Chad.
portschinskii (Radoszkowski, 1888) (*Mutilla*). – Russia: SET (south-east); Asia: ARM, AZ, KZ, TU (south-west).
punctata (Latreille, 1792) (*Mutilla*). – Europe: E, P, F (south), I, M, H, R, Croatia. North Africa: Egypt, Tunisia, Algeria, Morocco.
quadripunctata (Olivier, 1811) (*Mutilla*).
quadrimaculata (Lucas, 1849) (*Mutilla*).
melanothorax (André, 1901) (*Mutilla*). – North Africa: Algeria.
pusilla (Klug, 1835) (*Mutilla*). – Europe: AL, Croatia, GR, E; Asia: CY; North Africa: Algeria, Morocco.
nigrithorax (André, 1902) (*Mutilla*). – North Africa: Algeria, Morocco.
rubrocincta (Lucas, 1849) (*Mutilla*). – Europe: YU, AL, GR, I, F, E, P; North Africa: Algeria, Tunisia, Morocco.
insularis Schuster, 1949 (*Physetopoda*).
jacobi (Invrea, 1953) (*Smicromyrme*).
bitriangulata (Invrea, 1953) (*Smicromyrme*). – North Africa: Morocco.

- rubicula** (Invrea, 1953) (*Smicromyrme*). – North Africa: Morocco.
- rufosquamulata** (André, 1904) (*Mutilla*). – Europe: E, F (south).
- pronotalis** (Mercet, 1903) (*Mutilla*). – Europe: E.
- scutellaris** (Latreille, 1792) (*Mutilla*). – Europe: E, A, CH, I, Slovenia, Croatia, YU, F, D, CZ, SL, GR, R, H, ? Russia; Asia: ARM, TR; ? North Africa: Algeria, Tunisia.
- bimaculata** (Jurine, 1807) (*Mutilla*).
- subcomata** (Wesmael, 1852) (*Mutilla*).
- nigrescens** (André, 1901) (*Mutilla*). – Europe: F (south), CH, Croatia; ? Russia; North Africa: Algeria, Tunisia.
- nigricollis** (Hoffer, 1936) (*Smicromyrme*).
- castellaneta** Pagliano, 1983 (*Smicromyrme*).
- sericeiceps** (André, 1901) (*Mutilla*). – Europe: H, YU, I, F, P, E.
- moltonii** (Invrea, 1955) (*Smicromyrme*).
- similis** (Lelej, 1984) (*Paramyrme*). – Europe: U (Crimea); Asia: TR, ARM, AZ.
- trioma** (Invrea, 1955) (*Smicromyrme*). – Europe: I (Sardinia), F (Corsica).
- turgajensis** (Lelej, 1984) (*Smicromyrme*). – Russia: SET (south-east), South Ural; Asia: KZ, Mongolia (west).
- unicineta** (Lucas, 1849) (*Mutilla*). – North Africa: Algeria.

34. Genus *Promecilla* André, 1903

- primana** Skorikov, 1935 (*Promecilla*). – Asia: UZ, TJ.
- vinuta** (Nagy, 1972) (*Smicromyrme*). – Asia: Afghanistan.

35. Genus *Sinotilla* Lelej, 1995

- cyaneiventris** (André, 1896) (*Mutilla*). – Asia: China: NO (Neimenggu, Hebei, Shanxi), CE (Jiangsu, Jiangxi, Zhejiang).
- cyaneiventris cyaneiventris** (André, 1896). – Asia: China: NO (Hebei, Shanxi), CE (Jiangsu, Jiangxi, , Zhejiang).
- cyaneiventris gobiana** (Chen, 1957) (*Smicromyrme*). – Asia: China: NO (Neimenggu).
- pekiniana** (André, 1905) (*Mutilla*). – Asia: China: NO (Hebei, Shanxi), CE (Jiangsu), SE (Fujian).

36. Genus *Skorikovia* Ovchinnikov, 2002

- elongata** (Radoszkowski, 1885) (*Mutilla*). – Russia: SET; Asia: AZ, KZ, KY, TU, UZ, Iran (north), Afghanistan.
- pura** (André, 1901) (*Mutilla*).
- ambigua** (Skorikov, 1935) (*Smicromyrme*).
- baigacuma** (Skorikov, 1935) (*Smicromyrme*).

- gracilis* (Skorikov, 1935) (*Smicromyrme*).
kuznetzovi (Skorikov, 1935) (*Smicromyrme*).
pallipes (Lelej, 1985) (*Smicromyrme*). – Asia: Iran (east).
pliginskiji (Lelej, 1984) (*Smicromyrme*). – Europe: U (Crimea); Russia: SET, South Ural.
radoszkovskii (Skorikov, 1935) (*Ephutomma*). – Russia: SET (Daghestan); Asia: AZ, GE.
transcaucasica (Lelej, 1985) (*Smicromyrme*). – Asia: ARM, AZ.

37. Genus *Smicromyrme* Thomson, 1870

Subgenus *Astomyrme* Schwartz, 1984

- asiatica** Lelej, 1985 (*Smicromyrme*). – Asia: KZ, KY, UZ.
ausonia Invrea, 1950 (*Smicromyrme*). – Europe: F (Corsica), I, GR, U (Crimea), MO; Russia: SET, South Ural; Asia: KZ, TU (Kopet-Dag), CY.
sabatia Invrea, 1953 (*Smicromyrme*).
etrusca Invrea, 1950 (*Smicromyrme*). – Europe: I.
esterina Pagliano, 1983 (*Smicromyrme*). – Europe: I (Sicily).
inerma (Schwartz, 1984) (*Astomyrme*). – Europe: F (south).
massageta Lelej, 1985 (*Smicromyrme*). – Asia: KZ.
minuta Lelej, 1985 (*Smicromyrme*). – Asia: KZ, KY, UZ.
morawitzi Lelej, 1979 (*Smicromyrme*). – Asia: KZ (south), UZ, TU, TJ.
nikoskajae Lelej, 1985 (*Smicromyrme*). – Asia: KY, TJ.
opistomelas Invrea, 1950 (*Smicromyrme*). – Europe: I (Sardinia), E, North Africa: Morocco, Libya.
suberrata Invrea, 1957 (*Smicromyrme*). – Europe: E, F (south), I (south), M; North Africa: Egypt, Libya, Algeria, Morocco.
succedanea Ovchinnikov, 2002 (*Smicromyrme*). – Asia: KY.

Subgenus *Eremotilla* Lelej, 1985

- betpakdalensis** Lelej, 1971 (*Smicromyrme*). – Asia: KZ.
constans Ovchinnikov, 2002 (*Smicromyrme*). – Asia: KY.
coracina Lelej, 1985 (*Smicromyrme*). – Russia: South Ural, WS (Altai); Asia: KZ, KY, Mongolia (west).
gussakovskii Lelej, 1985 (*Smicromyrme*). – Asia: UZ, TJ.
inconstans (Radoszkowski, 1885) (*Mutilla*). – Asia: KZ, UZ, TU.
lehri Lelej, 1985 (*Smicromyrme*). – Asia: KZ, TU, UZ, TJ.
margiana Lelej, 1985 (*Smicromyrme*). – Asia: TU.
mesasiatica Lelej, 1985 (*Smicromyrme*). – Asia: UZ, TU.
nigriceps Nonveiller, 1959 (*Smicromyrme*). – Europe: U (Crimea), GR, YU, Macedonia; Russia: SET; Asia: GE, ARM, AZ, TU (Kopet-Dag), Iran (north).
orientalis Nonveiller, 1959 (*Smicromyrme*).

-
- maculata* Nonveiller, 1959 (*Smicromyrme*).
afasciata Nonveiller, 1972 (*Smicromyrme*).
pseudoorientalis Nonveiller, 1979 (*Smicromyrme*).
nonveilleri Invrea, 1954 (*Smicromyrme*). – Europe: R, Croatia, GR.
novaki Invrea, 1954 (*Smicromyrme*). – Europe: Croatia, R, GR.
dalmatina Invrea, 1954 (*Smicromyrme*).
perisii (Sichel et Radoszkowski, 1870) (*Mutilla*). – Europe: F (Corsica), I (Sardinia).
rufocapitata Skorikov, 1935 (*Smicromyrme*). – Asia: KZ, TJ.
schwarzi Suárez, 1975 (*Smicromyrme*). – Europe: U (south), GR; Asia: GE, AZ, TR.
tadzhikistanica Lelej, 1985 (*Smicromyrme*). – Asia: TJ.
tekensis Skorikov, 1935 (*Smicromyrme*). – Asia: KZ, KY, UZ, TU, TJ, China: NW (Xinjiang), Mongolia.
nigrocapitata Skorikov, 1935 (*Smicromyrme*).
triangularis (Radoszkowski, 1865) (*Mutilla*). – Europe: U; Russia: CET (Kasan), South Ural, SET.
verhoeffi Suárez, 1959 (*Smicromyrme*). – Europe: F (south).
vladani Nonveiller, 1972 (*Smicromyrme*). – Europe: GR (Crete).
infuscata Nonveiller, 1972 (*Smicromyrme*).
deminuta Nonveiller, 1972 (*Smicromyrme*).

Subgenus Erimyrme Lelej, 1985

- azerbaidzhanica** Lelej, 1985 (*Smicromyrme*). – Asia: AZ, TU (south-west).
bidenticulata Chen, 1957 (*Smicromyrme*). – Russia: FE (Primorskii krai); Asia: Mongolia (east), China: NO (Neimenggu, Shandong), CE (Jiangsu).
hasanensis Lelej, 1977 (*Smicromyrme*).
monopunctata Lelej, 1977 (*Smicromyrme*).
apunctata Lelej, 1977 (*Smicromyrme*).
borceai Nagy, 1968 (*Smicromyrme*). – Europe: R.
kurzenkoi Lelej, 1985 (*Smicromyrme*). – Asia: KZ (south-east), KY.
pulawskii Suárez, 1975 (*Smicromyrme*). – Europe: BG, GR (Rhodes); Asia TR.
ambigua Invrea, 1954 (*Smicromyrme*).
sicana (De Stefani, 1887) (*Mutilla*). – Europe: F, I, YU, Croatia, AL, GR, H, A, U, CZ, SL, R, TR; Russia: SET, South Ural, WS (Altai); Asia: TR, Syria.
septentrionalis Hoffer, 1936 (*Smicromyrme*).
gregory Hoffer, 1936 (*Smicromyrme*).
lutescens Invrea, 1954 (*Smicromyrme*).
stepposa Lelej, 1984 (*Smicromyrme*). – Europe: U (Crimea); Russia: SET; Asia: KZ (west).

Subgenus **Rhombotilla** Nagy, 1966

riparia Nagy, 1966 (*Smicromyrme*). – Europe: R.

Subgenus **Smicromyrme** Thomson, 1870

cephalotes Lelej, 1985 (*Smicromyrme*). – Asia: TJ.

iliensis Lelej, 1985 (*Smicromyrme*). – Asia: KZ (south-east), KY.

lampedusia Invrea, 1957 (*Smicromyrme*). – Europe: I (Lampedusa).

lewisi Mickel, 1935 (*Smicromyrme*). – Russia: ES (Irkutskaya oblast, Transbaicalia), FE (Amurskaya oblast, Khabarovskii krai, Primorskii krai, Sakhalin, Kunashir II.); Asia: Mongolia, China (Neimenggu, Hebei, Shandong, Jiangsu, Anhui, Jiangxi, Fujian), Korea, Japan (from Hokkaido to Tanegashima).

japonica (Cameron, 1900) (*Mutilla*).

cameroni (Yano, 1932) (*Mutilla*).

yanoi Mickel, 1935 (*Smicromyrme*).

nigricula Mickel, 1935 (*Smicromyrme*).

momidziyama Tsuneki, 1972 (*Smicromyrme*).

nikolajevi Lelej, 1985 (*Smicromyrme*). – Asia: KZ, KY, Mongolia (west).

ruficollis (Fabricius, 1793) (*Mutilla*). – Europe: U, I, F (Corsica), A, Croatia, BH, YU, Macedonia, GR, BG, E, F, M; Russia: WS (Altai); Asia: KZ, TR.

ruficollis ruficollis (Fabricius, 1793). – Europe: U, I, A, Croatia, BH, YU, Macedonia, GR, BG, E, F, M; Russia: WS (Altai); Asia: KZ, TR.

kiritshenkoae Skorikov, 1935 (*Smicromyrme*).

ruficollis ceresae Invrea, 1952 (*Smicromyrme*). – Europe: I (Sardinia), F (Corsica).

berlandi Invrea, 1953 (*Smicromyrme*).

rufipes (Fabricius, 1787) (*Mutilla*). – Europe: E (north), F, I (north, ? Sicily), Slovenia, Croatia, BH, YU, CH, A, D, H, CZ, SL, U, BE, S, SF; Russia: NET, CET, SET, South Ural, WS (Altai), ES (Yakutsk); Asia: KZ.

nigra (Rossi, 1792) (*Mutilla*).

ephippium (Rossi, 1792) (*Mutilla*).

sellata (Panzer, 1797) (*Mutilla*).

nigrita (Panzer, 1801) (*Mutilla*).

petiolaris (Fabricius, 1804) (*Mutilla*).

gibba (Baer, 1848) (*Mutilla*).

nemoralis (Baer, 1848) (*Mutilla*).

platiensis (De Stefani, 1897) (*Mutilla*). – Europe: I (Sicily).

fulvogastra (André, 1901) (*Mutilla*).

dusmeti (Mercet, 1905) (*Mutilla*). – Europe: E.

-
- nigricollaris* Skorikov, 1935 (*Smicromyrme*).
zavadili Hoffer, 1938 (*Smicromyrme*).
lidmilae Hoffer, 1938 (*Smicromyrme*).
kuehlhorni Invrea, 1963 (*Smicromyrme*).
globosa Franz, 1982 (*Smicromyrme*).
serta (Radoszkowski, 1885) (*Mutilla*). – Russia: South Ural.
strandii (Zavattari, 1913) (*Mutilla*). – Asia: China: NO (Shanxi), CE (Jiangxi, Zhejiang), SE (Fujian, Guangdong, Taiwan).
tristis Lelej, 1984 (*Smicromyrme*). – Europe: U; Russia: South Ural; Asia: KZ.
viktorovi Lelej, 1984 (*Smicromyrme*). – Russia: SET; Asia: AZ, GE.

Smicromyrme incertae sedis

- agusii** (Costa, 1884) (*Mutilla*). – Europe: I (Sardinia), F (Corsica).
balianii Invrea, 1932 (*Smicromyrme*). – Europe: I; North Africa: ? Morocco.
beaumonti Invrea, 1953 (*Smicromyrme*). – North Africa: Morocco.
bengasiana Invrea, 1932 (*Smicromyrme*). – North Africa: Libya.
berberica (André, 1902) (*Mutilla*). – North Africa: Algeria, Morocco.
melanosoma Suárez, 1958 (*Smicromyrme*). – North Africa: Morocco.
bletoni Giner, 1959 (*Smicromyrme*). – North Africa: Morocco.
bytinskii Invrea, 1965 (*Smicromyrme*). – Asia: Palestine.
rufula Invrea, 1965 (*Smicromyrme*).
ceballosi Suárez, 1959 (*Smicromyrme*). – Europe: E.
chinensis (Smith, 1855) (*Mutilla*). – Asia: China: CE (Jiangsu, Anhui).
castrensis Invrea, 1954 (*Smicromyrme*). – Europe: Croatia, R.
crassicornis (Maidl, 1922) (*Mutilla*).
cristinae Lo Cascio, 2000 (*Smicromyrme*). – Europe: GR.
damascena Invrea, 1962 (*Smicromyrme*). – Asia: Syria.
defixa Invrea, 1953 (*Smicromyrme*). – North Africa: Morocco.
ellipsifera (Gribodo, 1884) (*Mutilla*). – Africa: Egypt, Eritrea.
arsinoensis (André, 1896) (*Mutilla*).
elonica Invrea, 1965 (*Smicromyrme*). – Asia: Palestine.
eltihnica Invrea, 1960 (*Smicromyrme*). – North Africa: Egypt.
ferdinandi Invrea, 1953 (*Smicromyrme*). – Europe: E; North Africa: Morocco.
ferrugineiceps (André, 1902) (*Mutilla*). – Asia: Palestine.
ferrugineipes (André, 1910) (*Mutilla*). – Asia: Palestine; North Africa: Egypt, Libya.
gebatana Invrea, 1953 (*Smicromyrme*). – Asia: Palestine.
giarabubensis Invrea, 1932 (*Smicromyrme*). – North Africa: Libya.
gineri Invrea, 1953 (*Smicromyrme*). – North Africa: Morocco.
gridellii Invrea, 1960 (*Smicromyrme*). – North Africa: Egypt.
hierosolymitana Invrea, 1953 (*Smicromyrme*). – Asia: Palestine.
implicata Invrea, 1953 (*Smicromyrme*). – North Africa: Morocco.
ingauna Invrea, 1958 (*Smicromyrme*). – Europe: I, R.

- varinella* Invrea, 1960 (*Smicromyrme*).
innesi (André, 1893) (*Mutilla*). – North Africa: Egypt.
kurda (Skorikov, 1935) (*Smicromyrme*). – Asia: GE, ARM, AZ, Iran (north).
 subtunensis (Skorikov, 1935) (*Smicromyrme*).
 persa (Skorikov, 1935) (*Smicromyrme*).
longistigma Invrea, 1960 (*Smicromyrme*). – North Africa: Egypt.
lucens Invrea, 1963 (*Smicromyrme*). – North Africa: Egypt.
lybica Invrea, 1932 (*Smicromyrme*). – North Africa: Egypt, Libya, Morocco,
 Western Sahara.
 libyca Invrea, 1953 (*Smicromyrme*).
femoralis Invrea, 1953 (*Smicromyrme*). – North Africa: Morocco.
obscurata Invrea, 1953 (*Smicromyrme*). – North Africa: Morocco.
complecta Invrea, 1960 (*Smicromyrme*). – North Africa: Egypt.
fusca Invrea, 1932 (*Smicromyrme*). – North Africa: Libya.
invreai Giner, 1945 (*Smicromyrme*). – North Africa: Western Sahara.
mareotica Invrea, 1951 (*Smicromyrme*). – North Africa: Egypt, Libya.
mateui Suárez, 1953 (*Smicromyrme*). – North Africa: Morocco.
matritensis (Mercet, 1905) (*Mutilla*). – Europe: E.
matruhana Invrea, 1960 (*Smicromyrme*). – North Africa: Egypt.
mauromoustakisi Invrea, 1940 (*Smicromyrme*). – Asia: Palestine, CY.
melanolepis (Costa, 1884) (*Mutilla*). – Europe: I (Sardinia, Sicily), F (Corsica).
merctei (André, 1904) (*Mutilla*). – Europe: E.
metanotalis (André, 1902) (*Mutilla*). – Europe: E, P.
 metanotalis metanotalis (André, 1902). – Europe: E.
 metanotalis andradei Suárez, 1954 (*Smicromyrme*). – Europe: P.
 notomelanica Suárez, 1954 (*Smicromyrme*). – Europe: P.
mochii Invrea, 1962 (*Smicromyrme*). – Asia: Syria.
naefi Invrea, 1953 (*Smicromyrme*). – North Africa: Morocco.
negrei Suárez, 1965 (*Smicromyrme*). – Europe: E, P; North Africa: Morocco.
nuptura (Mercet, 1905) (*Mutilla*). – Europe: E.
 bofilli (Mercet, 1905) (*Mutilla*). – Europe: E.
pardoí Suárez, 1953 (*Smicromyrme*). – North Africa: Morocco.
partita (Klug, 1835) (*Mutilla*). – Europe: E, P, F (south), I, M, GR, YU; North
 Africa: Tunisia, Algeria, Morocco.
 unimaculata (Lucas, 1849) (*Mutilla*).
 hispanica (Sichel et Radoszkowski, 1870) (*Mutilla*).
partita fasciaticollis (Spinola, 1843) (*Mutilla*). – Europe: E, I (Sicily), M.
 cinereifrons (Costa, 1864) (*Mutilla*).
 intermedia (De Stefani, 1887) (*Mutilla*).
 intermedia (Costa, 1888) (*Mutilla*).
 destefanii (Dalla Torre, 1897) (*Mutilla*).
 costae (Dalla Torre, 1897) (*Mutilla*).
partita obscurithorax (André, 1902) (*Mutilla*). – Europe: E; Asia: Syria;

- North Africa: Algeria, Tunisia, Morocco.
- propodealis** Suárez, 1959 (*Smicromyrme*). – Europe: E, P.
- plantouriana** Schwartz, 1986 (*Smicromyrme*). – Europe: F (south).
- pluripicta** Invrea, 1965 (*Smicromyrme*). – Asia: Palestine.
- pouzdranensis** Hoffer, 1936 (*Smicromyrme*). – Europe: CZ.
- priesneri** Invrea, 1960 (*Smicromyrme*). – North Africa: Egypt.
- pyramidalorum** (André, 1910) (*Mutilla*). – Asia: Palestine; North Africa: Egypt.
- rectagonum** Invrea, 1953 (*Smicromyrme*). – North Africa: Morocco.
- dunalis** Invrea, 1953 (*Smicromyrme*). – North Africa: Morocco.
- rubidula** Invrea, 1953 (*Smicromyrme*). – North Africa: Morocco.
- rufithorax** Giner, 1959 (*Smicromyrme*). – North Africa: Morocco.
- [b**rufula** (André, 1901) (*Mutilla*). – "Arabia"]. [Not Palaearctic].
- schemeana** Invrea, 1953 (*Smicromyrme*). – Asia: Palestine.
- signata** (Klug, 1829) (*Mutilla*). – Africa: Egypt, Eritrea, Ethiopia, Sudan, Chad.
- subtilis** Invrea, 1953 (*Smicromyrme*). – North Africa: Morocco.
- sulcisia** Invrea, 1955 (*Smicromyrme*). – Europe: I (Sardinia).
- terricola** Nagy, 1972 (*Smicromyrme*). – Europe: GR.
- tetragona** Invrea, 1960 (*Smicromyrme*). – North Africa: Egypt.
- trinotata** (Costa, 1858) (*Mutilla*). – Europe: I; North Africa: ?Morocco.
- tumidula** Nagy, 1972 (*Smicromyrme*). – Europe: GR.
- turanica** (Morawitz, 1893) (*Mutilla*). – Europe: GR (Crete); Asia: Syria, TR,
 ARM, AZ, UZ, TJ, Afghanistan.
 arameana (André, 1902) (*Mutilla*).
 atrichorax (André, 1902) (*Mutilla*).
zavattarii Invrea, 1932 (*Smicromyrme*). – North Africa: Libya.

Tribe Petersenidiini Lelej, 1996

38. Genus Artiotilla Invrea, 1950

- Glossomyrme* Suárez, 1979.
- afghanica** (Suárez, 1979) (*Glossomyrme*). – Asia: Afghanistan.
- ariana** Lelej, 1980 (*Artiotilla*). – Asia: Afghanistan.
- biguttata** (Costa, 1858) (*Mutilla*). – Europe: I, Croatia, YU, Macedonia, GR,
 AL; Asia: AZ, TU, Iran, TR, CY, Palestine.
 dalmatica (André, 1893) (*Mutilla*).
 mervensis (Radoszkowski, 1893) (*Mutilla*).
 fertoni (Invrea, 1953) (*Smicromyrme*).
 palestinensis Invrea, 1965 (*Artiotilla*).

39. Genus Orientidia Lelej, 1996

- obscurilamina** (Chen, 1957) (*Smicromyrme*). – Asia: China (Shanghai, Jiangsu,
 Fujian).

40. Genus *Taiwanomyrme* Tsuneki, 1993

friekae (Zavattari, 1913) (*Mutilla*). – Asia: China: CE (Jiangsu, Anhui, Zhejiang), SE (Fujian, Taiwan).

Tribe *Trogaspidiini* Bischoff, 1920

Timullini Krombein, 1972.

41. Genus *Eotrogaspidia* Lelej, 1996

auroguttata (Smith, 1855) (*Mutilla*). – Asia: China: SW (Sichuan), CE (Jiangsu, Zhejiang, Hunan), SE (Fujian, Guangdong, Hainan, Taiwan, Hong Kong), Thailand (north).

42. Genus *Glossotilla* Bischoff, 1920

tricoloriformis Bischoff, 1920: (*Glossotilla*). – Africa: Egypt, Senegal.

43. Genus *Lobotilla* Bischoff, 1920

leucopyga (Klug, 1829) (*Mutilla*). – Africa: Egypt, Eritrea, from Sudan to Senegal.

nitidiventris (Klug, 1829) (*Mutilla*).

opaca (Lepeletier, 1845) (*Mutilla*).

44. Genus *Neotrogaspidia* Lelej, 1996

hammeri (Suárez, 1959) (*Trogaspidia*). – Asia: ARM, AZ, TU (south-west), CY, Palestine.

pustulata (Smith, 1873) (*Mutilla*). – Asia: Japan (Honshu, Kyushu, Tsushima, Yaku-shima, Tanega-shima, Amami-oshima, Bonin Is.), Korea, China (Sichuan, Jiangsu, Anhui, Zhejiang, Jiangxi, Hunan, Fujian, Guangdong, Taiwan).

insidiator (Smith, 1873) (*Mutilla*).

insidiator (Smith, 1874) (*Mutilla*).

petulans (Smith, 1874) (*Mutilla*).

japonica (Dalla Torre, 1897) (*Mutilla*).

elpinice (Mickel, 1933) (*Timulla*).

aestivalis (Hammer, 1949) (*Trogaspidia*).

45. Genus *Radoszkowskius* Ashmead, 1903

conversus (Chen, 1957) (*Trogaspidia*). – Asia: China (Shanghai, Anhui, Fujian, Taiwan).

oculata (Fabricius, 1804) (*Mutilla*). – Asia: China (Beijing, Jiangsu, Zhejiang, Hunan, Fujian, Taiwan, Guangdong, Yunnan, Hainan, Hong Kong), Vietnam, Laos, Thailand, Myanmar, Cambodia, Malaysia (Malay Peninsula).

46. Genus *Trogaspidia* Ashmead, 1899

arciformis Chen, 1957 (*Trogaspidia*). – Asia: China (Anhui, Fujian).

[castellana (Mercet, 1903) (*Mutilla*). – "Madrid", actually Luzon, Philippines].
[Not Palaearctic].

catanensis (Rossi, 1794) (*Mutilla*). – Europe: U (Crimea), H, R, Croatia, YU, Macedonia, I; Russia: SET; Asia: ARM, KZ (west); North Africa: Algeria.

ballioni (Radoszkowski, 1866) (*Mutilla*).

paripunctata (Sichel et Radoszkowski, 1870) (*Mutilla*).

ehrenbergi Bischoff, 1920 (*Trogaspidia*). – North Africa: Egypt.

fedtschenkoi (Radoszkowski, 1877) (*Mutilla*). – Asia: AZ, KZ (south), KY, UZ, TU, TJ.

transcaspica (Radoszkowski, 1888) (*Mutilla*).

binio (Radoszkowski, 1888) (*Mutilla*).

bisignata (Morawitz, 1890) (*Mutilla*).

rittneri (Dalla Torre, 1897) (*Mutilla*).

immaculata (Skorikov, 1935) (*Smicromyrme*).

floralis (Klug, 1829) (*Mutilla*). – Asia: Arabia, Yemen; Africa: Algeria, Libya, Egypt, Sudan, Senegal, Mali, Nigeria, Cameroon, Chad, Eritrea, Ethiopia, Somali.

interrupta (Olivier, 1811) (*Mutilla*).

divisa (Smith, 1855) (*Mutilla*).

aureiventris (Walker, 1871) (*Mutilla*).

klugiana (André, 1902) (*Mutilla*). – Asia: Iran, Yemen; Africa: Egypt, Eritrea, Ethiopia.

moslemita Bischoff, 1920 (*Trogaspidia*). – Asia: Arabia, Yemen; Africa: Egypt, Sudan.

fuscipennis (Fabricius, 1804) (*Mutilla*). – Asia: China: NO (Hebei), CE (Jiangsu, Anhui, Zhejiang), SE (Fujian, Guangdong).

hoffmanni (Mickel, 1933) (*Timulla*). – Asia: China: CE (Jiangsu), SE (Fujian, Hainan).

israeliana Invrea, 1965 (*Trogaspidia*). – Asia: Jordan.

kolthoffi Hammer, 1949 (*Trogaspidia*). – Asia: China: CE (Jiangsu).

lingnani (Mickel, 1933) (*Timulla*). – Asia: China (Jiangsu, Anhui, Zhejiang, Fujian, Hainan).

maritima Chen, 1957 (*Trogaspidia*). – Asia: China: CE (Jiangsu, Anhui, Zhejiang).

mesopotamica (Bischoff, 1914) (*Mutilla*). – Asia: Iraq.

- niveomaculata** (André, 1898) (*Mutilla*). – Asia: Palestine; Africa: Djibouti, Sudan.
- nodoa** (Mickel, 1933) (*Timulla*). – Asia: China: NO (Anhui), SE (Fujian, Guangdong, Hainan).
- pavesii** (Magretti, 1884) (*Mutilla*). – Africa: Egypt, Ethiopia.
- priesneri** Invrea, 1963 (*Trogaspidia*). – North Africa: Egypt.
- retinula** Chen, 1957 (*Trogaspidia*). – Asia: China: CE (Jiangsu), SE (Fujian, Taiwan).
- rhea** (Mickel, 1933) (*Timulla*). – Asia: China: CE (Anhui, Jiangsu, Zhejiang), SE (Fujian, Guangdong, Hainan).
- rhea rhea** (Mickel, 1933). – Asia: China (Jiangsu, Fujian, Hainan).
- rhea gaea** Chen, 1957 (*Trogaspidia*). – Asia: China (Anhui, Zhejiang, Fujian, Guangdong).
- rubripes** (André, 1901) (*Mutilla*). – Asia: Syria, Palestine.
- tricolor** (Klug, 1829) (*Mutilla*). – Africa: Egypt, Eritrea, Ethiopia.
- unguiculata** (Magretti, 1884) (*Mutilla*). – Asia: ? Palestine; Africa: Eastern Africa.

VII. Subfamily SPHAEROPTHALMINAE Schuster, 1949(1903)

Sphaeropthalmini Ashmead, 1903.

Cystomutillinae Invrea, 1964.

Tribe Sphaeropthalmini Schuster, 1949(1903)

47. Genus *Cystomutilla* André, 1896

- ruficeps** (Smith, 1855) (*Mutilla*). – Europe: GB, F, CH, A, I, SL, H, R, Croatia, AL, E, U; Russia: SET; North Africa: Algeria.
- erythrocephala* (Lucas, 1849) (*Mutilla*).
- parvula* (Kriechbaumer, 1896) (*Mutilla*).
- teranishii** Mickel, 1935 (*Cystomutilla*). – Asia: Korea, Japan (Hokkaido, Honshu).

[Genus *Dasymutilla* Ashmead, 1899]

[**nigripes** Fabricius, 1787 (*Mutilla*). "Habitat in Oriente", actually North America]. [Not Palaearctic].

VIII. Subfamily DASYLABRINAЕ Skorikov, 1935

48. Genus *Dasylabris* Radoszkowski, 1885

Allomutilla Ashmead, 1899.

Subgenus **Baltilla** Lelej, 1976

- adagirica** Invrea, 1953 (*Dasylabris*). – North Africa: Morocco.
- adversa** Skorikov, 1935 (*Dasylabris*). – Europe: U; Russia: SET, ES (Transbaikalia); Asia: Mongolia.
- kaszabi* Suárez, 1974 (*Dasylabris*).
- arabica** (Olivier, 1811) (*Mutilla*). – Asia: Arabia, Palestine, Iraq; North Africa: Morocco, Algeria, Egypt.
- ornata* (Klug, 1829) (*Mutilla*).
- detrita** (André, 1903) (*Mutilla*). – North Africa: Algeria, Morocco.
- sefrana** (André, 1903) (*Mutilla*). – North Africa: Algeria.
- univittata** (André, 1903) (*Mutilla*). – North Africa: Algeria.
- bicolor** (Pallas, 1771) (*Mutilla*). – Russia: SET (Daghestan, Astrachanskaya oblast); Asia: KZ, KY, UZ, China: NW (Xinjiang), Mongolia (west).
- albeola* (Pallas, 1773) (*Mutilla*).
- caspica* (Kolenati, 1846) (*Mutilla*).
- funeraria* (Erichson, 1849) (*Mutilla*).
- bergi* Skorikov, 1935 (*Dasylabris*).
- crassa** (Klug, 1829) (*Mutilla*). – Africa: Egypt, Chad.
- gobicola** Skorikov, 1935 (*Dasylabris*). – Russia: WS (Altai, Tuva); Asia: KZ (east), Mongolia, China: NW (Xinjiang, Neimenggu).
- funerula* Suárez, 1974 (*Dasylabris*).
- hazara** Lelej, 1980 (*Dasylabris*). – Asia: Afghanistan.
- kiritshenkoi** Skorikov, 1935 (*Dasylabris*). – Asia: Iran.
- koenigi** Radoszkowski, 1888 (*Dasylabris*). – Asia: KZ (south), UZ, TJ, TU, Iran (north).
- caecia* Skorikov, 1935 (*Dasylabris*).
- argyroloma* Skorikov, 1935 (*Dasylabris*).
- kozlovi** Skorikov, 1935 (*Dasylabris*). – Asia: Mongolia, China: NW (Gansu, Neimenggu).
- mesomelaena** Bischoff, 1921 (*Dasylabris*). – North Africa: Egypt.
- miranda** Lelej, 1976 (*Dasylabris*). – Asia: KZ, KY, TU, Afghanistan.
- olivieri** (Sichel et Radoszkowski, 1870) (*Mutilla*). – Asia: Palestine; North Africa: Tunisia, Algeria, Egypt.
- atrata* sensu Olivier, 1811 (*Mutilla*).
- pavlovskyi** Skorikov, 1935 (*Dasylabris*). – Asia: UZ, TU.
- przewalskii** Lelej, 1976 (*Dasylabris*). – Asia: KZ (south-east), China: NW (Xinjiang).
- tingitana** (André, 1903) (*Mutilla*). – North Africa: Morocco.

Subgenus **Craspedopyda** Lelej, 1976

Craspedopyda Skorikov, 1935.

desertorum (Radoszkowski, 1865) (*Mutilla*). – Asia: KZ, KY, UZ, TU, TJ, Afghanistan, Mongolia, China: NW (Xinjiang).

scissa (André, 1903) (*Mutilla*).

deficiens Skorikov 1935 (*Dasylabris*).

manderstiernii (Radoszkowski, 1865) (*Mutilla*). – Europe: U (Crimea), GR; Russia: SET; Asia: GE, ARM, AZ, TU, Iran (north), TR, Lebanon, Syria, CY.

manderstiernii manderstiernii (Radoszkowski, 1865). – Europe: U (Crimea); Russia: SET; Asia: GE, ARM, AZ, TU, Iran (north), TR, Lebanon, Syria.

manderstiernii cypria (Sichel et Radoszkowski, 1870) (*Mutilla*). – Europe: GR; Asia: CY, TR, Syria.

mauropsis Skorikov 1935 (*Dasylabris*).

canonota Skorikov 1935 (*Dasylabris*).

Subgenus **Dasylabris** Radoszkowski, 1885

andrei Mercet, 1903 (*Dasylabris*). – North Africa: Western Sahara, Morocco.

angelae Suárez, 1959 (*Dasylabris*). – North Africa: Canary Is.

atrata (Linnaeus, 1767) (*Mutilla*). – Europe: S, P; North Africa: Morocco, Algeria, Libya.

arenaria (Fabricius, 1787) (*Mutilla*).

nilotica (Klug, 1829) (*Mutilla*).

levida (Klug, 1835) (*Mutilla*).

oraniensis (Lucas, 1849) (*Mutilla*).

pseudocypria Invrea, 1940 (*Dasylabris*).

[*aureomaculata* (Sichel et Radoszkowski, 1870) (*Mutilla*), – "Crimea" (Ukraine), actually Africa].

bucharensis Skorikov, 1935 (*Dasylabris*). – Asia: TU, UZ, TJ, Afghanistan.

algira Skorikov, 1935 (*Dasylabris*).

canariensis Suárez, 1970 (*Dasylabris*). – North Africa: Canary Is.

cinerescens (Olivier, 1811) (*Mutilla*). – Asia: Iran.

crenata (Radoszkowski, 1865) (*Mutilla*). – Asia: KZ (south), KY, UZ, TJ, China: NW (Xinjiang).

egregia (Klug, 1835) (*Mutilla*). – Europe: E.

atriventris (André, 1903) (*Mutilla*). – Europe: E.

calonata (André, 1903) (*Mutilla*).

garciamercti Suárez, 1962 (*Dasylabris*). – North Africa: Western Sahara.

iberica Giner 1942 (*Dasylabris*). – Europe: F, E, P.

valverdei Suárez, 1970 (*Dasylabris*). – Europe: E.

nevadensis Suárez, 1965 (*Dasylabris*). – Europe: E.

-
- intermedia** Skorikov, 1935 (*Dasylabris*). – Russia: ES (south Buryatia); Asia: Afghanistan, Mongolia, China: NO (Neimenggu, Hebei, Shandong), CE (Jiangsu, Zhejiang), SE (Fujian), Korea.
- ypsilon* Chen, 1957 (*Dasylabris*).
- juxtarenaria** Skorikov, 1935 (*Dasylabris*). – North Africa: Tunisia, Algeria, Morocco.
- atratoria* Petersen 1988 (*Dasylabris*).
- kondarensis** Lelej, 1985 (*Dasylabris*). – Asia: TJ.
- kruegeri** Invrea, 1953 (*Dasylabris*). – North Africa: Libya.
- lybica** Invrea, 1940 (*Dasylabris*). – Africa: Morocco, Algeria, Tunisia, Libya, Chad.
- libyca* Invrea, 1953 (*Dasylabris*).
- trilineata* Invrea, 1953 (*Dasylabris*).
- maura** (Linnaeus, 1758) (*Mutilla*). – Europe: P, S, F, CH, I, M, Croatia, BH, YU, Macedonia, AL, GR, BG, H, TR, A, CZ, SL, D, PL, U; Russia: SET, South Ural, WS (Altai, south); Asia: KZ, KY, ARM, AZ, TU (Kopet-Dag), TR, Iran (north), ?Palestine.
- maura maura** (Linnaeus, 1758). – Europe: F, CH, I, M, Croatia, BH, YU, Macedonia, AL, GR, BG, H, TR, A, CZ, SL, D, PL, U (Crimea). Asia: GE, ARM, AZ.
- austriaca* (Panzer, 1799) (*Mutilla*).
- rubicans* (Lepeletier, 1845) (*Mutilla*).
- argenteofasciata* (Costa, 1858) (*Mutilla*).
- trisinuosa* (Costa 1886) (*Mutilla*).
- nigrociliata* Hoffer, 1938 (*Dasylabris*).
- maura sungora** (Pallas, 1773) (*Mutilla*). – Russia: SET, South Ural, WS (Altai, south); Asia: KZ, KY.
- quadrimaculata* (Radoszkowski, 1865) (*Mutilla*).
- nostras* Skorikov, 1935 (*Dasylabris*).
- argentata* Skorikov, 1935 (*Dasylabris*).
- argyrauxa* Skorikov, 1935 (*Dasylabris*).
- consueta* Skorikov, 1935 (*Dasylabris*).
- iberica* Skorikov, 1935 (*Dasylabris*).
- infuscata* Skorikov, 1935 (*Dasylabris*).
- quinquemaculata* Skorikov, 1935 (*Dasylabris*).
- tricincta* Skorikov, 1935 (*Dasylabris*).
- maura clausa** (Lepeletier, 1845) (*Mutilla*). – Europe: P, S, F (south), CH.
- maura armeniaca** (Kolenati, 1846) (*Mutilla*). – Russia: SET (Daghestan); Asia: ARM, AZ, TU (Kopet-Dag), TR, Iran (north), ? Palestine.
- mlokosewitzii* Radoszkowski, 1885 (*Dasylabris*).
- duplicolor* Skorikov, 1935 (*Dasylabris*).
- maura carinulata** (Dalla Torre, 1897) (*Mutilla*). – Europe: F (Corsica), I

- (Sardinia).
- carinata* (Sichel et Radoszkowski, 1870) (*Mutilla*).
- mitra** Lelej, 1980 (*Dasylabris*). – Asia: Afghanistan (east).
- mixta** (André, 1903) (*Mutilla*). – Asia: ARM, AZ, TU, Iran (north).
- cirrata* Skorikov, 1935 (*Dasylabris*).
- mongolica** (Radoszkowski, 1861) (*Mutilla*). – Asia: KZ (south and east), KY, UZ, TU, TJ, Iran (norh), Afghanistan (Gazni), Mongolia, China: NW (Xinjiang).
- exarenaria* Skorikov, 1935 (*Dasylabris*).
- turana* Skorikov, 1935 (*Dasylabris*).
- rufocephala** (André, 1903) (*Mutilla*). – Asia: Palestine; North Africa: Western Sahara, Tunisia, Morocco, Mauritania.
- pseudomongolica** Ovchinnikov, 2002 (*Dasylabris*). – Asia: KZ (south-east), KY.
- sarafschani** (Radoszkowski, 1877) (*Mutilla*). – Asia: KZ (south), KY, UZ, TU, Iran (north-east).
- specta* (André, 1903) (*Mutilla*).
- argyronota* Skorikov, 1935 (*Dasylabris*).
- siberica** (Christ, 1791) (*Mutilla*). – Russia: WS (Altai), ES (Tuva, south of Krasnoyarskii krai, Transbaicalia), FE (Amurskaya oblast, Primorskii krai); Asia: Mongolia, China: NO (Neimenggu, Shanxi, Hebei), CE (Jiangsu), ? Korea.
- rubrosignata* (Radoszkowski, 1865) (*Mutilla*).
- quadrimaculata* (Sichel et Radoszkowski, 1870) (*Mutilla*).
- quadripunctata* (Sichel et Radoszkowski, 1870) (*Mutilla*).
- insignata* Skorikov, 1935 (*Dasylabris*).
- incompleta* Skorikov, 1935 (*Dasylabris*).
- skorikovi** Lelej, 1971 (*Dasylabris*). – Asia: KZ (south), KY.
- wahrmani** (Invrea, 1965) (*StenoMutilla*). – Asia: Israel.
- zarudnyi** Lelej, 1985 (*Dasylabris*). – Asia: Iran (east).

Subgenus Inbaltilla Lelej, 1976

- anceps** (Radoszkowski, 1877) (*Mutilla*). – Asia: KZ (south), UZ, TU.
- mirifica* Skorikov, 1935 (*Dasylabris*).
- biblica** Invrea, 1950 (*Dasylabris*). – Europe: GR (Crete); Asia: Palestine.
- defecta** (André, 1903) (*Mutilla*). – Asia: Caucas, Turkestan; North Africa: Algeria, Egypt.
- erronea** Lelej, 1985 (*Dasylabris*). – Asia: ARM, AZ, KZ, KY, UZ, TJ.
- [ferrugineiventris** (André, 1903) (*Mutilla*). – Distribution: unknown]. Probably not Palaearctic.
- gussakovskii** Skorikov, 1935 (*Dasylabris*). – Asia: KZ, KY, UZ, TU, Iran (north).
- indra** Lelej, 1980 (*Dasylabris*). – Asia: Afghanistan (east).

- jacobsoni** Skorikov, 1935 (*Dasylabris*). – Asia: KZ, KY.
- lugubris** (Fabricius, 1804) (*Mutilla*). – Europe: E, P; North Africa: Morocco, Algeria, Tunisia, Egypt.
- elegans* (Klug, 1835) (*Mutilla*).
 - tristis* (Klug, 1835) (*Mutilla*).
 - decorata* (Sichel et Radoszkowski, 1870) (*Mutilla*).
 - decoratissima* (Dalla Torre, 1897) (*Mutilla*).
 - almeriensis* Suárez, 1952 (*Dasylabris*).
- lyuateyi** Invrea, 1953 (*Dasylabris*). – North Africa: Morocco.
- miogramma** Skorikov, 1935 (*Dasylabris*). – Europe: U (south), R.
- paupertina** Skorikov, 1935 (*Dasylabris*). – Asia: ARM, AZ, TU, UZ, KY, TJ.
- obscura* Skorikov, 1935 (*Dasylabris*).
 - ditescens* Skorikov, 1935 (*Dasylabris*).
 - medialis* Lelej, 1975 (*Dasylabris*).
- popovi** Skorikov, 1935 (*Dasylabris*). – Asia: ARM, Iran (north).
- regalis** (Fabricius, 1793) (*Mutilla*). – Europe: CZ, SL, A, H, YU, BG, U; Russia: SET, South Ural, WS (Altai); Asia: KZ.
- petiolata* (Baer, 1848) (*Mutilla*).
 - petiolaris* (Radoszkowski, 1865) (*Mutilla*).
 - concolora* (Radoszkowski, 1865) (*Mutilla*).
 - unipetiolaris* (Radoszkowski, 1866) (*Mutilla*).
 - triangulomaculata* Skorikov, 1935 (*Dasylabris*).
 - slovaca* Hoffer, 1938 (*Dasylabris*).
 - batai* Hoffer, 1938 (*Dasylabris*).
- scutila** Skorikov, 1935 (*Dasylabris*). – Asia: TR, ARM, GE, AZ.
- shelkovnikovi** Lelej, 1985 (*Dasylabris*). – Asia: ARM.
- armeniaca* Skorikov, 1935 (*Dasylabris*).
- stenothoracica** Skorikov, 1935 (*Dasylabris*). – Asia: KZ (south), KY, UZ, TU, TJ.
- paupera* Skorikov, 1935 (*Dasylabris*).
 - nigrifica* Skorikov, 1935 (*Dasylabris*).
 - argentata* Skorikov, 1935 (*Dasylabris*).
- vittata** (Olivier, 1811) (*Mutilla*). – Asia: Iran (north), Iraq, Syria.
- zimini** Skorikov, 1935 (*Dasylabris*). – Asia: KZ, UZ, TU, TJ.
- kopetdaga* Skorikov, 1935 (*Dasylabris*).
- zonsteini** Ovchinnikov, 2002 (*Dasylabris*). – Asia: KY, TJ.

49. Genus *Jaxartilla* Lelej, 1984

fedtshenkoi Lelej, 1984 (*Jaxartilla*). – Asia: KZ.

50. Genus *Orientilla* Lelej, 1979

tausignata (Chen, 1957) (*Stenomutilla*). – Asia: China: CE (Anhui, Hunan, Jianxi, Zhejiang), SE (Fujian, Guangdong).

variegata (Smith, 1855) (*Mutilla*). – Asia: China: CE (Jiangsu, Anhui, Hunan, Zhejiang), SE (Fujian, Guangxi, Hainan, Taiwan), Vietnam, ?Myanmar.
desponsa (Smith, 1855) (*Mutilla*).

51. Genus *Stenomutilla* André, 1896

Subgenus *Stenomutilla* André, 1896

acanthosterna Nonveiller, 1994 (*Stenomutilla*). – Asia: TR.
aethiops Nonveiller, 1994 (*Stenomutilla*). – North Africa: Libya.
anatolica Nonveiller, 1994 (*Stenomutilla*). – Asia: TR.
argentata (Villers, 1789) (*Mutilla*). – Europe: I, F (south), E, P.
rondanii (Spinola, 1843) (*Mutilla*).
austriaca (Lepeletier, 1845) (*Mutilla*).
aucta (Lepeletier, 1845) (*Mutilla*).
saundersivora Ferton, 1921 (*Stenomutilla*).
carinata Skorikov, 1935 (*Stenomutilla*).
melanica Nonveiller, 1994 (*Stenomutilla*).
bicornuta Nonveiller, 1994 (*Stenomutilla*). – Asia: CY.
bizonata (Smith, 1856) (*Mutilla*). – Europe: Croatia, AL, GR: Asia: Lebanon, Syria, Israel, Iran, TR, ARM.
bifasciata (Klug, 1829) (*Mutilla*).
shelkovnikovi Skorikov, 1935 (*Stenomutilla*).
collaris (Fabricius, 1787) (*Scolia*). – Europe: E, P, Gibraltar, I (Sicily); North Africa: Morocco, Tunisia, Algeria.
sabulosa (Klug, 1835) (*Mutilla*).
angustisquama Skorikov, 1935 (*Stenomutilla*).
erlandssoni Suárez, 1961 (*Stenomutilla*).
erubescens Invrea, 1965 (*StenoMutilla*). – Asia: Israel, Jordan.
hottentota (Fabricius, 1804) (*Mutilla*). – Europe: I (south, Sicily), M; North Africa: Morocco, Tunisia, Algeria.
barbara (Fabricius, 1793) (*Mutilla*).
luctuosa (Lucas, 1849) (*Mutilla*).
parens (Costa, 1860) (*Mutilla*).
propinqua Nonveiller, 1994 (*Stenomutilla*).
impressiventris Nonveiller, 1994 (*Stenomutilla*). – North Africa: Western Sahara.
invreai Nonveiller, 1994 (*Stenomutilla*). – North Africa: Tunisia.
osellai Nonveiller, 1994 (*Stenomutilla*). – Asia: TR.
quadricincta Giner, 1945 (*Stenomutilla*). – North Africa: Western Sahara.
rubripes Invrea, 1967 (*StenoMutilla*). – Asia: Israel, Jordan.
rubripes Invrea, 1965 (*Stenomutilla*).
rufinodis Giner, 1947 (*StenoMutilla*). – North Africa: Western Sahara.

- rungsi** Giner, 1959 (*Stenomutilla*). – North Africa: Morocco.
- sejugis** (Radoszkowski, 1885) (*Dasylabris*). – Asia: KZ (south), KY, UZ, TJ, Iran (north), Afghanistan.
- chorezmiensis* Skorikov, 1935 (*Stenomutilla*).
- infusca* Skorikov, 1935 (*Stenomutilla*).
- tabida** (Lucas, 1849) (*Mutilla*). – North Africa: Algeria.
- tetrazonia** Skorikov, 1935 (*Stenomutilla*). – Asia: KZ (south), TU, UZ, TJ, Mongolia.
- ferrugineocapitata* Skorikov, 1935 (*Stenomutilla*).

52. Genus *Tricholabiodes* Radoszkowski, 1885

- aegyptiacus** (Radoszkowski, 1876) (*Mutilla*). – North Africa: Egypt.
- asiaticus** Radoszkowski, 1885: (*Tricholabiodes*). – Asia: KZ (south), UZ, TU, TJ, Iran.
- bactrianus** Suárez, 1967 (*Tricholabiodes*). – Asia: TU, Iran.
- beludzhistanus** Lelej, 1985 (*Tricholabiodes*). – Asia: Iran.
- garamantis** Suárez, 1967 (*Tricholabiodes*). – North Africa: Algeria.
- israeliticus** Suárez, 1967 (*Tricholabiodes*). – Asia: Israel.
- mandibularis** Lelej, 1985 (*Tricholabiodes*). – Asia: Iran.
- marmarica** Invrea, 1932 (*Tricholabiodes*). – North Africa: Libya, Tunisia.
- niloticus** Suárez, 1967 (*Tricholabiodes*). – Africa: Egypt, Libya, Mauritania, Ethiopia, Sudan.
- nuristanus** Lelej, 1980 (*Tricholabiodes*). – Asia: Afghanistan (east).
- pallidicornis** Bischoff, 1920 (*Tricholabiodes*). – North Africa: Algeria, Tunisia, Libya, Western Sahara.
- pallidus** Lelej, 1985 (*Tricholabiodes*). – Asia: Iran.
- patrizii** Invrea, 1932 (*Tricholabiodes*). – North Africa: Libya (south), Chad.
- persicus** Suárez, 1967 (*Tricholabiodes*). – Asia: Iran
- saharicus** Suárez, 1967 (*Tricholabiodes*). – North Africa: Algeria, Morocco, Western Sahara, Mauritania, Libya, Chad.
- semistriatus** (Klug, 1829) (*Mutilla*). – Asia: Israel, Arabia, Oman; North Africa: Egypt, Libya, Eritrea, Somali, Cameroon.
- pedunculatus* (Klug, 1829) (*Mutilla*).
- palaestinensis** Suárez, 1967 (*Tricholabiodes*). – Asia: Israel.
- stenothorax** Suárez, 1967 (*Tricholabiodes*). – North Africa: Western Sahara, Morocco.
- sudanensis** Suárez, 1967 (*Tricholabiodes*). – Africa: Egypt, Libya, Eritrea, Chad, Mali, Cameroon.

IX. Subfamily EPHUTINAE Ashmead, 1903

Tribe Odontomutillini Lelej, 1983

53. Genus **Odontomutilla** Ashmead, 1899

sinensis (Smith, 1855) (*Mutilla*). – Asia: China: CE (Jiangsu, Anhui, Zhejiang), SE (Fujian).

54. Genus **Yamanetilla** Lelej, 1996

nipponica (Tsuneki, 1972) (*OdontoMutilla*). – Asia: Japan (Honshu).

quadruplex (Chen, 1957) (*Odontomutilla*). – Asia: China: CE (Jiangsu).

X. MUTILLIDAE INCERTAE SEDIS

Genus **Cretavus** Sharov, 1957

sibiricus Sharov, 1957 (fossil) (*Cretavus*).

Genus **Mutilla** Linnaeus, 1758

citigrada Gistel, 1857 (*Mutilla*). – Europe: P.

clythrae Rosenhauer, 1856 (*Mutilla*). – Europe: E.

defunctatrix Gistel, 1857 (*Mutilla*). – Europe: P.

dimidiata (Latreille, 1792) (*Mutilla*). – Europe: F.

interrupta Latreille, 1792 (*Mutilla*). – Europe: F (south).

irritabilis Smith, 1879 (*Mutilla*). – North Africa: Egypt.

lichtensteinii Tournier, 1889 (*Mutilla*). – Europe: F (south).

myrmecium Klug, 1817 (*Mutilla*). – Europe: I.

pertyana Gistel, 1857 (*Mutilla*). – Europe: P.

pulverifex Gistel, 1857 (*Mutilla*). – Europe: P.

pyriformis Gistel, 1857 (*Mutilla*). – Europe: I.

saltatrix Scopoli, 1763 (*Mutilla*). – Europe: I.

schmiedeknechti Bischoff, 1910 (*Mutilla*). – Europe: D

taurica Gistel, 1857 (*Mutilla*). – Europe: U (Crimea).

westerhauseri Gistel, 1857 (*Mutilla*). – Europe: I.

Genus **Myrmosa** Latreille, 1796

muscifex Gistel, 1857 (*Myrmosa*). – Europe: P.

psammodes Gistel, 1857 (*Myrmosa*). – Europe: P.

PALAEARCTIC SPECIES DESCRIBED OUTSIDE OF PALAEARCTIC REGION

Mutilla robusta Smith, 1855 "Hew Holland" actually is *Ronisia brutia* (Petagna, 1787) (Europe).

Physetopoda insularis Schuster, 1949 "Haiti" actually is *Physetopoda rubro-cincta* (Lucas, 1849) (Europe).

Mutilla dorsata Fabricius, 1798 "Habitat in Americae Insulia", actually "Genes" (Italy).

SPECIES ERRONEOUSLY DESCRIBED FROM PALAEARCTIC REGION

Mutilla caucasica Radoszkowski, 1865 "Caucasus" actually is a synonym of *Bothriomutilla ruginollis* (Westwood, 1843) (Australia).

Mutilla kuschakewitzi Radoszkowski, 1888 "Wernoje" (Kazakhstan) actually is a synonym of *Trogaspidia nigra* (Smith, 1858) (Papua New Guinea).

Mutilla castellana Mercet, 1903 "Madrid" actually is *Trogaspidia castellana* (Mercet, 1903) (Philippines).

Mutilla aureomaculata Sichel et Radoszkowski, 1870 "Crimea" (Ukraine) actually is a synonym of *Dasylabris merope* (Smith, 1855) (Africa).