

Kramerius 5

Digital library

Conditions of use

The Czech Academy of Sciences Library provides access to digitalised documents only for non-commercial, academic and study purposes and only for the personal requirements of library users. Some of the documents in the Digital Library of the Czech Academy of Sciences are subject to copyright. By using the digital library and the generation of a copy of part of a digitalised document the user undertakes to observe these conditions, which must comprise part of each copy made. No further copying of any kind of materials from the digital library is possible without the written permission of the Czech Academy of Sciences Library.

Main title: **Časopis Československé společnosti entomologické**

Publisher: **Nakladatelství ČSAV**

Published in: **1919-1952**

Periodical volume number: **28**

Periodical item date: **1931**

ISSN identifier: **0862-0245**

SICI identifier: **nezjištěn**

Page: **101**

NOVAE STOMINORUM¹⁾) FORMAE E AMERICA MERIDIONALI.

V. Lutshnik.

Tichonia peruana sp. n.

Piceo-nigra, parum nitida, elytris indistincte violaceo mican-
tibus, ore cum antennis pedibusque nigris, palpis rufescentibus. Ca-
pite lato, laevi, impressionibus frontalibus profundis. Oculis magnis,
convexit. Prothorace transverso, latitudine ad angulos basalis 1,2
breuiore, capite cum oculis multo latiore, basi parum emarginato,
antice fere recto, lateribus antice rotundato, latitudinem maximam
in medio attingente, basin sensim angustato, ante angulos basales
distincte sed leviter sinuato, his subacutis, angulis anticis rotundatis,
haud prominulis; disco medio convexo, linea media profunda
utrinque abbreviata, striga basali interna lata, externa oblitterata.
Elytris sat — latis, ellipticis, ad humeros thoracis basi haud la-
tioribus, latitudine 1,43 longioribus, lateribus multo rotundatis,
ante apicem sinuatis, apice acutis, profunde striatis, interstitiis
convexit. Sterno et abdome laevibus. Prosterni processu intercoxali
apice rotundato, haud marginato, segmento ultimo in ♀ utrinque
foveolis setigeris 2 instructo.

Long. 17 mm.

Peru: Chanchamyo. 1 ♀ in coll. m. Statura latiore, thorace
lateribus postice angustatis ante angulos basales sinuatis ab
speciebus hujus generis facile distinguenda.

Platysma (*Parhypates*) **irrequietum** sp. n. Majus, haud con-
vexum, piceo-nigrum, parum nitidum, antennis, palpis pedibusque
rufo-piceis. Capite lato, laeve, impressionibus frontalibus sat pro-
fundis, setis ocularibus utrinque 2. Prothorace transverso, capite
cum oculis latiore, cordato, anterius rotundato, latitudinem maxi-
mam ante medium attingente, basin versus angustato, ante an-
gulos posticos rectos sinuato, apice emarginato, angulis anticis
subprominulis, rotundatis, basi haud punctato, striga basali
haud profunda, linea media tenuiter impressa utrinque abbreviata,
setis lateralibus utrinque 3. Elytris latiusculis, lateribus rotundatis,
ad humeros thoracis basi distincte latioribus, ante apicem sinuatis,
dorso subplanatis, tenuiter striatis, striis subtilissime punctatis,
interstitiis planis, tertio ad striam tertiam punctis 3 instructo. Sub-
tus corpore laevi, prosterni processu intercoxali apice immarginato.

Long. 15 mm.

Chili. 1 specimen in coll. m. *Pl. gayi* Chaud. et *chalybicolor*
Chaud. affine, sed corpore haud metallico, lato, minus convexo
praesertim distinguendum.

Pl. (*Parhypates*) **profundestriatum** Motsch. ab. **expletum** ab.
n. Formae typicae simile, sed elytrorum interstitio tertio punctis
tribus distinctissimis notato.

Long. 12,5 mm.

Chili. 1 ♀ in coll. m.

¹⁾ Stomina = Pterostichina (cf. **Lutshnik**, Acta Societ. Entomol. Staurop., III., 1927, 51).