

INSS 1996–1499

2014 №1



Российский Журнал Биологических Инвазий

<http://www.sevin.ru/invasjour/>



Институт проблем экологии и эволюции
имени А.Н. Северцова
Российской Академии Наук

Российской академии наук
Институт проблем экологии и эволюции им. А.Н. Северцова (ИПЭЭ РАН)

Российский Журнал Биологических Инвазий

(ISSN – 1996–1499)

Основан в январе 2008 г.

Выходит 4 раза в год

Главный редактор
академик РАН Дгебуадзе Юрий Юлианович

Заместитель главного редактора
д.б.н., Петросян Варос Гарегинович

Ответственный секретарь
к.б.н., Дергунова Наталья Николаевна

Редакционная коллегия
к.б.н., В.В. Бобров, д.б.н., Ю.К. Виноградова, д.б.н., А.Ю. Звягинцев,
д.б.н., С.С. Ижевский, д.б.н., И.Н. Ильин, д.б.н., Крылов А.В.,
к.б.н., В.Ю. Масляков, к.б.н., О.В. Морозова, академик РАН, Д.С. Павлов,
д.б.н., А.Н. Пельгунов, д.б.н., Н.М. Пронин, к.б.н., Ю.В. Слынько,
д.б.н., И.В. Телеш, к.б.н., И.Ю. Фенева, к.б.н., Л.А. Хляп, д.б.н., Шиганова Т.А.,
д.б.н., Г.Х. Щербина

Тематика журнала

Теоретические вопросы биологических инвазий (теория, моделирование, результаты наблюдений и экспериментов): инвазионные коридоры, векторы инвазий, адаптации видов-вселенцев, уязвимость аборигенных экосистем, оценка риска инвазий, генетические, экологические, биологические, биогеографические и эволюционные аспекты влияния чужеродных видов на биологическое разнообразие биосистем различных уровней организации.

Мониторинг инвазионного процесса (сообщения о нахождении организмов за пределами естественного ареала, динамике расселения, темпах натурализации).

Методы, средства накопления, обработки и представления данных прикладных исследований (новые разработки, моделирование, результаты исследований) с применением фактографических и геоинформационных систем.

Использование результатов исследований биологических инвазий (методы и новые фундаментальные результаты) при изучении морских, пресноводных и наземных видов, популяций, сообществ и экосистем.

Контроль, рациональное использование и борьба с видами-вселенцами.

Индексирование журнала – SCOPUS, ПИНЦ, Google Scholar, Academic OneFile,
Summon by Serial Solutions, OCLC, CAB International, Global Health

Адрес: Россия, 119071, Москва, Ленинский проспект, д. 33.
тел. (495) 954-75-53; факс (495) 954-55-34;

Е-mail: invasjour@sevin.ru
<http://www.sevin.ru/invasjour/>

ALIEN BEETLE SPECIES IN THE REPUBLIC OF MOLDOVA: A REVIEW OF THEIR ORIGIN AND MAIN IMPACT

© 2014 Munteanu N.¹, Moldovan A.¹, Bacal S.², Toderas I.¹

¹ Laboratory of Systematics and Molecular Phylogeny, Institute of Zoology, Moldova Academy of Sciences, MD 2028, Chisinau, Academiei 1, Republic of Moldova, e-mail: munteanu_natalia_v@yahoo.com

² Laboratory of Entomology, Institute of Zoology, Moldova Academy of Sciences, MD 2028, Chisinau, Academiei 1, Republic of Moldova

Received September 15, 2013

A list of beetles considered to be of alien origin in the Republic of Moldova has been set up, with data on their origin, habitat, trophic range and economic impact. Altogether, 75 alien beetle species have been revealed, which represents 2.98% of local coleopteran fauna. The most numerous taxonomic groups are Curculionidae (30 species), Staphylinidae and Chrysomelidae (by 11). The majority of non-native beetles originates in Europe, Asia and Mediterranean region, while comparable less number of species originates in Africa and North America. From the total number of the investigated alien beetles 26.67% are agricultural, forest and storage pests.

Key words: alien species, beetles, Republic of Moldova.

Introduction

Invertebrates, and among them arthropods, represent the greatest part of alien animals and are of pronounced importance in agriculture, horticulture and forestry, with a major economic impact. Their activity can alter ecosystems structure and functioning and eventually lead to the extinction of native species, inducing biodiversity losses [Elton, 1958; Nentwig, 2008; Hulme et al., 2009]. Vulnerability of ecosystems to alien species is probably one of the least studied and most difficult questions. Knowledge of alien species effects and potential threats is still insufficient being crucial for managing the risks related to species transfer.

Establishment of first alien arthropods checklists in Europe started in 2002 and covered Austria [Essl, Rabitsch, 2002], Germany [Geiter et al., 2002], the Netherlands [Reemer, 2003], the Czech Republic [Šefrová, Laštůvka, 2005], the United Kingdom [Hill et al., 2005, Smith et al., 2007], Serbia and Montenegro [Glavendekić et al., 2005], Switzerland

[Kenis, 2005], Israel [Roll et al., 2007], Albania, Bulgaria and Macedonia [Tomov et al., 2009], and Russia [Maslyakov, Izhevsky, 2011].

The geographical position of the Republic of Moldova at the crossing point of three biogeographical zones (Central European, Eurasian and Mediterranean) along with the intense human activities makes it especially prone to alien species entry. Despite this, the problem of alien species on the investigated territory was not treated in the scientific literature. Available information contains some data on certain taxa or groups of beetles, their trophic preferences and distribution but it does not always contain information on their origin.

The main goal of this study was to determine which alien beetle species entered, spread and/or established in natural transitional, forest and agricultural ecosystems of the Republic of Moldova, its origin and impact, in order to better understand how to manage non-native beetles. Only in this manner the most

Table 1. Composition of alien beetle species in individual families; S – eusynanthropic, C – casual, N – naturalized non invasive, *N – naturalized post invasive, I – invasive.

Taxon	Spp. RM	Aliens	% of all	S	C	N	*N	I
Staphylinidae	321	11	3.42	–	–	–	11	–
Scarabaeidae	124	1	0.8	–	–	–	1	–
Dermestidae	14	1	7.1	–	–	–	1	–
Ptinidae	22	1	4.54	–	–	–	1	–
Coccinellidae	33	1	3.03	–	–	–	–	1
Tenebrionidae	52	5	9.61	5	–	–	–	–
Meloidae	26	1	3.84	–	–	–	1	–
Cerambycidae	125	2	1.6	–	1	–	1	–
Chrysomelidae	197	11	5.58	2	1	3	4	1
Brentidae	90	8	8.88	–	–	3	5	–
Dryophthoridae	5	3	60	2	1	–	–	–
Curculionidae	510	30	5.88	–	7	16	6	1
Total Coleoptera	2512	75	2.98	9	10	22	31	3

efficient methods to prevent new arrivals and/or disseminations, and reduce their impacts, can be proposed and implemented.

Material and methods

To initiate the first list of non-native beetle species in the Republic of Moldova, we reviewed professional published and unpublished information and available reports, particularly checklists of Miller, Zubowsky [1917], Medvedev, Shapiro [1957], Adashkevich [1972], Poiras [1991, 1994, 1998, 2006], Neculiseanu, Matalin [2000], Neculiseanu, Bacal [2005], Munteanu [2006], Bacal, Stan [2006], Bacal, Derunkov [2010], Derunkov, Bacal [2011]. Also, museum specimen collection of the Institute of Zoology, Academy of Sciences of Moldova and alien invasive species inventories websites were used.

Data were compiled into a preliminary list of non-native beetles documented in the country (either currently or historically). For each species we included information about its origin, first report, habitat, trophic preferences and host. Further, based on evidence in the literature, we identified a number of non-native species known as pests of agricultural crops, fruit trees, forests, ornamental plants and storage products. Distribution in alien species categories (eusynanthropic, casual, naturalized non invasive, naturalized post invasive and invasive) was performed [Šefrová, Laštůvka, 2005; Nentwig, Josefsson, 2010].

The taxonomy of the beetle species is based primarily on the works of Kryzhanovskij [1965] and Bouchard et al. [2011]. Biological and geographical classification was done according to concepts of Darlington [1966] and Banareescu [1970].

Results and discussions

In the Republic of Moldova about 2512 beetle species are known [Bacal et al., 2013]. From the total number, 75 ones (2.98%) represent alien species. Among them, 9 species (12%) create short or long-term populations only in heated spaces, with the ability to survive outside only during warm periods of the year, but not under winter conditions. A group of 10 species (13.33%) creates occasionally short-term populations in outdoor environments. The populations of these species are usually decimated by adverse weather conditions, surviving winter with difficulty or gradually disappearing for other reasons. Another 56 species (74.66%) can be considered naturalized; their populations survive for long time tolerating easily the local climatic conditions. Between these species, 22 (29.33% of aliens) ones remain non-invasive, other 31 (41.33%) species are post-invasive at present, and 3 species (4%) are invasive. The categories of aliens with the numbers of species are shown in the Table 1.

The greatest number of identified alien species (30) belongs to Curculionidae

family, followed by Staphylinidae, Chrysomelidae, Brentidae and Tenebrionidae. The largest proportion of species of alien origin is in Dryophthoridae (60%) and Tenebrionidae (9.61%), while for the rest it varies between 0.8–8.88 percent. By contrast, no alien species were documented in other 57 coleopteran families which occur in the Republic of Moldova.

The majority of revealed eusynanthropic alien species (5) are naturally found in tropical and subtropical regions; other two species are cryptogenic and by one are from North America and Asia. The group of casual species is represented by four European, three Asian and three Mediterranean species. Among 22 naturalized non-invasive species 8 ones originate in Eurasia and Mediterranean region, 5 species are Mediterranean, by three European and Eurasian, two Euro-Mediterranean and one North American species. Concerning naturalized post-invasive species the most of them (8) derive from Eurasia and Mediterranean region, five are Eurasian, four Asian, three European, by two African and Mediterranean, and one from America. This group also contains six cryptogenic

species. Invasive species originate from North America (2) and Asia (1). Detailed information on each species is presented in the Table 2.

From the total number of non-native beetles the majority originates in Europe, Asia and Mediterranean region and spreads with the progress of human civilization, expansion of agricultural crops and for other various reasons. Comparable less number of species originates in America; their dissemination is partly due to the human travel and trade activity, spontaneous spreading of those species is being almost impossible (the geographical barrier is difficult to pass without human intervention).

Alien species impact on economy and environment is a result of trophic and habitat preferences. Concerning habitats around 41.33% of the established alien beetle species colonize cultivated habitats. The second most populated are urban settlements (38.66%). Grasslands are also frequent habitats (28%). Woodlands and forest habitats are occupied by 20% of the non-native beetles. Heathlands, hedgerows and shrubs are populated by 5.33% of the species and only 4% occur in wetlands.

Table 2. Characteristics of alien beetle species in the Republic of Moldova.

Status of alien species and their occurrence: S – eusynanthropic, C – casual, N – naturalized non invasive, *N – naturalized post invasive, I – invasive; Habitats: B – coastal habitats; E – grasslands; F – heathlands, hedgerows and shrub plantations; G – woodlands and forests; H – cultivated habitats; J – urban settlements, N/A – data non available; Trophic range: Pr – Predator, De – Detritivorous, Ph – Phytophagous (herbiphagous (her), spermatophagous (spe), leaf/stem browser (lbw), root browser (rbw), root borer (rbo), phloeophagous (phl), saproxylophagous (sxp), xylomycetophagous (xmp), Un – Unknown.

Species	Origin	Status	Habitat	Trophic range	Host	Reference
Staphylinidae						
<i>Cilea silphoides</i> (Linnaeus, 1767)	Cryptogenic	*N	J	Pr	Cattle dung	Mihailov, 2012
<i>Atheta oblita</i> (Erichson, 1839)	Northern Europe	*N	G, J	Pr	Fungus, cattle dung	Adashkevich, 1972
<i>Carpelimus corticinus</i> (Gravenhorst, 1806)	Cryptogenic	*N	B, E	Ph	Floodplains, river banks, sand	Adashkevich, 1972
<i>C. gracilis</i> (Mannerheim, 1830)	Cryptogenic	*N	B, E	Pr	Floodplains, river banks, sand	Adashkevich, 1972
<i>C. pusillus</i> (Gravenhorst, 1802)	Cryptogenic	*N	B, E	Pr	Floodplains, river banks, sand	Adashkevich, 1972
<i>Lithocharis nigriceps</i> Kraatz, 1859	Asia-Tropical	*N	H, J	Pr	Compost	Neculiseanu, 1984
<i>L. ochracea</i> (Gravenhorst, 1802)	Eurasia	*N	J	Un	Dried vegetals	Miller, Zubowsky, 1917

<i>Gabrieus nigrutilus</i> (Gravenhorst, 1802)	Eurasia	*N	J	Un	Fungus	Yatsentkovsky, 1912
<i>Philonthus concinnus</i> (Gravenhorst, 1802)	Eurasia	*N	J	Pr	Fungus, cattle dung	Adashkevich, 1972
<i>P. discoideus</i> (Gravenhorst, 1802)	Northern Africa, Eurasia	*N	J	Pr	Dung	Adashkevich, 1972
<i>P. longicornis</i> Stephens, 1832	Eurasia	*N	J	Pr	Dung	Miller, Zubowsky, 1917
Scarabaeidae						
<i>Pleurophorus caesus</i> (Creutzer, 1796)	Eurasia, America	*N	E	De	Dung	Miller, Zubowsky, 1917
Dermestidae						
<i>Dermestes ater</i> De Geer, 1774	Cryptogenic	*N	J	De	Necrophagous	Miller, Zubowsky, 1917
Ptinidae						
<i>Ptinus fur</i> (Linnaeus, 1758)	Cryptogenic	*N	J	De	Waste, dried vegetals	Miller, Zubowsky, 1917
Coccinellidae						
<i>Harmonia axyridis</i> (Pallas, 1773)	Asia	I	H	Pr	Polyphagous, predator mainly aphids and coccids	Iazlovetchi, Sumenkova, 2013
Tenebrionidae						
<i>Tribolium castaneum</i> (Herbst, 1797)	Cryptogenic	S	J	De	Stored products	Neculiseanu, Bacal, 2005
<i>T. confusum</i> Jacquelin du Val, 1868	Africa	S	J	De	Stored products	Medvedev, Shapiro, 1957
<i>T. destructor</i> Uyttenboogaart, 1933	Tropical	S	J	De	Stored products	Neculiseanu, Bacal, 2005
<i>Alphitobius diaperinus</i> (Panzer, 1797)	Tropical, Subtropical	S	J, G	Pr, De	Residues, inhabitant of chicken houses, on faeces and wastes	Neculiseanu, Bacal, 2005
<i>Alphitophagus bifasciatus</i> (Say, 1823)	Cryptogenic	S	J, G	De	Residues, compost, rotten fruits, under bark old stumps	Neculiseanu, Bacal, 2005
Meloidae						
<i>Mylabris variabilis</i> (Pallas, 1781)	Eurasia	*N	E	Pr, Ph (her)	Adult floricolous, larvae parasite of Acrididae	Miller, Zubowsky, 1917
Cerambycidae						
<i>Rosalia alpine</i> (Linnaeus, 1758)	Central Europe, Alps	C	G, H, J	Ph (sxp)	<i>Fagus</i> , and other deciduous trees	Miller, Zubowsky, 1917
<i>Monochamus sutor</i> (Linnaeus, 1758)	Northern and Central Europe	*N	G	Ph (sxp)	<i>Picea</i> , <i>Pinus</i>	Baban, 2006
Chrysomelidae						
<i>Callosobruchus chinensis</i> (Linnaeus, 1758)	Asia-Temperate	C	H, J	Ph (spe)	Stored legumes	Munteanu, 2006
<i>Bruchus pisorum</i> (Linnaeus, 1758)	Asia-Temperate	S	H, J	Ph (spe)	<i>Lathyrus</i> , <i>Pisum</i> , <i>Vicia</i>	Medvedev, Shapiro, 1957
<i>B. rufimanus</i> Boheman, 1833	Africa	*N	H, J	Ph (spe)	<i>Phaseolus</i> , <i>Lathyrus</i> , <i>Lupinus</i> , <i>Pisum</i> , <i>Lens</i> , <i>Cicer</i> , <i>Vicia</i>	Medvedev, Shapiro, 1957
<i>B. signaticornis</i> Gyllenhal, 1833	Mediterranean region	*N	H, J	Ph (spe)	<i>Lathyrus</i> , <i>Lens</i> , <i>Vicia</i> seeds	Munteanu, 2009
<i>Bruchidius lividimanus</i> (Gyllenhal, 1833)	Mediterranean region	*N	H	Ph (spe)	<i>Genistea</i> , <i>Ononis</i> , <i>Cytisus</i> seeds	Munteanu, 2006
<i>Acanthoscelides obtectus</i> Say, 1831	America	S	H, J	Ph (spe)	<i>Phaseolus</i> seeds, wild and cultivated legumes outdoors	Medvedev, Shapiro, 1957

<i>A. pallidipennis</i> (Motschulsky, 1874)	North America	N	H, J	Ph (spe)	Fabaceae	Munteanu, 2006
<i>Cassida atrata</i> Fabricius, 1787	Central, South East Europe	N	H	Ph (her)	<i>Lamiaceae</i>	Calestru, 2003
<i>Crioceris asparagi</i> (Linnaeus, 1758)	Europe, Central Asia	N	H, J	Ph (her)	<i>Asparagus</i>	Medvedev, Shapiro, 1957
<i>Leptinotarsa decemlineata</i> Say, 1824	Northern and Central America	I	H	Ph (her)	<i>Solanum tuberosum</i> , other Solanaceae	Vorotyntseva, 1971
<i>Goniocetena fornicate</i> (Brüggemann, 1873)	Eastern Europe	*N	H	Ph (her)	<i>Medicago</i>	Medvedev, Shapiro, 1957
Brentidae						
<i>Aspidapion validum</i> (Germar, 1817)	Asia-Temperate	*N	H, F	Ph (spe)	<i>Alcea rosae</i>	Miller, Zubowsky, 1917
<i>A. radiolus</i> (Marsham, 1802)	Asia, Europe, Mediterranean region	*N	N/A	Ph (her)	Malvaceae	Medvedev, Shapiro, 1957
<i>Alocentron curvirostre</i> (Gyllenhal, 1833)	Asia-Temperate	*N	H, F	Ph (spe)	<i>Alcea rosae</i>	Solodovnikova, Talitskii, 1972
<i>Squamapion leucophaeatum</i> (Wencker, 1864)	Mediterranean region	N	E	Ph (spe)	No data	Solodovnikova, Talitskii, 1972
<i>Taeniapion rufulum</i> (Wencker, 1864)	Europe, Mediterranean region	N	E, F	Ph (her)	<i>Urtica dioica</i>	Poiras, 1994
<i>Rhopalapion longirostre</i> (Olivier, 1807)	Asia-Temperate	*N	H	Ph (spe)	<i>Alcea rosae</i>	Miller, Zubowsky, 1917
<i>Trichopterapion holosericeum</i> (Gyllenhal, 1833)	Asia, Europe, Mediterranean region	N	G	Ph (spe)	<i>Carpinus betulus</i> , <i>C. orientalis</i>	Poiras, 1998
<i>Ischnopterapion virens</i> (Herbst, 1797)	Asia, Europe, Mediterranean region	*N	N/A	Ph (her)	<i>Trifolium</i>	Poiras, 1994
Dryophthoridae						
<i>Sitophilus oryzae</i> (Linnaeus, 1763)	Asia-Tropical	S	J	Ph (spe)	Grain	Poiras, 1998
<i>S. granarius</i> Linnaeus, 1758	Africa	S	J	Ph (spe)	Stored grains, <i>Secale Triticum</i> , <i>Zea mays</i>	Miller, Zubowsky, 1917
<i>Acentrus histrio</i> Boheman, 1845	Mediterranean region	C	E	Ph (her)	<i>Glaucium</i>	Medvedev, Shapiro, 1957
Curculionidae						
<i>Lignyodes bischoffi</i> (Blatchley, 1916)	North America	I	G, H	Ph (spe)	<i>Fraxinus</i> , <i>Syringa</i>	Poiras, 1987
<i>Tychius caldaria</i> Dieckmann, 1986	Mediterranean region	N	E	Ph (her)	<i>Dorycnium suffruticosum</i> , <i>D. pentaphyllum</i>	Medvedev, Shapiro, 1957
<i>Rhynchaenus sparsus</i> Fahraeus, 1843	Mediterranean region	N	G	Ph (lbw)	<i>Quercus robur</i> , <i>Q. pubescens</i>	Poiras, 1991
<i>Baris janthina</i> Boheman, 1836	Asia, Europe, Mediterranean region	N	E	Ph (lbw)	<i>Caradaria draba</i>	Medvedev, Shapiro, 1957
<i>Baris prasina</i> (Boheman, 1836)	South Europe	C		Ph (lbw)	Brassicaceae	Poiras, 1998
<i>Rhinoncus pericarpus</i> (Linnaeus, 1758)	Asia, West Mediterranean region, Europe	*N	E, H	Ph (rbo)	<i>Rumex</i> , <i>Medicago</i> , <i>Melilotus</i>	Miller, Zubowsky, 1917
<i>Thamiochilus uniformis</i> (Gyllenhal, 1837)	Asia, Europe, Mediterranean region	N	E	Ph (her)	<i>Phlomis pungens</i>	Poiras, 1998

<i>Ranunculiphilus kuntzei</i> (Smreczynski, 1957)	Mediterranean region	N	E	Ph (her)	No Data	Poiras, 1998
<i>Otiorynchus albidus</i> Stierlin, 1861	Asia, West Mediterranean region, Europe	*N	E, H	Ph (lbw)	<i>Cerasus, Juglans, Populus, Tamarix, Tilia</i>	Ruscinski, 1937
<i>Trachyphloeus spinimanus</i> Germar, 1824	Europe, Asia	C	N/A	Ph (rbw)	<i>Cynodon</i>	Poiras, 1998
<i>Phyllobius canus</i> Gyllenhal, 1834	Mediterranean region, Asia	C	G, H	Ph (lbw)	Rosaceae, <i>Quercus robur</i>	Plugar, 1963
<i>Cycloderes canescens</i> (Rossi, 1792)	Mediterranean region	C	E	Ph (lbw)	<i>Plantago</i>	Poiras, 1998
<i>Sitona lineatus</i> (Linnaeus, 1758)	Asia, Europe, Mediterranean region	*N	H, J	Ph (rbw)	Fabaceae	Medvedev, Shapiro, 1957
<i>S. macularius</i> (Marsham, 1802)	Asia, Europe, Mediterranean region	*N	H, J	Ph (rbw)	<i>Trifolium</i>	Medvedev, Shapiro, 1957
<i>S. puncticollis</i> Stephens, 1831	Asia, Europe, Mediterranean region	*N	H	Ph (rbw)	<i>Trifolium, Melilotus</i>	Ruscinski, 1937
<i>Aparopion costatum</i> Fahraeus, 1843	Europe, Mediterranean region	C	E	Ph (her)	No data	Arnoldi et al., 1965
<i>Hypera nigrirostris</i> (Fabricius, 1775)	Asia, Europe, Mediterranean region	N	E, J	Ph (lbw)	<i>Ononis, Trifolium</i>	Medvedev, Shapiro, 1957
<i>H. postica</i> (Gyllenhal, 1813)	Asia, Europe, Mediterranean region	*N	H, J	Ph (lbw)	Fabaceae	Antonova, 1977
<i>H. pastinacae</i> (Rossi, 1790)	Asia, Europe, Mediterranean region	N	G, H	Ph (lbw)	<i>Pastinaca sativa, Daucus carota</i>	Medvedev, Shapiro, 1957
<i>H. contaminata</i> (Herbst, 1795)	Europe	N	E	Ph (lbw)	<i>Lathyrus tuberosus</i>	Ruscinski, 1937
<i>Rhabdorrhynchus menetriesi</i> Gyllenhal, 1842	East Mediterranean region	N	E	Ph (her)	No data	Poiras, 2001
<i>Lachnaeus horridus</i> Reitter, 1890	Asia	C	E	Ph (her)	<i>Inula</i>	Poiras, 1998
<i>Lixus astrachanicus</i> Faust, 1883	Asia	C	E	Ph (her)	No data	Poiras, 1998
<i>Pissodes castaneus</i> (De Geer, 1775)	Asia, Europe, Mediterranean region	N	G	Ph (phl)	<i>Pinus</i>	Poiras, 1991
<i>Scolytus laevis</i> Chapuis, 1873	Europe	N	G, H	Ph (phl)	<i>Ulmus</i>	Poiras, 2001
<i>S. rugulosus</i> (Ratzeburg, 1837)	Asia, Europe, Mediterranean region	N	H	Ph (phl)	Rosaceae trees	Miller, Zubowsky, 1917
<i>Xyleborinus saxesenii</i> (Ratzeburg, 1837)	Asia, Europe, Mediterranean region	N	H	Ph (xmp)	<i>Laurus, Pinus, Castanea</i>	Miller, Zubowsky, 1917
<i>Pteleobius kraatzii</i> (Eichhoff, 1864)	Europe, West Mediterranean region	N	H, G, F	Ph (phl)	<i>Ulmus</i>	Miller, Zubowsky, 1917
<i>Hylastes angustatus</i> (Herbst, 1793)	South Europe, Asia	N	G	Ph (phl)	<i>Pinus</i>	Stark, 1952
<i>H. ater</i> (Paykull, 1800)	Europe, Asia	N	G, H	Ph (phl)	<i>Pinus</i>	Miller, Zubowsky, 1917

According to larvae trophic preferences 54 species of investigated beetles are phytophagous, 12 are predators and 7 detritivorous; adults of *Alphitobius diaperinus* and *Mylabris variabilis* are being detritivorous and phytophagous, respectively. For remaining 2 species information on their diet is not available.

Most of the phytophagous beetles (29.09%) are herbivorous. Spermatophagous is the second most important feeding group (27.27%), followed by leaf/steam browsers (18.18%), phloeophagous (10.9%), root browsers (7.27%) and saproxylophagous (3.63%). The remaining (by 1.81%) are root borers and xylomycetophagous. From the total number of the investigated alien beetles 26.67% are agricultural, forest and storage pests. All non-native species occurring in closed spaces (9) are storage pests. Depending on their food bonds and population dynamics, post-invasive and invasive species have various degrees of importance, among them *Leptinotarsa decemlineata*, *Sitona lineatus* and *Rhopalapion longirostre* cause the biggest losses in agricultural crops yield [Vorotyntseva, 1971; Poiras, 2006; Munteanu et al., 2012], while *Lignyodes bischoffi* is one of the most dangerous forest pest [Poiras, 1987, 2006]. Of the alien species living under outdoor conditions, the species of casual occurrence and naturalized non-invasive species do not have great significance, because they are active only in short-terms or inhabit a very small territory.

Revealed alien predators belong to Staphylinidae and Coccinellidae families. Detritivorous species belong to Tenebrionidae, Ptinidae, Dermestidae and Scarabaeidae. Their influence on environment can be considered beneficial or neutral as long as they occur in small numbers and do not affect native species. A representative case of alien species expansion in the Republic of Moldova is *Harmonia axyridis* (Pall., 1773). The first unpublished information on this species, according to Timus and Stahi [2013], dates from 1970, which could be related to the

earlier release of this species for introduction in Ukraine [Katsoyannos et al., 1997]. Published data regarding establishment of *H. axyridis* in the Republic of Moldova appeared recently, [Iazlovetchi, Sumenkova, 2013; Timus, Stahi, 2013]. Within *H. axyridis* populations five forms were identified: *H. axyridis* var. *novemdecimsignata*, *H. a.* var. *siccoma*, *H. a.* var. *succinea*, *H. a.* var. *spectabilis* and *H. a.* var. *conspicua* [Timus, Stahi, 2013]. Iazlovetchi and Sumenkova [2013] mention that the species represented 42% of all coccinellids recorded in 2011, the proportion has reached 86% in 2012, and 90% in 2013, successfully eliminating autochthon coccinellids *Coccinella septempunctata* L. and *Adalia bipunctata* L.

The effects of an alien species entry in a new ecosystem can be diverse. Most significant is a direct competition with native species for food and space, an increasing abundance in the new environment until a complete replacement of native residents occurs. Despite that a large amount of attested alien beetles does not produce economic damages, it should not be ignored because non-invasive status does not mean that it will not become invasive. It could be just a matter of time and circumstances (biotic and abiotic conditions that may change) until it becomes invasive. That's why knowledge of species peculiarities and behavior in a new environment is extremely important for determining the factors that can lead to an alien species uncontrollable expansion and invasive statute modification.

Unfortunately, the hazards caused by alien species did not cause that much concern among scientists, nor did it attract public awareness as much as would have been expected [Hulme et al., 2009]. Conducted researches comprise too large regions and cannot solve the problems of restricted areas. In the Republic of Moldova, for example, this is the first study highlighting alien species. The location of the republic at the interference of biogeographical zones causes a continuous pressure of alien species that

are at the limits of their distribution area. Thus, detailed researches in this territory will allow the monitoring and forecasting of new invasions.

Acknowledgements

This study was supported by the institutional project, 11.817.08.13F „Biological invasions and their impact on biodiversity, structure and functioning of natural and anthropogenic ecosystems in Moldova”, Institute of Zoology, Academy of Sciences of Moldova. The authors thank anonymous reviewers for stimulation and critical comments. For additional consultations regarding species *Harmonia axyridis*, we are grateful to Mrs. Asea Timus.

References

- Adashkevich B. Useful entomofauna in the fields of Moldavia. Chisinau. Stiinta. 1972. 108 p.
- Antonova V. Pests of field crops. Chisinau. 1977. 115 p.
- Arnoldy L., Zaslavsky V.A., Ter-Minassian M.E. Family Curculionidae—weevils. Identification key of European part of the USSR. Vol. 2. (Ed. 89). Moscow; Leningrad. 1965. P. 485–621.
- Baban E. Diversity of beetles (Coleoptera: Carabidae, Silphidae, Scarabaeidae, Cerambycidae) in forest ecosystems of Central Moldavian Plateau. Summary of the PhD Thesis. Chisinau. 2006. 26 p.
- Bacal S., Derunkov A. Contributions to the knowledge of rove beetles (Coleoptera, Staphylinidae) from state nature reserve „Plaiul Fagului”, Republic of Moldova // Travaux du Museum National d'Histoire Naturelle "Grigore Antipa". 2010. Vol. LIII. P. 217–221.
- Bacal S., Munteanu N., Toderas I. Checklist of beetles (Insecta: Coleoptera) of the Republic of Moldova. Brukenthal. Acta Musei. 2013. Nr. VIII. 3. P. 415–450.
- Bacal S., Stan M. New contribution to the knowledge of staphylinids (Coleoptera: Staphylinidae) from the scientific landscape „Codrii Tigheciului” (Republic of Moldova). Oltenia Museum. Craiova. Studies and Reports. Life Sciences. 2006. Vol. XXII. P. 155–159.
- Banarescu P. Principles and problems of zoogeography. Bucuresti. 1970. 260 p.
- Bouchard P., Bousquet Y., Davies A.E., Alonso-Zarazaga M.A., Lawrence J.F., Lyal C.H.C., Newton A.F., Reid C.A.M., Schmitt M., Slipinski S.A., Smith A.B.T. Family-group names in Coleoptera (Insecta). ZooKeys. 2011.
- Calestru L. Chrysomelidae beetles (Coleoptera, Chrysomelidae) in the Republic of Moldova. Their biodiversity and importance. Chisinsu. 2003. Summary of the PhD thesis. 23 p.
- Darlington F. Zoogeography: the geographical distribution of animals. Moscow: Progress Publishers, 1966. 518 p.
- Derunkov A., Bacal S. Contributions to the knowledge of staphylinid fauna (Coleoptera: Staphylinidae) from the right side of the Dniester river. Oltenia Museum. Craiova. Studies and Reports. Life Sciences. 2011. Vol. 27. Nr. 1. P. 105–108.
- Elton C.S. The ecology of invasions by plants and animals. London. Methuen & Co. Ltd. 1958. 181 p.
- Essl F., Rabitsch W. Neobiota in Österreich. Umweltbundesamt. Wien. 2002. 432 p.
- Geiter O., Homma S., Kinzelbach R. Bestandsaufnahme und Bewertung von Neozoen in Deutschland. Berlin. 2002. Umweltbundesamt 25. P. 1–173.
- Glavendekić M., Mihajlović L., Petanović R. Introduction and spread of invasive mites and insects in Serbia and Montenegro // In: Alford D.V. & Backhaus G.F. (Eds) Abstracts of the Conference on Plant Protection and Plant Health in Europe, “Introduction and spread of invasives species”. Humboldt University. Berlin. Germany. 2005. P. 229–230.
- Hill M., Baker R., Broad G., Chandler P.J., Copp G.H. et al. Audit of non-native species in England. English Nature

- Research Reports 662, English Nature, Peterborough. 2005. 82 p.
- Hulme P., Pysek P., Nentwig W., Vila M. Will threat of biological invasions unite the European Union? Science. 2009. 324 p.
- Iazlovetchi I., Sumenkova V. The invasion of multicolored Asian ladybird *Harmonia axyridis* Pallas in the Republic of Moldova has happened // The environment. 2013. Nr. 2 (68). April. P. 19–28.
- Katsoyannos P., Kontodimas D.C., Stathas G.J., Tsartsalis C.T. Establishment of *Harmonia axyridis* on citrus and some data on its phenology in Greece // Phytoparasitica. 1997. Nr. 25. P. 183–191.
- Kenis M. Insects-Insecta // In: Wittenberg R. (Ed) An inventory of alien species and their threat to biodiversity and economy in Switzerland. Delémont. 2005. CABI Bioscience Switzerland. P. 131–212.
- Kryzhanovskij O.L. Insects of the European part of the USSR. Identification key. V. 2. Moscow. 1965. 668 p.
- Maslyakov V.Yu., Izhevsky S.S. Alien Phytophagous Insects Invasions in the European Part of Russia. IGRAS Moscow. 2011. 272 p.
- Medvedev S., Shapiro D. To the knowledge of beetle fauna (Coleoptera) of the Moldavian Soviet Socialist Republic and the neighboring regions of Ukraine, Scientific works on the IBF. 1957. Nr. 30. P. 173–206.
- Mihailov I. Fauna, ecology and biocenotic significance of staphylinid species (Coleoptera, Staphylinidae) in the Republic of Moldova. Summary of PhD thesis. Chisinau. 2012. 31 p.
- Miller E., Zubowsky N. Materials on the entomological fauna of Bessarabia. Proceedings of the Bessarabian Society Naturalists and Life Science Amateurs. 1917. Vol. 2. Part 1. Chisinau. Typography Bessarabian Provincial Board. P. 32–150.
- Munteanu N. New species of bruchids (Coleoptera, Bruchidae) for fauna of the Republic of Moldova. Scientific Annals of SUM. Chisinau. 2006. Nr. 3. P. 141–143.
- Munteanu N. Diversity of Bruchidae, Nemomychidae, Anthribidae, Rhynchitidae and Attelabidae beetles and their importance in natural and anthropogenic ecosystems in Republic of Moldova Chisinau. 2009. Summary of the PhD thesis. 24 p.
- Munteanu N., Malevanciuc N., Toderas L., Moldovan A., Bacal S. Distribution and biological peculiarities of species *Sitona lineatus* (Coleoptera: Curculionidae) in Republic of Moldova // Bulletin of Academy of Sciences of Moldova, Zoology. 2012. Nr. 3 (318). P. 131–138.
- Neculiseanu Z. Fauna and biology of rove beetles subfamilies Staphylininae and Paederinae (Coleoptera, Staphylinidae) in agrocenoses of Moldova. Chisinau. 1984. Summary of the PhD thesis, 25 p.
- Neculiseanu Z., Bacal S. Tenebrionids (Coleoptera, Tenebrionidae) of Republic of Moldova. Scientific Annals of SUM. Chisinau. 2005. Vol. 3. P. 203–205.
- Neculiseanu Z., Matalin A. A catalogue of the ground-beetles of the Republic of Moldova (Insecta, Coleoptera, Carabidae). Sophia. 2000. 164 p.
- Nentwig W. Biological Invasions. Ecological Studies 193. Heidelberg: Springer. 2008. 441 p.
- Nentwig W., Josefsson M. Chapter 1. Alien terrestrial arthropods of Europe. 2010. BioRisk Vol. Nr. 4(1). P. 5–9.
- Plugar S. Some results concerning study of entomofauna in oak forests of Moldova. Harmful entomofauna of Moldova and methods of control. Chisinau. 1963. P. 7–38.
- Poiras A. A new species of weevil *Lignyodes bischoffi* Blatchley (Coleoptera, Curculionidae) in USSR fauna. Bulletin of Zoology. ASM. Series: Biological and Chemical Sciences. Chisinau. 1987. Nr. 6. P. 64.
- Poiras A. New dendrophilous weevil species (Coleoptera: Attelabidae, Curculionidae) for fauna of Moldova // Bulletin of Zoology. ASM. Series:

- Biological and Chemical Sciences. Chisinau. 1991. Nr. 3. P. 73–74.
- Poiras A. New species of weevils for the fauna of Moldova (Coleoptera: Apionidae, Curculionidae) // Bulletin of Zoology. ASM. Series: Biological and Chemical Sciences. Chisinau. 1994. Nr. 1. P. 41–45.
- Poiras A. Catalogue of the weevils (Coleoptera, Curculionoidea) and their host plants in the Republic of Moldova. Sofia-Moscow Pensoft. 1998. 156 p.
- Poiras A. Addition to the catalogue of weevils of the Republic of Moldova. Diversity and ecology of animal world in natural and anthropized ecosystems. Chisinau. 2001. P. 112–116.
- Poiras A. Beetles of Curculionoidea superfamily (Insecta, Coleoptera) in Moldova, their biodiversity and importance. Chisinau. 2006. PhD Thesis. 300 p.
- Reemer M. Invasieve Arthropoda in Nederland: Een eerste inventarisatie. Stichting European Invertebrate Survey. Leiden. 2003. 63 p.
- Roll E., Dayan T., Simberloff D. Non-indigenous species in Israel and adjacent areas // Biological Invasions. 2007. Nr. 9. P. 629–643.
- Ruscinski A. Beitrage zur Curculionidenfauna von Bessarabien // Bulletin of Regional Museums of Basarabia. Chisinau. 1937. Nr. 8. P. 33–35.
- Šefrová H. Laštůvka Z. Catalogue of alien animal species in the Czech Republic. Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis. 2005. Nr. 53. P. 151–170.
- Smith R.M., Baker R.H., Malumphy C.P., Hockland S., Hammon R.P., Ostoják Starzewski J.C., Collins D.W. Recent non-native invertebrate plant pest establishments in Great Britain: origins, pathways, and trends // Agricultural and Forest Entomology. 2007. Nr. 9. P. 307–326.
- Solodovnikova V., Talitskii V. Some results of seed feeders' weevil fauna study of genus *Apion* Hbst (Coleoptera, Apionidae) in Moldova // Entomological Review. 1972. V 51. Nr. 4. P. 786–695.
- Stark V.N. Fauna of the USSR. Coleopterous insects. V. XXXI. Bark beetles. Edition of the Academy of Sciences of the USSR. Moscow; Leningrad. 1952. 534 p.
- Timus A., Stahi N. The acclimatization and trophic spectrum of *Harmonia axiridis* in the Republic of Moldova // International conference of zoologists "Actual problems of protection and sustainable use of the animal world diversity". Chisinau. 2013. P. 180–182.
- Tomov R., Trencheva K., Trenchev G., Cota E., Ramadhi A., Ivanov B., Naceski S., Papazova-Anakieva I., Kenis M. Non-indigenous insects and their threat to biodiversity and economy in Albania, Bulgaria and Republic of Macedonia. Sofia; Moscow: Pensoft publishers, 2009. 112 p.
- Vorotyntseva A.F. Biological features of the Colorado potato beetle (*Leptinotarsa decemlineata* Say) its importance as a pest of potatoes, eggplant, tomatoes in the central zone of Moldova. Chisinau. 1971. PhD Thesis. 150 p.
- Yatsentkovsky E. Materials on the entomological fauna of Bessarabia. Beetles, T. VIII, V. 2, Nr. 2. Proceedings of the Bessarabian Society Naturalists and Life Science Amateurs. 1912. P. 1–16.