

Latvijas Entomoloģijas biedrība
Entomological Society of Latvia

ISBN 9984-9768-0-7
ISBN 9984-9768-1-5

COMPENDIUM OF LATVIAN COLEOPTERA

VOLUME I

CHECK – LIST OF LATVIAN BEETLES (INSECTA: COLEOPTERA)

Second Edition

DMITRY TELNOV



RĪGA, 2004

INTRODUCTION

Dear BEETLEman,

Seven years have passed since the first complete list of Coleoptera recorded from Latvia was published (Telnov D., A.Barsevskis, F.Savich, F.Kovalevsky, S.Berdnikov, M.Doronin, R.Cibulskis, D.Ratniece, 1997. Check-List of Latvian Beetles (Insecta: Coleoptera). - *Mitteilungen des Internationalen Entomologischen Vereins, Supplementum 5*: 1-141). As is typical in entomology, many systematic changes have been made since that time, and many beetles new to the Latvian fauna have been discovered. So, it's time for a second edition of the "Check-List of Latvian Beetles (Insecta: Coleoptera)"!

The present work begins a new ongoing series of publications on Latvian Coleoptera, which I have called the "Compendium of Latvian Coleoptera". This series aims to introduce the Latvian beetle fauna, to compile all known data and to make them accessible to coleopterists and other interested people. After this List, the next volumes planned are "Bibliography on Latvian Beetles", several parts of "Ecology of Latvian Beetles", and, finally, "Distribution Maps of Latvian Beetles".

Many Coleopterists from Latvia and all over Europe have made contributions to this issue. I am highly indebted to all my friends and colleagues listed below in the chapters "Number of Coleoptera species according to families" & "Acknowledgements" for their support!

The Coleoptera of Latvia have been studied for more than 225 years, since 1778. The current check-list is based on a revision of all known published records of Latvian Coleoptera, as well as examination of major historical and recent collections stored in Latvia and all over Europe. These collections include those of Seidlitz, Gimmerthal, Spuris, Šmits, Lackschewitz, Mikutowicz, Brammanis and numerous others. In total, more than 400 bibliographical sources have been consulted.

In total, 105 families and 3434 species of Coleoptera have been confirmed for the Latvian fauna as of May 2004. Of course, this is not the final number, and we would predict the occurrence of ~4000 species of Coleoptera in Latvia. To compare: there are ~3000 species recorded from Estonia, ~2800 from Lithuania, ~3650 from Finland and ~4450 from Sweden.

Like any faunal monograph, this check-list is not constant, and will be updated continuously as the group's systematics advances. Hence, I am open to any comments, addenda and corrections.

DMITRY TELNOV

Chairman of Coleopterology
The Entomological Society of Latvia
Rīga, 2004.28.05.

INTERNET ADDRESS OF THIS ISSUE: <http://invertebrata.from.lv> -> under "Catalogue of Latvian Coleoptera"

CONTACT ADDRESS: Dmitry Telnov, Praulienas iela 4-35, LV-1021 Rīga, Latvia; E-mail: telnov@parks.lv

ORDERING INFORMATION: The Entomological Society of Latvia, c/o: Faculty of Biology, 4, Kronvalda Blvd., LV-1586 Rīga, Latvia / Lettland / Lettonie

CHOROTYPES OF LATVIA

In Latvia, several species of Coleoptera and other invertebrates reach the southernmost, northernmost, westernmost or easternmost extremes of their distributional area. The geographical position of Latvia on the borders of the boreal and broad-leaved forest zones on one side, and the forest and forest-steppe zones on the other, have made an excellent background for high beetle species diversity.

Except the large, sometimes “transcontinental” chorotypes, like the Holarctic (incl. circumpolar & boreomontane), Palaearctic (incl. North African-Eurasian, Eurasian), W Palaearctic, Asiatic-European, Sibero-European, and European chorotypes, the territory of Latvia includes the following other smaller chorotypes: Europeo-Mediterranean, N European, Central European, E European, Europeo-Caucasian, Europeo-British, Siberian-Baltic, and finally the Baltic chorotype.

ACKNOWLEDGEMENTS

I sincerely acknowledge all the authors of the first issue of this check-list, namely A.Barševskis (Daugavpils, Latvia), S.Berdnikov (Rīga, Latvia), R.Cibuļskis (Daugavpils), M.Dorонин (Rīga), F.Kovalevsky (Rīga), D.Ratniece (Saulkrasti, Latvia) and F.Savich (Rīga), who have all contributed their data towards the second edition.

My thanks to many of my colleagues from around the World. I sincerely thank everyone who helped me during the preparation of the current list, including M.A.Alonso-Zarazaga, F.Angelini, R.Angus, V.Assing, B.Büche, M.Danilevsky, L.Erbeling, R.Frieser, F.Fritzlar, M.Geisthardt, R.Gerstmeier, M.Gigli, M.Hartmann, J.Háva, F.Hebauer, M.Jäch, A.Kirejtshuk, B.Klausnitzer, R.Klinger, A.Kopetz, S.Kurbatov, A.Legalov, G.Liberti, I.Löbl, G.Lyubarsky, J.-M.Maes, M.Mandelshtam, O.Merkl, A.Napolov, N.Nikitsky, E.Palm, I.Plonski, D.Pollock, A.Pütz, J.Reibnitz, W.Rücker, M.Russell, G.Sama, M.Schülke, S.Skalicky, M.Sörensson, X.Vázquez, M.Volkovitsh, A.Weigel, H.Wendt, C.Wurst and D.Young.

I also deeply indebted to my friends V.Spuņģis (Faculty of Biology, Rīga), J.Gailis (Latvian Fund for Nature, Rīga), M.Kalniņš (Environmental State Inspectorate, Rīga), M.V.L.Barclay (The Natural History Museum, London), and M.Hartmann (Museum of Natural History, Erfurt) for consultations, access to current literature and other friendly support during the preparation of the current work.

My special loving thanks to my wife Kristina Greķe, and my children Alisa and Edwin for their assistance during the preparation of this work, and for the supportive atmosphere of the family.

LEGENDS

Taxa above the species-level are listed in accordance with their actual taxonomy, but species, subspecies and synonyms are ordered alphabetically. Names from suborder to tribe level are in **bold**, names of genera in ***bold italic***, and those of species and subspecies are in *italic*.

Synonyms are listed. It was not my intention to include the complete synonymy of all species, and it was left to the discretion of the compilers and consultants for each family which synonyms to include. Synonyms of adephagous Coleoptera are mostly taken from the „Catalogue of Palaearctic Coleoptera”, vol. 1: Archostemata-Myxophaga-Adephaga. Stenstrup, Apollo Books, pp. 1-819 (eds. I.Löbl, A.Smetana, 2003), but with some addenda and corrections.

Names that are unavailable according the ICZN (2000) are not listed.

Some Carabidae species mentioned for Latvia for the first time in vol. 1 of the Palaearctic Beetle catalogue, without further locality data, have not been included in to this check-list.

Several species are marked with a dash „—“ in the list. These are Coleopteran species mentioned (published) for the Latvian fauna, but which do not appear to occur here, and it is doubtful whether they ever occurred in Latvia. Some causes of such “dubious records” may include:

- misidentifications;
- species recorded for „Livland“ (or Livonia) with no precise locality. The southern part of the former Livland now belongs to Latvia, but the northern one is part of Estonia. If such species have not been recorded from the territory of Latvia since that time, such species cannot be reliably assigned to the Latvian fauna;
- unsuccessful introductions (especially some species of *Carabus* L.).

NUMBER OF COLEOPTERA SPECIES ACCORDING TO FAMILIES

Nr.	Family	Number of Species in LV	Compilers & Consultants
1.	Sphaeriusidae	1	D.Telnov
2.	Trachypachidae	1	D.Telnov
3.	Carabidae	326	M.Hartmann, D.Telnov
4.	Gyrinidae	11	D.Telnov
5.	Haliplidae	14	D.Telnov
6.	Noteridae	2	D.Telnov
7.	Dytiscidae	112	D.Telnov
8.	Helophoridae	12	R.Angus, D.Telnov
9.	Hydrophilidae	51	F.Hebauer, D.Telnov
10.	Hydrochidae	5	D.Telnov
11.	Spercheidae	1	M.Jäch, D.Telnov
12.	Georissidae	1	D.Telnov
13.	Sphaeritidae	1	D.Telnov
14.	Histeridae	58	L.Erbeling, D.Telnov
15.	Hydraenidae	13	M.Jäch, D.Telnov
16.	Ptiliidae	34	M.Sörensson, D.Telnov
17.	Leiodidae	36	F.Angelini, D.Telnov
18.	Colonidae	4	D.Telnov
19.	Cholevidae	20	A.Weigel, D.Telnov
20.	Platypsyllidae	1	D.Telnov
21.	Scydmaenidae	23	D.Telnov
•	Agyrtidae	(1)	D.Telnov
22.	Silphidae	19	J.Háva, D.Telnov
23.	Staphylinidae	612	V.Assing, M.Schülke, D.Telnov
24.	Scaphidiidae	8	I.Löbl, D.Telnov
25.	Pselaphidae	27	S.Kurbatov, D.Telnov
26.	Lucanidae	6	J.-M.Maes, D.Telnov
27.	Trogidae	2	A.Napolov, D.Telnov
28.	Geotrupidae	7	A.Napolov, D.Telnov
29.	Scarabaeidae	86	A.Napolov, D.Telnov
30.	Clambidae	4	D.Telnov
31.	Eucinetidae	1	D.Telnov
32.	Scirtidae	17	B.Klausnitzer, D.Telnov
33.	Dascillidae	1	D.Telnov
34.	Buprestidae	45	M.Gigli, M.Volkovitsh, D.Telnov
35.	Byrrhidae	13	A.Pütz, D.Telnov
36.	Elmidae	11	M.Jäch, D.Telnov
37.	Dryopidae	6	M.Jäch, D.Telnov
38.	Limnichidae	2	D.Telnov
39.	Heteroceridae	9	S.Skalicky, D.Telnov
40.	Psephenidae	1	M.Jäch, D.Telnov
41.	Elateridae	79	A.Napolov, C.Wurst, D.Telnov
42.	Eucnemidae	11	D.Telnov
43.	Throscidae	5	D.Telnov
44.	Lycidae	6	D.Telnov
45.	Drilidae	1	D.Telnov
46.	Lampyridae	2	M.Geisthardt, D.Telnov
47.	Cantharidae	47	A.Kopetz, D.Telnov
48.	Dermestidae	28	J.Háva, D.Telnov
49.	Bostrichidae	3	D.Telnov
50.	Lyctidae	1	D.Telnov

51.	Anobiidae	43	B.Büche, D.Telnov
52.	Ptinidae	12	D.Telnov
53.	Lymexylidae	3	D.Telnov
54.	Trogossitidae	7	D.Telnov
55.	Cleridae	9	R.Gerstmeier, D.Telnov
56.	Dasytidae	10	G.Liberti, I.Plonski, D.Telnov
57.	Malachiidae	14	D.Telnov
58.	Kateretidae	8	A.Kirejtshuk, D.Telnov
59.	Nitidulidae	86	A.Kirejtshuk, D.Telnov
60.	Monotomidae	16	D.Telnov
61.	Silvanidae	10	D.Telnov
62.	Cucujidae	2	D.Telnov
63.	Bothrideridae	1	D.Telnov
64.	Cryptophagidae	70	G.Lyubarsky, D.Telnov
65.	Languriidae	1	G.Lyubarsky, D.Telnov
66.	Erotylidae	8	A.Weigel, D.Telnov
67.	Cerylonidae	5	D.Telnov
68.	Byturidae	2	D.Telnov
69.	Laemophloeidae	4	D.Telnov
70.	Phalacridae	17	G.Lyubarskiy, D.Telnov
71.	Sphindidae	2	D.Telnov
72.	Corylophidae	8	D.Telnov
73.	Endomychidae	5	A.Weigel, D.Telnov
74.	Coccinellidae	61	B.Klausnitzer, D.Telnov
75.	Latridiidae	48	W.Rücker, D.Telnov
76.	Zopheridae	4	D.Telnov
77.	Mycetophagidae	14	N.Nikitsky, D.Telnov
78.	Ciidae	29	J.Reibnitz, D.Telnov
79.	Tetratomidae	5	N.Nikitsky, D.Telnov
80.	Melandryidae	22	N.Nikitsky, D.Telnov
81.	Mordellidae	13	B.Büche, D.Telnov
82.	Rhipiphoridae	2	D.Telnov
83.	Stenotrachelidae	1	D.Telnov
84.	Meloidae	6	M.Geisthardt, D.Telnov
85.	Oedemeridae	14	X.Vázquez, D.Telnov
86.	Pythidae	1	D.Pollock, D.Telnov
87.	Pyrochroidae	3	D.Young, D.Telnov
88.	Scaptiidae	1	B.Büche, D.Telnov
89.	Salpingidae	9	D.Pollock, D.Telnov
90.	Mycteridae	1	D.Pollock, D.Telnov
91.	Boridae	1	D.Pollock, D.Telnov
92.	Anthicidae	13	D.Telnov
93.	Aderidae	5	R.Klinger, D.Telnov
94.	Tenebrionidae	50	O.Merkl, D.Telnov
95.	Cerambycidae	115	M.Danilevsky, G.Sama, D.Telnov
96.	Bruchidae	9	H.Wendt, D.Telnov
97.	Chrysomelidae	329	F.Fritzlar, D.Telnov
98.	Nemonychidae	2	A.Legalov, D.Telnov
99.	Anthribidae	9	R.Frieser, D.Telnov
100.	Rhynchitidae	12	A.Legalov, D.Telnov
101.	Attelabidae	3	A.Legalov, D.Telnov
102.	Apionidae	68	M.A.Alonso-Zarazaga, M.Russell, D.Telnov
103.	Curculionidae	373	E.Palm, D.Telnov
104.	Scolytidae	68	M.Mandelshtam, D.Telnov
105.	Platypodidae	1	M.Mandelshtam, D.Telnov

Totaly: 105 families, 3434 species

SYSTEMATIC LIST OF COLEOPTERA RECORDED FOR THE FAUNA OF LATVIA

State: May, 2004.

Coleoptera

Myxophaga

Sphaeriiformia

Sphaerioidea

Sphaeriusidae Erichson, 1845

Sphaerius Waltl, 1838

acarooides Waltl, 1838

obsidianus (Kolenati, 1846)

Adephaga

Carabiformia

Caraboidea

Trachypachidae Thomson, 1857

Trachypachus Motschulsky, 1844

zetterstedti (Gyllenhal, 1827)

laticollis Motschulsky, 1864

transversalis Motschulsky, 1844

Note: Rare and insufficiently known species, known from single locality (E Latvia) but should be more widely distributed in old spruce forests of eastern and North-eastern Latvia.

Carabidae Latreille, 1802

Nebriinae Laporte de Castelnau, 1834

Nebriini Laporte de Castelnau, 1834

Leistus (*Leistus* Frölich, 1799)

ferrugineus (Linnaeus, 1758)

spinilabris (Panzer, 1797)

testaceus Frölich, 1799

piceus *piceus* Frölich, 1799

bosnicola P.W.J.Müller, 1937

fuscoaeneus (Panzer, 1803)

froehlichii (Duftschmid, 1812)

kaszabi Horvatovich, 1972

pseudoalpicola Maran, 1941

roubali Maran, 1941

terminatus (Panzer, 1793)

bructeri (Panzer, 1796)

nigripennis Wojas, 1993

praeustus (Fabricius, 1792)

rufescens (Fabricius, 1775)

Leistus (*Pogonophorus* Latreille, 1802)

rufomarginatus (Duftschmid, 1812)

italicus A.Fiori, 1913

ottomanus Schweiger, 1970

Note: This species is known from extreme SW part of Latvia (Pape area, Baltic Sea coast) situated at the border with Lithuania. North

border of main distribution area.

Nebria (*Boreonebria* Jeannel, 1937)

— nivalis *nivalis* (Paykull, 1798)

bifaria Mannerheim, 1853

changaica Horvatovich, 1973

femoralis Motschulsky, 1859

femorata Motschulsky, 1866

mollis Motschulsky, 1866

rufescens *rufescens* (Ström, 1768)

aethiops (Stephens, 1828)

arctica Dejean, 1826

attenuata Motschulsky, 1844

balbii Bonelli, 1810

baltica Motschulsky, 1866

besseri Fischer von Waldheim, 1828

curtulata Casey, 1924

dubia R.F.Sahlberg, 1844

geniculata Gredler, 1857

gyllenhali (Schönherr, 1806)

hyperborea Gyllenhal, 1827

impressa Newman, 1833

jamata Motschulsky, 1866

jockischii (Duftschmid, 1812)

maklini Munster, 1933

marshallana (Stephens, 1827)

subacuminata Chaudoir, 1843

Nebria (*Nebria* Latreille, 1802)

brevicollis (Fabricius, 1792)

barbara Chaudoir, 1843

cursor (O.F.Müller, 1776)

fuscata Bonelli, 1810

infida (Rossi, 1792)

lata Newman, 1833

obscurata J.R.Sahlberg, 1903

rufipes (Goeze, 1777)

rufomarginata Fischer von Waldheim, 1828

rugimarginata Marsham, 1802

sicula Preudhomme de Borre, Ragusa, 1882

uzokensis Obenberger, 1917

varicornis Newman, 1833

Nebria (*Paranebria* Jeannel, 1937)

livida *livida* (Linnaeus, 1758)

collaris (Thunberg, 1787)

lateralis (Fabricius, 1787)

sabulosa (Fabricius, 1787)

sibirica Csiki, 1902

Notiophilini Motschulsky, 1850

Notiophilus Duméril, 1806

aestuans Dejean, 1826

bigeminus Thomson, 1884

longipennis Putzeys, 1866

pusillus Waterhouse, 1833
aquaticus (Linnaeus, 1758)
blacki Edwards, 1913
dauricus Motschulsky, 1860
dybowskii J.L.Lomnicki, 1903
kroli J.L.Lomnicki, 1903
metallicus Waterhouse, 1833
parvulus Waterhouse, 1833
pristinus Friederichs, 1903
pusillus (Schreber, 1759)
semenovi Tschitschérine, 1903
semipunctatus (Fabricius, 1775)
spaethi Munster, 1922
strigifrons Baudi di Selve, 1864
biguttatus (Fabricius, 1779)
coerulescens Depoli, 1929
lateralis Motschulsky, 1864
latus Waterhouse, 1833
melanophthalmus Šlosser-Klekovski, 1877
nitidus Waterhouse, 1833
striatus Waterhouse, 1833
germinyi Fauvel, 1863
hypocrita Putzeys, 1866
stipraisi Barševskis, 1993
palustris (Duftschmid, 1812)
atavus Friederichs, 1903
brevis Waterhouse, 1833
davisii Waterhouse, 1833
decemstriatus Roubal, 1948
latior Waterhouse, 1833
newmanni Waterhouse, 1833
nitidulus Waterhouse, 1833
parallelus Waterhouse, 1833
tibialis Waterhouse, 1833
— ***substriatus*** Waterhouse, 1833
puncticollis Küster, 1848
subopacus Chaudoir, 1852
Loricinae Bonelli, 1810
Loricini Bonelli, 1810
Loricera (***Loricera*** Latreille, 1802)
pilicornis (*pilicornis* Fabricius, 1775)
aenea Latreille, 1806
alpina Heer, 1838
apennina Binaghi, 1942
rufilabris Motschulsky, 1845
semipunctata Eschscholtz, 1833
seticornis (O.F.Müller, 1776)
vinearum (Geoffroy, 1785)
Pelophilini Kavanaugh, 1996
Pelophila Dejean, 1826
borealis (Paykull, 1790)
angusticollis Motschulsky, 1860
arctica Dejean, 1826
californica Motschulsky, 1844
costata Ménétriés, 1851
dejeanii Dejean, 1826
elongata Mannerheim, 1823
eschscholtzii Mannerheim, 1823

gebleri Mannerheim, 1823
laevigata Motschulsky, 1844
marginata Mannerheim, 1823
ochotica R.F.Sahlberg, 1844
shermani Casey, 1913
ulkei G.H.Horn, 1870

Note: Scarce species with only a few known localities. South border of main distribution area.

Cicindelinae Latreille, 1802

Cicindelini Latreille, 1802

Cicindela Linnaeus, 1758

campestris (*campestris* Linnaeus, 1758)

affinis Fischer von Waldheim, 1820
austriaca Schrank, 1776
beuthini Schulz, 1908
bilineata Della Beffa, 1909
calabrica Mandl, 1944
coerulescens Schilsky, 1888
confluens Bremi-Wolf, 1856
connata Heer, 1838
desertorum Faldermann, 1836
destituta Srnka, 1890
dumetorum Motschulsky, 1840
exapicalis Codina, 1916
exsuturalis Beuthin, 1899
farellensis Graëlls, 1847
funebris Sturm, 1827
gebieni Barthe, 1908
humerosa Srnka, 1890
impunctata Westhoff, 1881
liturata Roeschke, 1891
marginepunctata Gistel, 1856
nigrescens Heer, 1838
palustris Beuthin, 1896
pontica Schaum, 1857
pseudopalustris Schulz, 1908
rubens I.Frivaldszky von Frivald, 1835
rufipennis Beuthin, 1885
saxeseni Endrulat, Tessien, 1854
semiapicalis Beuthin, 1896
sibirica Fischer von Waldheim, 1821
subapicalis Beuthin, 1899
tirolensis Schulz, 1908
vernalis Gistel, 1857
worleei Barthe, 1908

hybrida (*hybrida* Linnaeus, 1758)

apraca Stephens, 1827
beuthini Fuente, 1912
campestris Sulzer, 1761
conjugata Fuente, 1912
fasciata (Geoffroy, 1785)
inhumeralis Beuthin, 1885
integra Sturm, 1827
maculata DeGeer, 1774
melanostoma Schenkling, 1889
monasteriensis Westhoff, 1881
riparia Stephens, 1827
striatoscutellata Beuthin, 1885

- transdanubialis* Csiki, 1946
maritima maritima Dejean, 1822
baltica Motschulsky, 1844
finmarkica Mandl, 1936
intermedia Lengerken, 1909
obscura Schilsky, 1888
pseudomaritima Lengerken, 1912
tshemalensis Gebert, 1995

- sylvatica sylvatica* Linnaeus, 1758
caja Samko, 1938
immaculata Wanach, 1907
obotritica Schulz, 1909
pseudotypica Lutshnik, 1924
similis Westhoff, 1881
subinterrupta Beuthin, 1899

Cylindera (Cylindera Westwood, 1831)

- germanica germanica* (Linnaeus, 1758)
angustata (Motschulsky, 1844)
anthracina (Klug, 1834)
apricaria (Gistel, 1838)
berthouneai (Beuthin, 1904)
bipunctata (Kraatz, 1890)
bleusei (Beuthin, 1904)
boja (Schank, 1798)
coerulea (Herbst, 1806)
cyanea (Cristoforis, 1832)
inornata (Schilsky, 1888)
kunzei (Gistel, 1838)
laeta (Motschulsky, 1844)
oberthueri (Beuthin, 1904)
obscura (Fabricius, 1798)
obscurior (Schreber, 1759)
saphyrina (Barthe, 1922)
semiapicalis (Beuthin, 1905)
stevenii (Dejean, 1825)
subtruncata (Chaudoir, 1842)

Cylindera (Eugrapha Rivalier, 1950)

- arenaria viennensis* (Schrank, 1781)
adunca (Gravenhorst, 1807)
leucophthalma (Fischer von Waldheim, 1823)
padana (Schaum, 1856)
sinuata (Panzer, 1792)

Note: This species is known by a few specimens from SE part of Latvia (Daugavpils area). North border of main distribution area.

Carabinae Latreille, 1802

Carabini Latreille, 1802

Calosoma (Calosoma Weber, 1801)

- inquisitor inquisitor* (Linnaeus, 1758)
antiquum (Geoffroy, 1785)
batnense Lallemand, 1868
comanense Lapouge, 1930
funereum Ragusa, 1883
punctiventre Reiche, Saulcy, 1855
viridulum Kraatz, 1877

Note: Very rare species with only a few recent localities. Officially protected species in

Latvia.

- *sycophanta* (Linnaeus, 1758)
anthracinum (Lapouge, 1924)
habelmanni Schilsky, 1888
lapougei Breuning, 1927
nigroaeneum Polentz, 1937
purpuripenne Reitter, 1891
rapax Motschulsky, 1866
severum Chaudoir, 1850
smaragdynum Rossi, 1882

Calosoma (Campalita Motschulsky, 1866)

- europunctatum europunctatum* (Herbst, 1784)
funestum Géhin, 1885
herbsti (Gmelin, 1790)
nicolasi (Schuler, 1965)
sericeum (Fabricius, 1792)
tauricum (Motschulsky, 1850)

Note: Possible instinct species with no confirmed records in the last 150 years. North border of main distribution area.

Carabus (Archicarabus Seidlitz, 1887)

- nemoralis nemoralis* O.F.Müller, 1764
borealis Raynaud, 1973
canadensis Lapouge, 1908
cantalicus Jeanne, 1971
colasi Bourgin, 1963
fayardensis P.Machard, 1974
foetens (Voet, 1778)
krasae Roubal, 1903
miolansicus (Tarrier, 1965)
verdonensis Puisségur, 1961
waltersachi Mandl, 1984

Carabus (Carabus Linnaeus, 1758)

- arcensis arcensis* Herbst, 1784
aeratus Géhin, 1885
aereus Dejean, 1826
austriae Sokolár, 1907
borussicus Csiki, 1927
nigrescens Westhoff, 1881
nigrinopomeranus Rade, 1895
nigrinus Westhoff, 1881
noricus Sokolár, 1907
pomeranus Gmelin, 1790
seileri Heer, 1837
ziegleri Géhin, 1876

granulatus granulatus Linnaeus, 1758

- campestris* M.F.Adams, 1817
chalybeus (Voet, 1778)
confluens Fischer von Waldheim, 1828
forticostis Kraatz, 1878
fulvipes Géhin, 1885
haematomerus Kraatz, 1878
hibernicus Lindroth, 1956
hudsonicus Lapouge, 1925
parallelus Faldermann, 1836
parvicollis Kraatz, 1878
rubripes Géhin, 1876
wimmeli Schulz, 1900

menetriesi menetriesi Hummel, 1827

Note: Officially protected species in Latvia.

Carabus (Chaetocarabus) Thomson, 1875

intricatus intricatus Linnaeus, 1761

angustulus Haury, 1878

bohemicus Haury, 1881

carnicus Depoli, 1926

chionobatus Born, 1918

cyaneus Fabricius, 1776

eurynotus Lapouge, 1913

germanicus Semenov, 1898

giganteus Heer, 1837

gigas Heer, 1838

kaadeni Schaufuss, 1879

liburnicus Haury, 1881

macedoniens Jureček, 1928

marchali Raynaud, 1971

minor Haury, 1881

montenegrinus Kraatz, 1876

roeschkeanus Semenov, 1906

subrhodopensis Blumenthal, 1976

ulcerosus Lapouge, 1902

Note: Possible extinct species with no confirmed records in the last 150 years.

Carabus (Chrysocarabus) Thomson, 1875

— *smaragdinus smaragdinus* Fisch. von Waldheim, 1823

coreicus (G.Hauser, 1921)

— *splendens* Olivier, 1790

breiti Le Moult, 1913

carolinus Fabricius, 1792

cychricollis Lapouge, 1898

hemipterus Sirguey, 1931

kricheldorfii Le Moult, 1913

lapurdanus Lapouge, 1913

lineatopunctatus Lapouge, 1898

mascarauxi Lapouge, 1918

nereensis A.Nicolas, 1898

palesius Sirguey, 1931

pallens Lapouge, 1925

pinetorum Lapouge, 1910

pseudofaustulus Le Moult, 1912

viridis Dejean, 1826

Carabus (Eucarabus) Géhin, 1876

— *ulrichii ulrichii* Germar, 1824

comanensis Born, 1902

leuckarti Petri, 1885

pavlicheki Born, 1907

planitiae Csiki, 1929

podolicus Semenov, 1898

tigancanus Born, 1908

volhynicus Lutshnik, 1927

Carabus (Hemicarabus) Géhin, 1876

nitens Linnaeus, 1758

aquitaniensis Mollard, 1996

aureus DeGeer, 1774

cupreus Westhoff, 1881

fennicus Géhin, 1885

hookeri Nodier, 1821

interruptus Westhoff, 1881

marginatus (Voet, 1778)

niger Semenov, 1887

pulchellus Brüggemann, 1873

subnitens Reitter, 1896

Carabus (Limnocarabus) Géhin, 1876

clathratus clathratus Linnaeus, 1761

adspersus Gmelin, 1790

aeneopunctatus DeGeer, 1774

borealis Born, 1908

boreus Csiki, 1927

cupreus Schilsky, 1888

eversmanni Motschulsky, 1844

foveolatoseriatatus Reitter, 1896

jansoni Kraatz, 1890

longus (Voet, 1778)

multipunctatus Kraatz, 1890

Carabus (Megodontus) Solier, 1848

violaceus andrzejuscii Fischer von Waldheim, 1823

carbonatus Schaufuss, 1882

Carabus (Mesocarabus) Thomson, 1875

— *problematicus problematicus* Herbst, 1786

angustior Born, 1895

austriacus Sturm, 1815

clairvillei Lapouge, 1913

cyanescens Sturm, 1815

Carabus (Oreocarabus) Géhin, 1876

glabratus glabratus Paykull, 1790

angustatus Petri, 1912

extensus Kraatz, 1885

laevigatus L.G.Scriba, 1790

lapponicus Born, 1909

liebleri Bierig, 1918

punctatostatus Haury, 1885

sericeus Csiki, 1946

hortensis hortensis Linnaeus, 1758

alternans Kraatz, 1887

dofleini Csiki, 1927

duerckianus Ganglbauer, 1890

magdeleinei Lapouge, 1922

ostariensis Born, 1912

pillichi Bernau, 1914

rhodopensis Apfelbeck, 1904

starygradensis Born, 1912

striatus DeGeer, 1774

Carabus (Procrustes) Bonelli, 1810

coriaceus coriaceus Linnaeus, 1758

angusticollis Motschulsky, 1859

basilicatus (Born, 1902)

cansiglianus Bernau, 1914

capellae Depoli, 1929

cordicollis Motschulsky, 1866

gracilis Depoli, 1929

imminutus (Kraatz, 1879)

italianus Csiki, 1927

- italicus* Lapouge, 1925
occidentalis Born, 1907
pseudorugifer Sokolár, 1906
punctulatus (Voet, 1778)
spretus Dejean, 1826
sublineatus Géhin, 1885
tauricus (Ganglbauer, 1888)
- Carabus (Tachypus)** Weber, 1801
— *auratus auratus* Linnaeus, 1761
auratoides Reitter, 1896
brullei Géhin, 1885
catalaunicus Géhin, 1885
concyri P.Machard, 1973
confluentinus Bocklet, 1904
dufouranis Csiki, 1927
isignyensis P.Machard, 1977
labitsei Clément, 1904
laniarius Gistel, 1857
laticollaris Bocklet, 1904
navarricus Lapouge, 1925
obscuricornis Beuthin, 1885
perauratus Reitter, 1896
quadricostatus Bocklet, 1904
rotundatus Born, 1895
sulcatisimus Lapouge, 1898
sulcatus DeGeer, 1774
- cancellatus cancellatus* Illiger, 1798
dolosus Csiki, 1927
ingulensis Eidam, 1941
kniephofi Langenhan, 1913
kuennemanni Langenhan, 1913
kuntzeni Eidam, 1930
obscuriusculus Eidam, 1930
sajanensis Reitter, 1896
sarmaticus Bernau, 1913
setteli Langenhan, 1913
tuberculatus Dejean, 1826
wankae Sokolár, 1911
- Carabus (Tomocarabus)** Reitter, 1896
convexus convexus Fabricius, 1775
aemulis Fischer von Waldheim, 1823
antipai Panin, 1943
apenninus Depoli, 1927
borealis Géhin, 1885
bucciarellii Mandl, 1966
gibbus Brullé, 1835
hensleri Bodemeyer, 1922
kiskunensis Ádám, Merkl, 1986
longobardus Depoli, 1927
merklii Hopffgarten, 1878
oblongulus Motschulsky, 1866
paganettii (Born, 1905)
pseudogracilior Panin, 1943
pyrenaicola Csiki, 1927
pyrenaicus Born, 1907
simplicipennis Dejean, 1826
striolatus Fischer von Waldheim, 1823
turcomanicus Motschulsky, 1850
- Carabus (Trachycarabus)** Géhin, 1876
- *perrini planus* Géhin, 1885
campestris Fischer von Waldheim, 1820
parallelus Fischer von Waldheim, 1844
planatus Motschulsky, 1850
transcaucasicus Csiki, 1927
- Cychrini** Laporte de Castelnau, 1834
Cyhrus Fabricius, 1794
caraboides (Linnaeus, 1758)
alutaceus Sturm, 1826
angustatus Heer, 1837
coadunatus (DeGeer, 1774)
convexus Heer, 1837
costai Emery, 1872
elongatus Audinet-Serville, 1821
depressus Eidam, 1925
hoppei Ganglbauer, 1891
intermedius Westhoff, 1881
lapponicus Géhin, 1885
laticollis Jeanne, 1972
pehri Sokolár, 1907
prymnaeus Fischer von Waldheim, 1828
pygmaeus Chaudoir, 1837
pyrenaeus Kraatz, 1879
rostratus (Linnaeus, 1761)
sabaudus Fauvel, 1882
torulosus Fischer von Waldheim, 1823
zariqueyi Born, 1924
- Elaphrinae** Erichson, 1837
Elaphrini Erichson, 1837
Blethisa Bonelli, 1810
multipunctata multipunctata (Linnaeus, 1758)
curtula Motschulsky, 1844
- Elaphrus (Arctelaphrus)** Semenov, 1926
— *lapponicus lapponicus* Gyllenhal, 1810
elongatus Fischer von Waldheim, 1828
obscurior Kirby, 1837
violaceus Lutshnik, 1936
viridis Sparre-Schneider, 1888
- Elaphrus (Elaphroterus)** Semenov, 1895
angusticollis longicollis J.R.Sahlberg, 1880
jakowlevi Semenov, 1895
- Note: This species is known from a few specimens from the extreme SE part of Latvia (Piedruja area, river Daugava valley) situated at the border with Belarus'. North border of main distribution area.
- aureus aureus* P.W.J.Müller, 1821
littoralis Dejean, 1826
smaragdinus Reitter, 1887
- Elaphrus (Elaphrus)** Fabricius, 1775
riparius (Linnaeus, 1758)
baschkiricus Motschulsky, 1844
dilaticollis R.F.Sahlberg, 1844
latiusculus Motschulsky, 1850
paludosus Olivier, 1790
violaceomaculatus Motschulsky, 1845

Elaphrus (Neaelaphrus Hatch, 1951)	<i>seminiger</i> Reitter, 1913
<i>cupreus</i> Duftschmid, 1812	<i>strepens</i> Fischer von Waldheim, 1828
<i>arcticus</i> Dejean, 1826	<i>tibialis</i> Motschulsky, 1850
<i>uliginosus</i> Fabricius, 1792	
<i>impressifrons</i> Chaudoir, 1842	Scaritinae Bonelli, 1810
<i>laevisculptus</i> Bänninger, 1919	Clivinini Rafinesque, 1815
<i>purkynei</i> Obenberger, 1917	Clivina Latreille, 1802
Omophroninae Bonelli, 1810	<i>collaris</i> (Herbst, 1784)
Omophronini Bonelli, 1810	<i>contracta</i> (Geoffroy, 1785)
Omophron Latreille, 1802	<i>discicollis</i> Dejean, 1825
<i>limbatum</i> (Fabricius, 1776)	<i>discipennis</i> Dejean, 1825
<i>baenningeri</i> Krause, 1915	<i>elongata</i> Randall, 1838
<i>coccinelloides</i> (L.Petagna, 1819)	<i>gibbicollis</i> Dejean, 1825
<i>corcyreum</i> J.R.Sahlberg, 1903	<i>obliterata</i> Sumakov, 1909
<i>dubium</i> (Herbst, 1779)	<i>randalli</i> LeConte, 1857
<i>kanalense</i> Fauvel, 1882	<i>sanguinea</i> Dejean, 1825
<i>maculatipenne</i> Pic, 1901	<i>transcaucasica</i> Putzeys, 1866
<i>sokolari</i> Roubal, 1910	
<i>solskyi</i> Zaitzev, 1916	<i>fossor</i> <i>fossor</i> (Linnaeus, 1758)
Brachininae Bonelli, 1810	<i>arenaria</i> (Fabricius, 1792)
Brachinini Bonelli, 1810	<i>infuscata</i> Chaudoir, 1846
Brachinus (<i>Brachinus</i> Weber, 1801)	<i>meridionalis</i> J.Jacquet, 1938
<i>crepitans</i> (Linnaeus, 1758)	<i>ovipennis</i> Chaudoir, 1846
<i>altaicus</i> Motschulsky, 1864	<i>prolixa</i> Kiesenwetter, 1857
<i>atripennis</i> Motschulsky, 1864	
<i>fallax</i> Apfelbeck, 1904	Dyschiriini W.Kolbe, 1880
<i>fimbriolatus</i> Lucas, 1846	Dyschirius (<i>Dyschiriodes</i> Jeannel, 1941)
<i>flavosuturalis</i> Eichler, 1924	<i>aeneus</i> <i>aeneus</i> (Dejean, 1825)
<i>gracilis</i> Motschulsky, 1844	<i>aeratus</i> Stephens, 1828
<i>jeanneli</i> Razet, 1951	<i>aereus</i> (Ahrens, 1830)
<i>joenius</i> Patti, 1844	<i>ahenus</i> (Ahrens, 1830)
<i>obscuricornis</i> Brullé, 1834	<i>paludosa</i> (Ahrens, 1830)
<i>obscurus</i> Heer, 1837	<i>remotepunctatus</i> Putzeys, 1866
<i>scoteinus</i> Kolenati, 1845	<i>ruthenus</i> Motschulsky, 1849
<i>siculus</i> Patti, 1844	<i>stethogramme</i> Kolenati, 1845
<i>strepitans</i> Duftschmid, 1812	<i>tanaicensis</i> Motschulsky, 1844
<i>sulcatulus</i> Motschulsky, 1850	
Note: Scarce species with only a few confirmed records in the last 50 years. All known records are from southern part of Latvia.	<i>impunctipennis</i> Dawson, 1854
	<i>inermis</i> Dawson, 1854
	<i>laevistriatus</i> Fairmaire, Laboulbéné, 1854
	<i>lapponicus</i> Thomson, 1857
Brachinus (<i>Brachynidius</i> Reitter, 1919)	<i>intermedius</i> Putzeys, 1846
— <i>explodens</i> Duftschmid, 1812	<i>rufilabris</i> Motschulsky, 1864
<i>armoricanus</i> Razet, 1951	<i>sylvaticus</i> Thomson, 1857
<i>bombarda</i> Duftschmid, 1812	<i>similis</i> Petri, 1891
<i>brunnicornis</i> Motschulsky, 1864	
<i>chalybaeus</i> Motschulsky, 1864	<i>laeviusculus</i> Putzeys, 1846
<i>elongatus</i> Tournier, 1864	<i>nodifrons</i> Penecke, 1914
<i>fulviventris</i> Motschulsky, 1864	
<i>glabratus</i> Dejean, 1824	<i>neresheimeri</i> Wagner, 1915
<i>marginiventris</i> Motschulsky, 1864	
<i>monticola</i> Motschulsky, 1864	<i>nigricornis</i> Motschulsky, 1844
<i>nitidulus</i> Mulsant, Wachanru, 1852	<i>septentrionum</i> Munster, 1923
<i>obscuricornis</i> Ménétriés, 1832	
<i>palicari</i> Laporte de Castelnau, 1834	Note: This species is known only from single locality in Dunava area (E Latvia).
<i>prokschi</i> Mader, 1917	
<i>salicola</i> Motschulsky, 1864	<i>nitidus</i> <i>nitidus</i> (Dejean, 1825)
	<i>inermis</i> Curtis, 1831
	<i>thoracicus</i> (Duftschmid, 1812)
	<i>politus</i> <i>politus</i> (Dejean, 1825)
	<i>aureolus</i> Notman, 1920
	<i>elongatus</i> (Ahrens, 1830)
	<i>irkutensis</i> A.Fleischer, 1899
	<i>jenisseiensis</i> P.W.J.Müller, 1924
	<i>metallicus</i> (Ahrens, 1830)

- stronczynskii* Weissberg, 1924
- salinus*** *salinus* Schaum, 1843
klynstrai P.W.J.Müller, 1934
punctipennis Putzeys, 1846
- tristis*** Stephens, 1827
glypturus Bates, 1883
luedersi Wagner, 1915
unicolor Motschulsky, 1844
- Dyschirius*** (*Dyschirius* Bonelli, 1810)
angustatus (Ahrens, 1830)
jejunus Dawson, 1854
sabulicola (Lacordaire, 1835)
uliginosus Putzeys, 1846
- arenosus*** Stephens, 1827
niger (Ahrens, 1830)
riparius Mannerheim, 1844
septentrionalis Motschulsky, 1852
thoracicus (Rossi, 1790)
- digitatus*** (Dejean, 1825)
obscurus (Gyllenhal, 1827)
dilaticollis A.Fleischer, 1899
- Dyschirius*** (*Eudyschirius* Fedorenko, 1996)
globosus (Herbst, 1784)
gibbus (Fabricius, 1792)
glomerosus Bousquet, 1997
heydeni A.Fleischer, 1899
laevicollis (Ahrens, 1830)
maritimus (Boheman, 1849)
minimus (Ahrens, 1830)
muellieri Wagner, 1929
ragusae P.W.J.Müller, 1922
remotus (Marsham, 1802)
rotundicollis (Faldermann, 1835)
ruficollis Kolenati, 1845
tournieri Putzeys, 1866
— *rufipes* (Dejean, 1825)
- Broscinae*** Hope, 1838
Broscini Hope, 1838
Broscus Panzer, 1813
cephalotes (Linnaeus, 1758)
vulgaris Fischer von Waldheim, 1824
- Miscodera*** Eschscholtz, 1830
arctica (Paykull, 1798)
erythropus Motschulsky, 1844
hardyi Chaudoir, 1861
hypisia Schauberger, 1934
readii (Curtis, 1831)
- Trechinae*** Bonelli, 1810
Bembidiini Stephens, 1827
Asaphidion Des Gozis, 1886
austriacum Schweiger, 1975
- Note: This species is known by single record from Naujene area (SE Latvia). Probably, should be more widely distributed in whole of Latvia.
- curtum* *curtum* (Heyden, 1870)
Note: This species is only known by single record from Moricsala State Nature reserve (W Latvia). North border of main distribution area.
flavipes (Linnaeus, 1761)
impressum (Geoffroy, 1785)
quadripunctatum (Goeze, 1777)
- pallipes*** (Duftschmid, 1812)
Bembidion (*Actedium* Motschulsky, 1864)
pallidipenne (Illiger, 1802)
frisium Paul Meyer, 1938
- Bembidion*** (*Asioperyphus* Vysoky, 1986)
lunatum (Duftschmid, 1812)
koltzei Paul Meyer, 1919
ustum (Stephens, 1828)
- Bembidion*** (*Bembidion* Latreille, 1802)
humerale Sturm, 1825
bellum C.R.Sahlberg, 1827
pulchrum Gyllenhal, 1827
- quadrimaculatum*** *quadrimaculatum* (Linnaeus, 1761)
arragonense Wagner, 1926
coarctatum C.R.Sahlberg, 1827
formosum C.R.Sahlberg, 1827
pulchellum (Panzer, 1796)
quadriguttatum (Fabricius, 1775)
quadriguttatum Audinet-Serville, 1821
sibiricum Motschulsky, 1850
subglobosum (Rossi, 1792)
tenax Casey, 1918
- *quadripustulatum* *quadripustulatum*
Audinet-Serville, 1821
albomaculatum J.R.Sahlberg, 1900
- Bembidion*** (*Bembidionetolitzky* E.Strand, 1829)
— *tibiale* (Duftschmid, 1812)
cнемериум (Stephens, 1839)
вірідіаенум (Stephens, 1828)
- Bembidion*** (*Bracteon* Bedel, 1879)
argenteolum Ahrens, 1812
аметистинум P.Meier, 1899
азуреум (Gebler, 1833)
глабрускум Motschulsky, 1844
віренс Schilsky, 1908
- litorale* (Olivier, 1790)
каeruleum Krynicki, 1832
елеганс Germar, 1824
нігранс Barthe, 1912
палудосум (Panzer, 1794)
- velox* (Linnaeus, 1761)
гюнтері Seidlitz, 1887
импресум (Panzer, 1797)
семіцианум P.Meier, 1899
стріатум (Paykull, 1798)
- Bembidion*** (*Chlorodium* Motschulsky, 1864)
pygmaeum (Fabricius, 1792)

<i>bilunulatum</i> Bielz, 1852	<i>orichalcicum</i> (Stephens, 1828)
<i>chalcopterum</i> Dejean, 1831	<i>plumbbeum</i> (Motschulsky, 1844)
<i>fornicatum</i> Beck, 1817	<i>quatuordecimstriatum</i> Thomson, 1871
<i>orichalceum</i> (Panzer, 1796)	<i>velox</i> Erichson, 1837
<i>orichalcicum</i> (Illiger, 1798)	<i>Bembidion</i> (<i>Neja</i>) Motschulsky, 1864)
<i>serotinum</i> Hochhuth, 1871	<i>nigricorne</i> Gyllenhal, 1827
<i>Bembidion</i> (<i>Diplocampa</i>) Bedel, 1896)	<i>wahnschaffei</i> Heinemann, 1911
<i>assimile</i> Gyllenhal, 1810	<i>Bembidion</i> (<i>Nepha</i>) Motschulsky, 1864)
<i>castanopterum</i> (Stephens, 1828)	<i>illigeri</i> Netolitzky, 1914
<i>doris</i> (Stephens, 1828)	<i>kuesteri</i> Jakobson, 1906
<i>dubium</i> Heer, 1838	<i>Bembidion</i> (<i>Notaphus</i>) Dejean, 1821)
<i>fasciatum</i> Petri, 1912	<i>obliquum</i> Sturm, 1825
<i>guttula</i> (Duftschmid, 1812)	<i>fasciatum</i> (Motschulsky, 1844)
<i>haemorrhoidale</i> (Stephens, 1835)	<i>immaculatum</i> J.R.Sahlberg, 1875
<i>productum</i> Schilsky, 1889	<i>semipunctatum</i> (Donovan, 1806)
<i>spencii</i> (Stephens, 1828)	<i>accuratum</i> Casey, 1924
<i>viridanum</i> (Motschulsky, 1850)	<i>adustum</i> Schaum, 1860
<i>fumigatum</i> (Duftschmid, 1812)	<i>alternans</i> (Motschulsky, 1845)
<i>dejeani</i> Putzeys, 1846	<i>atrocyanum</i> Netolitzky, 1943
<i>stictum</i> (Stephens, 1828)	<i>elegantulum</i> R.F.Sahlberg, 1844
Note: Insufficiently known species, known only from the area of lake Lubānas (E Latvia).	<i>provoanum</i> Casey, 1918
<i>transparens</i> transparens (Gebler, 1829)	<i>rupestre</i> Dawson, 1854
<i>contaminatum</i> J.R.Sahlberg, 1875	<i>sturmii</i> (Duftschmid, 1812)
<i>edmontonense</i> Casey, 1924	<i>varium</i> (Olivier, 1795)
<i>fortestriatum</i> Mannerheim, 1852	<i>amoenulum</i> Péringuay, 1896
<i>sulcatum</i> (LeConte, 1848)	<i>bifasciatum</i> (Stephens, 1828)
<i>trepidum</i> (LeConte, 1848)	<i>dyrrhachii</i> De Monte, 1947
<i>Bembidion</i> (<i>Emphanes</i>) Motschulsky, 1850)	<i>flammulatum</i> (Duftschmid, 1812)
<i>azurescens</i> <i>azurescens</i> Dalla Torre, 1877	<i>heptapotamicum</i> Tschitschérine, 1895
<i>triste</i> Schilsky, 1888	<i>infuscatum</i> Schilling, 1847
<i>tenellum</i> tenellum Erichson, 1837	<i>majus</i> Gyllenhal, 1827
<i>atratum</i> Hornung, 1844	<i>marginicolle</i> Wollaston, 1864
<i>doris</i> (Duftschmid, 1812)	<i>mendacissimum</i> J.R.Sahlberg, 1913
<i>maeoticum</i> (Kolenati, 1845)	<i>nebulosum</i> (Stephens, 1828)
<i>pseudotenellum</i> Netolitzky, 1910	<i>rumelicum</i> Apfelbeck, 1902
<i>Bembidion</i> (<i>Eupetedromus</i>) Netolitzky, 1911)	<i>tenebrosum</i> (Motschulsky, 1844)
<i>dentellus</i> (Thunberg, 1787)	<i>Bembidion</i> (<i>Ocydromus</i>) Clairville, 1806)
<i>flammulatum</i> (Clairville, 1806)	<i>saxatile</i> <i>saxatile</i> Gyllenhal, 1827
<i>meridionale</i> (Kolenati, 1845)	<i>vectense</i> Fowler, 1887
<i>undulatum</i> Sturm, 1825	<i>Bembidion</i> (<i>Odontium</i>) LeConte, 1848)
<i>Bembidion</i> (<i>Metallina</i>) Motschulsky, 1850)	<i>striatum</i> (Fabricius, 1792)
<i>lampros</i> (Herbst, 1784)	<i>hamburgense</i> P.Meier, 1899
<i>acutum</i> (Marsham, 1802)	<i>maurum</i> Netolitzky, 1918
<i>annulosum</i> (Schuler, 1962)	<i>nigrescens</i> Schilsky, 1888
<i>boreonense</i> (Avon, 1994)	<i>orichalcicum</i> (Duftschmid, 1812)
<i>celer</i> (Fabricius, 1792)	<i>tschalikovi</i> Samko, 1932
<i>felixianum</i> Heer, 1837	<i>Bembidion</i> (<i>Paraprincipium</i>) Netolitzky, 1914)
<i>lithuanicum</i> (Motschulsky, 1850)	<i>ruficolle</i> (Panzer, 1796)
<i>pulchellum</i> (Marsham, 1802)	<i>volgense</i> Becker, 1872
<i>phygas</i> (Antoine, 1956)	<i>Bembidion</i> (<i>Peryphanes</i>) Jeannel, 1941)
<i>pygmaeum</i> (Paykull, 1798)	<i>deletum</i> <i>deletum</i> Audinet-Serville, 1821
<i>rufipes</i> (Paykull, 1790)	<i>alpinum</i> Dejean, 1831
<i>triste</i> (Fabricius, 1792)	<i>brunnipes</i> Sturm, 1825
<i>velocipes</i> (Rossi, 1790)	<i>heerii</i> Kelch, 1846
<i>properans</i> (Stephens, 1828)	<i>nitidulum</i> (Marsham, 1802)
<i>chalceum</i> (Stephens, 1828)	<i>reticulatum</i> Iablokoff-Khnzorian, 1976
	<i>rufipes</i> Gyllenhal, 1810

- stephensi stephensi* Crotch, 1866
affine Stephens, 1835
canadense Hayward, 1897
ferlani (Quezel, Verdier, 1952)
glandonense (J.Ochs, 1949)
heterocerum Thomson, 1868
marthae Reitter, 1908
- Bembidion (Peryphiolus*** Jeannel, 1941)
monticola monticola Sturm, 1825
fuscocrine Dejean, 1831
- Bembidion (Peryphus*** Dejean, 1821)
bruxellense Wesmael, 1835
elegans (Stephens, 1835)
femoratum Gyllenhal, 1827
- cruciatum polonicum* P.W.J.Müller, 1930
dissolutum Hellén, 1934
- femoratum femoratum* Sturm, 1825
caucasicum (Motschulsky, 1850)
concinnum (Stephens, 1828)
nigrothoracicum (Goeze, 1777)
quadriflerum (Geoffroy, 1785)
- tetracolum tetracolum* Say, 1823
humerale Heer, 1838
littorale (Olivier, 1795)
nactum Casey, 1918
omostigma Netolitzky, 1914
pseudoustulatum P.W.J.Müller, 1934
tetraspilotum (Stephens, 1835)
- Bembidion (Philochthus*** Stephens, 1828)
aeneum aeneum Germar, 1824
javeti Jacquel du Val, 1852
marinum Schiödte, 1841
- Note: Insufficiently known species, known from single locality in Ropaži area (Central Latvia). Probably, should be more widely distributed in suitable biotopes in whole of Latvia.
- biguttatus* (Fabricius, 1779)
fuscipes (Stephens, 1828)
guttula Audinet-Serville, 1821
obscurum Verhoeff, 1890
subfenestratum (Stephens, 1828)
vulneratum Dejean, 1831
- guttula guttula* (Fabricius, 1792)
binotatum (Stephens, 1828)
bipustulatum Redtenbacher, 1849
haemorrhoum (Stephens, 1828)
nigrescens Friederichs, 1903
sumatrense Louwerens, 1953
vittatum (Stephens, 1828)
- lunulatum* (Geoffroy, 1785)
bisignatum Audinet-Serville, 1821
toletanum Perris, 1864
- Note: This species is known from single locality in the Pape area (extreme SW Latvia). Probably, it should be more widely distributed
- in the western part of Latvia.
- mannerheimii* C.R.Sahlberg, 1827
grandicolle (Motschulsky, 1850)
unicolor Chaudoir, 1850
- neresheimeri neresheimeri* (P.W.J.Müller, 1929)
Note: Insufficiently known species, known from a single locality in Moricsala State Nature reserve (W Latvia). Should be more widely distributed in the western part of Latvia. North-eastern border of main distribution area.
- Bembidion (Phyla*** Motschulsky, 1844)
obtusum Audinet-Serville, 1821
gracile (Stephens, 1828)
immune (Stephens, 1828)
pusillum (Stephens, 1828)
- Note: This species is known from single locality in the Pape area (extreme SW Latvia). Probably, should be more widely distributed in the western part of Latvia.
- Bembidion (Principium*** Motschulsky, 1864)
punctulatum punctulatum Drapiez, 1820
aerosum Erichson, 1837
chlorophanum Sturm, 1825
stagnorum (Geoffroy, 1785)
striatum (Duftschmid, 1812)
velox Dawson, 1854
- Bembidion (Semicampa*** Netolitzky, 1910)
gilvipes Sturm, 1825
kollari Dejean, 1831
nigrum (Stephens, 1828)
- schueppelii* Dejean, 1831
- Bembidion (Sinechostictus*** Motschulsky, 1864)
— *ruficornis ruficornis* Sturm, 1825
albipes (Stephens, 1829)
brunnipes Dejean, 1831
rufipes (Illiger, 1901)
- Bembidion (Testedium*** Motschulsky, 1864)
bipunctatum bipunctatum (Linnaeus, 1761)
- Bembidion (Trepandedoris*** Netolitzky, 1918)
doris (Panzer, 1796)
aquaticum (Panzer, 1796)
aquatilis (Illiger, 1798)
assimile (Stephens, 1828)
- Bembidion (Trepaines*** Motschulsky, 1864)
articulatum (Panzer, 1796)
diluticornis Netolitzky, 1918
poecilum (Stephens, 1828)
subglobosum (Paykull, 1798)
- octomaculatum octomaculatum* (Goeze, 1777)
pictum (Duftschmid, 1812)
sturmii (Panzer, 1803)
- Note: Insufficiently known species, recorded only from the valley of the river Gauja valley

and its tributaries (Central and northern Latvia). Probably, it should be more widely distributed in suitable biotopes, especially in the Central part of Latvia.

Ocys Stephens, 1828

quinquestriatus *quinquestriatus* (Gyllenhal, 1810)

acrocilius (Beck, 1817)
currens Stephens, 1828
pumilio (Duftschmid, 1812)
virens (Audinet-Serville, 1821)

Porotachys Netolitzky, 1914

bisulcatus (Nicolai, 1822)

focki (Hummel, 1822)
frontalis (Hayward, 1900)
guerini (Gaubil, 1844)
latipennis (Sturm, 1825)
numidicus (Lucas, 1846)
silaceus (Dejean, 1831)

Tachys (**Paratachys**) Casey, 1918

bistriatus (Duftschmid, 1812)

elongatulus (Dejean, 1831)
flavus (Becker, 1871)
insularis Ragusa, 1875
minimus (Curtis, 1828)
minutissimus Stephens, 1828
parallelus Motschulsky, 1850
rufulus Rey, 1882
vandeli Mateu, Colas, 1954

Tachyta (**Tachyta**) Kirby, 1837

nana (Gyllenhal, 1810)

laticollis (Motschulsky, 1850)
minima (Duftschmid, 1812)
persica Morvan, 1971

Patrobini Kirby, 1837

Patrobus Dejean, 1821

assimilis Chaudoir, 1844

borealis Motschulsky, 1844
campestris Motschulsky, 1844
clavipes Thomson, 1859

atrorufus (Ström, 1768)

biharicus Maran, 1933
bulgaricus Roubal, 1928
danihelkai Maran, 1936
excavatus (Paykull, 1790)
lutshniki Roubal, 1928
meridionalis P.W.J.Müller, 1911
napoleonius Reiche, 1857
schaubergeri Roubal, 1928

septentrionis Dejean, 1828

alpinus Curtis, 1827
australis J.R.Sahlberg, 1875
bitschnaui Reitter, 1908
hyperboreus Dejean, 1828
labradorinus Casey, 1918
lacustris Motschulsky, 1844
lapponicus Chaudoir, 1844

minuens Casey, 1918

picicornis (Zetterstedt, 1828)
relictus Neresheimer, Wagner, 1927
rubripennis Thomson, 1857
serenus Gredler, 1863

Note: This species is known from only a few localities in eastern and South-eastern Latvia.

Pogonini Laporte de Castelnau, 1834

Pogonus (**Pogonus**) Dejean, 1821)

— *luridipennis* (Germar, 1823)
burrellii Curtis, 1824
flavipennis Dejean, 1828

Trechini Bonelli, 1810

Perileptus (**Perileptus**) Schaum, 1860)

areolatus *areolatus* (Creutzer, 1799)

acuticollis (Dufour, 1843)
angustior Baudi di Selve, 1864
apicalis Della Beffa, 1909
depressus (Stephens, 1828)
niger Heyden, 1880

Note: This species is known from single locality in the river Ogre valley (Central Latvia). It is probably more widely distributed, especially in Southern parts of Latvia. North border of main distribution area.

Blemus Dejean, 1821

discus *discus* (Fabricius, 1792)
mariae (Hummel, 1823)
unifasciatus (Panzer, 1796)

Trechoblemus Ganglbauer, 1891

micros (Herbst, 1784)

abdominalis (Motschulsky, 1844)
flavus (Sturm, 1825)
littoralis (Audinet-Serville, 1821)
pallidus Reitter, 1894
planatus (Duftschmid, 1812)
quadricollis (Putzeys, 1847)
sericeus (Fleischer, 1829)

Trechus (**Epaphius**) Leach, 1819

rivularis (Gyllenhal, 1810)

incilis Dawson, 1849
nigricornis Motschulsky, 1844
secalis *secalis* (Paykull, 1790)
laticollis Motschulsky, 1844
testaceus (Fabricius, 1792)

Trechus (**Trechus**) Clairville, 1806)

austriacus Dejean, 1831

angustus J.R.Sahlberg, 1908
gajaci Coiffait, 1973
platypterus Sturm, 1825
podolicus Ljovuschkin, 1962

Note: This species is known only from SE Latvia. North border of main distribution area.

obtusus *obtusus* Erichson, 1837

castanopterus Herr, 1837

- laevis* Stephens, 1835
obtusiooides Jeannel, 1927
renati Jeannel, 1922
tristis Stephens, 1828
- quadristriatus* (Schrank, 1781)**
- amaurocephalus* Kolenati, 1845
 - capitatus* (Geoffroy, 1785)
 - fuscipennis* Stephens, 1828
 - humeralis* Dejean, 1831
 - minutus* (Fabricius, 1792)
 - piciventris* Graëlls, 1858
 - politus* Faldermann, 1836
 - syriacus* Putzeys, 1870
 - tempestivus* (Panzer, 1799)
- rubens* (Fabricius, 1792)
- pallidus* Sturm, 1825
 - palpalis* (Duftschmid, 1812)
 - paludosus* (Gyllenhal, 1810)
 - tristis* (Schönherr, 1806)
- Harpalinae** Bonelli, 1810
- Chlaeniini** Brullé, 1834
- Callistus** Bonelli, 1810
- *lunatus lunatus* (Fabricius, 1775)
- bifasciatus* Reitter, 1912
 - bioculatus* Roubal, 1922
 - cruxpallidus* (Mandl, 1986)
 - eques* (Schrank, 1781)
 - plateosus* (Geoffroy, 1785)
 - sulzeri* (Brahm, 1790)
- Chlaenius (Agostenus)** F. von Waldheim, 1829)
- costulatus* (Motschulsky, 1859)
- illigeri* Crotch, 1871
- quadrисulcatus* (Paykull, 1790)
- caelatus* (Weber, 1801)
- Note: Very rare species, only known from a few localities in western, Central and eastern Latvia.
- sulcicollis* (Paykull, 1798)
- alternans* Chaudoir, 1856
 - gebleri* Ganglbauer, 1891
- Chlaenius (Chlaeniellus)** Reitter, 1908)
- kindermannii chrysothorax* Krynický, 1832
- cordicollis* Motschulsky, 1865
 - nitens* Faldermann, 1836
 - uliginosus* Hochhuth, 1871
- Note: Relict species, known only from Daugava river valley. North border of main distribution area.
- nigricornis* (Fabricius, 1787)
- coeruleocephalus* Motschulsky, 1844
 - erythropus* Fischer von Waldheim, 1828
 - fulgidus* Stephens, 1828
 - fuscitarsis* Schaum, 1857
 - geoffroae* (Brahm, 1790)
 - melanocornis* Dejean, 1826
- nitidulus* (Thunberg, 1784 nec Schrank, 1781)
- punctatus* Motschulsky, 1844
 - upsaliensis* (Gmelin, 1790)
 - viridis* Westhoff, 1881
- nitidulus* (Schrank, 1781)**
- cyanoides* Jeannel, 1942
 - jeani* Puel, 1925
 - schrankii* (Duftschmid, 1812)
- tibialis* Dejean, 1826
- lomnickyi* Patkiewicz, 1910
- tristis tristis* (Schaller, 1783)**
- anceps* Chaudoir, 1876
 - carbonarius* (Rossi, 1790)
 - douei* Peyron, 1858
 - holosericeus* (Fabricius, 1787)
 - rufipes* Šterba, 1910
- vestitus* (Paykull, 1790)**
- distinctus* Chaudoir, 1856
 - dubius* (Hoppe, 1796)
 - oreteus* Ragusa, 1881
- Cyclosomini** Laporte de Castelnau, 1834
- Masoreus** Dejean, 1821
- wetterhallii* *wetterhallii* (Gyllenhal, 1813)
- laticollis* (Sturm, 1825)
 - luxatus* (Audinet-Serville, 1821)
- Harpalini** Bonelli, 1810
- Anisodactylus (Anisodactylus)** Dejean, 1829)
- binotatus* (Fabricius, 1787)
- brevicollis* Chaudoir, 1844
 - calceatus* Stephens, 1835
 - spurcaticornis* Dejean, 1829
- *nemorivagus* (Duftschmid, 1812)
- atricornis* Stephens, 1835
 - atripes* Ganglbauer, 1900
 - austriacus* Dalla Torre, 1877
 - crouzeti* Puel, 1931
 - gilvipes* Dejean, 1829
 - nigricornis* (Stephens, 1835)
 - nigripes* Gersdorf, 1953
 - orientalis* Gautier des Cottes, 1872
 - porosus* Puel, 1931
 - propinquus* Ballion, 1871
- Anisodactylus (Pseudanisodactylus)** Noonan, 1973)
- signatus* (Panzer, 1796)
- katavensis* (Jedlička, 1957)
- Diachromus** Erichson, 1837
- germanus* (Linnaeus, 1758)
- bimaculatus* Joukl, 1905
 - exquisitus* Mulsant, Rey, 1868
 - rollei* Fleischer, 1907
- Harpalus (Harpalus)** Latreille, 1802)
- affinis* (Schrank, 1781)
- aeneolus* Reitter, 1900
 - aeneopiceus* Stephens, 1828
 - aenescens* Casey, 1884
 - aspromontis* Hille, 1914
 - assimilis* Dejean, 1829

<i>bifoveolatus</i> Küster, 1846	<i>rufus</i> Brüggemann, 1873
<i>borysthеникус</i> Krynicki, 1832	<i>froelichi</i> Sturm, 1818
<i>canonicus</i> Casey, 1884	<i>aimaki</i> Jedlička, 1968
<i>coerulescens</i> Schilsky, 1828	<i>regularis</i> Motschulsky, 1844
<i>concininus</i> Stephens, 1828	<i>rivularis</i> Motschulsky, 1860
<i>confinis</i> Stephens, 1828	<i>segnis</i> Dejean, 1829
<i>confusus</i> Dejean, 1829	<i>tardoides</i> Jedlička, 1965
<i>convictor</i> Casey, 1884	<i>hirtipes</i> (Panzer, 1796)
<i>dentatus</i> Stephens, 1828	<i>latus</i> Motschulsky, 1844
<i>elegans</i> Preller, 1862	<i>motschulskyi</i> Gemminger, Harold, 1868
<i>interstitialis</i> Gredler, 1863	<i>laevipes</i> Zetterstedt, 1828
<i>kubanicus</i> Jedlička, 1957	<i>ainus</i> Habu, Baba, 1963
<i>latus</i> (Goeze, 1777)	<i>alienus</i> LeConte, 1879
<i>limbopunctatus</i> Fuss, 1858	<i>baergi</i> Csiki, 1932
<i>lustralis</i> Casey, 1884	<i>cascadiensis</i> Hatch, 1949
<i>nigrinus</i> Schilsky, 1888	<i>egregious</i> Casey, 1914
<i>paganettii</i> Flach, 1907	<i>impressipennis</i> Motschulsky, 1844
<i>parumpunctatus</i> W.Kolbe, 1924	<i>instructus</i> Casey, 1924
<i>proteus</i> (Paykull, 1790)	<i>montivagus</i> Reitter, 1900
<i>pseudoaeneolus</i> Schauberger, 1930	<i>motschoulskyanus</i> Schauberger, 1928
<i>semipunctatus</i> Dejean, 1829	<i>nigripes</i> Letzner, 1885
<i>subcaeruleus</i> Stephens, 1828	<i>pimalicus</i> Casey, 1914
<i>transparens</i> Motschulsky, 1844	<i>quadripunctatus</i> Dejean, 1829
<i>viridiaeneus</i> Palisot de Beauvois, 1811	<i>rufimanus</i> LeConte, 1848
<i>viridis</i> Say, 1823	<i>sachalinensis</i> Matsumura, 1911
<i>viridis</i> Schilsky, 1888	<i>latus</i> (Linnaeus, 1758)
<i>viridulus</i> (Geoffroy, 1785)	<i>acuminatus</i> Stephens, 1835
<i>weiratheri</i> P.W.J.Müller, 1931	<i>flaviventris</i> Sturm, 1818
<i>anxius</i> (Duftschmid, 1812)	<i>fulvipes</i> (Fabricius, 1792)
<i>ambigenus</i> Reiche, 1853	<i>lateralis</i> Stephens, 1835
<i>amicus</i> Dufour, 1843	<i>limbatus</i> (Duftschmid, 1812)
<i>faber</i> Ménétriés, 1832	<i>metallescens</i> Rye, 1874
<i>femoralis</i> Stephens, 1828	<i>versus</i> Roubal, 1917
<i>friulanus</i> P.W.J.Müller, 1926	<i>ruficeps</i> Curtis, 1833
<i>lerati</i> Antoine, 1920	<i>rugulosus</i> Heer, 1837
<i>nigripes</i> Sturm, 1818	<i>surinamensis</i> (Fabricius, 1792)
<i>piger</i> (Duftschmid, 1812)	<i>luteicornis</i> (Duftschmid, 1812)
<i>sericeus</i> (Duftschmid, 1812)	<i>servus</i> Stephens, 1828
<i>tibialis</i> Audinet-Serville, 1821	<i>modestus</i> Dejean, 1829
<i>autumnalis</i> (Duftschmid, 1812)	<i>aestivus</i> (Duftschmid, 1812)
<i>impiger</i> (Duftschmid, 1812)	<i>flavitarsis</i> Dejean, 1829
<i>inunctus</i> Sturm, 1818	<i>picipennis</i> (Duftschmid, 1812)
<i>rufus</i> Schilsky, 1889	<i>multisetosus</i> Thomson, 1884
<i>seriepunctatus</i> Sturm, 1818	<i>progrediens</i> Schauberger, 1922
Note: This species is known from single locality in the extreme SE of Latvia (Ilgas area) at the border with Lithuania and Belarus'.	<i>pumilus</i> Sturm, 1818
<i>distinguendus</i> <i>distinguendus</i> (Duftschmid, 1812)	<i>barthei</i> Antoine, 1920
<i>aeneuoides</i> J.Jacquet, 1942	<i>fleischeri</i> Eichler, 1924
<i>coeruleus</i> Schilsky, 1888	<i>funestus</i> Audinet-Serville, 1821
<i>contemptoides</i> Puel, 1935	<i>nigripennis</i> Motschulsky, 1844
<i>contemptulus</i> Puel, 1935	<i>parvulus</i> (Ballion, 1878)
<i>nigricans</i> Schilsky, 1888	<i>rubripes</i> (Duftschmid, 1812)
<i>psittaceus</i> (Geoffroy, 1785)	<i>alpestris</i> Redtenbacher, 1849
<i>subcontemptus</i> Puel, 1935	<i>amoenus</i> Heer, 1837
<i>virens</i> Schilsky, 1888	<i>azurescens</i> Gyllenhal, 1827
<i>flavescens</i> (Piller, Mitterpacher, 1783)	<i>chloropterus</i> Stephens, 1828
<i>fabricii</i> Crotch, 1781	<i>glaberellus</i> Sturm, 1818
	<i>hyperboreus</i> Motschulsky, 1844
	<i>ignavus</i> Stephens, 1828

- marginellus* Stephens, 1828
munganasti Reitter, 1908
nigrocaeruleus Stephens, 1828
nobilitatus Faldermann, 1836
potisii Stephens, 1835
punctiger Stephens, 1828
rufipes Motschulsky, 1844
sobrinus Dejean, 1829
truncatus Rosenhauer, 1842
turkestanicus Csiki, 1932
viridulus Solsky, 1874
rufipalpis rufipalpis Sturm, 1818
annulicornis Stephens, 1828
atrocaeruleus Stephens, 1828
maculicornis Stephens, 1835
notatus Stephens, 1828
obscuricornis Stephens, 1828
rufitarsis (Duftschmid, 1812)
servus (Duftschmid, 1812)
bucculentus Kraatz, 1874
complanatus Sturm, 1818
dilatatus Motschulsky, 1844
eupatoriae Schauberger, 1932
maritimus Schneider, 1898
obscuricornis Motschulsky, 1844
sericeus Motschulsky, 1844
smaragdinus (Duftschmid, 1812)
discoideus Erichson, 1837
discriminatus J.Jacquet, 1942
duftschmidii Sturm, 1818
perplexus Gyllenhal, 1827
petisii (Duftschmid, 1812)
reiningi Schauberger, 1934
solitaris Dejean, 1829
chevrierii Heer, 1837
fuliginosus (Duftschmid, 1812)
germanicus Reitter, 1900
lapponicus Zetterstedt, 1828
tardus (Panzer, 1796)
amaroides Faldermann, 1836
angustior J.R.Sahlberg, 1875
jacqueti Puel, 1936
pueli Antoine, 1920
rufimanus (Marsham, 1802)
xanthopus winkleri Schauberger, 1923
sulcatus Faldermann, 1836
Harpalus (Pseudoophonus) Motschulsky, 1844
calceatus (Duftschmid, 1812)
calcitrarus Motschulsky, 1844
itosimanus (Habu, 1954)
nonsignatus (Krynicki, 1832)
griseus (Panzer, 1796)
reichei Desbrochers des Loges, 1867
rufipes (DeGeer, 1774)
cribripennis Chaudoir, 1842
fuscus (Gmelin, 1790)
nigrosericeus (Goeze, 1777)
pubescens (O.F.Müller, 1776)

- pulverulentus* (Rossi, 1790)
ruficornis (Fabricius, 1775)
sagowskii (Lutshnik, 1909)
sericeus (Geoffroy, 1785)
Harpalus (Semiophonus) Schauberger, 1933
signaticornis (Duftschmid, 1812)
debrunieri (Briel, 1964)
impuncticeps (Reitter, 1900)
janus Fairmaire, 1856
Note: This species is known from a single locality in Ķemeri National park (Central Latvia). North border of main distribution area.
Ophonus (Hesperophonus) Antoine, 1959
azureus (Fabricius, 1775)
agnatus Chaudoir, 1846
albarracinus (Wagner, 1926)
apterus Bedel, 1899
atrocyanus Chaudoir, 1842
boemicus Roubal, 1917
chlorophanus (Panzer, 1799)
cyanus Ballion, 1878
gallicus (Pater, 1939)
grottagliensis (Schauberger, 1927)
koniensis (Schauberger, 1927)
oberthuri (Pater, 1938)
ruficrus (Ménétriés, 1932)
supremus (Schauberger, 1927)
Note: Very rare and insufficiently known species, with no confirmed records in the last 100 years.
Ophonus (Metophonous) Bedel, 1897
laticollis Mannerheim, 1825
griseoides Reitter, 1900
punctatulus (Duftschmid, 1812)
puncticeps Stephens, 1828
angusticollis P.W.J.Müller, 1921
crebrior (Tschitschérine, 1902)
orientis (Schauberger, 1926)
Note: This species is known from only a few specimens from the extreme SE part of Latvia (Ilgas area) situated at the border with Lithuania and Belarus'. Probably should be more widely distributed in suitable biotopes in S and SE parts of Latvia.
puncticollis (Paykull, 1798)
foraminulosus (Marsham, 1802)
fuscicornis Rey, 1886
meridianus (Schauberger, 1926)
ovipennis Motschulsky, 1844
punctatissimus Stephens, 1828
rusticus (Geoffroy, 1785)
trentinus (Csiki, 1932)
rufibarbis (Fabricius, 1792)
antoineianus (Schauberger, 1929)
cribellum Stephens, 1828
imitans (Schauberger, 1926)

orienticola (Tschitschérine, 1902)
parcepunctatus Reitter, 1902
seladon (Schauberger, 1926)
subpunctatus Stephens, 1828

rupicola (Sturm, 1818)

gruardeti (Pater, 1938)
subcordatus (Dejean, 1829)
subferrugatus (Pater, 1938)
zigzag A.Costa, 1882

schaubergerianus Puel, 1937

Note: Insufficiently known species, recorded from a few localities in SE and Central Latvia.

***Acupalpus* (*Acupalpus* Latreille, 1829)**

dubius Schilsky, 1888

Note: This species is only common in the city of Rīga and its surroundings (Central Latvia).

exiguus Dejean, 1829

minutus (Motschulsky, 1844)
subtilis (Motschulsky, 1850)

flavicollis (Sturm, 1825)

luridus Dejean, 1829
nigriceps Dejean, 1829
pumilio (Schaum, 1857)

meridianus (Linnaeus, 1761)

anthracinus J.Jacquet, 1938
chevrolati (Gaubil, 1846)
clarissimus Roubal, 1930
cruciger (Fabricius, 1798)
geeri (Gmelin, 1790)
humeralis J.Jacquet, 1938
suturalis (Stephens, 1828)
suturatus (Geoffroy, 1785)
thoracicus (Schilsky, 1889)

parvulus (Sturm, 1825)

celer (Olivier, 1795)
confusus P.W.J.Müller, 1926
derelictus (Dawson, 1854)
discus Reitter, 1894
dorsalis (Fabricius, 1787)
gyllenhali (Thomson, 1859)
pallidipennis Fassati, 1949
proserpinae Ragusa, 1886
ruficapillus Matits, 1910
thomsoni (C.Roth, 1897)
vittatus (Graëlls, 1858)

suturalis *suturalis* Dejean, 1829

Note: Insufficiently known species, widely distributed only on the NE coast of the Rīga Gulf.

***Acupalpus* (*Ancylostria* Schauberger, 1930)**

interstitialis Reitter, 1884

flavicornis A.Fiori, 1903

Note: This species is only known from single locality in Piedruja area (extreme SE Latvia), situated at the border with Belarus'.

***Anthracus* Motschulsky, 1850**

consputus (Duftschmid, 1812)

bivittulus Reitter, 1893
cordicollis (Chaudoir, 1846)
ephippiger (Duftschmid, 1812)
leonhardi Schatzmayr, 1909
melanocephalus (Dejean, 1829)
reitteri Wanka, 1915
wimmeli Reitter, 1893

***Bradyceillus* (*Bradyceillus* Erichson, 1837)**

caucasicus (Chaudoir, 1846)

agonus Puel, 1914
alatus Schaubberger, 1922
carnicus Schweiger, 1969
collaris (Paykull, 1798)
imitator Schweiger, 1969
paganettii Schweiger, 1969
transalpinus Schaubberger, 1922
turcicus Schweiger, 1969

— *harpalinus* (Audinet-Serville, 1821)

apterus P.W.J.Müller, 1926
fulvus (Marsham, 1802)

Note: Record needs confirmation.

***ruficollis* (Stephens, 1828)**

circumcinctus (R.F.Sahlberg, 1834)
littoreus (J.Jacquet, 1936)
similis (Dejean, 1829)

***Dicheirotrichus* (*Trichocellus* Ganglbauer, 1891)**

cognatus (Gyllenhal, 1827)

axillaris (Mannerheim, 1853)
boreellus (Casey, 1914)
conflagratus (Mannerheim, 1853)
deutschii (C.R.Sahlberg, 1827)
lateralis (Casey, 1914)
longiusculus (Mannerheim, 1853)
marginicollis (Motschulsky, 1845)
monticola (Casey, 1914)
nitens (LeConte, 1858)
obscuritarsis (Motschulsky, 1844)
punctipennis (Casey, 1914)
ruficrus (Kirby, 1834)

placidus (Gyllenhal, 1827)

affinis (C.R.Sahlberg, 1827)
strandi (Lutshnik, 1936)

rufithorax (C.R.Sahlberg, 1827)

***Stenolophus* (*Stenolophus* Dejean, 1821)**

mixtus (Herbst, 1784)

humeralis Mulsant, Godart, 1860
intermedius A.Fiori, 1914
pallidus Motschulsky, 1844
sinuatus Motschulsky, 1844
vespertinus (Panzer, 1796)
ziegleri (Panzer, 1809)

teutonus (Schrank, 1781)

aegeus P.W.J.Müller, 1921
anglicus Schiödte, 1861
confluens Kolenati, 1845
hirticornis Krynicki, 1829

- Lebiini** Bonelli, 1810
- Cymindis** (*Cymindis*) Latreille, 1806)
- angularis* Gyllenhal, 1810
- lonae* Wagner, 1927
 - lunaris* (Duftschmid, 1812)
 - rectangula* (Motschulsky, 1850)
 - sibirica* V.E.Jakovlev, 1891
- humeralis* (Geoffroy, 1785)
- designata* Reiche, 1863
 - dianae* (Fabricius, 1792)
 - elongata* Heer, 1841
 - hungarica* Csiki, 1908
 - humerosa* (Quensel, 1806)
 - hybrida* Heer, 1841
 - marmorae* Gené, 1839
 - rubrotestacea* Fleischer, 1892
 - sequistriata* (Panzer, 1789)
 - servillei* Solier, 1835
- Cymindis** (*Tarsostinus*) Motschulsky, 1864)
- macularis* Fischer von Waldheim, 1824
- crucifera* Reitter, 1923
 - fasciolata* Schilsky, 1888
 - fenestrata* Schilsky, 1888
- Cymindis** (*Tarulus*) Bedel, 1906)
- vaporariorum* (Linnaeus, 1758)
- asahiensis* Habu, Baba, 1962
 - basalis* Gyllenhal, 1810
 - dilatipennis* (Motschulsky, 1865)
 - gebleri* (Motschulsky, 1850)
 - humeralis* (Paykull, 1790)
 - immaculata* Dejean, 1829
 - intricata* Motschulsky, 1844
 - mollis* (Ström, 1768)
 - punctata* Dejean, 1825
 - subarctica* Kanô, 1933
- Demetrias** (*Aetophorus*) Schmidt-Göbel, 1846)
- imperialis* (Germar, 1824)
- interruptus* (Schilsky, 1888)
 - ruficeps* Schaum, 1857
 - rufus* Ragusa, 1892
- Demetrias** (*Demetrias*) Bonelli, 1810)
- atricapillus* (Linnaeus, 1758)
- angulatus* Motschulsky, 1844
 - confusus* Heer, 1838
 - elongatus* (Duftschmid, 1812)
 - erythrocephalus* Buxson, 1901
 - fulvus* (Geoffroy, 1785)
 - obscurus* Stephens, 1835
 - obtusus* Motschulsky, 1864
 - punctatostriatus* Motschulsky, 1864
 - sagitta* Cope, 1870
- Note: Very rare species with only a few confirmed records within the last 100 years.
- monostigma* Samouelle, 1819
- apicalis* Motschulsky, 1844
 - unipunctatus* (Germar, 1824)
- Calodromius** Reitter, 1905
- spilotus** (Illiger, 1798)
- basalis* (Schilsky, 1888)
 - championi* (Bedel, 1907)
 - geminatus* (Haldeman, 1843)
 - nigricornis* (Brullé, 1834)
 - quadrinotatus* (Panzer, 1799)
- Dromius** (*Dromius*) Bonelli, 1810)
- agilis* (Fabricius, 1787)
- bimaculatus* Dejean, 1825
 - caucasicus* Semenov, 1900
 - rufus* Dufour, 1820
 - rufescens* Hope, 1838
 - truncatus* (Fabricius, 1792)
- fenestratus* (Fabricius, 1794)
- arcticus* (Olivier, 1795)
 - biguttatus* (Clairville, 1806)
 - colon* (Thunberg, 1787)
- quadraticollis* A.Morawitz, 1862
- breviceps* Bates, 1883
 - cordicollis* Vorbringer, 1898
 - longulus* J.Frivaldzsky, 1884
 - oberti* Motschulsky, 1864
- quadrimaculatus* (Linnaeus, 1758)
- cracoviensis* Mazur, 1936
 - maxillosus* (Geoffroy, 1785)
- schneideri* Crotch, 1871
- marginellus* (Fabricius, 1794)
- Microlestes** Schmidt-Göbel, 1846
- maurus* *maurus* (Sturm, 1827)
- angustatus* (Stephens, 1839)
 - glabratus* (Stephens, 1827)
 - pygmaeus* (Heer, 1838)
- minutulus* (Goeze, 1777)
- femoralis* (Marsham, 1802)
 - glabratus* (Duftschmid, 1812)
 - longulus* (Reitter, 1901)
 - minutus* (Geoffroy, 1785)
- Paradromius** (*Manodromius*) Reitter, 1905)
- linearis* *linearis* (Olivier, 1795)
- praeustus* (Steven, 1809)
 - punctatostriatus* (Duftschmid, 1812)
- Paradromius** (*Paradromius*) Fowler, 1887)
- longiceps* (Dejean, 1826)
- capitalis* (Fairmaire, 1858)
 - muelleri* (Cadamuro Morgante, 1958)
- Philorhizus** Hope, 1838
- notatus* (Stephens, 1827)
- amplipennis* (Roubaud, 1928)
 - nigriventris* (Thomson, 1857)
 - oblitus* (Boieldieu, 1859)
- sigma* (Rossi, 1790)
- amurensis* (Reitter, 1887)
 - bepennifer* (Babington, 1835)
 - fasciatus* (Paykull, 1790)
 - sturmi* (Babington, 1835)
- Lebia** (*Lamprias*) Latreille, 1802)

chlorocephala (J.J.Hoffmann, 1803)
chrysocephala (Motschulsky, 1864)
insignicolor Roubal, 1937
micans Des Gozis, 1873
palustris E.Jacquet, 1887
cyanoccephala cyanoccephala (Linnaeus, 1758)
annulata Brullé, 1834
femoralis Chaudoir, 1844
formosa Comolli, 1837
geniculata Mannerheim, 1837
jeanneli Pater, 1939
nigritarsis (Stephens, 1827)
pilosella (Motschulsky, 1862)
subovata (Motschulsky, 1864)
violaceipennis (Motschulsky, 1862)

***Lebia* (*Lebia*) Latreille, 1802)**
cruxminor cruxminor (Linnaeus, 1758)
caucasica Motschulsky, 1844
errata (Rossi, 1790)
minor Ménétriés, 1848
nigricollis Gené, 1839
nigripes Dejean, 1825
sexmaculata Patkiewicz, 1922

Syntomus Hope, 1838
ai Barševskis, 1993

Note: This species is only known from the type locality in the Kauguri area (Rīga Gulf dunes, Jūrmala, Central Latvia). The state of this species is in dispute.

foveatus (Geoffroy, 1785)
foveola (Gyllenhal, 1810)
punctatellus (Duftschmid, 1812)

— *pallipes* (Dejean, 1825)
humeralis (Motschulsky, 1844)
pallidipes (Bedel, 1913)
tibialis (Motschulsky, 1844)

truncatellus (Linnaeus, 1761)
angelescui (Marcu, 1937)
impunctatus (Motschulsky, 1844)
sibiricus (Motschulsky, 1844)
tartarus (Bates, 1878)

Licinini Bonelli, 1810

***Badister* (*Badister*) Clairville, 1806)**
bullatus (Schrank, 1798)
ancora Ménétriés, 1832
crucifer (Geoffroy, 1785)
lunulatus Polentz, 1932
microcephalus Stephens, 1835
striatoquatermaculatus (Goeze, 1777)

lacertosus Sturm, 1815
binotatus Fischer von Waldheim, 1828

meridionalis Puel, 1925
antoinei Puel, 1925
kineli Makolski, 1952
suturalis Stephens, 1827

unipustulatus Bonelli, 1813

cephalotes Dejean, 1826
episternalis Gerhardt, 1888
pici Schatzmayr, 1909
teutonoides Preudhomme de Borre, 1882
trapezicollis Apfelbeck, 1904
unipunctatus Jacquelin du Val, 1857

***Badister* (*Trimorphus*) Stephens, 1828)**
dorsiger (Duftschmid, 1812)
xanthomus Chaudoir, 1844

sodalis (Duftschmid, 1812)
confinis (Stephens, 1828)
humeralis Bonelli, 1813
luteidorsis Roubal, 1917
scapularis (Stephens, 1828)

***Badister* (*Baudia*) Ragusa, 1884)**
collaris Motschulsky, 1844
anomalus (Perris, 1866)
gladiator Apfelbeck, 1904
striatulus V.Hansen, 1944
pseudopeltatus De Wailly, 1949

dilatatus Chaudoir, 1837
grafi Reitter, 1913

peltatus peltatus (Panzer, 1796)
chalybeus Sturm, 1824
corruscus Fischer von Waldheim, 1828
erro (Newman, 1838)
maeoticus Kolenati, 1845
ponticus Motschulsky, 1845

***Licinus* (*Licinus*) Latreille, 1802)**
depressus (Paykull, 1790)
cossyphoides (Duftschmid, 1812)
helveticus Heer, 1838

Odacanthini Laporte de Castelnau, 1834

***Odacantha* (*Odacantha*) Paykull, 1798)**
melanura (Linnaeus, 1767)
angustata (Fabricius, 1787)
puncticollis Mannerheim, 1844

Oodini LaFerté-Sénectère, 1851

Oodes Bonelli, 1810

gracilis Villa, Villa, 1833
gracilior Lambert, 1853
parallelogrammus Motschulsky, 1858
parallelus Motschulsky, 1858
similis Chaudoir, 1837

helopiooides helopiooides (Fabricius, 1792)
fiorii Porta, 1923
thessalonicensis Schatzmayr, 1909

Panagaeini Bonelli, 1810

***Panagaeus* (*Panagaeus*) Latreille, 1802)**
bipustulatus (Fabricius, 1755)
communimacula A.Fiori, 1883
conjunctus Everts, 1893
putzeysi Preudhomme de Borre, 1878
quadripustulatus Sturm, 1815

cruxmajor (Linnaeus, 1758)
elongatus Chaudoir, 1842

- equestris* (Geoffroy, 1785)
nobilis (Goeze, 1777)
trimaculatus Dejean, 1826
vicus Gory, 1833
- Perigonini** Horn, 1881
- Perigona (Treichicus)** LeConte, 1853
- nigriceps* (Dejean, 1831)
atriceps (Fairmaire, 1869)
fimicola (Wollaston, 1854)
jansoniana (Wollaston, 1858)
japonica (Bates, 1873)
pusilla (Péringuier, 1896)
tachyoides Bates, 1883
- Platynini** Bonelli, 1810
- Agonum (Agonum)** Bonelli, 1810
- afrum* (Duftschmid, 1812)
dolosum Puel, 1938
elevatum Puel, 1938
emarginatum (Gyllenhal, 1827)
falsum Puel, 1938
prominulum Puel, 1938
tenuimarginatum Leoni, 1907
- *gisellae* Csiki, 1931
angustatum Dejean, 1828
- chalconotum* Ménétriés, 1832
archangelicum (J.R.Sahlberg, 1875)
sahlbergii (Chaudoir, 1850)
- Note: This species is known only from SE part of Latvia.
- dolens* (C.R.Sahlberg, 1827)
latipenne Dejean, 1828
molestem Motschulsky, 1844
olivaceum Dejean, 1828
shimoyamai (Habu, 1974)
subtile (Motschulsky, 1844)
tarsatum (Zetterstedt, 1828)
triste Dejean, 1828
- duftschmidi* J.Schmidt, 1994
moestum (Duftschmid, 1812)
- ericeti* (Panzer, 1809)
bifoveolatum (C.R.Sahlberg, 1827)
fulgens Davis, 1832
- hypocrita* (Apfelbeck, 1904)
lugubre (Duftschmid, 1812)
- impressum* (Panzer, 1796)
splendidulum (Motschulsky, 1844)
- *longicornis* Chaudoir, 1846
holdhausi Apfelbeck, 1904
- lugens* (Duftschmid, 1812)
longipenne Mannerheim, 1844
- marginatum* (Linnaeus, 1758)
agrorum (Geoffroy, 1785)
flavocinctum Suffrian, 1854
lucorum (Geoffroy, 1785)
pretiosum (Friederichs, 1903)
viridinitidum (Goeze, 1777)
viridipunctatum (Goeze, 1777)
- muelleri* (Herbst, 1784)
chalibaeum (Gradl, 1881)
clandestinum Sturm, 1824
melletii Heer, 1837
parumpunctatum (Fabricius, 1792)
plicicolle Nicolai, 1822
tibiale Dejean, 1828
- sexpunctatum* (Linnaeus, 1758)
assimile (Motschulsky, 1865)
auraticolle Westhoff, 1881
cervini (Dupuis, 1911)
duodecimpunctatus (O.F.Müller, 1776)
mattosi (Paolino de Oliveira, 1882)
montanum Heer, 1837
motschulskyi Reitter, 1891
- versutum* Sturm, 1824
laeve Gyllenhal, 1827 nec Dejean, 1828
laeve Dejean, 1828 nec Gyllenhal, 1827
- viduum* (Panzer, 1796)
abnormicolle Puel, 1938
basale Puel, 1938
benoiti (Antoine, 1920)
brueggemanni (Verhoeff, 1891)
coerulescens Motschulsky, 1844
fallax (Apfelbeck, 1904)
limbicolle Puel, 1938
metallescens Dejean, 1828
obscurum (Paykull, 1790)
unicolor (Dejean, 1828)
vernale (Paykull, 1798)
- Agonum (Europhilus)** Chaudoir, 1859
- *consimile* (Gyllenhal, 1810)
borealis (Motschulsky, 1865)
- fuliginosum* (Panzer, 1809)
castaneipenne Motschulsky, 1844
convexum Gebler, 1829
minutum Motschulsky, 1844
pullum Stephens, 1828
pusillum Stephens, 1828
putridum (Zetterstedt, 1828)
simpsoni Stephens, 1828
striatum Stephens, 1828
- gracile* Sturm, 1824
atratum Stephens, 1828
latvicus (Barsevskis, 1993)
oblongum Dejean, 1831
patruelle (Gemminger, Harold, 1868)
- micans* (Nicolai, 1822)
cursitor Stephens, 1828
inauratum Mannerheim, 1825
- munsteri* (Hellén, 1835)
- Note: This species is known from only a few records from large peat-bogs in eastern Latvia.
- piceum* (Linnaeus, 1758)
canellipes Motschulsky, 1844
fuscipenne Nicolai, 1822
lutescens (Panzer, 1796)
pelidnum (Herbst, 1784)

- picipes* (Fabricius, 1787)
scitulum Dejean, 1828
thoreyi thoreyi Dejean, 1828
affine Stephens, 1828
chivense Lutshnik, 1934
lehmanni Chaudoir, 1837
longicolle Lacordaire, 1835
longulum Motschulsky, 1844
melanocephalum (Dejean, 1828)
puellum Dejean, 1828
stavopolitanum Lutshnik, 1934
Agonum (Platynomicrus) Casey, 1920
gracilipes (Duftschmid, 1812)
elongatum Fischer von Waldheim, 1823
Anchomenus (Anchomenus) Bonelli, 1810
dorsalis (Pontoppidan, 1763)
bicolor (Geoffroy, 1785)
cyanicollis Gebler, 1841
discophorus Chaudoir, 1842
infuscatus Chevrolat, 1854
malyi (Obenberger, 1914)
marchicus (Herbst, 1784)
maroccanus (Antoine, 1941)
prasinus (Thunberg, 1784)
thunbergi (Gmelin, 1790)
violaceus (Thunberg, 1784)
viridanus (Fabricius, 1787)
viridifulvus (Goeze, 1777)
viridis (Gmelin, 1790)
Olisthopus Dejean, 1828
rotundatus rotundatus (Paykull, 1790)
rotundicollis (Marsham, 1802)
vafer (Duftschmid, 1812)
Oxypselaphus Chaudoir, 1843
obscurus (Herbst, 1784)
oblongus (Fabricius, 1792)
pallidulus Chaudoir, 1843
taeniatus (Panzer, 1796)
Paranchus Lindroth, 1974
albipes (Fabricius, 1796)
antennatus (Gautier des Cottes, 1860)
flavipes (Geoffroy, 1785)
oblongus (Fabricius, 1792)
pallipes (Fabricius, 1801)
pavidus (Panzer, 1796)
ruficornis (Goeze, 1777)
sordidus (Marsham, 1802)
Platynus (Batenus) Motschulsky, 1865
livens (Gyllenhal, 1810)
bipunctatus (Sturm, 1824)
memnonius (Nicolai, 1822)
mundus (Germar, 1836)
mannerheimii (Dejean, 1828)
melas (Gebler, 1848)
morio (Gebler, 1847)
octofoveolatus (Mäklin, 1857)
- Note: Insufficiently known species.

- Platynus (Platynus)** Bonelli, 1810
assimilis (Paykull, 1790)
angusticollis (Fabricius, 1801)
corsicus (Tournier, 1860)
distinctus (Chaudoir, 1843)
junceus (Scopoli, 1763)
piceus (Dejean, 1831)
putzeysi (Preudhomme de Borre, 1882)
rufipes (Dejean, 1828)
krynickii (Sperk, 1835)
collaris (Motschulsky, 1844)
laticollis (Motschulsky, 1844)
magnicollis (Motschulsky, 1865)
nigerrimus (Chaudoir, 1837)
uliginosus (Erichson, 1837)
longiventris (Mannerheim, 1825)
impressipennis (Schilsky, 1891)
microthorax Fischer von Waldheim, 1829
Sericoda Kirby, 1837
quadripunctata (DeGeer, 1774)
ambigua (Mäklin, 1857)
cuprata (Sturm, 1824)
foveolata (Illiger, 1801)
nigrosericans (Heller, 1923)
octocola (Mannerheim, 1853)
- Note: This species is included on the list of indicator species of natural forest habitats.
- Pterostichini** Bonelli, 1810
Abax (Abax) Bonelli, 1810
— *parallelepipedus parallelepipedus* (Piller, Mitterpacher, 1783)
anuanus Depoli, 1939
athelinus Depoli, 1939
baldensis Depoli, 1939
bodonensis Depoli, 1939
dolomiticola Depoli, 1939
eukanensis Schatzmayr, 1944
tridentinus Depoli, 1939
zoldanus Depoli, 1939
- Poecilus (Poecilus)** Bonelli, 1810
cupreus cupreus (Linnaeus, 1758)
affinis (Sturm, 1824)
beryllinus Preller, 1862
brandisi Reitter, 1908
calabrus Flach, 1907
graecus Heyden, 1887
hoeltzerae (Lutshnik, 1916)
iridicolor (Westhoff, 1882)
prasinocupreus (Wagner, 1926)
puncticeps (Thomson, 1867)
rufifemoratus Stephens, 1828
viridis Preller, 1862
- lepidus lepidus* (Leske, 1785)
galiberti (Puel, 1932)
silvaticus Preller, 1862
virens (O.F.Müller, 1776)
viridanus Preller, 1862

- punctulatus* (Schaller, 1783)
crassicornis (Fischer von Waldheim, 1842)
punctatocollis Gemminger, Harold, 1868
puncticollis Motschulsky, 1884
- versicolor* (Sturm, 1824)
cupreoides (Heer, 1837)
janae (Kult, 1943)
medius Dejean, 1828
metallicus (Scopoli, 1763)
pauciseta (Thomson, 1867)
planicollis Motschulsky, 1860
punctatostriatus Stephens, 1828
subcyanus Preller, 1862
tenebricosus (Westhoff, 1881)
viridicolor (Westhoff, 1881)
- Pterostichus (Adelosia** Stephens, 1835)
- macer* *macer* (Marsham, 1802)
anachoreta (Ménétriés, 1832)
moniticola Nicolai, 1822
picimanus (Duftschmid, 1812)
siagonicus (Ménétriés, 1849)
sibiricus (Gebler, 1829)
thessalicus Schatzmayr, 1943
- Pterostichus (Argutor** Dejean, 1821)
- vernalis* (Panzer, 1796)
biimpressus (Fuss, 1858)
crenatus (Duftschmid, 1812)
degorsi (Croissandeau, 1893)
distinguendus (Hochhuth, 1871)
marijanovi (Lutshnik, 1922)
rotundicollis (Sturm, 1824)
rufomarginatus (Curtis, 1827)
ruthenus (Motschulsky, 1866)
sedulus (Dejean, 1828)
- Pterostichus (Bothriopterus** Chaudoir, 1835)
- *adstrictus* Eschscholtz, 1823
arcticus (Reiche, 1857)
borealis (Zetterstedt, 1828)
bulwerii (Stephens, 1828)
fugax A.Morawitz, 1862
orinomus (Stephens, 1828)
seriepunctatus Mannerheim, 1843
vireus (Dejean, 1828)
- oblongopunctatus oblongopunctatus* (F., 1787)
foveolatus (Gebler, 1841)
hafniensis (Gmelin, 1790)
octopunctatus Stephens, 1828
- quadrifoveolatus* Letzner, 1852
angustatus (Duftschmid, 1812)
octopunctatus Apfelbeck, 1904
- Pterostichus (Cheporus** Latreille, 1829)
- *burmeisteri burmeisteri* (Heer, 1838)
gundermanni Hubenthal, 1908
metallicus (Fabricius, 1792)
palae C.Müller, 1900
virens Schilsky, 1888
viridinitidus Pic, 1893
zebianus Schuberger, 1927

- Pterostichus (Eosteropus** Tschitschérine, 1902)
- aethiops* (Panzer, 1796)
cognatus (Stephens, 1835)
concinnus (Curtis, 1827)
glaber Vorbringer, 1897
monticola (Crotch, 1871)
vorbringeri Csiki, 1930
- Pterostichus (Melanius** Bonelli, 1810)
- aterrimus aterrimus* (Herbst, 1784)
- Pterostichus (Morphnosoma** Lutshnik, 1915)
- melanarius melanarius* (Illiger, 1798)
affinis (Stephens, 1828)
alternans (Carret, 1897)
arvernus (Sirguey, 1924)
ater (C.R.Sahlberg, 1827)
corrugatus (J.R.Sahlberg, 1880)
cribricollis (Motschulsky, 1850)
furvis (C.R.Sahlberg, 1827)
hyperoides (Gautier des Cottes, 1869)
nemoralis Heer, 1837
nigerrimus (Sturm, 1824)
pennatus (Dejean, 1828)
quadripunctatus (Sirguey, 1928)
rugicollis (Stephens, 1828)
sulcatus (Stephens, 1835)
- Pterostichus (Phonias** Des Gozis, 1886)
- diligens* (Sturm, 1824)
assimilis (Hochhuth, 1871)
barthei (Jeannel, 1949)
boreellus (J.R.Sahlberg, 1871)
heyrieri (Sturm, 1824)
kutensis Poppius, 1905
maeoticus (Motschulsky, 1850)
minutus (Motschulsky, 1866)
orcinulus Poppius, 1906
politus (Heer, 1837)
pullus (Gyllenhal, 1827)
rotundicollis (Duftschmid, 1812)
- *ovoideus ovoideus* (Sturm, 1824)
eruditus (Dejean, 1828)
interstinctus (Sturm, 1824)
- Pterostichus (Panzer, 1796)**
- difficilis* (Chaudoir, 1846)
dilatatus (Motschulsky, 1850)
erythropus (Marsham, 1802)
gagates (Duftschmid, 1812)
intermedius (Dejean, 1828)
lectulus Reitter, 1888
maritimus (Jeannel, 1942)
nigriceps (Sturm, 1824)
pygmaeus (Sturm, 1818)
pyrenaeus (Jeannel, 1942)
solers (Sturm, 1824)
wasastjerna (J.R.Sahlberg, 1875)
- Pterostichus (Platysma** Bonelli, 1810)
- niger niger* (Schaller, 1783)
baerii Motschulsky, 1845
bulgaricus Obenberger, 1917

bulgarus Csiki, 1930
clavipes (L.G.Scriba, 1790)
frischii (Herbst, 1784)
nigrostriatus (DeGeer, 1774)
scotus (Jeannel, 1942)
striatus (Rossi, 1792)
Pterostichus (***Pseudomaseus*** Chaudoir, 1838)
anthracinus anthracinus (Illiger, 1798)
corpulentus (Gebler, 1829)
crassipes (Ménétriés, 1832)
maurus (Fabricius, 1798)
gracilis gracilis (Dejean, 1828)
gottwaldi Jedlička, 1966
guentheri (Sturm, 1824)
rotundicollis (Stephens, 1835)
minor minor (Gyllenhal, 1827)
brunneus (Sturm, 1824)
laevigatus (Stephens, 1828)
nigrita (Paykull, 1790)
confluens (Panzer, 1796)
costulatus (Motschulsky, 1850)
excavatus (Boudier, 1844)
longibasis Jedlička, 1964
rufifemoratus (Stephens, 1828)
rhaeticus Heer, 1837
Stomis Clairville, 1806
pumicatus (Panzer, 1796)
tenuis (Marsham, 1802)
Sphodrini Laporte de Castelnau, 1834
Calathus (***Calathus*** Bonelli, 1810)
fuscipes fuscipes (Goeze, 1777)
cisteloides (Panzer, 1793)
frigidus (Fabricius, 1801)
leporinus (Geoffroy, 1785)
planipennis Germar, 1824
rossii Depoli, 1929
violatus Germar, 1824
Calathus (***Neocalathus*** Ball, Nègre, 1972)
ambiguus ambiguus (Paykull, 1790)
alatus Verhoeff, 1891
chevrolatii Gautier des Cottes, 1867
dilutus Chaudoir, 1842
fucus (Fabricius, 1792)
rufangulus (Marsham, 1802)
rufipes (Fabricius, 1792)
cinctus Motschulsky, 1850
erythroderus Gemminger, Harold, 1868
ruficollis Gautier des Cottes, 1868
 Note: Insufficiently known species, recorded from a few localities in SE and Central Latvia. Probably, should be more widely distributed in suitable biotopes in whole of Latvia.
erratus erratus (C.R.Sahlberg, 1827)
crocopus Stephens, 1828
erratoambiguus Verhoeff, 1891
fulvipes (Gyllenhal, 1810)

lasserrei Heer, 1838
rubidicollis Verhoeff, 1891
melanocephalus melanocephalus (Linnaeus, 1758)
alpinus Dejean, 1828
apicalis Newman, 1833
melanotus Putzeys, 1873
noricus K.Daniel, J.Daniel, 1891
nubigena Haliday, 1838
obscuricollis Chaudoir, 1837
ochropterus (Duftschmid, 1812)
parisiensis Gautier des Cottes, 1867
tarsalis J.R.Sahlberg, 1875
tricolor Reitter, 1887
micropterus (Duftschmid, 1812)
barthei Puel, 1939
borealis Motschulsky, 1850
elongatus Dejean, 1828
glabripennis Sturm, 1824
maculicornis Maran, 1935
microcephalus Dejean, 1828
sibiricus Gebler, 1841
mollis mollis (Marsham, 1802)
maritimus Schilsky, 1894
 Note: Insufficiently known species, probably should be widely distributed in suitable biotopes in SSE Latvia.
Laemostenus (***Pristonychus*** Dejean, 1828)
terricola terricola (Herbst, 1784)
castillanus (Coiffait, 1956)
cyanescens (Fairmaire, 1862)
episcopalalis (Drapiez, 1819)
inaequalis (Panzer, 1796)
lithuanicus (Motschulsky, 1850)
marginatus (Descourtilz, 1827)
reichenbachii (Schaufuss, 1861)
sardeus (Dejean, 1828)
silvaticus (Coiffait, 1956)
subaequalis (Schaufuss, 1865)
subcyaneus (Illiger, 1801)
subterraneus (Dejean, 1831)
torressalai (Schaufuss, 1861)
Sphodrus Clairville, 1806
leucophthalmus (Linnaeus, 1758)
armeniacus Osculati, 1844
indus Chaudoir, 1852
obsoletus (Rossi, 1790)
planus (Fabricius, 1792)
siculus Motschulsky, 1865
spiniger (Paykull, 1790)
 Note: Very rare species with only a few confirmed records within the last 50 years.
Synuchus (***Synuchus*** Gyllenhal, 1810)
vivalis vivalis (Illiger, 1798)
alpinus Heer, 1838
impiger (Panzer, 1809)
marginicollis (Duftschmid, 1812)
nivalis (Panzer, 1796)

- sivalis* (Motschulsky, 1865)
vehemensis Gistel, 1857
- Zabrini** Bonelli, 1810
- Amara** (*Amara* Bonelli, 1810)
- aenea* (DeGeer, 1774)
atra Stephens, 1828
devillei Jeannel, 1942
devincta Casey, 1918
maculipes Grimmer, 1841
palanda Jedlička, 1957
palustris Gistel, 1857
persica Chaudoir, 1842
sincera Lutshnik, 1933
sororcula Tschitschérine, 1895
thisbe Antoine, 1951
trivialis (Gyllenhal, 1810)
vogesiaca Bourgeois, 1898
- anthobia* Villa, Villa, 1833
krekichi Schatzmayr, 1909
- Note: This species is known by a few specimens from the Ilgas area (extreme SE Latvia) situated at the border with Lithuania and Belarus'.
- communis* (Panzer, 1797)
aemiliana A.Fiori, 1903
alpicola Heer, 1837
atrocoerulea Sturm, 1825
aubryi Schuler, 1964
dalei Rylands, 1841
ferrea Sturm, 1825
impressa Motschulsky, 1844
latescens Stephens, 1828
makolskii Roubal, 1923
manevali Jeannel, 1942
nigrita Chaudoir, 1844
udensis Jedlička, 1957
vagabunda (Duftschmid, 1812)
viatica Motschulsky, 1844
- convexior* Stephens, 1828
continua Thomson, 1873
obtusa Stephens, 1828
puncticollis Rylands, 1841
- curta* Dejean, 1828
aeruginosa Kolenati, 1845
brunnicornis Heer, 1837
cyanocnemis Thomson, 1857
ovalis Mulsant, Rey, 1861
- eurynota* (Panzer, 1796)
acuminata (Paykull, 1798)
aerata Stephens, 1828
exsculpta Hausmann, 1866
raymondi Putzeys, 1870
- famelica* Zimmermann, 1832
bidula Antoine, 1944
contrusa Schiödte, 1841
- familiaris* (Duftschmid, 1812)
cursor Strum, 1825

- elegans* Rylands, 1841
humilis Casey, 1918
levis Strum, 1825
perplexa Dejean, 1828
- littorea* Thomson, 1857
bogatshevi Lutshnik, 1934
cyanescens Baliani, 1940
dubia Motschulsky, 1844
kodymi Jedlička, 1936
motschulskyi Heyden, 1880
ovata Motschulsky, 1844
tockoensis Jedlička, 1936
wuki Matcha, 1922
- Note: This species is known from only a few specimens from the extreme SE part of Latvia (Ilgas area) situated at the border with Lithuania and Belarus'.
- lucida* (Duftschmid, 1812)
chobauti Puel, 1924
erythropa Stephens, 1828
gemina Zimmermann, 1932
heydeni Schatzmayr, 1909
septentrionalis Curtis, 1840
- lunicollis* Schiödte, 1837
arsenjevi Lutshnik, 1935
assimilis Chaudoir, 1844
carriana Casey, 1924
fatica Roubal, 1922
graculoides Jeannel, 1942
inepta LeConte, 1855
limbata Schiödte, 1837
marquettensis Casey, 1918
poecilloides Heer, 1837
zaisani Jedlička, 1964
- nitida* Sturm, 1825
consobrina Mäklin, 1880
guentheri J.R.Sahlberg, 1887
formosa Schiödte, 1837
laticollis Stephens, 1828
sinuata Motschulsky, 1844
- ovata* (Fabricius, 1792)
adamantina Kolenati, 1845
aequatorialis Baliani, 1939
audax Gistel, 1857
incisa Grimmer, 1841
kaulbacki Baliani, 1938
melanochlora Lutshnik, 1933
pratensis Sturm, 1825
sarsi Munster, 1927
similatoides Jedlička, 1934
tristis Baliani, 1937
trivialis (Duftschmid, 1812)
- pulpani* Kult, 1949
pseudocommunis Burakowski, 1957
- similata* (Gyllenhal, 1810)
agilis Rylands, 1841
curvicrus Thomson, 1871
depressa Zimmermann, 1832

<i>maghrebica</i> Antoine, 1941	<i>nivina</i> (Gistel, 1857)
<i>obsoleta</i> (Duftschmid, 1812)	<i>obscuricornis</i> (Motschulsky, 1859)
<i>spreta</i> Dejean, 1831	<i>punctulata</i> Dejean, 1828
<i>bipartita</i> Motschulsky, 1860	<i>rugulosa</i> (Gistel, 1857)
<i>tibialis</i> (Paykull, 1798)	<i>septentrionalis</i> Schiödte, 1837
<i>obscura</i> Motschulsky, 1844	<i>sibirica</i> Csiki, 1929
<i>viridis</i> (Duftschmid, 1812)	<i>torva</i> Lutshnik, 1933
Amara (Bradytus Stephens, 1827)	<i>trophina</i> (Gistel, 1857)
<i>apricaria</i> (Paykull, 1790)	<i>xenechthum</i> (Gistel, 1857)
<i>analis</i> (Fabricius, 1792)	Note: Insufficiently known species, known only from a few localities.
<i>armena</i> (Motschulsky, 1844)	
<i>baschkirica</i> (Schweiger, 1966)	<i>infima</i> (Duftschmid, 1812)
<i>convexilabris</i> Schiödte, 1837	<i>baeckmanni</i> Lutshnik, 1933
<i>paphlagoniensis</i> (Schweiger, 1966)	<i>granaria</i> Dejean, 1828
<i>parallela</i> (Chaudoir, 1842)	<i>ingenua</i> (Duftschmid, 1812)
<i>petri</i> Hummel, 1825	<i>liberta</i> (Duftschmid, 1812)
<i>putzeysi</i> G.H.Horn, 1875	<i>ruficornis</i> Dejean, 1828
<i>putzeysiana</i> Csiki, 1929	<i>subaenea</i> Sturm, 1825
<i>pygmea</i> Couper, 1865	<i>municipalis</i> <i>municipalis</i> (Duftschmid, 1812)
<i>rhaetica</i> Heer, 1837	<i>aerumnosa</i> Lutshnik, 1933
<i>uludagensis</i> (Schweiger, 1966)	<i>compos</i> Fischer von Waldheim, 1829
<i>consularis</i> (Duftschmid, 1812)	<i>dilatata</i> (Motschulsky, 1844)
<i>crassa</i> Stephens, 1828	<i>elevata</i> (Motschulsky, 1844)
<i>nigra</i> (Chaudoir, 1837)	<i>laevigata</i> (Motschulsky, 1844)
<i>patrata</i> Schiödte, 1841	<i>melancholica</i> Schiödte, 1837
<i>regismontium</i> (Schweiger, 1966)	<i>modesta</i> Dejean, 1828
<i>uralensis</i> (Motschulsky, 1844)	<i>quadrifoveolata</i> (Ballion, 1878)
<i>crenata</i> Dejean, 1828	<i>quadrifossulata</i> Csiki, 1929
<i>crenatostriata</i> (Chaudoir, 1846)	<i>praetermissa</i> (C.R.Sahlberg, 1827)
<i>fulva</i> (O.F.Müller, 1776)	<i>grandicollis</i> Zimmermann, 1832
<i>concolor</i> (Olivier, 1795)	<i>indivisa</i> Putzeys, 1871
<i>iridipennis</i> Heer, 1837	<i>oreophila</i> Zimmermann, 1832
<i>majuscula</i> Chaudoir, 1850	<i>monticusta</i> Iablokoff-Khnzorian, 1957
Amara (Celia Zimmermann, 1832)	<i>pallens</i> Sturm, 1825
<i>bifrons</i> (Gyllenhal, 1810)	<i>rufocincta</i> (C.R.Sahlberg, 1827)
<i>discrepans</i> Stephens, 1828	<i>solitaria</i> (Gistel, 1857)
<i>kleini</i> (Schuler, 1864)	<i>transcaucasica</i> Lutshnik, 1935
<i>longiceps</i> Seidlitz, 1887	Amara (Curtonotus Stephens, 1827)
<i>oricalcia</i> (O.F.Müller, 1776)	<i>aulica</i> (Panzer, 1796)
<i>zimmermanni</i> Heer, 1837	<i>abruzzensis</i> Schauberger, 1923
<i>brunnea</i> (Gyllenhal, 1810)	<i>caucasica</i> (Motschulsky, 1844)
<i>amplicollis</i> (Mannerheim, 1853)	<i>lithuanica</i> (Motschulsky, 1844)
<i>lapponica</i> (C.R.Sahlberg, 1827)	<i>convexuscula</i> (Marsham, 1802)
<i>mongolica</i> (Jedlička, 1966)	<i>bohemica</i> Fassati, 1944
<i>sahlbergi</i> (Zetterstedt, 1838)	<i>intermedia</i> (Motschulsky, 1844)
<i>cursitans</i> Zimmermann, 1832	Note: Insufficiently known species, widely distributed in coastal habitats along the Baltic Sea and Rīga Gulf, and with only a few localities inland.
<i>arnoldii</i> Lutshnik, 1934	<i>gebleri</i> Dejean, 1831
<i>bernhardti</i> (Schuler, 1965)	<i>circassica</i> Reitter, 1888
<i>cerdanica</i> (Aubry, 1970)	<i>helleri</i> Gredler, 1868
<i>fuscicornis</i> Zimmermann, 1832	<i>uralensis</i> (Motschulsky, 1850)
<i>oculata</i> Helliesen, 1912	— torrida (Panzer, 1796)
<i>ougsburgeri</i> Heer, 1837	<i>albertana</i> (Casey, 1924)
<i>properans</i> Zimmermann, 1832	<i>biarcuata</i> (Casey, 1924)
<i>erratica</i> (Duftschmid, 1812)	<i>brevilabris</i> (Kirby, 1837)
<i>affinis</i> (Motschulsky, 1850)	
<i>caucasicola</i> Lutshnik, 1935	
<i>cupricolor</i> Nakane, 1963	
<i>gracula</i> Heer, 1837	

- brevipennis* (Casey, 1924)
californica (Motschulsky, 1859)
consueta Fall, 1926
cylindrica (LeConte, 1878)
dura (Casey, 1924)
eschscholtzi (Chaudoir, 1837)
holmbergi (Putzeys, 1866)
hudsonica Hayward, 1908
infausta (LeConte, 1855)
johnsahlbergi Csiki, 1929
labradorensis (Casey, 1918)
melanogastrica Dejean, 1828
nairica Iablokoff-Khnzorian, 1964
picipes (Motschulsky, 1844)
pulla Jedlička, 1957
reflexa (Putzeys, 1966)
ruficornis J.R.Sahlberg, 1880
rufimana (Kirby, 1837)
rufimana (Motschulsky, 1844)
scrutata (Casey, 1918)
somnolenta (Putzeys, 1866)
striolata (Putzeys, 1866)
tenuestriata Baliani, 1943
turanica Jedlička, 1957
- Amara (Paracelia)** Bedel, 1899
quenseli silvicola Zimmermann, 1832
maritima Schiödte, 1837
- Amara (Percosia)** Zimmermann, 1832
equestris equestris (Duftschmid, 1812)
laticollis (Motschulsky, 1850)
mancipium (Duftschmid, 1812)
marginata (Stephens, 1831)
nobilis Sturm, 1825
patricia (Duftschmid, 1812)
plebeja (Duftschmid, 1812)
redikorzevi Lutshnik, 1934
tomoricensis P.W.J.Müller, 1923
- Amara (Zezea)** Csiki, 1929
plebeja (Gyllenhal, 1810)
barnevillei Croissandieu, 1893
brisouti Csiki, 1929
cylindrella Schiödte, 1837
lapidicola Heer, 1837
lenticularis Schiödte, 1837
noctivaga Hochhuth, 1871
punctibasis Jedlička, 1957
tridens A.Morawitz, 1862
varicolor Heer, 1838
- strenua* Zimmermann, 1832
vectensis Dawson, 1849
- *tricuspidata* Dejean, 1831
domascena (Reiche, 1876)
diversa Matits, 1911
impunctata Reiche, Saulcy, 1855
matitsi Roubal, 1917
- Zabrus (Zabrus)** Clairville, 1806
— *tenebrioides tenebrioides* (Goeze, 1777)
fossor (Fabricius, 1781)
- gibbosus* (Marsham, 1802)
gibbus (Fabricius, 1794)
hispanus Freude, 1988
obesus Stephens, 1828
piger (Geoffroy, 1785)
tenebrosus (Fabricius, 1801)
- Dytisciformia**
- Gyrinoidea**
- Gyrinidae** Latreille, 1810
- Gyrininae** Latreille, 1810
- Gyrinus (Gyrinulus)** Zaitzev, 1908
minutus Fabricius, 1798
kirbii Marsham, 1802
ohbayashii Sato, 1985
- Gyrinus (Gyrinus)** O.F.Müller, 1764
aeratus Stephens, 1835
edwardsi Sharp, 1914
thomsoni Zaitzev, 1908
- caspicus* Ménétriés, 1832
angustatus Aubé, 1838
celox Schiödte, 1841
elongatus Aubé, 1838
- Note: This species is known only by small series of individuals, collected in SE part of Latvia. North border of main distribution area.
- distinctus* Aubé, 1838
aegyptiacus Régimbart, 1883
atlanticus Régimbart, 1892
fairmairei Régimbart, 1883
niloticus Waltl, 1838
- marinus* Gyllenhal, 1808
aeneus Stephens, 1828
anthracinus Sturm, 1836
dorsalis Gyllenhal, 1808
epipleuralis Munster, 1925
lembus Schiödte, 1841
- natator* (Linnaeus, 1758)
marginatus Germar, 1824
mergus Ahrens, 1812
wanckowiczi Régimbart, 1883
- opacus* C.R.Sahlberg, 1819
blairi Omer-Cooper, 1931
indicus Aubé, 1838
lecontei Omer-Cooper, 1930
- paykulli* G.Ochs, 1927
pullatus Zaitzev, 1908
- substriatus* Stephens, 1828
cercurus Schiödte, 1841
corpulentus Schatzmayr, 1903
fowleri Omer-Cooper, 1930
oblitus Sharp, 1914
siculus Régimbart, 1882
- *suffrani* W.Scriba, 1855
- Orectochilus (Orectochilus)** Dejean, 1833
villosus villosum O.F.Müller, 1776
aquatica (Modeer, 1776)

- modeeri* (Marsham, 1802)
seidlitzi Jakobson, 1908

Haliploidea

Haliplidae Kirby, 1837

Brychius Thomson, 1859

elevatus (Panzer, 1793)

 - cristatus* J.R.Sahlberg, 1875
 - intermedius* P.W.J.Müller, 1908
 - norvegicus* Munster, 1922
 - obenbergeri* Riha, 1952
 - rossicus* Semenov, 1898

Note: Officially protected species in Latvia.

Haliplus (*Haliplidus* Guignot, 1928)

confinis Stephens, 1828

 - halberti* Bullock, 1928
 - helliesenii* Munster, 1922
 - lineatus* Aubé, 1836
 - pallens* Fowler, 1887
 - varius* Bold, 1868

obliquus (Fabricius, 1787)

 - albinus* Reitter, 1911
 - amoenus* (Olivier, 1795)
 - mulleri* Reitter, 1912

Haliplus (*Haliplus* Latreille, 1802)

fluviatilis Aubé, 1836

furcatus Seidlitz, 1887

heydeni Wehncke, 1875

 - foveostriatus* Thomson, 1884
 - multipunctatus* Wehncke, 1875
 - pedemontanus* A.Fiori, 1904
 - romanus* A.Fiori, 1904

immaculatus Gerhardt, 1877

interjectus H.Lindberg, 1937

lineolatus Mannerheim, 1844

 - brownianus* Sharp, 1913
 - browni* Sharp, 1913
 - laevigatus* Falkenström, 1936
 - nomax* Balfour-Browne, 1877
 - schaumi* Solsky, 1867
 - transversus* Thomson, 1870

ruficollis (DeGeer, 1774)

 - affinis* Stephens, 1828
 - impressus* (Fabricius, 1787)
 - marginepunctatus* (Panzer, 1793)

wehnkei Gerhardt, 1877

 - asiaticus* E.Scriba, 1905
 - borealis* Gerhardt, 1877
 - holsticus* Reitter, 1908
 - sibiricus* Motschulsky, 1860

Haliplus (*Liaphlus* Guignot, 1928)

flavicollis Sturm, 1834

 - expallidus* Zaitzev, 1908
 - pallidus* Semenov, 1904

fulvus (Fabricius, 1801)

 - carlittensis* Régimbart, 1901
 - interpunctatus* (Marsham, 1802)

lapponum Thomson, 1856

niger Seidlitz, 1887

pyreneus Delarouzée, 1857

salinarius Wallis, 1933

spareschneideri Munster, 1922

subguttatus Crotch, 1873

unicolor A.Fiori, 1914

unicolor Munster, 1922

variegatus Sturm, 1834

 - galileus* J.R.Sahlberg, 1913
 - leopardinus* J.R.Sahlberg, 1900
 - obsoletus* Westhoff, 1881
 - subnubilis* Babington, 1836
 - transvolgensis* Semenov, 1904

Peltodytes Régimbart, 1901

 - *caesus* (Duftschmid, 1805)
 - delhermi* (Mazéret, 1923)
 - levantinus* (J.R.Sahlberg, 1903)
 - quadrimaculatus* (Drapiez, 1820)

Dytiscoidea

Noteridae Thomson, 1860

Noterinae Thomson, 1860

Noterini Thomson, 1860

Noterus Clairville, 1806

clavicornis (DeGeer, 1774)

 - convexusculus* Reiche, Saulcy, 1855
 - semipunctatus* (Fabricius, 1792)
 - sparsus* (Marsham, 1802)

crassicornis (O.F.Müller, 1776)

 - capricornis* (Herbst, 1784)
 - geeri* Leach, 1817
 - simulator* Porta, 1923

Dytiscidae Leach, 1815

Agabinae Thomson, 1867

Agabini Thomson, 1867

Agabus (*Acatodes* Thomson, 1859)

clypealis (Thomson, 1867)

 - scholzi* W.Kolbe, 1916

confinis (Gyllenhal, 1808)

 - longulus* (LeConte, 1878)
 - ovoideus* (Crotch, 1873)

congener (Thunberg, 1794 nec Paykull, 1798)

 - congener* (Paykull, 1798 nec Thunberg, 1794)
 - daisetsuzanus* Kamiya, 1938
 - foveolatus* Mulsant, Godart, 1860
 - finkii* Seidlitz, 1887
 - venturi* Bertolini, 1870

fuscipennis (Paykull, 1798)

 - eversmanni* Ballion, 1855
 - fossarum* (Germar, 1824)
 - obscurior* (J.R.Sahlberg, 1875)

pseudoclypealis Scholtz, 1933

 - haraldi* H.Lindberg, 1933

setulosus (J.R.Sahlberg, 1895)

sturmii (Gyllenhal, 1808)

 - fallax* Munster, 1932

- goedelii* (Villa, Villa, 1833)
- Agabus*** (*Agabus* Leach, 1817)
- labiatus* (Brahm, 1791)
 - assimilis* (Sturm, 1834)
 - femoralis* (Paykull, 1798)
 - femoralis* (Gravenhorst, 1807)
 - hochhuti* Zaitzev, 1908
 - impressus* (Zoubkoff, 1833)
 - transcaucasicus* Zaitzev, 1927
 - uliginosus* (Linnaeus, 1761)
 - aeratus* (Stephens, 1828)
 - dispar* Bold, 1849
 - reichei* Aubé, 1837
 - uliginosus* O.F.Müller, 1776
 - undulatus* (Schrink, 1776)
 - abbreviatus* (Fabricius, 1787)
 - imperfectus* P.Meier, 1899
 - interruptus* Schilsky, 1888
 - pictus* P.Meier, 1899
 - ruficeps* (Ménétriés, 1832)
 - Agabus*** (*Gaurodytes* Thomson, 1859)
 - affinis* (Paykull, 1798)
 - branchiatus* (Babington, 1832)
 - guttulus* (Schönherz, 1808)
 - biguttulus* (Thomson, 1867)
 - boreellus* (J.R.Sahlberg, 1871)
 - ovalis* (J.R.Sahlberg, 1877)
 - bipustulatus* (Linnaeus, 1767)
 - abdominalis* (A.Costa, 1847)
 - acuductus* (Marsham, 1802)
 - alpestris* Heer, 1837
 - alpicola* Zaitzev, 1927
 - alticola* Brunéau de Mire, Legros, 1963
 - bipustulatus* (O.F.Müller, 1776)
 - callosus* (Thomson, 1884)
 - carbonarius* (Fabricius, 1801)
 - dolomitanus* Scholz, 1935
 - falcozi* Guignot, 1932
 - immaculatus* (Schrink, 1781)
 - kiesenwetterii* Seidlitz, 1887
 - latus* (Gebler, 1841)
 - luctuosus* (Geoffroy, 1785)
 - maurus* (A.Zimmermann, 1919)
 - peyerimhoffi* Brunéau de Mire, Legros, 1963
 - picipennis* J.R.Sahlberg, 1903
 - pyrenaeus* Fresneda, Hernando, 1989
 - regalis* Petri, 1903
 - remotus* J.R.Sahlberg, 1913
 - sexualis* Reiche, 1857
 - snowdonius* (Newman, 1833)
 - solieri* Aubé, 1837
 - guttatus* (*guttatus* (Paykull, 1798)
 - nigripes* (A.Costa, 1847)
 - picinus* (Marsham, 1802)
 - septemstriatus* (J.R.Sahlberg, 1875)
 - severior* Des Gozis, 1912
 - signatus* (Grimmer, 1841)
 - styriacus* Sharp, 1882

- nebulosus* (Forster, 1771)
- abdominalis* Ragusa, 1887
 - bipunctatus* (Fabricius, 1787)
 - immaculatus* (Gschwendtner, 1927)
 - mixtus* Guignot, 1949
 - naevius* (Gmelin, 1790)
 - nigromaculatus* (Goeze, 1777)
 - notatus* (Bergsträsser, 1778)
 - pratensis* Schaufuss, 1882
 - ragusai* Zaitzev, 1908
 - rugosipennis* Guignot, 1932
 - tesselatus* (Geoffroy, 1785)
 - paludosus* (Fabricius, 1801)
 - marginalis* Sharp, 1882
 - pallidipennis* (Laporte de Castelnau, 1835)
 - politus* (Marsham, 1802)
 - striolatus* (Gyllenhal, 1808)
 - costatus* Gerhardt, 1910
 - rectus* (Babington, 1841)
 - unguicularis* (Thomson, 1867)
 - Ilybius*** Erichson, 1832
 - aenescens* Thomson, 1870
 - kiesenwetteri* Kraatz, 1872
 - angustior* (Gyllenhal, 1808)
 - ater* (DeGeer, 1774 nec Panzer, 1796)
 - ater* (Panzer, 1796 nec DeGeer, 1774)
 - ungularis* (LeConte, 1862)
 - chalconatus* (Panzer, 1796)
 - aterrimus* (Stephens, 1828)
 - fuscoaeneus* (Régimbart, 1878)
 - nigroaeneus* (Marsham, 1802)
 - erichsoni* Gemminger, Harold, 1868
 - cincticollis* Mäklin, 1881
 - lutosus* (Crotch, 1873)
 - fenestratus* (Fabricius, 1781)
 - aeneus* (Panzer, 1796)
 - ciliatus* (Olivier, 1791)
 - prescotti* (Mannerheim, 1829)
 - fuliginosus* (*fuliginosus* (Fabricius, 1792)
 - lacustris* (Panzer, 1796)
 - pirinicus* V.G.Guéorguiev, 1957
 - guttiger* (Gyllenhal, 1808)
 - immunis* (Stephens, 1828)
 - kiesenwetteri* Wehncke, 1872
 - quadrinotatus* (Stephens, 1828)
 - montanus* (Stephens, 1828)
 - melanocornis* A.Zimmermann, 1915
 - neglectus* (Erichson, 1837)
 - quadriguttatus* (Lacordaire, 1835)
 - obscurus* (Marsham, 1802 nec Panzer, 1794)
 - sexdentatus* Schiödte, 1841
 - similis* Thomson, 1856
 - ovatus* Hochhuth, 1871
 - subaeneus* Erichson, 1837
 - badenii* Wehncke, 1872
 - chalybeatus* Thomson, 1860
 - lapponicus* Scholz, 1917

<i>suffusus</i> Crotch, 1873	<i>suturalis</i> (McLeay, 1825)
<i>viridiaeneus</i> Crotch, 1873	<i>annamita</i> Régimbart, 1899
<i>subtilis</i> Erichson, 1837	<i>australis</i> (Aubé, 1838)
<i>altaicus</i> Gebler, 1848	<i>birmanicus</i> Vazirani, 1970
<i>wasastjernae</i> (C.R.Sahlberg, 1824)	<i>bramardi</i> Guignot, 1942
<i>kenaiensis</i> (Fall, 1926)	<i>chinensis</i> Falkenström, 1936
<i>palustris</i> (Wallis, 1926)	<i>dispar</i> Régimbart, 1899
<i>Platambus</i> Thomson, 1859	<i>flaviventris</i> Schilsky, 1908
<i>maculatus</i> (Linnaeus, 1758)	<i>hypochlorus</i> Des Gozis, 1911
<i>biocellatus</i> (O.F.Müller, 1776)	<i>montrouzieri</i> (Lucas, 1860)
<i>cantalicus</i> Pic, 1912	<i>morneri</i> Falkenström, 1937
<i>caucasicus</i> Zaitzev, 1927	<i>neoguineensis</i> V.B.Guéorguiev, Rocchi, 1993
<i>escalerai</i> Régimbart, 1900	<i>pulverosus</i> (Stephens, 1828)
<i>glacialis</i> (Graëlls, 1858)	<i>punctatus</i> (Geoffroy, 1785)
<i>graellsii</i> (Gemminger, Harold, 1868)	<i>regimbarti</i> A.J.Jakovlev, 1896
<i>hebraicus</i> (Geoffroy, 1785)	<i>ruficollis</i> Schilsky, 1908
<i>inornatus</i> Schilsky, 1888	<i>rufimanus</i> (White, 1846)
<i>ornatus</i> (Herbst, 1784)	<i>sharpi</i> A.I.Jakovlev, 1896
<i>pulchellus</i> (Heer, 1839)	<i>suturellus</i> (Harris, 1828)
<i>Colymbetini</i> Erichson, 1837	<i>semilineatus</i> A.Fleischer, 1898
<i>Colymbetes</i> Clairville, 1806	<i>subopacus</i> (Mannerheim, 1853)
<i>fuscus</i> (Linnaeus, 1758)	<i>zimmermanni</i> Wallis, 1933
<i>affinis</i> Stein, 1881	<i>Copelatinae</i> Van den Branden, 1885
<i>semirufus</i> A.Zimmermann, 1921	<i>Copelatini</i> Van den Branden, 1885
<i>transversestriatus</i> (DeGeer, 1774)	<i>Copelatus</i> Erichson, 1832
<i>paykulli</i> Erichson, 1837	<i>haemorrhoidalis</i> (Fabricius, 1787)
<i>striatus</i> (Linnaeus, 1758)	<i>agilis</i> (Fabricius, 1792)
<i>bogemannii</i> (Gyllenhal, 1813)	<i>oblongus</i> (Illiger, 1801)
<i>Rhantus</i> (<i>Nartus</i>) Zaitzev, 1907	<i>ruficollis</i> (Schaller, 1783)
<i>grapii</i> (Gyllenhal, 1808)	<i>schalleri</i> (Gmelin, 1790)
<i>niger</i> (Lacordaire, 1835)	<i>Dytiscinae</i> Leach, 1815
<i>Rhantus</i> (<i>Rhantus</i>) Dejean, 1833)	<i>Aciliini</i> Thomson, 1867
<i>bistriatus</i> (Bergsträsser, 1778)	<i>Acilius</i> (<i>Acilius</i>) Leach, 1817)
<i>aberratus</i> Gemminger, Harold, 1868	<i>canaliculatus</i> (Nicolai, 1822)
<i>nigropunctatus</i> Motschulsky, 1860	<i>caliginosus</i> Curtis, 1825
<i>pericaspius</i> Guignot, 1956	<i>dispar</i> Lacordaire, 1835
<i>exsoletus</i> (Forster, 1771)	<i>kotulae</i> Ulanowski, 1884
<i>adspersus</i> (Panzer, 1796)	<i>laevisulcatus</i> Motschulsky, 1845
<i>collaris</i> (Paykull, 1798)	<i>subimpressus</i> Motschulsky, 1845
<i>insolatus</i> (Gebler, 1829)	<i>sulcatoaffinis</i> (Hummel, 1822)
<i>melanopterus</i> (Zetterstedt, 1828)	<i>sulcipennis</i> (Zetterstedt, 1824)
<i>nigriventris</i> Newberry, Sharp, 1915	<i>xantischius</i> Des Gozis, 1911
<i>oculatus</i> (Herbst, 1784)	<i>sulcatus</i> (Linnaeus, 1758)
<i>frontalis</i> (Marsham, 1802)	<i>blancki</i> Peyerimhoff, 1927
<i>flavicollis</i> (Eschscholtz, 1818)	<i>brevis</i> Aubé, 1837
<i>gyllenhalii</i> (Laporte de Castelnau, 1840)	<i>fasciatus</i> (DeGeer, 1774)
<i>punctatus</i> (Hope, 1795)	<i>punctatus</i> (Scopoli, 1763)
<i>semicirculatus</i> O.Schneider, 1898	<i>scopolii</i> (Gmelin, 1790)
<i>suturalis</i> Lacordaire, 1835	<i>scoticus</i> Stephens, 1828
<i>vermicularis</i> Fauvel, 1889	<i>tomentosus</i> Motschulsky, 1845
<i>virgulatus</i> (Illiger, 1806)	<i>varipes</i> Stephens, 1828
<i>incognitus</i> Scholz, 1927	<i>Graphoderus</i> Dejean, 1833
<i>latitans</i> Sharp, 1882	<i>austriacus</i> (Sturm, 1834)
<i>notaticollis</i> (Aubé, 1837)	<i>bilineatus</i> (DeGeer, 1774)
<i>infuscatus</i> (Erichson, 1837)	Note: Officially protected species in Latvia.
<i>nigriventris</i> Motschulsky, 1860	<i>cinereus</i> (Linnaeus, 1758)
<i>vibicicollis</i> (Hochhuth, 1846)	<i>intermedius</i> (Westhoff, 1881)

- simulator* (Westhoff, 1882)
taeniatus (Rossi, 1792)
zonatus verrucifer (C.R.Sahlberg, 1824)
bertolinii Seidlitz, 1887
flavotestaceus J.R.Sahlberg, 1875
piciventris Thomson, 1868
rosenbergeri Seidlitz, 1887
sahlbergii Seidlitz, 1887
thomsonii Seidlitz, 1887
- Cybistrini** Sharp, 1882
- Cybister* (*Melanectes*)** Brinck, 1945
- lateralimarginalis* *lateralimarginalis* (DeGeer, 1774)
- dispar* (Rossi, 1790)
 - dissimilis* (Rossi, 1792)
 - glaber* (Bergsträsser, 1778)
 - herbeti* (Peytoureau, 1894)
 - intricatus* (Schaller, 1783)
 - jordanis* (Reiche, Saulcy, 1855)
 - lepidus* Apetz, 1852
 - lusitanicus* Sharp, 1882
 - pallidus* (Ménétriés, 1832)
 - roeselii* (Fuessly, 1775)
 - tataricus* (Gebler, 1829)
 - velutus* (Schrank, 1798)
 - virens* (O.F.Müller, 1776)
 - virescens* (Gmelin, 1790)
- Dytiscini** Leach, 1815
- Dytiscus*** Linnaeus, 1758
- circumcinctus* Ahrens, 1811
- angustatus* Curtis, 1826 nec Stephens, 1828
 - angustatus* Stephens, 1828 nec Curtis, 1826
 - anxius* Mannerheim, 1843
 - circumscriptus* Lacordaire, 1835
 - confusus* Motschulsky, 1860
 - dubius* Gyllenhal, 1827
 - flavocinctus* Hummel, 1823
 - fuscostriatus* Motschulsky, 1859
 - ooligbukii* Kirby, 1837
- circumflexus* Fabricius, 1810
- dubius* Audinet-Serville, 1830
 - flavomaculatus* Curtis, 1826
 - flavoscutellatus* Latreille, 1804
 - kunstleri* Peytoureau, 1894
 - perplexus* Lacordaire, 1835
- dimidiatus* Bergsträsser, 1778
- lapponicus* *lapponicus* Gyllenhal, 1808
- latissimus* Linnaeus, 1758
- amplissimus* O.F.Müller, 1776
 - anastomozanus* Well, 1781
- Note: Officially protected species in Latvia.
- marginalis* *marginalis* Linnaeus, 1758
- circumductus* Audinet-Serville, 1830
 - conformis* Kunze, 1818
 - semicostatus* Reineck, 1921
 - semistriatus* Linnaeus, 1758
 - submarginatus* Stephens, 1828
- totomarginalis* DeGeer, 1774
- semisulcatus*** O.F.Müller, 1776
- expectatus* Peyerimhoff, 1905
 - frischii* Bergsträsser, 1778
 - laevis* Engert, 1911
 - maurus* Schaufuss, 1883
 - porcatus* Thunberg, 1794
 - punctatus* Olivier, 1795
 - punctulatus* Fabricius, 1776
 - stagnalis* Geoffroy, 1785
- Hydaticini** Sharp, 1882
- Hydaticus* (*Hydaticus*)** Leach, 1817
- aruspe* Clark, 1864
- americanus* Sharp, 1882
 - laevipennis* Thomson, 1867
 - laeviusculus* Poppius, 1906
 - modestus* Sharp, 1882
 - rugosus* Poppius, 1905
- seminiger* (DeGeer, 1774)
- cinctus* (O.F.Müller, 1776)
 - danicus* (Gmelin, 1790)
 - hybneri* (Fabricius, 1787)
 - parapleurus* (Marsham, 1802)
 - sexpunctatus* (Drapiez, 1819)
- transversalis* *transversalis* (Pontoppidan, 1763)
- degeneratus* Westhoff, 1882
 - inderiensis* Zaitzev, 1915
 - interruptemaculatus* Gerhardt, 1899
 - punctipennis* Thomson, 1867
 - sahlbergi* Zaitzev, 1905
- Hydroporinae** Aubé, 1836
- Bidessini** Sharp, 1882
- Bidessus*** Sharp, 1882
- grosssepunctatus* Vorbringer, 1907
- unistriatus* (Goeze, 1777)
- bisulcatus* (Curtis, 1840)
 - monostriatus* (Geoffroy, 1785)
 - nanus* (Gmelin, 1790)
 - opuscus* Guignot, 1932
 - subrufulus* Schneider, 1898
 - unistriatus* (Schrank, 1781)
- Hydroglyphus*** Motschulsky, 1853
- geminus* (Fabricius, 1787)
- corsicus* (O.Schneider, 1903)
 - inmaculatus* (Báguena Corella, 1935)
 - monaulacus* (Drapiez, 1820)
 - nitens* (Falkenström, 1939)
 - obscurus* (J.R.Sahlberg, 1903)
 - ocellatus* (Báguena Corella, 1935)
 - pectinatus* (Báguena Corella, 1935)
 - pusillus* (Fabricius, 1781)
 - pygmaeus* (Olivier, 1795)
 - symbolum* (Kolenati, 1845)
 - trifidus* (Panzer, 1795)
- hamulatus* (Gyllenhal, 1813)
- Hydroporini** Aubé, 1836
- Deronectes*** Sharp, 1882

latus (Stephens, 1829)
castaneus (Heer, 1837)
ovatus (Sturm, 1835)
pyrenaeus (Schaefer, 1930)

Note: Officially protected species in Latvia.

Graptodytes Seidlitz, 1887
granularis (Linnaeus, 1767)
funestus (Schilsky, 1888)
istrus P.W.J.Müller, 1926
pseudopictus Zaitzev, 1928
rhodanensis Schaefer, 1952
suturalis (Müller, 1821)
unilineatus (Schrink, 1781)

pictus (Fabricius, 1787)
arcuatus (Panzer, 1795)
cruciatus (Schilsky, 1890)
flexuosus (Marsham, 1802)
fortiterpunctatus Guignot, 1932

Hydroporus Clairville, 1806
angustatus Sturm, 1835
acuminatus Sturm, 1835
brevis R.F.Sahlberg, 1834
discretus Fairmaire, Brisout de Barneville, 1859
alpestris Falkenström, 1939
cyprius Régimbart, 1878
lundbladi Falkenström, 1939
maurus Sharp, 1882
peschetti Guignot, 1930
ponticus Zaitzev, 1927
sublaevis Falkenström, 1922
taticus Kinel, 1949

elongatulus Sturm, 1835
eugeniae Zaitzev, 1909
lenensis Poppius, 1905

erythrocephalus (Linnaeus, 1758)
deplanatus (Gyllenhal, 1826)
derelictus Clark, 1862
faroensis Mjöberg, 1917
sericeus (Eschscholtz, 1818)
subcostatus Gerhardt, 1909

fuscipennis Schaum, 1868
criniticoxis Larson, 1975
flavipennis Munster, 1932
piceus Sturm, 1835
puberulus Mannerheim, 1853

glabriusculus Aubé, 1838
incognitus Sharp, 1869
discedens Régimbart, 1877
ellypticus Petri, 1912
seidlitzii Gerhardt, 1899

longicornis Sharp, 1871
parallelus Sharp, 1869

melanarius Sturm, 1835
montanus Helliesen, 1890
monticola Sharp, 1869
munsterii Helliesen, 1890
obscuricornis J.R.Sahlberg, 1921

ruficornis Zetterstedt, 1838
memnonius Nicolai, 1822
castaneus Aubé, 1838
incertus Aubé, 1838
insularis Sharp, 1882
jugularis Babington, 1832
niger Sturm, 1835
revelierei Sharp, 1882
subelongatus Stephens, 1835

morio Aubé, 1838
atriceps Crotch, 1870
caliginosus LeConte, 1850
caminarius Motschulsky, 1860
lugubris Motschulsky, 1845
melancholicus Motschulsky, 1860
scaphiformis Sharp, 1871
valliger Hellen, 1929
watanabei Takizawa, 1933

neglectus Schaum, 1845
incrassatus Thomson, 1871

nigellus Mannerheim, 1853
bungei Zaitzev, 1910
mtiula Zaitzev, 1927
obscuripes Motschulsky, 1860
pyrenaeus Wehncke, 1872
sieversi J.R.Sahlberg, 1910
tungus Zaitzev, 1910

nigrita (Fabricius, 1792)
bisbiguttatus J.R.Sahlberg, 1921
convexior Seidlitz, 1887
glabellus Thomson, 1867
monilicornis J.R.Sahlberg, 1875
obovatus J.R.Sahlberg, 1880
subalpinus Thomson, 1871
trivialis Stephens, 1828

obscurus Sturm, 1835
hockingi Larson, 1975

palustris (Linnaeus, 1761)
apicalis Schilsky, 1888
buresi Maran, 1938
cambriensis Stephens, 1835
cristatomarginatus (Goeze, 1777)
fimbratus (Gmelin, 1790)
fuscorufus Munster, 1927
lituratus (Fabricius, 1781)
proximus Stephens, 1828
sexpustulatus (Fabricius, 1776)
styriacus Seidlitz, 1887
tinctus Clark, 1862
variegatus (Geoffroy, 1785)

planus (Fabricius, 1781)
corsicus Wehncke, 1872
frisius Verhoeff, 1891
fuscatus Stephens, 1828
fusculus (Schrink, 1781)
holosericeus (Marsham, 1802)
humeralis (Marsham, 1802)
lividus (Geoffroy, 1785)
nigricans (Schrink, 1781)

- nigriceps* Preller, 1862
pallidescens Seidlitz, 1887
sordidus (Herbst, 1784)
- pubescens*** (Gyllenhal, 1808)
antidotes Sharp, 1882
caliginosus Stephens, 1839
estrellensis Schaufuss, 1882
fulvipennis Munster, 1932
habelmanni Wehncke, 1876
roseni A.Zimmermann, 1919
scaphula Sturm, 1835
scopularis Schiödte, 1841
- rufifrons*** (O.F.Müller, 1776)
duftschmidti Rye, 1872
intermedius J.R.Sahlberg, 1880
piceus Stephens, 1828
rufifrons (Duftschmid, 1805)
- striola*** (Gyllenhal, 1827)
ambiguus Aubé, 1838
subtonsus LeConte, 1855
vittula Erichson, 1837
- tesselatus*** (Drapiez, 1819)
chrysostomus Des Gozis, 1914
nigricollis Desbrochers des Loges, 1871
pueli Barthe, 1916
xanthopus Stephens, 1835
- tristis*** (Paykull, 1798)
ampliceps J.R.Sahlberg, 1921
ruficapillus Mannerheim, 1852
varians LeConte, 1850
- umbrosus*** (Gyllenhal, 1808)
minutus Stephens, 1828
- Nebrioporus*** (*Nebrioporus* Régimbart, 1906)
assimilis (Paykull, 1798)
affinis (Sturm, 1835)
frater (Kunze, 1818)
hyperboreus (Gyllenhal, 1826)
- depressus*** (Fabricius, 1775)
boristhenicus (Hochhuth, 1871)
latescens (Falkenström, 1932)
neuhoffii (Cederjelm, 1798)
pallidus (Heer, 1837)
- Porhydrus*** Guignot, 1845
lineatus (Fabricius, 1775)
ovalis (Marsham, 1802)
parvulus (Herbst, 1784)
pygmaeus (Fabricius, 1792)
quadrilineatus (Drapiez, 1819)
velox (O.F.Müller, 1776)
- Scarodytes*** Des Gozis, 1914
halensis halensis (Fabricius, 1787)
amoenus (J.R.Sahlberg, 1908)
areolatus (Duftschmid, 1805 nec Grav., 1807)
areolatus (Gravenhorst, 1807 nec Duft., 1805)
ibericus (Régimbart, 1901)
nubilus (Gmelin, 1790)
reticulosus Falkenström, 1939
- Sphrodytes*** Des Gozis, 1914
dorsalis (Fabricius, 1787)
dubius (Melsheimer, 1844)
eliasi (Wirén, 1968)
figuratus (Gyllenhal, 1826)
maeklini (Zaitzev, 1910)
marginalis (Schilsky, 1889)
punctatus (O.F.Müller, 1776)
quadricolor (Gmelin, 1790)
rufifrons (Fabricius, 1792)
sibiricus (Mäklin, 1881)
- Hygrotini*** Portevin, 1929
Hygrotus (*Coelambus* Thomson, 1860)
impressopunctatus *impressopunctatus* (Schaller, 1783)
alternans (Gravenhorst, 1807)
decemlineatus (Mannerheim, 1853)
flavicans (Gmelin, 1790)
lineatus (O.F.Müller, 1776)
lineellus (Gyllenhal, 1808)
opus (Falkenström, 1932)
ovalis (Thunberg, 1784)
picipes (Fabricius, 1787)
punctatus (Marsham, 1802)
similis (Kirby, 1837)
- marklini*** (Gyllenhal, 1813)
astur (Sharp, 1882)
awajewi (A.I.Jakovlev, 1899)
canadensis (Fall, 1919)
lutzi (Reitter, 1908)
mistellus (Marseul, 1882)
mixtus (Motschulsky, 1860)
pallens (Aubé, 1838)
steppensis (Branden, 1885)
virgo (Fall, 1919)
- parallelogrammus*** (Ahrens, 1812)
consobrinus (Kunze, 1818)
kaszabi (V.B.Guéorguiev, 1970)
punctum (Gebler, 1829)
- polonicus*** (Aubé, 1842)
sahlbergi (Sharp, 1882)
unguicularis (J.R.Sahlberg, 1880)
- Hygrotus*** (*Hygrotus* Stephens, 1828)
decoratus (Gyllenhal, 1810)
unicolor Schilsky, 1888
- inaequalis*** *inaequalis* (Fabricius, 1776)
dalmatinus A.Zimmermann, 1915
minor (A.Costa, 1847)
parvulus (Fabricius, 1792)
uhagonii Seidlitz, 1887
- versicolor*** (Schaller, 1783)
affinis Stephens, 1828
collaris (Panzer, 1795)
confluens Munster, 1932
recurvus (Marsham, 1802)
reticulatus (Fabricius, 1792)
semilineatus A.Zimmermann, 1915

Hyhydrini Sharp, 1882	<i>aequalis</i> Thomson, 1868
Hyhydrus Illiger, 1802	<i>aeneus</i> (DeGeer, 1774)
<i>ovatus</i> (Linnaeus, 1761)	Note: Record needs confirmation.
<i>fabricii</i> Cristoforis, 1832	
<i>ferrugineus</i> (Linnaeus, 1767)	
<i>gibbus</i> (Fabricius, 1776)	
<i>grossus</i> (O.F.Müller, 1776)	
<i>ovalis</i> (Illiger, 1798)	
<i>sphaericus</i> (DeGeer, 1774)	
<i>variegatus</i> Stephens, 1828	
Laccornini Wolfe, Roughley, 1990	
Laccornis Des Gozis, 1914	
<i>oblongus</i> (Stephens, 1835)	
<i>bohemani</i> (Thomson, 1856)	
<i>nitidus</i> (Sturm, 1835)	
Laccophilinae Gistel, 1856	
Laccophilini Gistel, 1856	
Laccophilus Leach, 1815	
<i>hyalinus</i> <i>hyalinus</i> (DeGeer, 1774) nec Marsham, 1802)	
<i>cyprioticus</i> Mouchamps, 1956	
<i>hyalinus</i> (Marsham, 1802 nec DeGeer, 1774)	
<i>interruptus</i> (Panzer, 1795)	
<i>marmoratus</i> (Geoffroy, 1785)	
<i>marmoreus</i> (Olivier, 1795)	
<i>minutus</i> (Linnaeus, 1758)	
<i>cimicoides</i> (O.F.Müller, 1776)	
<i>gibbus</i> (Gmelin, 1790)	
<i>gilvus</i> (O.F.Müller, 1776)	
<i>melanophthalmos</i> (Geoffroy, 1785)	
<i>obscurus</i> (Panzer, 1795)	
<i>striatus</i> Depoli, 1930	
<i>variolosus</i> (Herbst, 1784)	
<i>virescens</i> (Brahm, 1791)	
<i>poecilus</i> Klug, 1834	
<i>latrix</i> Pic, 1896	
<i>parumpunctatus</i> O.Schneider, 1903	
<i>ponticus</i> Sharp, 1822	
<i>variegatus</i> (Germar, Kaulfuss, 1812)	
Polyphaga	
Staphyliniformia	
Hydrophiloidea	
Helophoridae Leach, 1815	
Helophorus (<i>Embleurus</i>) Hope, 1838)	
<i>nubilus</i> Fabricius, 1776	
<i>costatus</i> (Goeze, 1777)	
<i>linearis</i> Kuwert, 1886	
<i>lineellus</i> Kuwert, 1886	
<i>meridionalis</i> (Motschulsky, 1853)	
<i>mesopotamiae</i> Kuwert, 1886	
<i>striatus</i> (Geoffroy, 1785)	
Helophorus (<i>Kyphohelophorus</i>) Kuwert 1886)	
<i>tuberculatus</i> Gyllenhal, 1808	
<i>baicalicus</i> (Motschulsky, 1860)	
<i>scaber</i> LeConte, 1850	
Helophorus (<i>Helophorus</i>) Fabricius, 1775)	
	<i>aqualis</i> Thomson, 1868
	<i>aeneus</i> (DeGeer, 1774)
	<i>aquaticus</i> (Linnaeus, 1758)
	<i>frigidus</i> Graëlls, 1847
	<i>grandis</i> Illiger, 1798
	Helophorus (<i>Rhopalohelophorus</i>) Kuwert, 1886)
	<i>brevipalpis</i> Bedel, 1881
	<i>brevipalpis</i> d'Orchymont, 1927 nec Motschulsky, 1860
	<i>bulbipalpis</i> Kuwert, 1890
	<i>creticus</i> Kiesenwetter, 1858
	<i>insignis</i> Rey, 1885
	<i>mixtus</i> Rey, 1885
	<i>flavipes</i> Fabricius, 1792
	<i>arcuatus</i> Mulsant, 1844
	<i>balticus</i> Kuwert, 1886
	<i>monticola</i> Rey, 1885
	<i>phalleterus</i> Sharp, 1916
	<i>planicollis</i> Thomson, 1870
	<i>shetlandicus</i> Kuwert, 1890
	<i>sphagnicola</i> Hardy, 1871
	<i>viridicollis</i> Stephens, 1828
	<i>granularis</i> (Linnaeus, 1761)
	<i>brevicollis</i> Thomson, 1868
	<i>minimus</i> Kuwert, 1887 nec Fabricius, 1792
	<i>pygmaeus</i> Fabricius, 1792
	<i>griseus</i> Herbst, 1793
	<i>affinis</i> Sharp, 1916
	<i>minutus</i> Fabricius, 1775
	<i>nanus</i> Sturm, 1835
	<i>strigifrons</i> Thomson, 1868
	<i>championi</i> Sharp, 1915
	Hydrophilidae Latreille, 1802
	Hydrophilinae Latreille, 1802
	Berosini Mulsant, 1844
	Berosus (<i>Berosus</i>) Leach, 1817)
	<i>luridus</i> (Linnaeus, 1761)
	<i>chalcaspis</i> (Eschscholtz, 1818)
	<i>fuscus</i> (DeGeer, 1774)
	<i>signaticollis</i> (Charpentier, 1825)
	Chaetarthriini Bedel, 1881
	Chaetarthria Stephens, 1835
	<i>seminulum</i> (Herbst, 1797)
	<i>picea</i> Hochhuth, 1871
	Anacaenini Hansen, 1991
	Anacaena Thomson, 1859
	<i>globulus</i> (Paykull, 1798)
	<i>glabricollis</i> (Schaufuss, 1869)
	<i>limbata</i> (Fabricius, 1792)
	<i>carinata</i> Thomson, 1870
	<i>foveolata</i> (Haworth, 1807)
	<i>lutescens</i> (Stephens, 1829)
	<i>ambigua</i> (Rey, 1885)
	<i>immatura</i> Abeille de Perrin, 1901
	<i>nitida</i> (Heer, 1841)

- variabilis* Sharp, 1870
- Laccobiini** Bertrand, 1954
- Laccobius** Erichson, 1837
- bipunctatus* (Fabricius, 1775)
 - alutaceus* Thomson, 1868
 - emmeryanus* Rottenberg, 1874
- colon* (Stephens, 1829)
 - biguttatus* Gerhardt, 1877
 - mongolicus* Gentili, 1973
- minutus* (Linnaeus, 1758)
 - chrysomelinus* (O.F.Müller, 1776)
 - dermestoides* (Forster, 1771)
 - globosus* Heer, 1841
 - perla* (Geoffroy, 1785)
- striatulus* (Fabricius, 1801)
 - belgicus* Gentili, 1979
 - bucciarelli* Gentili, 1974
 - maculiceps* Gerhardt, 1877
 - nigriceps* Thomson, 1853
 - purpurascens* Newbery, 1908
- Hydrophilini** Latreille, 1802
- Helochares** Mulsant, 1844
- obscurus* (O.F.Müller, 1776)
 - erythrocephalus* (Fabricius, 1792)
 - subcompressus* Rey, 1885
 - variegatus* Herbst, 1797
- Enochrus** Thomson, 1859
- affinis* (Thunberg, 1794)
- bicolor* (Fabricius, 1792)
 - atricornis* (Kuwert, 1888)
 - chrysomelinus* (Fabricius, 1787)
 - ferrugineus* (Küster, 1849)
 - flavus* (Kuwert, 1888)
 - fulvus* (Marsham, 1802)
 - grisescens* (Gyllenhal, 1827)
 - levanderi* (J.R.Sahlberg, 1892)
 - maritimus* (Thomson, 1853)
 - mediterraneus* (J.R.Sahlberg, 1900)
 - sahlbergi* (Kuwert, 1888)
 - sternospina* (Kuwert, 1888)
 - unguidebilis* (Kuwert, 1888)
 - yukinoae* (Matsui, 1994)
- coarctatus* (Gredler, 1863)
 - suturalis* (Sharp, 1872)
- fuscipennis* Thomson, 1884
- melanocephalus* (Olivier, 1792)
 - atricapillus* (Stephens, 1829)
- ochropterus* (Marsham, 1802)
 - frontalis* (Erichson, 1837)
 - nigricans* (Zetterstedt, 1838)
- quadripunctatus* (Herbst, 1797)
 - minutus* (Paykull, 1798)
 - rectus* (J.R.Sahlberg, 1875)
 - ytenensis* (Sharp, 1915)
- testaceus* (Fabricius, 1801)
- Cymbiodyta** Bedel, 1881
- marginella* (Fabricius, 1792)
- carbonarius* (Kuwert, 1892)
- marginatus* (Dejean, 1833)
- ovalis* (Thomson, 1853)
- Hydrobius** Leach, 1815
- fuscipes* (Linnaeus, 1758)
 - aeneus* Dejean, 1833 nec Solier, 1833
 - aeneus* Solier, 1833 nec Dejean, 1833
 - angustatus* (Villa, Villa, 1838)
 - aquaticus* (Linnaeus, 1761)
 - arcadius* Brullé, 1832
 - chalconotus* Stephens, 1829
 - gyrinoides* (Schrank, 1781)
 - insculptus* LeConte, 1855
 - levigatus* Laporte de Castelnau, 1840
 - ovalis* Dejean, 1833
 - plicrus* Thomson, 1884
 - regularis* LeConte, 1855
 - rottenbergii* Gerhardt, 1872
 - scarabaeoides* (Linnaeus, 1758)
 - seriatus* LeConte, 1855
 - subrotundus* Stephens, 1829
 - substriatus* Curtis, 1829
- Hydrochara** Berthold, 1827
- caraboides* (Linnaeus, 1758)
 - nigricornis* (DeGeer, 1774)
 - scarabaeoides* (Schrank, 1781)
 - scrobiculatus* (Panzer, 1799)
 - smaragdinus* (Bach, 1856)
 - subaeneus* (Motschulsky, 1845)
 - substriatus* (Sturm, 1843)
- flavipes* (Steven, 1808)
 - profanifuga* Semenov, 1900
- Hydropilus** Geoffroy, 1762
- aterrimus* Eschscholtz, 1822
 - africanus* (Kuwert, 1893)
 - morio* (Cristoforis, Jan, 1832)
- piceus* (Linnaeus, 1758)
 - angustior* Rey, 1885
 - ruficornis* DeGeer, 1774
 - turkestanus* (Kuwert, 1893)
 - viridicollis* Redtenbacher, 1844
- Sphaeridiinae** Latreille, 1802
- Coelostomatini** Heyden, 1891
- Coelostoma** Brullé, 1835
- orbiculare* (Fabricius, 1775)
 - breve* (Sharp, 1874)
 - brevitarse* (Heyden, 1870)
 - dalmatinum* (Küster, 1848)
 - immaculatum* (Rossi, 1790)
 - minor* (Sharp, 1872)
 - pilula* (O.F.Müller, 1776)
 - punctatum* (Goeze, 1777)
- Megasternini** Mulsant, 1844
- Cercyon** (*Paracercyon*) Seidlitz, 1888
- analis* (Paykull, 1798)
 - acutum* Stephens, 1829

<i>apicale</i> Stephens, 1829	<i>nubilipenne</i> Stephens, 1835
<i>aquaticum</i> Stephens, 1829	<i>pulchellus</i> Heer, 1841
<i>falli</i> Winters, 1944	<i>testaceus</i> Stephens, 1829 in part
<i>maculatum</i> Melsheimer, 1844	<i>troglodytes</i> Dejean, 1836
Cercyon (Paracycreon d'Orchymont, 1942)	<i>ustulatum</i> Stephens, 1829
<i>laminatus</i> Sharp, 1873	<i>vicinale</i> Walker, 1859
<i>sharpi</i> Harold, 1878	
Cercyon (Dicyrtocercyon Ganglbauer, 1904)	obsoletus (Gyllenhal, 1808)
<i>ustulatus</i> (Preyssler, 1790)	<i>lugubris</i> auct. nec (Olivier, 1790)
<i>haemorrhoidalis</i> (Fabricius, 1792) nec	
Fabricius, 1775)	pygmaeus (Illiger, 1801)
<i>haemorrhoum</i> (Gyllenhal, 1808)	<i>apicalis</i> (Say, 1823)
<i>xanthorhoeum</i> Stephens, 1829	<i>conspurcatus</i> (Sturm, 1807)
Cercyon (Cercyon Leach, 1817)	<i>erythropus</i> Stephens, 1829
<i>bifenestratus</i> Kuster, 1851	<i>ferrugineus</i> (Herbst, 1792)
<i>palustre</i> Thomson, 1853	<i>fuscescens</i> Stephens, 1829
<i>convexusculus</i> Stephens, 1829	<i>merdarius</i> (Sturm, 1807)
<i>alni</i> Vogt, 1969	<i>stercorator</i> Stephens, 1829
<i>granarius</i> Erichson, 1837	quisquilius (Linnaeus, 1761)
<i>haemorrhoidalis</i> (Fabricius, 1775 nec F., 1792)	<i>dispar</i> (Paykull, 1798)
<i>aquatalis</i> , Donisthorpe, 1932	<i>flavipennis</i> Küster, 1848
<i>basilaris</i> Notman, 1920	<i>flavus</i> (Marsham, 1802)
<i>erythropterus</i> Mulsant, 1844	<i>minimus</i> (Scopoli, 1763)
<i>femoralis</i> Stephens, 1829	<i>xanthopterum</i> (Laicharting, 1781)
<i>flavipes</i> (Fabricius, 1792)	sternalis (Sharp, 1918)
<i>glabratus</i> (Melsheimer, 1806)	<i>pumilio</i> (Sharp, 1918)
<i>infuscatus</i> Stephens, 1829	terminatus (Marsham, 1802)
<i>nigricollis</i> (Say, 1825)	<i>fimetarium</i> Wollaston, 1854
<i>obscurus</i> (Fabricius, 1792)	<i>lepidum</i> Wollaston, 1864
<i>picinus</i> (Marsham, 1802)	<i>nanum</i> Melsheimer, 1844
<i>punctulatus</i> Dejean, 1836	<i>plagiatum</i> Erichson, 1837
<i>similis</i> (Marsham, 1802)	<i>scutellaris</i> Stephens, 1829
<i>striatus</i> (Turton, 1802)	tristis (Illiger, 1801)
<i>suturalis</i> Stephens, 1829	<i>boletophagus</i> (Marsham, 1802)
<i>impressus</i> (Sturm, 1807)	<i>convexior</i> Stephens, 1829 in part
<i>piceus</i> (Marsham, 1802)	<i>convexus</i> Stephens, 1829
<i>xanthocephalum</i> Stephens, 1829	<i>laevigatus</i> Stephens, 1829
<i>lateralis</i> (Marsham, 1802)	unipunctatus (Linnaeus, 1758)
<i>icrematus</i> Notman, 1920	<i>cordiger</i> (Herbst, 1784)
<i>rufus</i> (Sturm, 1826)	<i>dispar</i> (Paykull, 1798)
<i>littoralis</i> (Gyllenhal, 1808)	<i>janssoni</i> Nyholm, 1952
<i>binotatus</i> Stephens, 1829	Megasternum Mulsant, 1844
<i>dilatatus</i> Stephens, 1829	concinnum (Marsham, 1802)
<i>ruficornis</i> Stephens, 1829	<i>acutum</i> (Stephens, 1829)
<i>marinus</i> Thomson, 1853	<i>aquaticum</i> (Stephens, 1829)
<i>aquaticus</i> Dejean, 1833	<i>boletophagum</i> (Marsham, 1802) nec Marsham,
<i>melanocephalus</i> (Linnaeus, 1758)	1802
<i>nigripes</i> (Marsham, 1802)	<i>calabricum</i> Kuwert, 1890
<i>atricapillus</i> (Marsham, 1802)	<i>castaneum</i> (Heer, 1841)
<i>atriceps</i> Stephens, 1829	<i>caucasicum</i> Kuwert, 1890
<i>atriceps</i> Gemminger, Harold, 1868	<i>contaminatum</i> (Stephens, 1829)
<i>bimaculatus</i> Stephens, 1829	<i>convexior</i> (Stephens, 1829)
<i>centrimaculatus</i> (Sturm, 1807)	<i>immune</i> (Stephens, 1829)
<i>inustus</i> (Marsham, 1802)	<i>obscurum</i> (Marsham, 1802)
<i>laevis</i> (Marsham, 1802)	<i>pusillum</i> (Kuwert, 1890)
<i>mundus</i> Melsheimer, 1844	<i>stercorarium</i> (Marsham, 1802)
<i>nigriceps</i> Motschulsky, 1863	<i>testaceum</i> (Stephens, 1829)
	Cryptopleurum Mulsant, 1844
	crenatum (Kugelann, 1794)
	<i>vaucherii</i> Tournier, 1867

- minutum* (Fabricius, 1775)
sordidus (Marsham, 1802)
- Sphaeridiini** Latreille, 1802
- Sphaeridium** Fabricius, 1775
- bipustulatum* Fabricius, 1781
haemorrhous (Schrank, 1781)
testudinarius (Geoffroy, 1785)
- lunatum* Fabricius, 1792
- scarabaeoides* (Linnaeus, 1758)
crenatum Palisot de Beauvois, 1817
pictum Ménétriés, 1832
quadrimaculatum (Schrank, 1781)
quadrimaculatum Küster, 1845
striolatum Heer, 1841
testudinarium (DeGeer, 1774)
- Hydrochidae** Thomson, 1859
- Hydrochus** Leach, 1817
- angustatus* Germar, 1824
- brevis* (Herbst, 1793)
- carinatus* Germar, 1824
- elongatus* (Schaller, 1783)
- ignicollis* Motschulsky, 1860
elongatus auct. nec (Schaller, 1783)
- Spercheidae** Erichson, 1837
- Spercheus** Illiger, 1798
- emarginatus* (Schaller, 1783)
luridus Mathieu, 1858
sordidus (Marsham, 1802)
texanus Spangler, 1961
verrucosus (Marsham, 1802)
- Georissidae** Laporte de Castelnau, 1840
- Georissus** (*Georissus*) Latreille, 1809
- crenulatus* (Rossi, 1794)
- Histeroidea**
- Sphaeritidae** Shuckard, 1839
- Sphaerites** Duftschmid, 1805
- glabratus* (Fabricius, 1792)
politus Ohara, 1994 nec Mannerheim, 1846
- Histeridae** Gyllenhal, 1808
- Histerinae** Gyllenhal, 1808
- Holeoptini** Hope, 1840
- Holepta** Paykull, 1811
- plana* (Sulzer, 1776)
- Platysomatini** Bickhardt, 1914
- Platysoma** Leach, 1817
- compressum* (Herbst, 1783)
- deplanatum* (Gyllenhal, 1808)
- elongatum* (Thunberg, 1787)
oblongum Fabricius, 1792
- Cylistix** Marseul, 1857
- angustata* (Hoffmann, 1803)
ferrugineum (Thunberg, 1794 nec Olivier, 1789)
- lineare* (Erichson, 1834)
- Eblisia** Lewis, 1889
- minor* (Rossi, 1792)
frontale (Paykull, 1798)
- Histerini** Gyllenhal, 1808
- Margarinotus** Marseul, 1853
- bipustulatus* (Schrank, 1781)
- brunneus* (Fabricius, 1775)
cadaverinus (Hoffmann, 1803)
impressus (Fabricius, 1798)
- carbonarius* Hoffmann, 1803
- marginatus* (Erichson, 1834)
- merdarius* (Hoffmann, 1803)
- neglectus* (Germar, 1813)
- obscurus* (Kugelann, 1792)
stercorarius (Hoffmann, 1803)
- purpurascens* (Herbst, 1792)
- striola* (C.Sahlberg, 1819)
succicola (Thomson, 1862)
- terricola* (Germar, 1824)
- ventralis* (Marseul, 1854)
- Hister** Linnaeus, 1758
- bissexstriatus* Fabricius, 1801
- funestus* Erichson, 1834
- helluo* Truqui, 1852
- illigeri* Duftschmid, 1805
- quadrimaculatus* Linnaeus, 1758
- quadrinotatus* L.G.Scriba, 1790
- unicolor* Linnaeus, 1758
- Atholus** Thomson, 1859
- bimaculatus* (Linnaeus, 1758)
- corvinus* (Germar, 1817)
- duodecimstriatus* (Schrank, 1781)
- Hetaeriinae** Marseul, 1857
- Hetaerius** Erichson, 1834
- ferrugineus* (Olivier, 1789)
- Dendrophilinae** Reitter, 1909
- Dendrophilini** Reitter, 1909
- Dendrophilus** Leach, 1817
- corticalis* (Paykull, 1798)
punctatus (Herbst, 1792 nec O.F.Müller, 1776)
- pygmaeus* (Linnaeus, 1758)
- Paromalini** Reitter, 1909
- Carcinops** Marseul, 1855
- pumilio* (Erichson, 1834)
quatuordecimstriatus (Stephens, 1835)
- Platyomalus** Cooman, 1948
- complanatus* (Panzer, 1797)
- Paromalus** Erichson, 1834
- flavicornis* (Herbst, 1792)
- parallelepipedus* (Herbst, 1792)
- Abraeinae** McLeay, 1819

Abraeini Marseul, 1857	<i>rugifrons</i> (Paykull, 1798)
Abraeus Leach, 1817	Staphylinoidea
<i>perpusillus</i> (Marsham, 1802)	Hydraenidae Mulsant, 1844
<i>globosus</i> (Hoffmann, 1803)	Ochthebiinae Thomson, 1859
Plegaderini Portevin, 1929	Ochthebius Leach, 1815
Plegaderus Erichson, 1834	<i>marinus</i> (Paykull, 1798)
<i>caesus</i> (Herbst, 1792)	<i>minimus</i> (Fabricius, 1792)
<i>saucius</i> Erichson, 1834	<i>impressus</i> (Marsham, 1802)
<i>vulneratus</i> (Panzer, 1797)	<i>remotus</i> Reitter, 1887
Acritini Wenzel, 1944	Note: This species is known by single record from Murjāni area (Central Latvia) only. North border of main distribution area.
Acritus LeConte, 1853	Hydraeninae Mulsant, 1844
<i>homoeopathicus</i> Wollaston, 1857	Hydraena Kugelann, 1794
<i>minutus</i> (Herbst, 1792)	<i>excisa</i> Kiesenwetter, 1849
<i>nigricornis</i> (Hoffmann, 1803)	<i>gracilis</i> Germar, 1824
Note: Insufficiently known species, registered from only a few localities.	<i>palustris</i> Erichson, 1837
Teretriini Bickhardt, 1916	<i>pulchella</i> Germar, 1824
Teretrius Erichson, 1834	<i>flavipes</i> Sturm 1836
<i>fabricii</i> Mazur, 1972	<i>reyi</i> Kuwert, 1888
<i>picipes</i> (Fabricius, 1792 nec Olivier, 1789)	<i>riparia</i> Kugelann, 1794
Saprininae Blanchard, 1845	Limnebius Leach, 1815
Gnathoncus Jacquelin du Val, 1858	<i>crinifer</i> Rey, 1885
<i>buyssoni</i> Auzat, 1917	<i>papposus</i> Mulsant, 1844
<i>nannetensis</i> Marseul, 1868	<i>parvulus</i> (Herbst, 1797)
<i>rotundatus</i> (Kugelann, 1792)	<i>truncatulus</i> Thomson, 1851
<i>nanus</i> (L.G.Scriba, 1790 nec Piller, Mtpp., 1783)	<i>truncatellus</i> (Thunberg, 1794)
<i>punctulatus</i> Thomson, 1862	Ptiliidae Erichson, 1845 / Motschulsky, 1845
Myrmetes Marseul, 1862	Ptenidium Erichson, 1845
<i>paykulli</i> Kanaar, 1979	<i>formicetorum</i> Kraatz, 1851
<i>piceus</i> (Paykull, 1809 nec Marsham, 1802)	<i>myrmecophilum</i> (Motschulsky, 1845 nec Allibert, 1844)
Saprinus Erichson, 1834	<i>gressneri</i> Erichson, 1845
<i>aeneus</i> (Fabricius, 1775)	<i>intermedium</i> Wankowicz, 1869
<i>immundus</i> (Gyllenhal, 1827)	<i>laevigatum</i> Erichson, 1845
<i>planiusculus</i> Motschulsky, 1849	<i>longicorne</i> Fuss, 1848
<i>cuspidatus</i> Ihssen, 1949	<i>brisoutii</i> Matthews, 1872
<i>rugifer</i> (Paykull, 1809)	<i>nitidum</i> (Heer, 1841)
<i>quadristriatus</i> auct. nec (Thunberg, 1794)	<i>pusillum</i> (Gyllenhal, 1808)
<i>semistriatus</i> (L.G.Scriba, 1790)	<i>apicale</i> Erichson, 1845
<i>subnitescens</i> Bickhardt, 1909	<i>corpulentum</i> Lucas, 1846
<i>tenuistrius sparustus</i> Solsky, 1875	<i>turgidum</i> Thomson, 1855
<i>virescens</i> (Paykull, 1798)	Actidium Matthews, 1868
Chalcionellus Reichardt, 1932	<i>boudieri</i> (Allibert, 1844)
<i>decemstriatus</i> (Rossi, 1792)	<i>transversale</i> (Erichson, 1845)
<i>conjugens</i> (Paykull, 1798)	Oligella Motschulsky, 1869
Note: Very rare and insufficiently known species.	<i>foveolata</i> (Allibert, 1844)
Hypocacculus Bickhardt, 1916	<i>clandestina</i> (Haliday, 1855)
<i>rufipes</i> (Kugelann, 1792)	<i>excavata</i> (Erichson, 1845)
Hypococcus Thomson, 1867	Micridium Motschulsky, 1869
<i>metallicus</i> (Herbst, 1792)	<i>halidaii</i> (Matthews, 1868)
<i>rugiceps</i> (Duftschmid, 1805)	Millidium Motschulsky, 1855

<i>minutissimum</i> (Ljungh, 1804)	Sogdini Lopatin, 1961
<i>trisulcatum</i> (Aubé, 1833)	
Ptilium Gyllenhal, 1827	Hydnobius Schmidt, 1841
<i>exaratum</i> (Allibert, 1844)	<i>punctatus</i> (W.Sturm, 1807)
<i>canaliculatum</i> Erichson, 1845	<i>punctatissimus</i> (Stephens, 1829)
<i>myrmecophilum</i> (Allibert, 1844)	
<i>haemorrhoidale</i> Boheman, 1844	Leiodini Fleming, 1821
<i>inquilinum</i> Erichson, 1845	Leiodes Latreille, 1796
Euryptilium Matthews, 1872	<i>ciliaris</i> (W.Schmidt, 1841)
<i>gillmeisteri</i> Flach, 1889	<i>dubia</i> (Kugelann, 1794)
Ptiliola Haldeman, 1849	<i>ferruginea</i> (Fabricius, 1787)
<i>brevicolle</i> (Matthews, 1860)	<i>ovalis</i> (W.Schmidt, 1841)
<i>kunzei</i> (Heer, 1841)	<i>fracta</i> (Seidlitz, 1874)
<i>depressa</i> (Motschulsky, 1845)	<i>rahaetica</i> auct. nec (Erichson, 1845)
Ptiliolum Flach, 1888	<i>gyllenhali</i> Stephens, 1829
<i>sahlbergi</i> (Flach, 1888)	<i>parvula</i> (R.F.Sahlberg, 1833)
Pteryx Matthews, 1858	<i>picea</i> (Panzer, 1797)
<i>suturalis</i> (Heer, 1841)	<i>polita</i> (Marsham, 1802)
<i>flavicornis</i> (Mäklin, 1846)	<i>calcarata</i> (Erichson, 1845)
<i>mutabilis</i> Matthews, 1858	<i>silesiaca</i> (Kraatz, 1852)
Nephanes Thomson, 1859	<i>triepkii</i> (W.Schmidt, 1841)
<i>titan</i> (Newman, 1834)	Liocyrtusa Daffner, 1982
<i>abbreviatella</i> (Heer, 1841)	<i>minuta</i> (Ahrens, 1812)
<i>curta</i> (Allibert, 1844)	Cyrtusa Erichson, 1842
Smicrus Matthews, 1872	<i>subtestacea</i> (Gyllenhal, 1813)
<i>filicornis</i> (Fairmaire, Laboulbène, 1855)	Colenis Erichson, 1842
Baeocrara Thomson, 1859	<i>immunda</i> (Sturm, 1807)
<i>japonica</i> (Matthews, 1884)	Agaricophagus W.Schmidt, 1841
Acrotrichis Motschulsky, 1848	<i>cephalotes</i> W.Schmidt, 1841
<i>atomaria</i> (DeGeer, 1774)	Agathidiini Westwood, 1838
<i>flavicornis</i> (Waltl, 1838)	Anisotoma Panzer, 1797
<i>hellenica</i> (Matthews, 1889)	<i>axillaris</i> Gyllenhal, 1810
<i>brevipennis</i> (Erichson, 1845)	<i>castanea</i> (Herbst, 1792)
<i>dispar</i> (Matthews, 1865)	<i>glabra</i> (Kugelann, 1794)
<i>nana</i> A.Strand, 1946	<i>humeralis</i> (Fabricius, 1792)
<i>fascicularis</i> (Herbst, 1793)	<i>orbicularis</i> (Herbst, 1792)
<i>grandicollis</i> (Mannerheim, 1844)	Liodopria Reitter, 1909
<i>intermedia</i> (Gillmeister, 1845)	<i>serricornis</i> (Gyllenhal, 1813)
<i>carpathica</i> Fleischer, 1923	Amphicyllus Erichson, 1845
<i>montandoni</i> (Allibert, 1844)	<i>globiformis</i> (R.F.Sahlberg, 1833)
<i>longicornis</i> (Mannerheim, 1844)	<i>globus</i> (Fabricius, 1792)
<i>parva</i> Rosskothen, 1935	Agathidium Panzer, 1797
<i>sericans</i> (Heer, 1841)	<i>atrum</i> (Paykull, 1798)
<i>bovina</i> (Motschulsky, 1845)	<i>badium</i> Erichson, 1845
<i>depressa</i> (Gillmeister, 1845)	<i>confusum</i> Brisout de Barneville, 1863
<i>picicornis</i> sensu Heer, 1841 nec (Mannh., 1843)	<i>convexum</i> Sharp, 1866
<i>sitkaensis</i> (Motschulsky, 1845)	<i>piceum</i> Erichson, 1845 nec Melsheimer, 1844
<i>fratercula</i> sensu Motschulsky, 1845 nec	<i>haemorrhoum</i> Erichson, 1845
<i>(Matthews, 1878)</i>	<i>laevigatum</i> Erichson, 1845
<i>strandi</i> Sundt, 1958	<i>marginatum</i> Sturm, 1807
<i>thoracica</i> (Waltl, 1838)	<i>nigripenne</i> (Fabricius, 1792)
Leiodidae Fleming, 1821	<i>seminulum</i> (Linnaeus, 1758)
Leiodinae Fleming, 1821	<i>plagiatum</i> (Gyllenhal, 1910)

<i>pulchellum</i> Wankowicz, 1869	<i>umbrina</i> (Erichson, 1837)
<i>rotundatum</i> (Gyllenhal, 1827)	Sciodrepoides Hatch, 1933
<i>sphaerulum</i> Reitter, 1898	<i>fumatus</i> (Spence, 1815)
<i>seminulum</i> (Linnaeus, 1758)	<i>ambiguus</i> (Heer, 1838)
<i>varians</i> Beck, 1817	<i>scitulus</i> (Erichson, 1837)
Colonidae Horn, 1800	<i>umbrinus</i> (Thomson, 1862)
Colon Herbst, 1797	<i>watsoni</i> (Spence, 1815)
<i>bidentatum</i> (C.R.Sahlberg, 1822)	<i>agilis</i> (Fabricius, 1801)
<i>rufescens</i> Kraatz, 1850	<i>curticornis</i> (Fairmaire, 1877)
<i>serripes</i> (C.R.Sahlberg, 1822)	<i>fumatus</i> (Erichson, 1837)
<i>viennense</i> (Herbst, 1797)	Catops Paykull, 1798
Cholevidae Kirby, 1837	<i>fuscus</i> (Panzer, 1794)
Ptomaphaginae Jeannel, 1911	<i>festinans</i> Spence, 1815
Ptomaphagus Illiger, 1798	<i>rufescens</i> Fabricius, 1801
<i>subvilllosus</i> (Goeze, 1777)	<i>sericeus</i> Paykull, 1798
<i>carpaticus</i> Jeannel, 1934	grandicollis Erichson, 1837
<i>nigriclavus</i> Guillebeau, 1891	<i>morio</i> (Fabricius, 1792)
<i>rufiventris</i> Pic, 1913	<i>dissimulator</i> Spence, 1815
<i>sericeus</i> Panzer, 1801	<i>fornicatus</i> J.R.Sahlberg, 1817
<i>silphoides</i> Marsham, 1802	<i>lapponicus</i> J.R.Sahlberg, 1889
<i>tarbensis</i> Reitter, 1884	<i>laticollis</i> J.R.Sahlberg, 1889
<i>truncatus</i> Illiger, 1802	<i>sericeus</i> Gyllenhal, 1827
<i>varicornis</i> (Rosenhauer, 1847)	<i>substriatus</i> Reitter, 1884
<i>ruthenus</i> Reitter, 1890	<i>sunicus</i> Iablokoff-Khnzorian, 1963
Nemadinae Jeannel, 1936	nigricans (Spence, 1815)
Nemadus Thomson, 1867	<i>flavicornis</i> Thomson, 1867
<i>colonoides</i> (Kraatz, 1851)	<i>fornicatus</i> Stephens, 1830
Cholevinae Kirby, 1837	<i>hervei</i> Guillebeau, 1891
Cholevini Kirby, 1837	<i>longipennis</i> Chaudoir, 1845
Nargus Thomson, 1867	<i>major</i> Kraatz, 1852
<i>wilkini</i> (Spence, 1815)	<i>marginicollis</i> Reitter, 1884
<i>flavescens</i> (Dejean, 1833)	<i>suburratus</i> Des Gozis, 1886
<i>istrianus</i> Depoli, 1915	<i>nigriclavus</i> Gerhardt, 1900
<i>praecox</i> (Erichson, 1837)	<i>dorni</i> Reitter, 1913
Choleva Latreille, 1796	tristis (Panzer, 1793)
<i>agilis</i> (Illiger, 1798)	<i>abdominalis</i> Rosenhauer, 1847
<i>adusta</i> Reitter, 1896	<i>dichrous</i> Munster, 1912
<i>gausapata</i> Stephens, 1830	<i>leachi</i> Spence, 1815
<i>gomphosata</i> Stephens, 1829	<i>montivagus</i> Heer, 1838
<i>jailensis</i> Jeannel, 1923	<i>nigrita</i> Sturm, 1839
<i>testacea</i> Latreille, 1897	<i>spencei</i> Stephens, 1830
<i>angustata</i> (Fabricius, 1781)	<i>univestis</i> Guillebeau, 1891
<i>cisteloides</i> Britten, 1918	<i>ventricosus</i> Reitter, 1885
<i>pyraenaica</i> Jeannel, 1922	Apocatops Zwick, 1968
<i>vaulogeri</i> Pic, 1922	<i>nigritus</i> (Erichson, 1837)
<i>elongata</i> (Paykull, 1798)	<i>affinis</i> (Stephens, 1830)
<i>lucidicollis</i> Reitter, 1887	<i>caliginosus</i> (Stephens, 1830)
<i>rufescens</i> Illiger, 1798	<i>marshami</i> (Spence, 1815)
<i>nivalis</i> (Kraatz, 1856)	<i>morio</i> (Paykull, 1798)
<i>breiti</i> Jeannel, 1923	<i>soror</i> (Newman, 1933)
<i>oblonga</i> Latreille, 1807	Fissocatops Zwick, 1968
<i>intermedia</i> Kraatz, 1852	<i>westi</i> (Krogerus, 1931)
Catopini Chaudoir, 1845	Platypyllidae Ritsema, 1869
Dreposcia Jeannel, 1922	Platypyllinae Ritsema, 1869
	Platypyllus Ritsema, 1869
	<i>castoris</i> Ritsema, 1869

- Scydmaenidae** Leach, 1815
Scydmaeninae Leach, 1815
Eutheiini Casey, 1897
Eutheia Stephens, 1830
linearis Mulsant, 1861
scydmaenoides Stephens, 1830
Cephenniini Reitter, 1882
Cephennium Müller, Kunze, 1822
laticolle Aubé, 1842
majus Reitter, 1882
thoracicum Müller, Kunze, 1822
Cyrtoscydmini Schaufuss, 1889
Nevraphes Thomson, 1859
angulatus (Müller, Kunze, 1822)
elongatus (Müller, Kunze, 1822)
Scydmoraphes Reitter, 1891
helvolus (Schaum, 1844)
sparshalli (Denny, 1825)
Stenichnus Thomson, 1859
collaris (Müller, Kunze, 1822)
foveola Rey, 1888
godarti (Latreille, 1806)
scutellaris (Müller, Kunze, 1822)
Microscydmus Saulcy, Croissandieu, 1893
minimus (Chaudoir, 1845)
Euconnus Thomson, 1859
claviger (Müller, Kunze, 1822)
denticornis (Müller, Kunze, 1822)
fimetarius (Chaudoir, 1845)
hirticollis (Illiger, 1798)
maklinii (Mannerheim, 1844)
wetterhallii (Gyllenhal, 1813)
Scydmaenini Leach, 1815
Scydmaenus Latreille, 1802
hellwigii (Herbst, 1792)
rufus Müller, Kunze, 1822
tarsatus Müller, Kunze, 1822
Agyrtidae Thomson, 1859
Pterolomatinae Thomson, 1862 sensu Perkovskij, 2000
Pterolomatini Thomson, 1862
Pteroloma Gyllenhal, 1827
-forsstroemii (Gyllenhal, 1810)
brunnea (Fischer von Waldheim, 1828)
Silphidae Latreille, 1807
Nicrophorinae Kirby, 1837
Nicrophorini Kirby, 1837
Nicrophorus Fabricius, 1775
interruptus Stephens, 1830
fossor Erichson, 1837
- germanicus* (Linnaeus, 1758)
Note: This very rare species is only known from a few localities in western Latvia. North border of main distribution area.
humator (Gleditsch, 1767)
investigator Zetterstedt, 1824
sepultor Charpentier, 1825
vespillo (Linnaeus, 1758)
vespilloides Herbst, 1783
Silphinae Latreille, 1807
Necrodes Leach, 1815
littoralis (Linnaeus, 1758)
Thanatophilus Leach, 1815
dispar (Herbst, 1793)
rugosus (Linnaeus, 1758)
sinuatus (Fabricius, 1775)
Oiceoptoma Leach, 1815
thoracica (Linnaeus, 1758)
Aclypea Reitter, 1884
opaca (Linnaeus, 1758)
undata (O.F.Müller, 1776)
Dendroxena Motschulsky, 1858
quadrimaculata (Scopoli, 1772)
quadripunctata (Linnaeus, 1761 nec L., 1758)
Note: Officially protected species in Latvia.
Silpha Linnaeus, 1758
carinata Herbst, 1783
Note: Insufficiently known species, probably rare, recorded from only a few localities.
obscura Linnaeus, 1758
tristis Illiger, 1798
Phosphuga Leach, 1817
atrata (Linnaeus, 1758)
Staphylinidae Latreille, 1802
Staphylininae Latreille, 1802
Staphylinini Latreille, 1802
Erichsonius Fauvel, 1874
cinerascens (Gravenhorst, 1802)
Gabrius Stephens, 1833
appendiculatus Sharp, 1910
subnigritulus Reitter, 1909
breviventer Sperk, 1835
coxalus (Hochhuth, 1871)
hublei Coiffait, Seegers, 1985
pennatus Sharp, 1910
nigritulus (Gravenhorst, 1802)
osseticus (Kolenati, 1846)
vernalis (Gravenhorst, 1806)
splendidulus (Gravenhorst, 1802)
trossulus (Nordmann, 1837)
velox Sharp, 1910

Rabigus Mulsant, Rey, 1876
tenuis (Fabricius, 1792)
Bisnius Leach, 1829
cephalotes (Gravenhorst, 1802)
fimetarius (Gravenhorst, 1802)
nigriventris (Thomson, 1867)
puella (Nordmann, 1837)
sordidus (Gravenhorst, 1802)
subuliformis (Gravenhorst, 1802)
 fuscus (Gravenhorst, 1802)
Philonthus Curtis, 1829
addendus Sharp, 1867
 — *aerosus* Kiesenwetter, 1851
albipes (Gravenhorst, 1802)
alpinus Eppelsheim, 1875
atratus (Gravenhorst, 1802)
carbonarius (Gravenhorst, 1810)
 varius (Gyllenhal, 1810)
coerulescens Boisduval, Lacordaire, 1835
cognatus (Stephens, 1832)
 fuscipennis (Mannerheim, 1830)
concinnus (Gravenhorst, 1802)
coprophilus (Jarrige, 1949)
corvinus Erichson, 1839
cruentatus (Gmelin, 1790)
cyanipennis (Fabricius, 1792)
debilis (Gravenhorst, 1802)
decorus (Gravenhorst, 1802)
discoideus (Gravenhorst, 1802)
ebeninus (Gravenhorst, 1802)
fumarius (Gravenhorst, 1806)
intermedius (Boisduval, Lacordaire, 1835)
laevicollis (Boisduval, Lacordaire, 1835)
laminatus (Creutzer, 1799)
lepidus (Gravenhorst, 1802)
longicornis (Stephens, 1832)
mannerheimi Fauvel, 1869
marginatus (Ström, 1768)
micans (Gravenhorst, 1802)
micantoides (G.Benick, Lohse, 1956)
montivagus Heer, 1838
Note: Record needs confirmation.
nigrita (Gravenhorst, 1806)
nitidulus (Gravenhorst, 1802)
nitidus (Fabricius, 1787)
parvicornis (Gravenhorst, 1802)
 agilis (Gravenhorst, 1806)
politus (Linnaeus, 1758)
punctus (Gravenhorst, 1802)
quisquiliarius (Gyllenhal, 1810)
 puthzi (Korge, 1964)

rectangulus Sharp, 1874
rotundicollis (Ménétriés, 1832)
rubripennis (Stephens, 1832)
 fulvipes (Fabricius, 1793)
rufipes (Stephens, 1832)
 immundus (Gyllenhal, 1810)
sanguinolentus (Gravenhorst, 1802)
splendens (Fabricius, 1792)
succicola Thomson, 1860
temporalis Mulsant, Rey, 1853
tenuicornis Mulsant, Rey, 1853
 carbonarius (Gravenhorst, 1810)
umbratilis (Gravenhorst, 1802)
varians (Paykull, 1789)
ventralis (Gravenhorst, 1802)
 — *virgo* Gravenhorst, 1802
Neobisnius Ganglbauer, 1895
 — *procerulus* (Gravenhorst, 1802)
villosulus (Stephens, 1833)
Remus Holme, 1837
sericeus Holme, 1837
Dinothenarus Thomson, 1858
pubescens (DeGeer, 1774)
Ontholestes Ganglbauer, 1895
haroldi Eppelsheim, 1882
murinus (Linnaeus, 1758)
 dieckmanni Smetana, 1958
tesselatus (Geoffroy, 1785)
Emus Samouelle, 1819
hirtus (Linnaeus, 1758)
Platydracus Thomson, 1858
chalcocephalus (Fabricius, 1801)
fulvipes (Scopoli, 1763)
latebricola (Gravenhorst, 1806)
stercorarius (Olivier, 1795)
Staphylinus Linnaeus, 1758
caesareus Cederhjelm, 1798
dimidiaticornis Gemminger, 1851
erythropterus Linnaeus, 1758
Ocyphus Samouelle, 1819
aeneocephalus (DeGeer, 1774)
brunnipes (Fabricius, 1781)
fulvipennis Erichson, 1840
fuscatus (Gravenhorst, 1802)
nitens (Schrank, 1781)
 nero (Faldermann, 1835)
ophthalmicus ophthalmicus (Scopoli, 1763)
picipennis picipennis (Fabricius, 1792)
 — *sericeicollis* Ménétriés, 1832
 cupreus (Rossi, 1790)
Tasgius Stephens, 1829

ater (Gravenhorst, 1802)
globulifer (Geoffroy, 1785)
melanarius Heer, 1839
pedator (Gravenhorst, 1802)
Creophilus Samouelle, 1819
maxillosus (Linnaeus, 1758)
Heterothops Stephens, 1829
dissimilis (Gravenhorst, 1802)
praevius Erichson, 1840
quadripunctulus (Gravenhorst, 1806)
Euryporus Erichson, 1839
picipes (Paykull, 1800)
Velleius Samouelle, 1819
dilatatus (Fabricius, 1787)
 Note: Officially protected species in Latvia;
 also included on the list of indicator species of
 natural forest habitats.
Quedius Stephens, 1829
boops (Gravenhorst, 1802)
brevicornis Thomson, 1860
brevis Erichson, 1840
cinctus (Paykull, 1790)
cruentus (Olivier, 1795)
curtipennis Bernhauer, 1908
fulgidus (Fabricius, 1793)
 assimilis (Nordmann, 1837)
fuliginosus (Gravenhorst, 1802)
levicollis (Brullé, 1832)
 tristis (Gravenhorst, 1802)
limbatus (Heer, 1839)
 limbatoides Coiffait, 1963
longicornis Kraatz, 1857
lucidulus Erichson, 1840
maurus (C.R.Sahlberg, 1830)
mesomelinus mesomelinus (Marsham, 1802)
microps (Gravenhorst, 1847)
molochinus (Gravenhorst, 1806)
nigriceps Kraatz, 1857
nitipennis (Stephens, 1833)
ochripennis (Ménétriés, 1832)
paradisianus (Heer, 1839)
 Note: Record needs confirmation.
picipes (Mannerheim, 1830)
plagiatus (Mannerheim, 1843)
plancus Erichson, 1840
riparius Kellner, 1843
scitus (Gravenhorst, 1806)
umbrinus Erichson, 1839
 pseudoumbrinus Lohse, 1958
vexans Eppelsheim, 1881
xanthopus Erichson, 1839

Acylophorus Nordmann, 1837
glaberrimus (Herbst, 1784)
Xantholinini Erichson, 1839
Leptacinus Erichson, 1839
batychrus (Gyllenhal, 1827)
formicetorum Märkel, 1841
intermedius Donisthorpe, 1936
pusillus (Stephens, 1833)
 linearis (Gravenhorst, 1802)
 $-\text{othioides}$ Baudi di Selve, 1869
sulcifrons (Stephens, 1833)
 ops Coiffait, 1956
Phacophallus Coiffait, 1956
parumpunctatus (Gyllenhal, 1827)
Gauropterus Thomson, 1856
fulgidus (Fabricius, 1787)
Gyrohypnus Mannerheim, 1830
angustatus Stephens, 1833
 scoticus (Joy, 1913)
fracticornis (O.F.Müller, 1776)
Nudobius Thomson, 1860
lentus (Gravenhorst, 1806)
Hypnogyra Casey, 1906
glabra (Nordmann, 1837)
 angularis (Ganglbauer, 1895)
Xantholinus Dejean, 1821
distans distans Mulsant, Rey, 1853
laevigatus Jakobson, 1847
 clairei Coiffait, 1956
linearis (Olivier, 1795)
longiventris Heer, 1839
 sejugatus G.Benick, 1953
relucens (Gravenhorst, 1806)
tricolor (Fabricius, 1787)
Othiini Thomson, 1859
Othius Stephens, 1839
angustus Stephens, 1833
punctulatus (Goeze, 1777)
subuliformis Stephens, 1832
 myrmecophilus Kiesenwetter, 1843
Atrecus Jacquelain du Val, 1856
affinis (Paykull, 1789)
longiceps (Fauvel, 1872)
pilicornis (Paykull, 1790)
Paederinae Fleming, 1821
Paederus Fabricius, 1775
brevipennis Lacordaire, 1835
fuscipes Curtis, 1826
littoralis Gravenhorst, 1802
riparius (Linnaeus, 1758)
Astenus Stephens, 1833

<i>gracilis</i> (Paykull, 1789)	<i>beszedesi</i> Reitter, 1914
<i>procerus</i> (Gravenhorst, 1806)	<i>bluehweissi</i> Scheerpeltz, 1936
<i>filiformis</i> (Latreille, 1806)	
<i>pulchellus</i> (Heer, 1839)	<i>Euaesthetus</i> Gravenhorst, 1806
Rugilus Samouelle, 1819	<i>bipunctatus</i> (Ljungh, 1804)
<i>erichsoni</i> (Fauvel, 1867)	<i>ruficapillus</i> Lacordaire, 1835
<i>fragilis</i> (Gravenhorst, 1806)	Steninae McLeay, 1825
<i>angustatus</i> (Geoffroy, 1785)	Stenus Latreille, 1796
<i>scutellatus</i> (Motschulsky, 1858)	<i>ageus</i> Casey, 1884
<i>orbiculatus</i> (Paykull, 1789)	<i>gerhardti</i> L.Benick, 1915
<i>rufipes</i> Germar, 1836	<i>argus</i> Gravenhorst, 1802
<i>similis</i> (Erichson, 1839)	<i>ater</i> Mannerheim, 1830
Sunius Stephens, 1829	<i>atratus</i> Erichson, 1839
<i>melanocephalus</i> (Fabricius, 1792)	<i>bifoveolatus</i> Gyllenhal, 1827
Pseudomedon Mulsant, Rey, 1877	<i>biguttatus</i> (Linnaeus, 1758)
<i>obscurellus</i> (Erichson, 1840)	<i>bimaculatus</i> Gyllenhal, 1810
<i>obsoletus</i> (Nordmann, 1837)	<i>binotatus</i> Ljungh, 1804
Lithocaris Dejean, 1833	<i>bohemicus</i> Machulka, 1947
<i>ochracea</i> (Gravenhorst, 1802)	<i>boops</i> Ljungh, 1804
Scopaeus Erichson, 1840	<i>brevipennis</i> Thomson, 1851
<i>laevigatus</i> (Gyllenhal, 1827)	<i>brunnipes</i> Stephens, 1833
<i>pusillus</i> Kiesenwetter, 1843	<i>calcaratus</i> W.Scriba, 1864
Lathrobium Gravenhorst, 1802	<i>canaliculatus</i> Gyllenhal, 1827
<i>brunnipes</i> (Fabricius, 1792)	<i>carbonarius</i> Gyllenhal, 1827
<i>dilutum</i> Erichson, 1839	<i>cautus</i> Erichson, 1839
<i>microps</i> L.Benick, 1942	<i>vafellus</i> Erichson, 1839
<i>elongatum</i> (Linnaeus, 1767)	<i>cicindeloides</i> (Schaller, 1783)
<i>fovulum</i> Stephens, 1833	<i>circularis</i> Gravenhorst, 1802
<i>fulvipenne</i> Gravenhorst, 1806	<i>clavicornis</i> (Scopoli, 1763)
<i>muelleri</i> Bernhauer, 1899	<i>comma</i> LeConte, 1863
<i>impressum</i> Heer, 1841	<i>bipunctatus</i> Erichson, 1839
<i>filiforme</i> Gravenhorst, 1806	<i>crassus</i> Stephens, 1833
<i>laevipenne</i> Heer, 1839	<i>eumerus</i> Kiesenwetter, 1850
<i>longulum</i> Gravenhorst, 1802	<i>europaeus</i> Puthz, 1966
<i>patris</i> G.Benick, 1950	<i>cautus</i> Puthz, 1966 nec Erichson, 1839
<i>pallidipenne</i> Hochhuth, 1851	<i>excubitor</i> Erichson, 1839
<i>ripicola</i> Czwalina, 1888	<i>rossicus</i> Bernhauer, 1903
<i>rufipenne</i> Gyllenhal, 1813	<i>flavipalpis</i> Thomson, 1860
<i>volgense</i> Hochhuth, 1851	<i>flavipes</i> Stephens, 1833
<i>geminum</i> Kraatz, 1857	<i>fornicatus</i> Stephens, 1833
Lobrathium Mulsant, Rey, 1878	<i>fulvicornis</i> Stephens, 1833
<i>multipunctum</i> (Gravenhorst, 1802)	<i>fuscipes</i> Gravenhorst, 1802
Tetartopeus Czwalina, 1888	<i>geniculatus</i> Gravenhorst, 1806
<i>quadratus</i> (Paykull, 1789)	<i>guttula</i> P.W.J.Müller, 1821
<i>scutellaris</i> (Nordmann, 1837)	<i>humilis</i> Erichson, 1839
<i>terminatus</i> (Gravenhorst, 1802)	<i>impressus</i> Germar, 1824
<i>zetterstedti</i> (Rye, 1872)	<i>icanus</i> Erichson, 1839
Ochthephilum Stephens, 1829	<i>incrassatus</i> Erichson, 1839
<i>fracticorne</i> (Paykull, 1800)	<i>juno</i> (Paykull, 1789)
Euasthetinae Thomson, 1859	<i>latifrons</i> Erichson, 1839
Edaphus Motschulsky, 1857	<i>longipes</i> Heer, 1839

Note: North border of main distribution area.

- longitarsis* Thomson, 1851
lustrator Erichson, 1839
melanarius Stephens, 1833
melanopus (Marsham, 1802)
mendicus Erichson, 1840
Note: North border of main distribution area.
morio Gravenhorst, 1806
nanus Stephens, 1833
nigritulus Gyllenhal, 1827
nitens Stephens, 1833
ochropus Kiesenwetter, 1858
 erichsoni Rye, 1864
opticlus Gravenhorst, 1806
pallipes Gravenhorst, 1802
pallitarsis Stephens, 1833
palposus Zetterstedt, 1828
proditor Erichson, 1839
providus Erichson, 1839
 rogeri Kraatz, 1857
pubescens Stephens, 1833
pumilio Erichson, 1839
pusillus Stephens, 1833
 exiguus Erichson, 1840
ruralis Erichson, 1840
scabriculus J.R.Sahlberg, 1876
scrutator Erichson, 1840
similis (Herbst, 1784)
sylvester Erichson, 1839
tarsalis Ljungb., 1804
umbratilis Casey, 1884
Dianous Samouelle, 1819
coerulescens (Gyllenhal, 1810)
Oxyporinae Fleming, 1821
Oxyporus Fabricius, 1775
mannerheimii Gyllenhal, 1827
Note: Officially protected species in Latvia.
maxillosus Fabricius, 1792
rufus (Linnaeus, 1758)
Proteininae Erichson, 1839
Megarthrus Curtis, 1829
denticollis (Beck, 1817)
depressus (Paykull, 1789)
hemipterus (Illiger, 1794)
nitidulus Kraatz, 1858
Proteinus Latreille, 1796
atomarius Erichson, 1840
brachypterus (Fabricius, 1792)
laevigatus Hochhuth, 1871
Omaliini McLeay, 1825
Eusphalerum Kraatz, 1857
- lapponicum* (Mannerheim, 1830)
longipenne (Erichson, 1839)
luteum (Marsham, 1802)
 ophthalmicum (Paykull, 1800)
minutum (Fabricius, 1792)
primulae (Stephens, 1834)
Acrulia Thomson, 1858
inflata (Gyllenhal, 1813)
Pycnoglypta Thomson, 1858
lurida (Gyllenhal, 1813)
Phyllodrepa Thomson, 1859
floralis (Paykull, 1789)
ioptera (Stephens, 1834)
linearis (Zetterstedt, 1828)
 scabriuscula (Kraatz, 1857)
melanocephala (Fabricius, 1787)
nigra (Gravenhorst, 1806)
Hapalaraea Thomson, 1858
pygmaea (Paykull, 1800)
Omalium Gravenhorst, 1802
caesum Gravenhorst, 1806
excavatum Stephens, 1834
rivulare (Paykull, 1789)
Phloeostiba Thomson, 1858
lapponica (Zetterstedt, 1838)
plana (Paykull, 1792)
Phloeonomus Heer, 1839
pusillus (Gravenhorst, 1806)
sjoeberti Strand, 1937
Anthophagini Thomson, 1859
Xylodromus Heer, 1839
brunneipennis (Stephens, 1832)
depressus (Gravenhorst, 1802)
Orochares Kraatz, 1858
angustatus (Erichson, 1840)
Deliphrum Erichson, 1839
tectum (Paykull, 1789)
Deliphrosoma Reitter, 1909
macrocephalum (Eppelsheim, 1873)
 pechlaneri Lohse, 1963
Note: North border of main distribution area.
Anthobium Samouelle, 1819
atrocephalum (Gyllenhal, 1827)
melanocephalum (Illiger, 1794)
Olophrum Erichson, 1839
assimile (Paykull, 1800)
boreale (Paykull, 1792)
 florae Scheerpeltz, 1929
piceum (Gyllenhal, 1810)
Acidota Stephens, 1829
crenata (Fabricius, 1792)

<i>cruentata</i> Mannerheim, 1830	<i>laqueatus</i> (Marsham, 1802)
<i>Lesteva</i> Latreille, 1796	<i>piceus</i> (Linnaeus, 1767)
<i>longoelytrata</i> (Goeze, 1777)	<i>politus</i> Erichson, 1840
<i>pubescens</i> Mannerheim, 1830	<i>sculptus</i> Gravenhorst, 1806
<i>Geodromius</i> Redtenbacher, 1856	<i>Anotylus</i> Thomson, 1859
<i>plagiatus</i> (Fabricius, 1798)	<i>affinis</i> (Czwalina, 1870)
<i>Anthophagus</i> Gravenhorst, 1802	<i>clypeonitens</i> (Pandellé, 1867)
<i>angusticollis</i> (Mannerheim, 1830)	<i>complanatus</i> (Erichson, 1839)
<i>caraboides</i> (Linnaeus, 1758)	<i>chilensis</i> Coiffait, Saiz, 1968
<i>omalinus</i> Zetterstedt, 1828	<i>ixellensis</i> (Dvorak, 1954)
<i>Coryphiini</i> Portevin, 1929	<i>fairmairei</i> (Pandellé, 1867)
<i>Coryphium</i> Stephens, 1834	<i>hamatus</i> Fairmaire, Laboulbéné, 1856
<i>angusticolle</i> Stephens, 1834	<i>insecatus</i> (Gravenhorst, 1806)
<i>letzneri</i> Schwarz, 1873	<i>inustus</i> (Gravenhorst, 1806)
Oxytelinae Fleming, 1821	<i>excavatus</i> (Motschulsky, 1857)
Coprophilini Erichson, 1839	<i>maritimus</i> (Thomson, 1861)
Syntomium Curtis, 1828	<i>perrisi</i> (Fauvel, 1862)
<i>aeneum</i> (P.W.J.Müller, 1821)	<i>nitidulus</i> (Gravenhorst, 1802)
<i>Deleaster</i> Erichson, 1839	<i>borealis</i> (Motschulsky, 1860)
<i>dichrous</i> (Gravenhorst, 1802)	<i>rugifrons</i> (Hochhuth, 1849)
<i>Coprophilus</i> Latreille, 1829	<i>rugosus</i> (Fabricius, 1775)
<i>striatulum</i> (Fabricius, 1792)	<i>sculpturatus</i> (Gravenhorst, 1806)
<i>Manda</i> Blackwelder, 1952	<i>tetracarinatus</i> (Block, 1799)
<i>mandibularis</i> (Gyllenhal, 1827)	<i>Platystethus</i> Mannerheim, 1830
<i>Oxytelini</i> Fleming, 1821	<i>arenarius</i> (Geoffroy, 1785)
<i>Carpelimus</i> Kirby, 1819	<i>cornutus</i> (Gravenhorst, 1802)
<i>bilineatus</i> Stephens, 1834	<i>nitens</i> (R.F.Sahlberg, 1832)
<i>corticinus</i> (Gravenhorst, 1806)	<i>nodifrons</i> Mannerheim, 1830
<i>despectus</i> (Baudi di Selve, 1869)	<i>Bledius</i> Samouelle, 1819
<i>elongatulus</i> (Erichson, 1839)	<i>atricapillus</i> Williams, 1929
<i>erichsoni</i> (Sharp, 1871)	<i>praetermissus</i> Williams, 1929
<i>augustae</i> (Bernhauer, 1901)	<i>crassicollis</i> Lacordaire, 1835
— <i>exiguus</i> (Erichson, 1839)	<i>cribricollis</i> Heer, 1839
<i>foveolatus</i> (R.F.Sahlberg, 1832)	<i>dama jutlandensis</i> Herman, 1986
— <i>fuliginosus</i> (Gravenhorst, 1802)	<i>defensus</i> Fauvel, 1879
<i>gracilis</i> (Mannerheim, 1830)	<i>denticollis</i> Fauvel, 1872
<i>graciliformis</i> Konzelmann, Lohse, 1981	<i>dissimilis</i> Erichson, 1840
<i>heidenreichi</i> (L.Benick, 1934)	<i>erraticus</i> Erichson, 1839
<i>impressus</i> (Lacordaire, 1835)	<i>femoralis</i> (Gyllenhal, 1827)
<i>lindrothi</i> (Palm, 1942)	<i>fergussoni</i> Joy, 1912
<i>manchuricus subtilicornis</i> (Roubal, 1946)	<i>arenarius</i> (Paykull, 1800)
<i>pusillus</i> (Gravenhorst, 1802)	<i>filipes</i> Sharp, 1911
<i>lindbergi</i> (Scheerpeltz, 1937)	<i>frater</i> Kraatz, 1858
<i>rivularis</i> (Motschulsky, 1860)	<i>gallicus</i> (Gravenhorst, 1806)
<i>subterraneus</i> Smetana, 1960	<i>fracticornis</i> (Paykull, 1790)
<i>Aploderus</i> Stephens, 1833	<i>littoralis</i> Heer, 1839
<i>caelatus</i> (Gravenhorst, 1802)	<i>longulus</i> Erichson, 1839
— <i>caesus</i> (Erichson, 1839)	<i>nanus</i> Erichson, 1840
<i>Oxytelus</i> Gravenhorst, 1802	<i>opus</i> (Block, 1799)
<i>fulvipes</i> Erichson, 1839	<i>subsinuatus</i> Mulsant, Rey, 1878
<i>intricatus</i> Erichson, 1840	<i>pallipes</i> (Gravenhorst, 1806)
	<i>larseni</i> Hansen, 1940

- subterraneus* Erichson, 1839
talpa (Gyllenhal, 1810)
terebrans (Schiödte, 1866)
tibialis Heer, 1839
tricornis (Herbst, 1784)
vilis Mäklin, 1876
- Phloeochaerinae** Erichson, 1839
Phloeocharis Mannerheim, 1830
subtilissima Mannerheim, 1830
- Habrocerinae** Mulsant, Rey, 1875
Habrocerus Erichson, 1839
capillaricornis (Gravenhorst, 1806)
- Trichophyinae** Thomson, 1859
Trichophya Mannerheim, 1830
pilicornis (Gyllenhal, 1810)
- Tachyporinae** McLeay, 1825
Bolitobiini Horn, 1877
Mycetoporus Mannerheim, 1830
clavicornis (Stephens, 1832)
dispersus Schülke, Kocian, 2000
glaber (Sperk, 1835)
lepidus (Gravenhorst, 1806)
 brunneus (Marsham, 1802)
longulus Mannerheim, 1830
— nigricollis (Stephens, 1835)
 splendens (Marsham, 1802)
punctus (Gravenhorst, 1806)
rufescens (Stephens, 1832)
Ischnosoma Stephens, 1829
bergrothi (Hellén, 1925)
longicorne (Mäklin, 1847)
splendidum (Gravenhorst, 1806)
- Bryoporus** Kraatz, 1857
rugipennis (Pandellé, 1869)
- Lordithon** Thomson, 1859
exoletus (Erichson, 1839)
lunulatus (Linnaeus, 1760)
pulchellus (Mannerheim, 1830)
speciosus (Erichson, 1839)
thoracicus (Fabricius, 1776)
trimaculatus (Fabricius, 1793)
trinotatus (Erichson, 1839)
- Bolitobius** Leach, 1819
castaneus (Stephens, 1832)
cingulatus Mannerheim, 1830
- Parabolitobius** Li, Zhao, Sakei, 2000
formosus (Gravenhorst, 1806)
- Tachyporini** McLeay, 1825
Sepedophilus Gistel, 1856
bipunctatus (Gravenhorst, 1802)
- *bipustulatus* (Gravenhorst, 1802)
constans (Fowler, 1888)
 stoeckli (Lokay, 1913)
immaculatus (Stephens, 1832)
littoreus (Linnaeus, 1758)
pedicularius (Gravenhorst, 1802)
testaceus (Fabricius, 1793)
wankowiczi Pandellé, 1969
- Tachyporus** Gravenhorst, 1802
abdominalis (Fabricius, 1781)
atriceps Stephens, 1832
chrysomelinus (Linnaeus, 1758)
corpulentus J.R.Sahlberg, 1876
hypnorum (Fabricius, 1775)
nitidulus (Fabricius, 1781)
obtusus (Linnaeus, 1767)
pallidus Sharp, 1871
 scutellaris Rye, 1871
- pusillus* Gravenhorst, 1806
 macropterus Stephens, 1832
pulchellus Mannerheim, 1843
quadriscopulatus Pandellé, 1869
ruficollis Gravenhorst, 1802
scitulus Erichson, 1839
solutus Erichson, 1839
transversalis Gravenhorst, 1806
- Lamprinodes** Luze, 1901
saginatus (Gravenhorst, 1806)
- Lamprinus** Heer, 1839
erythropterus (Panzer, 1796)
- Tachinus** Gravenhorst, 1802
bipustulatus (Fabricius, 1793)
corticinus Gravenhorst, 1802
elongatus Gyllenhal, 1810
 ochsi Coiffait, 1954
- fimetarius* Gravenhorst, 1802
humeralis Gravenhorst, 1802
laticollis Gravenhorst, 1802
lignorum (Linnaeus, 1758)
marginatus (Fabricius, 1793)
marginellus (Fabricius, 1781)
pallipes (Gravenhorst, 1806)
proximus Kraatz, 1855
rufipennis Gyllenhal, 1810
signatus Gravenhorst, 1802
 rufipes auct. nec Linnaeus, 1758
- subterraneus* (Linnaeus, 1758)
- Coproporus** Kraatz, 1858
heterocerus Fauvel, 1898
- Cilea** Jacquelin du Val, 1856
silphoides (Linnaeus, 1767)

Aleocharinae Fleming, 1821	
Aleocharini Fleming, 1821	
Aleochara Gravenhorst, 1802	
<i>bilineata</i> Gyllenhal, 1810	
<i>binotata</i> Kraatz, 1856	
<i>bipustulata</i> (Linnaeus, 1761)	
<i>brevipennis</i> Gravenhorst, 1806	
<i>curtula</i> (Goeze, 1777)	
<i>fumata</i> Gravenhorst, 1802	
<i>inconspicua</i> Aubé, 1850	
<i>intricata</i> Mannerheim, 1830	
<i>laevigata</i> Gyllenhal, 1810	
<i>lanuginosa</i> Gravenhorst, 1802	
<i>moerens</i> Gyllenhal, 1827	
<i>sanguinea</i> (Linnaeus, 1758)	
<i>sparsa</i> Heer, 1839	
<i>tristis</i> Gravenhorst, 1806	
Tinotus Sharp, 1883	
<i>morion</i> (Gravenhorst, 1802)	
Oxypodini Thomson, 1859	
Oxypoda Mannerheim, 1830	
<i>abdominalis</i> Mannerheim, 1830	
<i>acuminata</i> (Stephens, 1832)	
<i>alternans</i> (Gravenhorst, 1802)	
<i>brachyptera</i> (Stephens, 1832)	
<i>difficilis</i> Roubal, 1931	
<i>brevicornis</i> (Stephens, 1832)	
<i>carnica</i> Scheerpeltz, 1956	
<i>umbrata</i> (Gyllenhal, 1810)	
<i>elongatula</i> Aubé, 1850	
— <i>exoleta</i> Erichson, 1839	
<i>bavarica</i> Scheerpeltz, 1956	
<i>huetheri</i> Scheerpeltz, 1956	
<i>flavicornis</i> Kraatz, 1856	
<i>amoena</i> Fairmaire, Laboulbène, 1856	
<i>formiceticola</i> Märkel, 1841	
<i>funebris</i> Kraatz, 1856	
<i>haemorrhoa</i> Mannerheim, 1830	
<i>opaca</i> (Gravenhorst, 1802)	
<i>togata</i> Erichson, 1837	
<i>vittata</i> Märkel, 1842	
Hygropora Kraatz, 1856	
<i>cunctans</i> (Erichson, 1837)	
Ocyusa Kraatz, 1856	
<i>maura</i> (Erichson, 1837)	
Calodera Mannerheim, 1830	
<i>aethiops</i> (Gravenhorst, 1802)	
<i>nigrita</i> Mannerheim, 1830	
<i>riparia</i> Erichson, 1837	
<i>rubens</i> (Erichson, 1837)	
Parocyusa Bernhauer, 1902	
	<i>rubicunda</i> (Erichson, 1837)
	Ischnoglossa Kraatz, 1856
	<i>prolixa</i> (Gravenhorst, 1802)
	Dexiogya Thomson, 1858
	<i>corticina</i> (Erichson, 1837)
	Thiasophila Kraatz, 1856
	<i>angulata</i> (Erichson, 1837)
	Cratarea Thomson, 1858
	<i>suturalis</i> (Mannerheim, 1830)
	Haploglossa Kraatz, 1856
	<i>gentilis</i> (Märkel, 1844)
	— <i>marginalis</i> (Gravenhorst, 1806)
	<i>picipennis</i> (Gyllenhal, 1827)
	<i>villosula</i> (Stephens, 1832)
	<i>pulla</i> (Gyllenhal, 1827)
	Ocalea Erichson, 1837
	<i>badia</i> Erichson, 1837
	<i>picata</i> (Stephens, 1832)
	Ilyobates Kraatz, 1856
	<i>nigricollis</i> (Paykull, 1800)
	<i>hoelzeli</i> Scheerpeltz, 1947
	Amarochara Thomson, 1858
	<i>forticornis</i> (Lacordaire, 1835)
	Phloeopora Erichson, 1837
	<i>corticalis</i> (Gravenhorst, 1802)
	Dinarda Samouelle, 1819
	<i>dentata</i> (Gravenhorst, 1806)
	<i>hagensii</i> Wasmann, 1889
	<i>maerkelii</i> Kiesenwetter, 1843
	Meotica Mulsant, Rey, 1873
	<i>filiformis</i> (Motschulsky, 1860)
	<i>apicalis</i> G.Benick, 1954
	<i>bohemica</i> G.Benick, 1954
	<i>capitalis</i> Mulsant, Rey, 1873
	<i>ermischi</i> G.Benick, 1954
	<i>foveolata</i> G.Benick, 1954
	<i>globulosa</i> G.Benick, 1954
	<i>hoelzeli</i> G.Benick, 1954
	<i>orbicularis</i> G.Benick, 1954
	<i>soniae</i> Bondroit, 1913
	Apimela Mulsant, Rey, 1874
	<i>mulsanti</i> (Ganglbauer, 1895)
	<i>pallens</i> (Mulsant, Rey, 1852)
	Athetini Casey, 1910
	Dasygnypeta Lohse, 1974
	— <i>velata</i> (Erichson, 1837)
	Thinonoma Thomson, 1859
	<i>atra</i> (Gravenhorst, 1806)
	Ischnopoda Stephens, 1835
	<i>coarctata</i> (Erichson, 1837)
	<i>constricta</i> (Erichson, 1837)
	<i>leucopus</i> (Marsham, 1802)

- scitula* (Erichson, 1837)
Brachyusa Mulsant, Rey, 1874
concolor (Erichson, 1839)
Schistoglossa Kraatz, 1856
gemina (Erichson, 1837)
viduata (Erichson, 1837)
Aloconota Thomson, 1858
debilicornis (Erichson, 1839)
gregaria (Erichson, 1839)
languida (Erichson, 1837)
Amidobia Thomson, 1858
talpa (Heer, 1841)
Liogluta Thomson, 1858
longiuscula (Gravenhorst, 1802)
microptera Thomson, 1867
 oblongiuscula (Sharp, 1869)
Geostiba Thomson, 1858
circellaris (Gravenhorst, 1806)
Dadobia Thomson, 1856
immersa (Erichson, 1837)
Plataraea Thomson, 1858
brunnea (Fabricius, 1798)
Atheta Thomson, 1858
aeneicollis (Sharp, 1869)
basicornis (Mulsant, Rey, 1852)
britteni Joy, 1913
brunneipennis (Thomson, 1852)
cadaverina (Brisout de Barnevile, 1860)
castanoptera (Mannerheim, 1830)
crassicornis (Fabricius, 1792)
 repanda (Mulsant, Rey, 1874)
cauta (Erichson, 1837)
celata (Erichson, 1837)
dadopora Thomson, 1867
debilis (Erichson, 1837)
elongatula (Gravenhorst, 1802)
euryptera (Stephens, 1832)
flavipes (Gravenhorst, 1806)
 wasserburgeri Bernhauer, 1931
fungi (Gravenhorst, 1806)
 beskidica (Paenik, 1999)
 foreistica (Paenik, 1999)
 otryticia (Paenik, 1999)
– *fungicola* (Thomson, 1852)
gagatina (Baudi di Selve, 1848)
graminicola (Gravenhorst, 1806)
grisea (Thomson, 1852)
hygrobia (Thomson, 1856)
laevana (Mulsant, Rey, 1852)
longicornis (Gravenhorst, 1802)
luridipennis (Mannerheim, 1830)
- *luteipes* (Erichson, 1837)
malleus Joy, 1913
 hygrobia auct. nec (Thomson, 1856)
melanaria (Mannerheim, 1830)
melanocera (Thomson, 1856)
nigricornis (Thomson, 1852)
nigritula (Gravenhorst, 1802)
obtusangula Joy, 1913
occulta (Erichson, 1837)
orbata (Erichson, 1837)
orphana (Erichson, 1837)
parca (Mulsant, Rey, 1874)
picipes (Thomson, 1856)
pilicornis (Thomson, 1852)
ravilla (Erichson, 1839)
setigera (Sharp, 1869)
sodalis (Erichson, 1837)
subsinuata (Erichson, 1839)
subtilis (W.Scriba, 1866)
terminalis (Gravenhorst, 1806)
triangulum (Kraatz, 1856)
trinotata (Kraatz, 1856)
volans (W.Scriba, 1859)
zosterae (Thomson, 1856)
Dinarea Thomson, 1858
aequata (Erichson, 1837)
angustula (Gyllenhal, 1810)
 hubenthali (Heymes, 1929)
linearis (Gravenhorst, 1802)
Lypocorrhe Thomson, 1859
anceps (Erichson, 1837)
Nehemitropia Lohse, 1971
lividipennis (Mannerheim, 1830)
 sordida (Marsham, 1802)
Acrotona Thomson, 1859
aterrima (Gravenhorst, 1802)
muscorum (Brisout de Barnevile, 1860)
parvula (Mannerheim, 1830)
pygmaea (Gravenhorst, 1806)
sylvicola (Kraatz, 1856)
Amischa Thomson, 1858
analis (Gravenhorst, 1802)
Thamiarea Thomson, 1858
cinnamomea (Gravenhorst, 1802)
Falagriini Mulsant, Rey, 1873
Falagria Samouelle, 1819
caesa (Erichson, 1837)
sulcatula (Gravenhorst, 1806)
Cordalia Jacobs, 1925
obscura (Gravenhorst, 1802)

Anaulacaspis Ganglbauer, 1895
nigra (Gravenhorst, 1802)
Lomechusini Fleming, 1821
Drusilla Samouelle, 1819
canaliculata (Fabricius, 1787)
Zyras Stephens, 1835
cognatus (Märkel, 1842)
collaris (Paykull, 1800)
funestus (Gravenhorst, 1806)
humeralis (Gravenhorst, 1802)
laticollis (Märkel, 1844)
limbatus (Paykull, 1789)
lugens (Gravenhorst, 1802)
Lomechusa Gravenhorst, 1806
emarginata (Paykull, 1789)
paradoxa (Gravenhorst, 1806)
Homalotini Heer, 1839
Gyrophaena Mannerheim, 1830
affinis Mannerheim, 1830
angustata (Stephens, 1832)
bihamata Thomson, 1867
boleti (Linnaeus, 1758)
congrua Erichson, 1837
fasciata (Marsham, 1802)
gentilis Erichson, 1839
joyi Wendler, 1924
minima Erichson, 1837
nana (Paykull, 1800)
poweri Crotch, 1866
Encephalus Kirby, 1832
complicans Kirby, 1832
Bolitochara Mannerheim, 1830
lucida (Gravenhorst, 1802)
mulsanti Sharp, 1875
obliqua Erichson, 1837
pulchra (Gravenhorst, 1806)
 lunulata (Paykull, 1789)
Leptusa Kraatz, 1856
fumida (Erichson, 1839)
norvegica Strand, 1941
pulchella (Mannerheim, 1830)
ruficollis (Erichson, 1839)
Euryusa Erichson, 1837
castanoptera Kraatz, 1856
Anomagnathus Solier, 1849
cuspidatus (Erichson, 1839)
Homalota Mannerheim, 1830
plana (Gyllenhal, 1810)
Cyphaea Fauvel, 1863
curtula (Erichson, 1837)

Placusini Mulsant, Rey, 1871
Placusa Erichson, 1837
atrata (Mannerheim, 1830)
incompleta Sjöberg, 1934
tachyporoides (Waltl, 1838)
Autaliini Thomson, 1859
Autalia Samouelle, 1819
impressa (Olivier, 1795)
 brevicornis Blair, 1944
puncticollis Sharp, 1864
rivularis (Gravenhorst, 1802)
Hygronomini Thomson, 1859
Hygronoma Erichson, 1837
dimidiata (Gravenhorst, 1806)
Hypocyphtini Laporte de Castelnau, 1835
Holobus Solier, 1849
flavicornis (Boisduval, Lacordaire, 1835)
Oligota Mannerheim, 1830
inflata (Mannerheim, 1830)
pusillima (Gravenhorst, 1806)
Cypha Samouelle, 1819
discoidea (Erichson, 1839)
imitator (Luze, 1902)
laeviuscula (Mannerheim, 1830)
seminulum (Erichson, 1839)
Myllaenini Ganglbauer, 1895
Myllaena Erichson, 1837
brevicornis (Matthews, 1838)
dubia (Gravenhorst, 1806)
intermedia Erichson, 1837
minuta (Gravenhorst, 1806)
Gymnusini Heer, 1839
Gymnusa Gravenhorst, 1806
brevicollis (Paykull, 1800)
Deinopsini Sharp, 1883
Deinopsis Matthews, 1838
erosa (Stephens, 1832)
Micropeplinae Leach, 1815
Micropeplus Latreille, 1809
caelatus Erichson, 1839
porcatus (Paykull, 1789)
staphylinoides (Marsham, 1802)
Scaphidiidae Latreille, 1807
Scaphidiinae Latreille, 1807
Scaphidium Olivier, 1790
quadrivirgatum Olivier, 1790
Scaphisominae Casey, 1893
Scaphisoma Leach, 1815
agaricinum (Linnaeus, 1758)
 major J.R.Sahlberg, 1889

- petzi* Csiki, 1908
podoces Lundblad, 1952
pulicarium (Rossi, 1792)
punctipenne Petz, 1905
assimile assimile Erichson, 1845
balcanicum Tamanini, 1954
boleti boleti (Panzer, 1793)
robustior Pic, 1905
boreale Lundblad, 1952
inopinatum Löbl, 1967
subalpinum subalpinum Reitter, 1881
Pselaphidae Latreille, 1802
Euplectinae LeConte, 1861
Bibloporus Thomson, 1859
bicolor (Denny, 1825)
Bibloplectus Reitter, 1882
ambiguus (Reichenbach, 1816)
pusillus (Denny, 1825)
spinosus Raffray, 1914
Meliceria Raffray, 1898
traegardhi Palm, 1938
Note: This species is known from a few specimens from extreme SE part of Latvia situated at the border with Lithuania and Belarus'. Probably should be more widely distributed in old broad-leaved forests of Central and eastern Latvia.
Euplectus Leach, 1817
duponti Aubé, 1833
karsteni (Reichenbach, 1816)
kirbii revelierei Reitter, 1884
— nanus (Reichenbach, 1816)
punctatus Mulsant, 1861
sanguineus Denny, 1825
signatus (Reichenbach, 1816)
Trimium Aubé, 1833
brevicorne (Reichenbach, 1816)
Batriscinae Reitter, 1882
Batrisodes Reitter, 1882
venustus (Reichenbach, 1816)
Goniacerinae Reitter, 1882
Bythinini Raffray, 1890
Bryaxis Kugelann, 1794
bulbifer (Reichenbach, 1816)
curtisii (Leach, 1817)
nigripennis (Aubé, 1844)
puncticollis (Denny, 1825)
gracilipes (Raffray, 1914)
validus (Aubé, 1844)
Tychini Raffray, 1904
Tychus Leach, 1817
- niger* (Paykull, 1800)
Brachyglutini Raffray, 1904
Rybaxis Saulcy, 1876
longicornis (Leach, 1817)
sanguinea Leach, 1817 nec (Linnaeus, 1758)
Brachygluta Thomson, 1859
fossulata (Reichenbach, 1816)
haematica sinuata (Aubé, 1833)
Reichenbachia Leach, 1826
juncorum (Leach, 1817)
Trissemus Jeannel, 1949
impressus (Panzer, 1803)
Pselaphinae Latreille, 1802
Pselaphini Latreille, 1802
Pselaphaulax Reitter, 1909
dresdensis (Herbst, 1792)
Pselaphus Herbst, 1792
heisei Herbst, 1792
Tyrini Reitter, 1882
Tyrus Aubé, 1833
mucronatus (Panzer, 1803)
Clavigerinae Leach, 1815
Claviger Preussler, 1790
testaceus Preussler, 1790
Note: Insufficiently known species.
Scarabaeiformia
Scarabaeoidea
Lucanidae Latreille, 1806
Syndesinae McLeay, 1819
Sinodendron Schneider, 1791
cylindricum (Linnaeus, 1758)
Lucaninae Latreille, 1806
Lucanus Scopoli, 1763
cervus cervus (Linnaeus, 1758)
americanus Hope, Westwood, 1845
antennatus Rey, 1890
armiger Herbst, 1790
capitalis Möllenkamp, 1910
capra Olivier, 1789
capreolus Fuessly, 1775
cornutus (Dale, 1693)
dorcas O.F.Müller, 1776
grandis Haworth, 1807
hircus Herbst, 1790
inermis Marsham, 1802
longipennis Von Rothenburg, 1900
lusitanicus Hope, Westwood, 1845
major Burmeister, 1847
microcephalus Mulsant, 1842
nigrinus March, 1931
ornatus Bernau, 1929
validus Möllenkamp, 1912
vicinus Hope, Westwood, 1845

Note: The last known record is from the end of the 1960s from Ieriķi area (Gaujas National park, Central Latvia). There are also some unconfirmed records from W and E Latvia. North border of main distribution area. Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.

Dorcus McLeay, 1819

- parallelipipedus* (Linnaeus, 1758)
 - bipunctatus* (Pontoppidan, 1765)
 - bituberculatus* McLeay, 1819 nec Parry, 1864
 - bituberculatus* Parry, 1864 nec McLeay, 1819
 - capra* (Panzer, 1797)
 - dama* (O.F.Müller, 1776)
 - immaturus* Mulsant, 1842
 - infractus* (Bergsträsser, 1778)
 - punctatus* (Pontoppidan, 1763)
 - tuberculatus* McLeay, 1819

Note: Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.

Platycerus O.F.Müller, 1764

- caprea* (DeGeer, 1774)

Note: This species is included on the list of indicator species of natural forest habitats.

- caraboides caraboides* (Linnaeus, 1758)
 - aeneus* (Weise, 1883)
 - azurea* Dalla Torre, 1789
 - carniolicus* (Roubaud, 1919)
 - coeruleescens* (Udman, 1753)
 - cribratus* (Mulsant, Rey, 1863)
 - virescens* Mulsant, 1842
 - viridiaeneus* Mulsant, 1842
 - viridicollis* (Depoli, 1917)
 - viridis* Dalla Torre, 1789

Note: This species is included on the list of indicator species of natural forest habitats.

Ceruchus McLeay, 1819

- chrysomelinus* (Hochenwarth, 1785)
 - piceus* (Bonsdorff, 1785)
 - silesiacus* (Mergele, Dejean, 1837)
 - sylvicola* (Mulsant, 1842)
 - tarandus* (Panzer, 1789)
 - tenebrioides* (Latreille, 1807)
 - tenebroides* (Fabricius, 1787)

Note: Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.

Trogidae McLeay, 1819

- Trox** Fabricius, 1775
 - sabulosus* (Linnaeus, 1758)
 - scaber* (Linnaeus, 1767)

Geotrupidae Latreille, 1806

Bolboceratinae Mulsant, 1842

- Odonteus** Samouelle, 1772
 - armiger* (Scopoli, 1772)
- Geotrupinae** Latreille, 1806
 - Anoplotrupes** Jekel, 1865
 - stercorosus* (L.G.Scriba, 1791)
 - sylvaticus* (Panzer, 1798)
 - Geotrupes** (**Geotrupes** Latreille, 1796)
 - mutator* (Marsham, 1802)

Note: Very rare species, known only from a few records during the last 100 years, mostly in the Eastern part of Latvia. North border of main distribution area.

- spiniger* (Marsham, 1802)
- stercorarius* (Linnaeus, 1758)

Trypocopris Motschulsky, 1859

- vernalis vernalis* (Linnaeus, 1758)

Typhaeus (**Typhaeus** Leach, 1815)

- *typhoeus* (Linnaeus, 1758)

Lethrinae Mulsant, Rey, 1871

Lethrus (**Lethrus** Scopoli, 1777)

- apterus* (Laxmann, 1770)

Note: Scarce species with no confirmed records in last 100 years. North border of main distribution area.

Scarabaeidae Latreille, 1802

Scarabaeinae Latreille, 1802

Ateuchetus Bedel, 1892

- *laticollis* (Linnaeus, 1767)

Copris Geoffroy, 1762

- lunaris* (Linnaeus, 1758)

Caccobius Thomson, 1859

- schreberi* (Linnaeus, 1767)

Onthophagus (**Palaeonthophagus** Zunino, 1979)

- coenobita* (Herbst, 1783)

Note: Very rare and insufficiently known species with no confirmed records within the last 100 years.

fracticornis (Preyssler, 1790)

gibbulus (Pallas, 1781)

- austriacus* (Panzer, 1793)

nuchicornis (Linnaeus, 1758)

ovatus (Linnaeus, 1767)

vacca (Linnaeus, 1767)

Note: Very rare and insufficiently known species with no confirmed records in the last 100 years.

- *vitulus* (Fabricius, 1776)

Onthophagus (**Onthophagus** Latreille, 1802)

- *taurus* (Schreber, 1759)
- Sisyphus*** Latreille, 1807
- *schaefferi* (Linnaeus, 1758)
- Aegialiinae*** Lacordaire, 1856
- Aegialia*** Latreille, 1807
- arenaria* (Fabricius, 1787)
- sabuleti* (Panzer, 1797)
- rufa* (Fabricius, 1792 nec DeGeer, 1778)
 - rufina* Silfverberg, 1977
 - spissipes* LeConte, 1878

Note: Insufficiently known species with only a few confirmed records in the last 50 years.

- Aphodiinae*** Leach, 1815
- Aphodiini*** Leach, 1815
- Aphodius* (*Otophorus*)** Mulsant, 1842
- haemorrhoidalis* (Linnaeus, 1758)
- Aphodius* (*Teuchestes*)** Mulsant, 1842
- fossor* (Linnaeus, 1758)
 - silvaticus* (Ahrens, 1812)
- Aphodius* (*Eupleurus*)** Mulsant, 1842
- subterraneus* (Linnaeus, 1758)
- Aphodius* (*Colobopterus*)** Mulsant, 1842
- erraticus* (Linnaeus, 1758)
- Aphodius* (*Ammoecius*)** Mulsant, 1842
- brevis* Erichson, 1848
 - elevatus* (Paykull, 1798)
- Aphodius* (*Plagiogonus*)** Mulsant, 1842
- putridus* (Geoffroy, 1785)
 - arenarius* (Olivier, 1789 nec Fabricius, 1787)
 - rhododactylus* (Marsham, 1802)
- Aphodius* (*Calamosternus*)** Motschulsky, 1859
- granarius* (Linnaeus, 1767)
- Aphodius* (*Liothorax*)** Motschulsky, 1859
- niger* (Panzer, 1797)
- plagiatus* (Linnaeus, 1767)
- Aphodius* (*Labarrus*)** Mulsant, Rey, 1869
- *lividus* (Olivier, 1789)
- Aphodius* (*Trichonotulus*)** Bedel, 1911
- scrofa* (Fabricius, 1787)
- Aphodius* (*Euorodalus*)** Dellacasa, 1983
- coenosus* (Panzer, 1798)
 - tristis* (Zenker, 1801)
- Aphodius* (*Esymus*)** Mulsant, Rey, 1869
- merdarius* (Fabricius, 1775)
- pusillus* (Herbst, 1789)
- Aphodius* (*Phalacronothus*)** Motschulsky, 1859
- *biguttatus* Germar, 1824
 - sanguinolentus* (Panzer, 1797 nec Herbst, 1783)
 - tirolensis* Rosenhauer, 1847
- *quadrimaculatus* (Linnaeus, 1761)
- Aphodius* (*Nialis*)** Mulsant, Rey, 1869

- *varians* Duftschmid, 1805
 - bimaculatus* (Fabricius, 1787 nec Laxmann, 1770)
- Aphodius* (*Acanthobodilus*)** Dellacasa, 1983
- immundus* Creutzer, 1799
- Aphodius* (*Parammoeclius*)** Seidlitz, 1891
- *gibbus* Germar, 1817
- Aphodius* (*Nimbus*)** Mulsant, Rey, 1869
- *contaminatus* (Herbst, 1783)
- *obliteratus* Panzer, 1823
- Aphodius* (*Chilothorax*)** Motschulsky, 1859
- conspurcatus* (Linnaeus, 1758)
- distinctus* (O.F.Müller, 1776)
 - inquinatus* (Herbst, 1783)
- melanostictus* Schmidt, 1840
- paykulli* Bedel, 1908
 - tessulatus* (Paykull, 1798 nec Laicharting, 1781)
- pictus* Sturm, 1805
- sticticus* (Panzer, 1798)
 - equestris* (Panzer, 1798)
- Aphodius* (*Nobius*)** Mulsant, Rey, 1869
- *serotinus* (Panzer, 1799)
- Aphodius* (*Eudolus*)** Mulsant, Rey, 1869
- quadriguttatus* (Herbst, 1783)
- Aphodius* (*Acrossus*)** Mulsant, 1842
- bimaculatus* (Laxmann, 1770)
- Note: Very rare species with no confirmed records in the last 50 years.
- depressus* (Kugelann, 1792)
- luridus* (Fabricius, 1775)
- rufipes* (Linnaeus, 1758)
- Aphodius* (*Planolinus*)** Mulsant, Rey, 1869
- borealis* Gyllenhal, 1827
- uliginosus* Hardy, 1847
 - fasciatus* (Olivier, 1789 nec Linnaeus, 1758)
 - foetidus* (Fabricius, 1792 nec Herbst, 1783)
 - putridus* (Herbst, 1789 nec Geoffroy, 1785)
 - tenellus* Hardy, 1847
- Aphodius* (*Aphodius*)** Illiger, 1798
- fimetarius* (Linnaeus, 1758)
- foetens* (Fabricius, 1787)
 - aestivalis* Stephens, 1839
- foetidus* (Herbst, 1783)
 - scybalaria* (Olivier, 1789 nec Fabricius, 1781)
- Aphodius* (*Agoliinus*)** A.Schmidt, 1913
- piceus* Gyllenhal, 1808
- Aphodius* (*Agrilinus*)** Mulsant, Rey, 1869
- ater* (DeGeer, 1774)
- nemoralis* Erichson, 1848
- ardoei* Landin, 1954
- sordidus* (Fabricius, 1775)
- rufus* (Moll, 1782)
- scybalaria* (Fabricius, 1781)

— *lapporum* Gyllenhal, 1806
Aphodius (*Bodilus*) Mulsant, Rey, 1869
ictericus (Laicharting, 1781)
nitidulus (Fabricius, 1792)
lugens Creutzer, 1799
Aphodius (*Melinopterus*) Mulsant, 1842
— *consputus* Creutzer, 1799
prodromus (Brahm, 1790)
punctatosulcatus Sturm, 1805
sabulicola Thomson, 1868
sphaelatetus (Panzer, 1798)
Aphodius (*Pubinus*) Mulsant, Rey, 1869
tomentosus (O.F.Müller, 1776)
lutarius (Fabricius, 1792)
Aphodius (*Amidorus*) Mulsant, Rey, 1869
obscurus (Fabricius, 1792)
Euheptaulacus Dellacasa, 1983
sus (Herbst, 1783)
Note: Very rare and insufficiently known species with only few records in the last 50 years.
villosus (Gyllenhal, 1806)
Note: Very rare and insufficiently known species with only few records in the last 50 years.
Heptaulacus Mulsant, 1842
testudinarius (Fabricius, 1775)
Note: Very rare and insufficiently known species. North border of main distribution area
Oxyomus Dejean, 1821
sylvestris (Scopoli, 1763)
Psammodiini Mulsant, 1842
Psammodius Fallén, 1807
asper (Fabricius, 1775)
sulcicollis (Illiger, 1802)
Diastictus Mulsant, 1842
vulneratus (Sturm, 1805)
Note: Very rare species known only from a few specimens from the Cekule area (Central Latvia). North border of main distribution area.
Pleurophorus Mulsant, 1842
caesus (Creutzer, 1796)
Note: This species is known only from a single record, from the Cekule area (Central Latvia). North border of main distribution area.
Rhyssemus Mulsant, 1842
germanus (Linnaeus, 1767)
asper auct. nec. (Fabricius, 1775)
Note: North border of main distribution area.
Sericinae Hope, 1837
Serica McLeay, 1819

brunnea (Linnaeus, 1758)
fulva (DeGeer, 1774)
fulvescens (Geoffroy, 1785)
Maladera (*Maladera*) Mulsant, Rey, 1871
holosericea (Scopoli, 1772)
berolinensis (Herbst, 1786)
chrysomelina (Schrank, 1781)
chrysomelooides (Schrank, 1781)
fumosa (Geoffroy, 1785)
fusca Mulsant, 1842
lausellata (Geoffroy, 1785)
pellucida (Schönherr, 1817)
pellucidula (Sulzer, 1776)
sulzeri (Fuessly, 1775)
variabilis (Fabricius, 1776)
Note: North border of main distribution area.
Omaloplia (*Omaloplia*) Schönherr, 1817
ruricola (Fabricius, 1775)
atrata (Geoffroy, 1785)
disca (Mulsant, 1842)
floricola (Laicharting, 1781)
humeralis (Fabricius, 1775)
immarginata (Mulsant, 1842)
nigromarginata (Herbst, 1786)
obscura (Mulsant, 1842)
Melolonthinae Samouelle, 1819
Amphimallon (*Amphimallon*) Berthold, 1827
solstitiale (*solstitiale*) Linnaeus, 1758
aurantiacus Mulsant, Rey, 1842
autumnalis (Geoffroy, 1785)
falleni (Gyllenhal, 1817)
fulvicollis Mulsant, 1842
lateralis Mulsant, Rey, 1842
maldipesi Chevrolat, 1902
montivagus Halbherr, 1892
ochraceus (Knoch, 1801)
subsulcatus Faldermann, 1835
suturalis Mulsant, 1842
Melolontha (*Melolontha*) Fabricius, 1775
hippocastani (*hippocastani*) Fabricius, 1801
aethiops Westhoff, 1884
albicans Stierlin, 1862
amasicola Westhoff, 1884
coronata Mulsant, 1842
discoidalis Dalla Torre, 1879
fuscicollis Kraatz, 1885
metzleri Westhoff, 1884
mulsanti Bedel, 1911
nig Mulsant, 1842
pectoralis Megerle, 1812
pectoralis Westhoff, 1885
picea Wanach, 1906
rex Dalla Torre, 1879
schwarzii Kraatz, 1888
suturalis Kraatz, 1888
tibialis Mulsant, 1842
vulgaris Olivier, 1789
vulgaris Herbst, 1789

***melolontha* (Linnaeus, 1758)**

- albida* Mulsant, 1842 nec Redtenbacher, 1849
- albida* Redtenbacher, 1849 nec Mulsant, 1842
- asiatica* Brenske, 1900
- discicollis* Mulsant, 1842
- femoralis* Kraatz, 1885 nec Reitter, 1885
- funesta* Westhoff, 1884
- humeralis* Westhoff, 1884
- luctuosa* Westhoff, 1884
- lugubris* Mulsant, 1842
- majalis* Moller, 1785
- marginata* Kraatz, 1888
- melanopus* Westhoff, 1884
- nigra* Kellner, 1877
- nigritarsis* Pfanneberg, 1905
- obscuripes* Westhoff, 1884
- pulcherrima* Dalla Torre, 1879
- redtenbacheri* Dalla Torre, 1912
- ruficeps* Kraatz, 1887
- ruficollis* Mulsant, 1842
- scapularis* Westhoff, 1884
- vulgaris* Fabricius, 1775

***Polyphylla* (*Polyphylla* Harris, 1841)**

- fullo*** (Linnaeus, 1758)
 - boryi* (Brullé, 1832)
 - luctuosa* Mulsant, 1842
 - marmorata* Mulsant, 1842

Note: Very rare species known only from a few specimens from Southern (both Eastern & Western) parts of Latvia. One unconfirmed record also from NW Latvia (Sītere National park). North border of main distribution area.

Rutelinae McLeay, 1819***Anomala* Samouelle, 1819**

- dubia*** (Scopoli, 1763)
 - aenea* (DeGeer, 1774)
 - dubia* (Herbst, 1790)
 - julii* (Fabricius, 1792)
 - frischii* (Fabricius, 1775)

***Anisoplia* (*Anisoplia* Dejean, 1821)**

- *agricola*** (Poda, 1761)
 - cracifera* (Herbst, 1790)
 - conjuncta* Schilsky, 1888
 - cyathigera* (Scopoli, 1763)
 - delesa* Schilsky, 1888

***Chaetopteroplus* Medvedev, 1949**

- *segetum*** (*segetum* (Herbst, 1783))
 - campestris* (Herbst, 1783)
 - fruticola* (Fabricius, 1775)
 - rasa* (Zoubkoff, 1833)
 - straminea* (Brullé, 1832)
 - velutina* (Erichson, 1847)

***Phyllopertha* Stephens, 1830**

- horticola*** (Linnaeus, 1758)
 - adiaphora* (Poda, 1761)
 - cyancephala* Mulsant, 1842
 - descordans* Meissner, 1907

- macularis* Mulsant, 1842
- maculata* Dalla Torre, 1879
- metallica* Dalla Torre, 1879
- nigra* Dalla Torre, 1879
- perrisi* Mulsant, 1842
- rufipes* Dalla Torre, 1879
- rufiventris* Westwood, 1881
- ursulatipennis* Villa, 1885
- viridicollis* (DeGeer, 1774)
- zea* Reitter, 1903

Hopliinae Latreille, 1829***Hoplia* (*Decamera* Mulsant, 1842)**

- praticola*** Duftschmid, 1805
 - bosnica* Reitter, 1903
 - palustris* Heer, 1842
 - rupicola* Mulsant, 1842

Note: Known from a few individuals from two localities in Central and SW Latvia. North border of main distribution area.

***Hoplia* (*Hoplia* Illiger, 1803)**

- *argentea*** (Poda, 1761 nec Marsham, 1802)
 - ambigua* Mulsant, 1842
 - deflorata* Mulsant, 1842
 - farinosa* (Linnaeus, 1761)
 - glauxa* Mulsant, 1842
 - griseola* Rey, 1890
 - karamani* Reitter, 1893
 - micans* Mulsant, 1842
 - rufolutea* Mulsant, 1842
 - sedicolor* Mulsant, 1842
 - squamosa* (Fabricius, 1792)
 - sublutea* Mulsant, 1842
 - viridula* Mulsant, 1842

***graminicola* (Fabricius, 1792)**

- argentea*** (Marsham, 1802 nec Poda, 1761)
- farinosa* Herbst, 1890 nec (Linnaeus, 1761)
- pulverulenta* (Fabricius, 1775)

***parvula* Krynicki, 1832**

- pollinosa*** Erichson, 1847 nec Krynicki, 1832
- pulverosa* Küster, 1853

zaitzevi* Jakobson, 1914*Dynastinae McLeay, 1819**

- Oryctes* (*Oryctes* Illiger, 1798)**
- nasicornis*** (*nasicornis* (Linnaeus, 1758))
 - punctipennis* Minck, 1915

Cetoniinae Leach, 1815***Oxythyrea* Mulsant, 1842**

- funesta*** (Poda, 1761)
 - albopunctata* (DeGeer, 1774)
 - consobrina* (Villa, Villa, 1833)
 - deleta* Mulsant, 1842
 - funeraria* (Geoffroy, 1785)
 - greeni* (Donovan, 1807)
 - nigrominuta* Seabra, 1905
 - stictica* (Schaefer, 1766)
 - viridana* Seabra, 1905

***Tropinota* Mulsant, 1842**

hirta (Poda, 1761)

heyrovskyi (Obenberger, 1917)
nigrina Mulsant, 1842
subfasciata Mulsant, 1842
tonsa (Burmeister, 1842)
vestita (Say, 1825)

Note: Very rare species with no records in the last 50 years. North border of main distribution area.

Cetonia Fabricius, 1775

aurata aurata (Linnaeus, 1758)
aenea (Fuessly, 1778)
chlorochrysa (Schröter, 1776)
cupreofulgens Mulsant, 1842
kalichi Mikšić, 1959
maculata L.G.Scriba, 1790
nobilis Schrank, 1798
piligera Mulsant, 1842
praeclera Mulsant, 1842
purpurata Heer, 1841
smaragda (DeGeer, 1774)
variabilis Preyssler, 1790
viridicorusca Voet, 1779

Liocola Thomson, 1859

lugubris (Herbst, 1786)
aenea (L.G.Scriba, 1790)
aeruginea (Herbst, 1790)
marmorata (Fabricius, 1792)
punctatoundulata (Petagna, 1819)
quercus (Schrank, 1788)

Note: Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.

Cetonischema Reitter, 1898

— *aeruginosa* (Drury, 1770)
aurata (Fuessly, 1782)
aureocuprea (Mulsant, 1842)
germania (Voet, 1779)
fastuosa (Fabricius, 1792)
frischii (Schrank, 1798)
immaculata (L.G.Scriba, 1790)
ignea Reitter, 1898
miribella Reitter, 1898
nudiventris (Germar, 1824)
smaragda (Brahm, 1790)
speciosissima (Scopoli, 1786)
superba (Villers, 1798)
viridis (Fuessly, 1786)

Potosia Mulsant, Rey, 1871

cuprea metallica (Herbst, 1782)
albohirta (Kraatz, 1883)
excisa (Fieber, 1831)
floricola (Herbst, 1790)
freyi Mikšić, 1958
mulsanti Reitter, 1898
ottomana (Blanchard, 1842)
rubrocuprea (Mulsant, 1842)

sexundata (Petagna, 1819)

sternohirta Seidlitz, 1888

— *ungarica ungarica* (Herbst, 1790)

rubrocuprea Medvedev, 1947
simplex (Schilsky, 1888)
viridis (Fabricius, 1792)

Valginae Mulsant, 1842***Valgus*** L.G.Scriba, 1790***hemipterus*** (Linnaeus, 1758)

rufosquamatus (Dalla Torre, 1879)
variegatus (Scopoli, 1763)

Note: North border of main distribution area.

Trichiinae Fleming, 1821***Osmoderma*** Le Peletier, Aud.-Serville, 1825***eremita lassallei*** Barraud, Tauzin, 1991

coriarium DeGeer, 1774
eremiticum Knoch, 1901
sociale Horn, 1871
variabile Linnaeus, 1758

Note: Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.

Gnorimus Le Peletier, Audinet-Serville, 1825

nobilis (Linnaeus, 1758)
auratus (Schrank, 1781)
cupidatus (Fabricius, 1787)
purpureus Depoli, 1912
viridulus (DeGeer, 1774)

Note: Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.

variabilis (Linnaeus, 1758)

albopunctatus (DeGeer, 1774)
cursor (Voet, 1769)
decempunctatus (Schrank, 1798)
nigricollis Mulsant, 1842
octopunctatus (Fabricius, 1775)

Note: Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.

Trichius Fabricius, 1787***fasciatus*** (Linnaeus, 1758)

abbraviatus Mulsant, 1842
abdominalis Mulsant, 1842
abruptus Rossi, 1882
albohirtus Reitter, 1892
andersoni Bogdanov-Katkov, 1913
beckersi Geilenkeiser, 1892
bimaculatus Gebler, 1830
bipunctatus Kraatz, 1891
borealis Schulze, 1910
commutatus Rossi, 1882
confluens Mulsant, 1842
divisus Mulsant, 1842
dubius Mulsant, 1842
erichsoni Rossi, 1882

fabricii Rossi, 1882
fennicus Tesař, 1935
hieroglyphicus Jesatko, 1935
interruptus Mulsant, 1842
jureceki Jesatko, 1935
krogerusi Tesař, 1935
kueni Friese, 1896
lindbergi Tesař, 1935
lineaticollis Kraatz, 1891
linnei Rossi, 1882
lojoensis Tesař, 1935
merkli Roubal, 1917
noui Pellet, 1871
obenbergeri Jesatko, 1935
obliquus Mulsant, 1842
omissus Tesař, 1935
prolongatus Mulsant, 1842
pseudosibiricus Schulze, 1910
pulchellus Rossi, 1882
quadrimaculatus Kraatz, 1891
rangnovi Schulze, 1910
reitteri Kraatz, 1891
sahlbergi Schulze, 1910
scutellaris Kraatz, 1891
sekerai Tesař, 1935
sibiricus Rossi, 1882
sterbai Tesař, 1935
succinctus Fabricius, 1787
suecicus Tesař, 1935
vseteckai Tesař, 1935
vulgaris Rossi, 1882

Elateriformia**Scirtoidea**

Clambidae Fischer von Waldheim, 1821

Clambus Fischer von Waldheim, 1821

armadillo (DeGeer, 1774)

gibbulus (LeConte, 1850)

radula Endrödy-Younga, 1960

pubescens Redtenbacher, 1849

punctulum (Beck, 1817)

borealis Strand, 1946

Eucinetidae Lacordaire, 1857

Eucinetus Germar, 1818

haemorrhoidalis (Germar, 1818)

haemorrhous (Duftschmid, 1825)

Scirtidae Fleming, 1821

Elodes Latreille, 1796

minuta (Linnaeus, 1767)

pseudominuta Klausnitzer, 1971

tricuspis Nyholm, 1985

Microcara Thomson, 1859

testacea (Linnaeus, 1767)

bohemani Mannerheim, 1844

Cyphon Paykull, 1799

coarctatus Paykull, 1799

hilaris Nyholm, 1944

kongsbergensis Munster, 1924

laevipennis Tournier, 1868

phragmiteticola Nyholm, 1955

ochraceus Stephens, 1830

padi (Linnaeus, 1758)

palustris Thomson, 1855

pubescens (Fabricius, 1792)

punctipennis Sharp, 1873

ruficeps Tournier, 1868

variabilis (Thunberg, 1787)

nigriceps Kiesenwetter, 1860

Prionocyphon Redtenbacher, 1858

serricornis (P.W.J.Müller, 1821)

Scirtes Illiger, 1807

hemisphaericus (Linnaeus, 1767)

orbicularis (Panzer, 1793)

Dascilloidea

Dascillidae Lacordaire, 1857

Dascillinae Lacordaire, 1857

Dascillini Lacordaire, 1857

Dascillus Latreille, 1796

cervinus (Linnaeus, 1758)

Buprestoidea

Buprestidae Leach, 1815

Chrysochroinae Laporte de Castelnau, 1835

Chrysochroini Laporte de Castelnau, 1835

Chalcophora Dejean, 1833

mariana *mariana* (Linnaeus, 1758)

deaurata (Voet, 1806)

hiulca (Pallas, 1782)

Note: Officially protected species in Latvia;
also included on the list of indicator species of
natural forest habitats.

Poecilonotini Jakobson, 1911

Poecilonota Eschscholtz, 1829

variolosa *variolosa* (Paykull, 1799)

conspersa (Gyllenhal, 1801)

plebeja (Olivier, 1790 nec Gory, Laporte de
Castelnau, 1837)

plebeja (Gory, Laporte de Castelnau, 1837 nec
Olivier, 1790)

rustica (Herbst, 1787 nec Linnaeus, 1758)

setulosa Fleischer, 1896

tenebrionis (Schaefer, 1766 nec L., 1758)

tremulae Abeille de Perrin, 1896

Note: This species is included on the list of
indicator species of natural forest habitats.

Dicercini Gistel, 1848

Dicerca (*Dicerca*) Eschscholtz, 1829

aenea *aenea* (Linnaeus, 1758)

austriaca (Schrank, 1781)

carniolica (Fabricius, 1801)

cuprea (Rossi, 1790)

<i>oxyptera</i> (Pallas, 1781)	
<i>reticulata</i> (Fabricius, 1793)	
<i>scabrosa</i> Mannerheim, 1837	
<i>subrugosa</i> (Paykull, 1799)	
<i>acuminata</i> (Pallas, 1782 nec DeGeer, 1774)	
<i>calcarata</i> (Fabricius, 1801)	
<i>furcata</i> (Thunberg, 1787)	
Note: This species is included on the list of indicator species of natural forest habitats.	
<i>alni alni</i> (Fischer von Waldheim, 1823)	
<i>berolinensis</i> (Paykull, 1799 nec Herbst 1779)	
<i>calcarata</i> Mannerheim, 1837 nec (F., 1801)	
<i>fagi</i> Laporte de Castelnau, Gory, 1837	
<i>marianna</i> (DeGeer, 1774 nec Linnaeus, 1758)	
Note: Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.	
Dicerca (Argante Gistel, 1834)	
<i>moesta</i> (Fabricius, 1792)	
<i>quadrilineata</i> (Herbst, 1801)	
Note: This species is included on the list of indicator species of natural forest habitats.	
Buprestinae Leach, 1815	
Buprestini Leach, 1815	
Buprestis (Buprestis Linnaeus, 1758)	
<i>haemorrhoidalis haemorrhoidalis</i> Herbst, 1780	
<i>punctata</i> Fabricius, 1787	
<i>quadristigma</i> Herbst, 1801	
<i>novemmaculata novemmaculata</i> Linnaeus, 1767	
<i>flavomaculata</i> Fabricius, 1787	
<i>flavopunctata</i> DeGeer, 1774	
<i>maculata</i> Fabricius, 1781	
<i>maculosa</i> Gmelin, 1788	
<i>major</i> Olivier, 1790	
<i>octomaculata</i> Pallas, 1806	
<i>tetrastichon</i> Linnaeus, 1758	
Note: This species is included on the list of indicator species of natural forest habitats.	
<i>octoguttata octoguttata</i> Linnaeus, 1758	
<i>albopunctata</i> Schaefer, 1776	
Note: This species is included on the list of indicator species of natural forest habitats.	
<i>rustica rustica</i> Linnaeus, 1758	
<i>lata</i> Sulzer, 1776	
<i>violacea</i> DeGeer, 1774	
Anthaxiini Gory, Laporte de Castelnau, 1839	
Anthaxia (Anthaxia Eschscholtz, 1829)	
<i>manca</i> (Linnaeus, 1767)	
<i>bistriata</i> (Fabricius, 1775)	
<i>mancatula</i> Abeille de Perrin, 1900	
<i>rubinus</i> (Geoffroy, 1785)	
Note: Very rare species with no confirmed records in the last 50 years.	
<i>nitidula</i> (Linnaeus, 1758)	

<i>cyanipennis</i> Gory, 1841	
<i>grabowskyi</i> Obenberger, 1912	
<i>laeta</i> (Schaefer, 1766)	
<i>millefolii</i> Abeille de Perrin, 1894 nec (Fabricius, 1801)	
<i>styria</i> (Voet, 1806)	
Note: Very rare species with no confirmed records in the last 50 years.	
<i>salicis</i> (Fabricius, 1776)	
<i>amplexus</i> Rey, 1891	
<i>croesus</i> (Villers, 1789)	
<i>hirticollis</i> Rey, 1891	
Note: Very rare species with no confirmed records in the last 50 years.	
Anthaxia (Melanthaxia Rikhter, 1949)	
<i>godeti</i> Gory, Laporte de Castelnau, 1839	
<i>granulata</i> Küster, 1850	
<i>raclinae</i> Baudon, 1959	
<i>submontana</i> Obenberger, 1930	
<i>helvetica</i> Stierlin, 1868	
<i>nigrocyanea</i> Rey, 1891	
<i>tirolensis</i> Obenberger, 1913	
<i>quadripunctata</i> (Linnaeus, 1758)	
<i>angulata</i> Küster, 1850	
<i>angulicollis</i> Küster, 1850	
<i>punctata</i> (Ponza, 1805)	
<i>quadriimpressa</i> Motschulsky, 1844	
<i>similis</i> Saunders, 1871	
<i>morio</i> (Fabricius, 1792)	
Note: This species is included on the list of indicator species of natural forest habitats.	
Melanophilini Bedel, 1921	
Melanophila Eschscholtz, 1829	
<i>acuminata</i> (DeGeer, 1774)	
<i>acuta</i> (Gmelin, 1788)	
<i>anthaxoides</i> Marquet, 1876	
<i>appendiculata</i> (Fabricius, 1792)	
<i>assimilis</i> LeConte, 1850	
<i>immaculata</i> Mannerheim, 1837	
<i>longipes</i> (Say, 1823)	
<i>morio</i> (Paykull, 1799)	
<i>obscurata</i> Lewis, 1893	
<i>opaca</i> LeConte, 1859	
<i>pecchiolii</i> Gory, Laporte de Castelnau, 1839	
<i>rugata</i> LeConte, 1857	
Note: This species is included on the list of indicator species of natural forest habitats.	
Phaenops Dejean, 1833	
<i>cyanea</i> (Fabricius, 1775)	
<i>azurea</i> (Olivier, 1790)	
<i>calybaea</i> (Villers, 1789)	
<i>clypeata</i> (Paykull, 1799)	
<i>sibirica</i> Pic, 1918	
<i>tarda</i> (Fabricius, 1792)	
<i>transbaicalica</i> Obenberger, 1924	
Chrysobothrini Laporte de Castelnau, Gory, 1838	

- Chrysobothris*** (*Chrysobothris*) Eschscholtz, 1829
- affinis affinis* (Fabricius, 1794)
 - assimulans* Schreibers, 1843
 - chrysostigma* (Geoffroy, 1785 nec L., 1758)
 - congener* (Paykull, 1799)
 - inaequalis* Reitter, 1895
 - materculae* Holynski, 1975
 - reitteri* Kerremans, 1900
 - sibirica* Obenberger, 1916
 - tristis* Abeille de Perrin, 1900
 - chrysostigma chrysostigma* (Linnaeus, 1758)
 - kerremansi* Abeille de Perrin, 1893
 - quadraticollis* Abeille de Perrin, 1894
 - Agrilinae*** Laporte de Castelnau, 1835
 - Coraebini*** Bedel, 1921
 - Coraebus*** Gory, Laporte de Castelnau, 1839
 - elatus elatus* (Fabricius, 1787)
 - aeruginosus* (Latreille, 1804)
 - lampsanae* (Bonelli, 1804)
 - metallicus* Gory, Laporte de Castelnau, 1839
 - pruinosus* Küster 1846
 - sinuatus* (Panzer, 1796)
 - subfasciatus* Küster 1846
 - undatus* (Fabricius, 1775)
 - pruni* (Panzer, 1796)
 - quadrifasciatus* (Rossi, 1790)
 - Agrilinae*** Laporte de Castelnau, 1835
 - Agrilini*** Laporte de Castelnau, 1835
 - Agrilus*** Curtis, 1825
 - angustulus* (Illiger, 1803)
 - angustatus* Petri, 1885
 - laetefrons* Mannerheim, 1837
 - olivaceus* (Gyllenhal, 1808)
 - pavidus* Gory, Laporte de Castelnau, 1837
 - rugicollis* Ratzeburg, 1839
 - viridis* Stephens, 1830 nec (Linnaeus, 1758)
 - ater ater* (Linnaeus, 1767)
 - biguttatus* (Rossi, 1790 nec Fabricius, 1776)
 - sexguttatus* (Brahm, 1790)
 - auricollis auricollis* Kiesenwetter, 1857
 - betuleti* (Ratzeburg, 1837)
 - impressicollis* Marseul, 1865 nec (Gory, 1841)
 - biguttatus* (Fabricius, 1776)
 - caeruleoviolaceus* (Thomson, 1864)
 - octoguttatus* (Geoffroy, 1785)
 - pannonicus* (Piller, Mitterpacher, 1783)
 - subfasciatus* Ménétriés, 1832
- Note: This species is included on the list of indicator species of natural forest habitats.
- convexicollis* Redtenbacher, 1849
 - cuprescens cuprescens* Ménétriés, 1832
 - altaicola* Obenberger, 1935
 - aurichalceus* Redtenbacher, 1849
 - chrysoderes* Bétis, 1914
 - communis* Obenberger, 1924

- egenus* Abeille de Perrin, 1895 nec (Gory, 1841)
- epistomalis* Abeille de Perrin, 1897
- fagi* Weiss, 1914
- krasai* Obenberger, 1924
- kuznecovi* Obenberger, 1935
- kuznecovinus* Obenberger, 1936
- lacrymans* Abeille de Perrin, 1897
- mokrzeckii* Obenberger, 1927
- obtusus* Abeille de Perrin, 1897
- politus* Weiss, 1914
- proximus* Rey, 1891 nec (Saunders, 1871)
- rubicola* Abeille de Perrin, 1897
- cyanescens*** (Ratzeburg, 1837)
 - amabilis* Gory, Laporte de Castelnau, 1837
 - amethystinus* (Schönherr, 1817 nec Oliver, 1790)
 - bidentulus* Ganglbauer, 1889
 - coeruleus* (Rossi, 1790 nec Thunberg, 1789)
 - cyaneus* (Lacordaire, Boisduval, 1835 nec Rossi, 1790)
 - fissifrons* Abeille de Perrin, 1897
 - kyselyi* Obenberger, 1924
 - sulcaticeps* Abeille de Perrin, 1869
- hyperici*** (Creutzer, 1789)
 - catalonicus* Pochon, 1964
 - elatus* Méquignon, 1907 nec (Fabricius, 1787)
 - hofferi* Obenberger, 1936
 - modestulus* Semenov, 1895
 - prasinus* Mulsant, 1863
 - sulcifer* Bétis, 1914
- integerrimus*** (Ratzeburg, 1839)
 - cupreus* Redtenbacher, 1849
- laticornis*** (Illiger, 1803)
 - aceris* Chevrolat, 1837
 - asperrimus* Marseul, 1865
 - laticollis* Kiesenwetter, 1857
 - minusculus* Marseul, 1865
 - reyi* Bauduer, 1878
 - scaberrimus* Ratzeburg, 1837
- mendax*** Mannerheim, 1837
 - faldermanni* Gory, Laporte de Castelnau, 1837
- Note: This species is included on the list of indicator species of natural forest habitats.
- paludicola*** Krogerus, 1922
- pratensis pratensis*** Ratzeburg 1837
 - linearis* (Paykull, 1799)
 - praeclarus* Krogerus, 1922
 - pseudocoeruleus* Obenberger, 1930
 - roberti* Chevrolat, 1837
 - strigicollis* Rey, 1891
- pseudocyanus*** Kiesenwetter, 1857
 - bialowiezaensis* Gutowski, 1993
 - foveola* Roubal, 1917 nec (Rey, 1891)
 - obrucevi* Obenberger, 1936
 - roubalianus* Obenberger, 1936
- subauratus subauratus*** (Gebler, 1833)
 - auripennis* Gory, Laporte de Castelnau, 1837
 - coryli* Ratzeburg, 1839

- sulcicollis* Lacordaire, 1835
cyanus (Rossi, 1790)
elongatus Saunders, 1871
elongatus (Herbst, 1780)
maksimirus Stiller, 1918
sahlbergi Mannerheim, 1837
tenuis Ratzeburg, 1839
viridis Gory, Laporte de Castelnau, 1839 nec
(Linnaeus, 1758)
suvorovi Obenberger, 1935
populneus (Schaefer, 1946)
viridis viridis (Linnaeus, 1758)
atra Kiesenwetter, 1859
aubei Gory, Laporte de Castelnau, 1837
bicolor Redtenbacher, 1849
caprea Chevrolat, 1838
darwini Wollaston, 1857
distinguendus Gory, Laporte de Castelnau, 1837
fagi Ratzeburg, 1839
filiformis (Herbst, 1801)
ignota Schaefer, 1946
linearis (Panzer, 1789)
littlei Curtis, 1840
mixtulus Obenberger, 1924
nocivus Ratzeburg, 1839
obtusus Obenberger, 1925
purpureicolor Pic, 1918
quercinus Redtenbacher, 1849
rosaceus (Scopoli, 1763)
rudis Abeille de Perrin, 1897
serraticornis (Scopoli, 1763)
viridipennis Gory, Laporte de Castelnau, 1837
- Aphanisticini** Jacquelin du Val, 1863
Aphanisticus Latreille, 1829
pusillus (Olivier, 1790)
emarginatus (Fallén, 1802 nec Olivier, 1790)
latus Dejean, 1836
lineola Germar, 1834
- Trachydini** Laporte de Castelnau, 1835
Trachys Fabricius, 1801
minutus (Linnaeus, 1758)
reflexiformis Obenberger, 1916
supraviolacea Thomson, 1864
- troglodytes* Gyllenhal, 1817
aeneus Mannerheim, 1837
cribratus Rey, 1891
foveicollis Rey, 1891
laevicollis Kiesenwetter, 1857
pumilus Laporte de Castelnau, Gory, 1839
puncticeps Abeille de Perrin, 1900
subglastra Rey, 1891
- Byrrhoidea**
Byrrhidae Latreille, 1804
Byrrhinae Latreille, 1804
Simplocariini Mulsant, Rey, 1869
Simplocaria (*Simplocaria* Stephens, 1829)
— *acuminata* Erichson, 1847
- metallica* (Sturm, 1807)
semistriata (Fabricius, 1794)
Morychini El Moursy, 1961
Morychus Erichson, 1846
aeneus (Fabricius, 1775)
modestus Kiesenwetter, 1850
- Pedilophorini** Casey, 1912
Lamprobyrrhulus Ganglbauer, 1902
nitidus (Schaller, 1783)
- Byrrhini** Latreille, 1804
Cytulus Erichson, 1846
sericeus (Forster, 1771)
auricomus (Duftschmid, 1825)
- Byrrhus** (*Byrrhus* Linnaeus, 1767)
arietinus arietinus Steffahny, 1843
fasciatus fasciatus (Forster, 1771)
pilula pilula (Linnaeus, 1758)
pustulatus pustulatus (Forster, 1771)
- Porcinolus** Mulsant, Rey, 1869
murinus (Fabricius, 1794)
- Syncalyptinae** Mulsant, Rey, 1869
Chaetophora Kirby, Spence, 1823
spinosa (Rossi, 1794)
- Curimopsis** (*Curimopsis* Ganglbauer, 1902)
paleata (Erichson, 1846)
setigera (Illiger, 1798)
— *setosa* (Waltl, 1838)
- Elmidae** Curtis, 1830
Elmina Curtis, 1830
Elmini Curtis, 1830
Elmis Latreille, 1798
aenea (P.W.J.Müller, 1806)
maugetii Latreille, 1802
obscura (P.W.J.Müller, 1806)
- Esolus** Mulsant, Rey, 1872
angustatus (P.W.J.Müller, 1821)
- Oulimnius** Des Gozis, 1886
tuberculatus (P.W.J.Müller, 1806)
- Limnius** Illiger, 1802
muelleri (Erichson, 1847)
- Note: Record needs confirmation. This species is often confused with *L. intermedius* Fairmaire, 1881.
- opacus* P.W.J.Müller, 1806
- Note: This species known from a single individual from the Cekule area (Central Latvia).
- volckmari* (Panzer, 1793)
ceconii Reitter, 1906
- Normandia** Pic, 1900

- nitens* (P.W.J.Müller, 1817)
Riolus Mulsant, Rey, 1872
cupreus (P.W.J.Müller, 1806)
Macronychini Mulsant, Rey, 1872
Macronychus Müller 1806
quadrituberculatus P.W.J.Müller, 1806
 Note: Several records from central, northern, eastern and South-eastern parts of Latvia.
 Should be more widely distributed.
- Dryopidae** Billberg, 1820
Dryopinae Billberg, 1820
Dryops (*Dryops*) Olivier, 1791
anglicanus Edwards, 1909
auriculatus (Geoffroy, 1785)
rudolfi Rusek, 1973
ernesti Des Gozis, 1886
griseus (Erichson, 1847)
luridus (Erichson, 1847)
similaris Bollow, 1936
Dryops (*Yrdops*) Steffan, 1961)
nitidulus (Heer, 1841)
- Limnichidae** Erichson, 1846
Limnichus Latreille, 1829
pygmaeus (Sturm, 1807)
sericeus (Duftschmid, 1825)
- Heteroceridae** McLeay, 1825
Heterocerinae McLeay, 1825
Heterocerus Fabricius, 1792
fenestratus (Thunberg, 1784)
asiaticus Nomura, 1958
completus Hatch, 1965
fausti Reitter, 1879
laevigatus Panzer, 1794
latus Grouvelle, 1915
marshami Stephens, 1828
multimaculatus Motschulsky, 1853
pusillus Waltl, 1839
siculus Kuwert, 1890
tonkinensis Grouvelle, 1896
- flexuosus* Stephens, 1828
apfelbecki Kuwert, 1890
arenarius Kiesenwetter, 1851
corsicus Rey, 1890
damryi Kuwert, 1890
dentifasciatus Kuwert, 1890
fausti Reitter, 1879
femoralis Krynický, 1832
hamifer Gené, 1836
heydeni Kuwert, 1890
maculosus Fairmaire, 1868
marginatus Gyllenhal, 1808
senegalensis Fairmaire, 1894
- fusculus* Kiesenwetter, 1843
oblongus Kuwert, 1890
- pulchellus* Kuwert, 1890
similis Kuwert, 1890
marginatus (Fabricius, 1787)
scutellaris Guillebeau, 1897
sulcatus Kuwert, 1890
obsoletus Curtis, 1828
Augyles Schiödte, 1866
hispidulus (Kiesenwetter, 1843)
intermedius (Kiesenwetter, 1843)
maritimus (Guerin-Meneville, 1844)
basalis (Mamitzia, 1929)
britannicus (Kuwert, 1890)
burchanensis (Schneider, 1896)
nanus (Gené, 1836)
pruiniosus (Kiesenwetter, 1851)
- Psephenidae** Lacordaire, 1857
Eubria Germar, 1818
palustris Germar, 1818
- Elateroidea**
- Elateridae** Leach, 1815
Agyrpninae Lacordaire, 1857
Agyrpnini Lacordaire, 1857
Agyrpnus Eschscholtz, 1829
murinus (Linnaeus, 1758)
Pyrophorinae Candéze, 1863
Adelocerini Bulyssón, 1893
Lacon Laporte de Castelnau, 1836
lepidopterus (Panzer, 1801)
 Note: Very rare and insufficiently known species.
— querceus (Herbst, 1784)
Danosoma Thomson, 1859
conspersus (Gyllenhal, 1808)
fasciatus (Linnaeus, 1758)
Negastriinae Nakane, 1953
Negastrius Thomson, 1859
pulchellus (Linnaeus, 1761)
sabulicola (Bohemian, 1853)
Zorochros Thomson, 1859
minimus (Lacordaire, 1835)
dermestoides (Herbst, 1806 nec L., 1767)
dufouri (Bulyssón, 1851)
- Oedostethus** LeConte, 1853
quadripustulatus (Fabricius, 1792)
tenuicornis (Germar, 1824)
Denticollinae Stein, Weise, 1877
Denticollini Stein, Weise, 1877
Athous Eschscholtz, 1829
— bicolor (Goeze, 1777)
longicollis (Olivier, 1790)
haemorrhoidalis (Fabricius, 1801)
subfuscus (O.F.Müller, 1767)

vittatus (Fabricius, 1792)

Hemicrepidius Germar, 1839

hirtus (Herbst, 1784)

niger (Linnaeus, 1758)

Crepidophorus Rosenhauer, 1847

mutilatus Rosenhauer, 1847

Diacanthous Reitter, 1905

undulatus (DeGeer, 1774)

Note: This species is included on the list of indicator species of natural forest habitats.

Stenagostus Thomson, 1859

rufus (DeGeer, 1774)

Note: Officially protected species in Latvia.

Denticollis Piller, Mitterpacher, 1783

borealis (Paykull, 1800)

Note: Scarce and insufficiently known species. South border of main distribution area. This species is included on the list of indicator species of natural forest habitats.

linearis (Linnaeus, 1758)

rubens Piller, Mitterpacher, 1783

Note: Officially protected species in Latvia.

Cidnopus Thomson, 1859

aeruginosus (Olivier, 1790)

minutus (Linnaeus, 1758)

parvulus (Panzer, 1799)

pilosus (Leske, 1785)

Hypnoidus Dillwyn, 1829

riparius (Fabricius, 1792)

rivularius (Gyllenhal, 1808)

consobrinus (Mulsant, Guillebeau, 1855)

Note: Scarce species. South border of main distribution area.

Prosternini Gistel, 1856

Ctenicera Latreille, 1829

– *cuprea* (Fabricius, 1775)

pectinicornis (Linnaeus, 1758)

Liotrichus Kiesenwetter, 1863

affinis (Paykull, 1800)

Orithales Kiesenwetter, 1863

serraticornis (Paykull, 1800)

Actenicerus Kiesenwetter, 1858

sjaelandicus (O.F.Müller, 1764)

Prosternon Latreille, 1834

tessellatum (Linnaeus, 1758)

holosericeum (Olivier, 1790)

Anostirus Thomson, 1859

castaneus castaneus (Linnaeus, 1758)

purpureus (Poda, 1761)

Note: Very rare and insufficiently known

species.

Aplotarsus Stephens, 1830

incanus (Gyllenhal, 1827)

quercus (Gyllenhal, 1808)

Selatosomus Stephens, 1830

aeneus (Linnaeus, 1758)

cruciatus (Linnaeus, 1758)

latus (Fabricius, 1801)

Note: Insufficiently known species with only a few records in the last 50 years.

– *melancholicus* (Fabricius, 1798)

Paraphotistus Kishii, 1966

impressus (Fabricius, 1792)

nigricornis (Panzer, 1799)

Calambus Thomson, 1859

bipustulatus (Linnaeus, 1767)

Elaterinae Leach, 1815

Megapenthini Gurjeva, 1973

Procraerus Reitter, 1905

tibialis (Lacordaire, 1835)

Apedini Gistel, 1856

Apedus Dejean, 1833

– *aethiops* (Lacordaire, 1835)

scrofa Germar, 1844

balteatus (Linnaeus, 1758)

cardinalis (Schiödte, 1865)

Note: Rare and insufficiently known species with only a few records in the last 50 years.

Should be more widely distributed.

cinnabarinus (Eschscholtz, 1829)

elegantulus (Schönherr, 1817)

elongatulus (Fabricius, 1787)

erythrogonus (P.W.J.Müller, 1821)

Note: This species is included on the list of indicator species of natural forest habitats.

hjorti (Rye, 1905)

– *mixtus* (Herbst, 1806)

nigrinus (Herbst, 1784)

nigroflavus (Goeze, 1777)

pomonae (Stephens, 1830)

pomorum (Herbst, 1784)

brigittae Bouwer, 1890

ferrugatus (Lacordaire, 1835)

robustus Bouwer, 1890

praeustus (Fabricius, 1792)

sanguineus (Linnaeus, 1758)

sanguinolentus (Schrank, 1776)

suecicus Palm, 1976

borealis Palm, 1947 nec (Paykull, 1800)

tristis (Linnaeus, 1758)

Elaterini Leach, 1815

- Sericus** Eschscholtz, 1829
brunneus (Linnaeus, 1758)
Melanotini Candéze, 1859
Melanotus Eschscholtz, 1829
castanipes (Paykull, 1800)
crassicollis (Erichson, 1841)
punctolineatus (Pelerin, 1829)
 niger (Fabricius, 1792 nec Linnaeus, 1758)
villosus (Geoffroy, 1785)
 erythropus (Gmelin, 1790)
 rufipes (Herbst, 1784 nec Goeze, 1777)
- Agriotini** Laporte de Castelnau, 1840
Agriotes Eschscholtz, 1829
gallicus Boisduval, Lacordaire, 1835
lineatus (Linnaeus, 1767)
obscurus (Linnaeus, 1758)
 — *pilosellus* (Schönherr, 1817)
 elongatus Burysson, 1893
 pilosus (Panzer, 1795)
sputator (Linnaeus, 1758)
 — *ustulatus* (Schaller, 1783)
- Ectinus** Eschscholtz, 1829
aterrimus (Linnaeus, 1761)
- Dalopius** Eschscholtz, 1829
marginatus (Linnaeus, 1758)
- Adrastus** Eschscholtz, 1829
 — *limbatus* (Fabricius, 1776)
- pallens* (Fabricius, 1792)
 nitidulus (Marsham, 1802 nec Fabricius, 1787)
- Synaptus** Eschscholtz, 1829
filiformis (Fabricius, 1781)
- Cardiophorinae** Candéze, 1859
Cardiophorus Eschscholtz, 1829
asellus Erichson, 1840
ebeninus (Germar, 1824)
gramineus (Scopoli, 1763)
rufipes (Goeze, 1777)
ruficollis (Linnaeus, 1758)
- Dicronychus** Brullé, 1832
cinereus (Herbst, 1784)
Note: Very rare
equiseti (Herbst, 1784)
equisetoides Lohse, 1976
 — *rubripes* (Germar, 1824)
- Paracardiophorus** Schwarz, 1895
 — *musculus* (Erichson, 1840)
- Lissominae** Laporte de Castelnau, 1840
Drapetes Dejean, 1821
mordelloides (Host, 1789)
 biguttatus (Piller, Mitterpacher, 1783 nec Fabricius, 1776)
- cinctus* (Panzer, 1796)
- Eucnemidae** Eschscholtz, 1829
Melasinae Fleming, 1821
Hylocharini Jacquelin du Val, 1859
Hylochares Latreille, 1834
cruentatus (Gyllenhal, 1808)
- Dirhaginae** Reitter, 1911
Microrhagus (*Microrhagus*) Dejean, 1833
lepidus Rosenhauer, 1847
 lindbergi (Palm, 1958)
pygmaeus (Fabricius, 1792)
- Rhacopus** Hampe, 1855
sahlbergi (Mannerheim, 1823)
- Clypeorhagus** Olexa, 1975
clypeatus (Hampe, 1850)
- Xylobiinae** Reitter, 1911
Otho Lacordaire, 1857
spondyloides (Germar, 1818)
- Xylophilus** Mannerheim, 1823
corticalis (Paykull, 1800)
 humeralis (Dufour, 1851)
- Hylis** Des Gozis, 1886
foveicollis (Thomson, 1874)
 fleischeri (Olexa, 1954)
- olexai* (Palm, 1955)
procerulus (Mannerheim, 1823)
- Eucneminae** Eschscholtz, 1829
Eucnemis Ahrens, 1812
capucina Ahrens, 1812
- Throscidae** Laporte de Castelnau, 1840
Aulonothroscus Horn, 1890
brevicollis (Bonvouloir, 1859)
- Trixagus** Kugelann, 1794
carinifrons (Bonvouloir, 1859)
dernestoides (Linnaeus, 1767)
elateroides (Heer, 1841)
exul (Bonvouloir, 1859)
- Cantharoidea**
- Lycidae** Laporte de Castelnau, 1840
Erotinae LeConte, 1881
Erotini LeConte, 1881
Dictyoptera Latreille, 1829
aurora (Herbst, 1784)
erythroptera Baudi di Selve, 1871
- Pyropterus** Mulsant, 1838
nigroruber (DeGeer, 1774)
 affinis (Paykull, 1799)
- Platycis** Thomson, 1859
minuta (Fabricius, 1787)
- Platerodini** Kleine, 1928
- Lopheros** LeConte, 1884

rubens (Gyllenhal, 1817)
Calochrominae Lacordaire, 1857
Lygistopterus Dejean, 1833
sanguineus (Linnaeus, 1758)
Drilidae Blanchard, 1845
Drilus Olivier, 1790
concolor Ahrens, 1812
Lampyridae Latreille, 1817
Lampyris Geoffroy, 1762
noctiluca (Linnaeus, 1767)
Phosphaenus Laporte de Castelnau, 1833
hemipterus (Geoffroy, 1762)
Cantharidae Imhoff, 1856
Cantharinae Imhoff, 1856
Podabrus Westwood, 1838
alpinus (Paykull, 1798)
Ancistronycha Märkel, 1852
cyanipennis (Faldermann, 1835)
violacea (Paykull, 1798 nec Thunberg, 1784)
Cantharis Linnaeus, 1758
decipiens Baudi di Selve, 1871
figurata Mannerheim, 1843
fulvicollis Fabricius, 1792
fusca Linnaeus, 1758
lateralis Linnaeus, 1758
livida Linnaeus, 1758
nigra (DeGeer, 1774)
bicolor Herbst, 1784 nec Linnaeus, 1763
thoracica (Olivier, 1790)
nigricans (O.F.Müller, 1776)
obscura Linnaeus, 1758
oculata (Gyllenhal, 1810)
paludosa Fallén, 1807
pellucida Fabricius, 1792
pulicaria Fabricius, 1781
quadripunctata (O.F.Müller, 1776)
rufa Linnaeus, 1758
rustica Fallén, 1807
Rhagonycha Eschscholtz, 1830
atra (Linnaeus, 1767)
elongata (Fallén, 1807)
fulva (Scopoli, 1763)
fugax Mannerheim, 1843
scopoli Gredler, 1866
lignosa (O.F.Müller, 1764)
limbata Thomson, 1864
testacea (Linnaeus, 1758)
Absidia Mulsant, 1863
rufotestacea (Letzner, 1845)
schoenherri (Dejean, 1837)
pilosa (Paykull, 1798 nec Scopoli, 1763)

Silis Dejean, 1825
nitidula (Fabricius, 1792)
Crudosilis Kasantsev, 1994
ruficollis (Fabricius, 1775)
rugicollis (Gebler, 1833)
Malthininae LeConte, 1861
Malthinus Latreille, 1806
balteatus Suffrian, 1851
biguttatus (Linnaeus, 1758)
biguttulus (Paykull, 1800)
frontalis (Marsham, 1802)
punctatus (Geoffroy, 1785)
flaveolus (Herbst, 1786)
Malthodes Kiesenwetter, 1852
brevicollis (Paykull, 1798)
crassicornis (Mäklin, 1846)
debilis Kiesenwetter, 1852
dimidiaticollis (Rosenhauer, 1847)
europaeus Wittmer, 1970
fibulatus Kiesenwetter, 1852
flavoguttatus Kiesenwetter, 1852
fuscus (Waltl, 1838)
guttifer Kiesenwetter, 1852
marginatus (Latreille, 1806)
maurus (Laporte de Castelnau, 1840)
minimus (Linnaeus, 1758)
pumilus (Brebisson, 1835)
atomus Thomson, 1864
spathifer Kiesenwetter, 1852
Cucujiformia
Bostrichoidea
Dermestidae Latreille, 1807
Dermestinae Latreille, 1807
Dermestini Latreille, 1807
Dermestes (*Dermestes* Linnaeus, 1758)
ater DeGeer, 1774
cadaverinus Fabricius, 1775
lardarius Linnaeus, 1758
Dermestes (*Dermestinus* Zhantiev, 1967)
erichsoni Ganglbauer, 1904
tessellatus Erichson, 1848
frischii Kugelann, 1792
vulpinus Herbst in Jablonsky, 1792
gyllenhali *gyllenhali* Laporte de Castelnau, 1840
atomarius Erichson, 1846
laniarius *laniarius* Illiger, 1801
maculatus DeGeer, 1774
vulpinus Fabricius, 1781
murinus *murinus* Linnaeus, 1758
— mustelinus Erichson, 1846
Note: All previous references are invalid and

- relate to *D. undulatus* (Brahm).
undulatus (Brahm, 1790)
- Megatominae** Leach, 1815
- Attagenini** Casey, 1900
- Attagenus** (*Attagenus*) Latreille, 1802
- brunneus* Faldermann, 1835
dalmatinus Küster, 1847
sordidus Heer, 1841
- pellio* (Linnaeus, 1758)
- schaefferi schaefferi* (Herbst, 1792)
- smirnovi* Zhantiev, 1973
- unicolor unicolor* (Brahm, 1791)
megatoma (Fabricius, 1798)
piceus (Olivier, 1790 nec Thunberg, 1781)
- Megatomini** Ganglbauer, 1904
- Trogoderma** Dejean, 1821
- angustum* (Solier, 1849)
- glabrum* (Herbst, 1783)
nigrum (Herbst, 1797)
- variabile* Ballion, 1878
- versicolor* (Creutzer, 1799)
- Globicornis** Latreille, 1829
- emarginata* (Gyllenhal, 1808)
marginata (Paykull, 1798 nec Thunberg, 1781)
- Megatoma** Herbst, 1792
- undata undata* (Linnaeus, 1758)
- Ctesiini** Rees, 1943
- Ctesias** Stephens, 1830
- serra* (Fabricius, 1792)
- Anthrenini** Casey, 1900
- Anthrenus** (*Anthrenus*) O.F.Müller, 1764
- pimpinellae pimpinellae* Fabricius, 1775
- scrophulariae scrophulariae* (Linnaeus, 1758)
- Anthrenus** (*Florilinus*) Mulsant, Rey, 1868
- museorum* (Linnaeus, 1761)
- olgae* Kalik, 1946
- Anthrenus** (*Helocerus*) Mulsant, Rey, 1868
- fuscus* Olivier, 1789
claviger (Erichson, 1846)
- polonicus* Mroczkowski, 1951
- Trinodinae** Casey, 1900
- Trinodini** Casey, 1900
- Trinodes** Dejean, 1821
- hirtus* (Fabricius, 1781)
- Bostrichidae** Latreille, 1802
- Dinoderinae** Thomson, 1863
- Rhyzopertha** Stephens, 1830
- dominica* (Fabricius, 1792)
- Stephanopachys** Waterhouse, 1888
- linearis* (Kugelann, 1792)
- Note: Insufficiently known species, known
- only from a few localities in old and middle-aged boreal forests. Officially protected species in Latvia.
- *substriatus* (Paykull, 1800)
- Note: Officially protected species in Latvia.
- Bostrichinae** Latreille, 1802
- Bostrichus** O.F.Müller, 1764
- capucinus* (Linnaeus, 1758)
- Note: Very rare species with no confirmed records in the last 100 years.
- Lyctidae** Billberg, 1920
- Lyctus** Fabricius, 1792
- linearis* (Goeze, 1777)
- Note: Very rare and insufficiently known species.
- Anobiidae** Kirby, 1837
- Hedobiinae** Mulsant, Rey, 1868
- Ptinomorphus** Mulsant, Rey, 1868
- imperialis* (Linnaeus, 1767)
- Dryophilinae** LeConte, 1861
- Dryophilus** Chevrolat, 1832
- pusillus* (Gyllenhal, 1808)
- Xestobium** Motschulsky, 1845
- rufovillosum* (DeGeer, 1774)
- Ernobiinae** Pic, 1912
- Ernobius** Thomson, 1859
- abietinus* (Gyllenhal, 1808)
- abietis* (Fabricius, 1792)
- angusticollis* (Ratzeburg, 1847)
tabidus Kiesenwetter, 1877
- explanatus* (Mannerheim, 1843)
- kiesenwetteri* Schilsky, 1898
- Note: Record needs confirmations.
- longicornis* (Sturm, 1837)
densicornis (Mulsant, Rey, 1863)
- mollis* (Linnaeus, 1758)
- nigrinus* (Sturm, 1837)
- pini* (Sturm, 1837)
- Anobiinae** Kirby, 1837
- Stegobium** Motschulsky, 1860
- paniceum* (Linnaeus, 1758)
- Gastrallus** Jacquelin du Val, 1860
- immarginatus* (P.W.J.Müller, 1821)
- Anobium** Fabricius, 1775
- punctatum* (DeGeer, 1774)
- Cacotemnus** LeConte, 1861
- rufipes* (Fabricius, 1792)
- thomsoni* Kraatz, 1881
- Hemicoelus** LeConte, 1861
- nitidus* Fabricius, 1792

Note: Probably correct name for this species is <i>canaliculatus</i> (Thomson, 1863).	Ptinus Linnaeus, 1767
Microbregma Seidlitz, 1889	<i>bicinctus</i> Sturm, 1837
<i>emarginatum</i> (Duftschmid, 1825)	<i>clavipes</i> Panzer, 1792
Hadrobregmus Thomson, 1859	<i>testaceus</i> Olivier, 1790 nec Thunberg, 1784
<i>confusus</i> (Kraatz, 1881)	Note: Introduced species.
<i>denticollis</i> (Creutzer, 1796)	<i>dubius</i> Sturm, 1837
<i>pertinax</i> (Linnaeus, 1758)	<i>fur</i> (Linnaeus, 1758)
Priobium Motschulsky, 1845	<i>pilosus</i> P.W.J.Müller, 1821
<i>carpini</i> (Herbst, 1793)	<i>raptor</i> Sturm, 1837
Ptilininae Shuckard, 1840	<i>rufipes</i> Olivier, 1790
Ptilinus Müller, 1764	<i>sexpunctatus</i> Panzer, 1795
<i>fuscus</i> Geoffroy, 1785	<i>subpilosus</i> Sturm, 1837
<i>pectinicornis</i> (Linnaeus, 1758)	<i>villiger</i> Reitter, 1884
Xyletininae Gistel, 1856 / Thomson, 1859?	<i>balticus</i> Iablokoff-Khnzorian, Karapetian, 1991
Xyletinus Latreille, 1809	Lymexyloidea
<i>ater</i> (Creutzer, 1796)	Lymexylidae Fleming, 1821
<i>fabyensis</i> Lundblad, 1949	Hylecoetinae Böving, Craighead, 1931
<i>groenblomi</i> Y.Kangas, 1955	Hylecoetus Latreille, 1806
<i>laticollis</i> (Duftschmid, 1825)	<i>dermestoides</i> (Linnaeus, 1761)
<i>longitarsis</i> Jansson, 1942	<i>flabellicornis</i> (Schneider, 1791)
<i>pectinatus</i> (Fabricius, 1792)	Lymexylinae Fleming, 1821
<i>planicollis</i> Lohse, 1957	Lymexylon Fabricius, 1775
<i>suecicus</i> Lundberg, 1972	<i>navale</i> (Linnaeus, 1758)
Lasioderma Stephens, 1835	Note: This species is included on the list of indicator species of natural forest habitats.
<i>serricorne</i> (Fabricius, 1792)	Cleroidea
Dorcatominae Thomson, 1859	Trogossitidae Latreille, 1802
Stagetus Wollaston, 1861	Peltinae Kirby, 1837
<i>borealis</i> Israelsson, 1971	Peltini Latreille, 1807
Dorcatoma Herbst, 1792	Peltis O.F.Müller, 1764
<i>chrysomelina</i> Sturm, 1837	<i>grossa</i> (Linnaeus, 1758)
<i>dresdensis</i> Herbst, 1792	Note: This species is included on the list of indicator species of natural forest habitats.
<i>flavicornis</i> (Fabricius, 1792)	Ostoma Laicharting, 1781
<i>janssoni</i> Bueche, Lundberg, 2001	<i>ferruginea</i> (Linnaeus, 1758)
<i>punctulata</i> Mulsant, Rey, 1864	Thymalus Latreille, 1802
<i>robusta</i> Strand, 1938	<i>limbatus</i> (Fabricius, 1787)
<i>substriata</i> Hummel, 1829	Note: This species is included on the list of indicator species of natural forest habitats.
<i>serra</i> (Panzer, 1796 nec Fabricius, 1792)	Lophocaterinae Crowson, 1964
Caenocara Thomson, 1859	Lophocaterini Crowson, 1964
<i>affinis</i> (Sturm, 1837)	Grynocharis Thomson, 1859
<i>bovistae</i> (Hoffmann, 1803)	<i>oblonga</i> (Linnaeus, 1758)
Anitys Thomson, 1863	Note: This species is included on the list of indicator species of natural forest habitats.
<i>rubens</i> (Hoffmann, 1803)	Calitinae Reitter, 1922
<i>cognata</i> Mulsant, Rey, 1864	Gymnochilini Lacordaire, 1854
Ptinidae Latreille, 1802	Calitys Thomson, 1859
Ptininae Latreille, 1802	<i>scabra</i> (Thunberg, 1784)
Ptinini Latreille, 1802	
Niptus Boieldieu, 1856	
<i>hololeucus</i> (Faldermann, 1836)	
Epauloecus Mulsant, Rey, 1868	
<i>unicolor</i> (Piller, Mitterpacher, 1783)	

Note: Very rare species. This species is included on the list of indicator species of natural forest habitats.

Trogossitinae Latreille, 1802

Nemozoma Latreille, 1804

elongatum (Linnaeus, 1761)

Note: Very rare and insufficiently known species.

Tenebroides Piller, Mitterpacher, 1783

mauritanicus (Linnaeus, 1758)

Cleridae Latreille, 1802

Tillinae Leach, 1815

Tillus Olivier, 1790

elongatus (Linnaeus, 1758)

ater (Fabricius, 1792)

aterrimus Eschscholtz, 1818

coeruleus (Voet, 1806)

marchiae (Gmelin, 1788)

niger Ragusa, 1896

ruficollis (Herbst, 1784)

Clerinae Latreille, 1802

Opilo Latreille, 1802

mollis (Linnaeus, 1758)

cruciger (Geoffroy, 1785)

fuscofasciatus (DeGeer, 1775)

Note: Very rare species. Note: This species is included on the list of indicator species of natural forest habitats.

Thanasimus Latreille, 1806

femoralis (Zetterstedt, 1828)

pectoralis (Fuss, 1863)

rufipes (Brahm, 1797 nec DeGeer, 1775)

substriatus Heyden, 1864

formicarius (Linnaeus, 1758)

assimilis Méquignon, 1948

brevicollis (Spinola, 1844)

fasciatus (Geoffroy, 1785)

hilaris Buxson, 1901

laterufus Corporaal, 1931

schaeferi Méquignon, 1949

Trichodes Herbst, 1792

apiarius (Linnaeus, 1758)

angusticollis Pic, 1902

apicita Spinola, 1844

apivore Walckenaer, 1802

corallinus Ménétriés, 1832

crassipedarius Chevrolat, 1876

elegans Spinola, 1844

georgianus Chevrolat, 1874

pannonicus Spinola, 1844

suturalis Trella, 1924

Korynetinae Laporte de Castelnau, 1840

Korynetes Herbst, 1792

caeruleus (DeGeer, 1775)

cyanellus Dejean, 1837

geoffroyanus (Laicharting, 1783)

meridionalis Obenberger, 1916

nigrocoeruleus (Geoffroy, 1799)

obenbergeri Jansson, 1936

unicolor Chevrolat, 1876

violaceus (Linnaeus, 1758)

Necrobia Olivier, 1795

ruficollis (Fabricius, 1775)

rufipes (DeGeer, 1775)

amethystina Stephens, 1832

aspera Walker, 1858

cupreonitens Lauffer, 1905

dermestoides (Piller, Mitterpacher, 1783)

flavipes (Klug, 1842)

foveicollis Schenkling, 1900

glabra Jurin apud Champollion, 1814

mumiарum Hope, 1834

pilifera Reitter, 1894 (female)

reticulata (Klug, 1842)

violacea (Linnaeus, 1758)

angustata (Faldermann, 1835)

chalybea (Sturm, 1837)

dalmatina (Obenberger, 1916)

errans Melsheimer, 1846

jablanicensis (Obenberger, 1916)

quadra (Marsham, 1802)

Dasytidae Laporte de Castelnau, 1840

Rhadalinae LeConte, 1861

Aplocnemus Stephens, 1830

nigricornis (Fabricius, 1792)

chalybaeus (Germar, 1824)

femoralis (Illiger, 1807)

punctatus (Germar, 1824)

variolatus A.Costa, 1882

vierteli Schilsky, 1894

tarsalis (C.R.Sahlberg, 1822)

küsteri Schilsky, 1894

rufitarsis (C.R.Sahlberg, 1822)

Dasytinae Laporte de Castelnau, 1840

Danacaeinae Thomson, 1859

Danacaea Laporte de Castelnau, 1836

— *pallipes* (Panzer, 1793)

ambigua Prochazka, 1894 nec Mulsant, Rey, 1868

Dasytinae Jakobson, 1911

Dasytes Paykull, 1799

cyaneus (Fabricius, 1775)

caeruleus (DeGeer, 1774 nec Linnaeus, 1758)

fusculus (Illiger, 1801)

femoralis Krynicki, 1832

pallipes Faldermann, 1832

niger (Linnaeus, 1761)

palustris Fiori, 1906

villosum (Olivier, 1790)

obscurus Gyllenhal, 1813

<i>letzneri</i> Weise, 1887	<i>Anthocomus</i> Erichson, 1840
<i>rugipennis</i> Thomson, 1869	<i>equestris</i> (Fabricius, 1781)
<i>plumbeus</i> (O.F.Müller, 1776)	<i>fasciatus</i> (Linnaeus, 1758)
<i>atrides</i> Deville, 1908	<i>rufus</i> (Herbst, 1784)
<i>coerulescens</i> Schilsky, 1898	<i>coccineus</i> (Schaller, 1783 nec Linnaeus, 1761)
<i>coxalis</i> Mulsant, Rey, 1868	
<i>flavipes</i> (Fabricius, 1781)	Laiini Jakobson, 1913
<i>nigrinus</i> Schilsky, 1898	Paratinus Abeille de Perrin, 1891
<i>nigrofemoralis</i> Schilsky, 1898	<i>femoralis</i> (Erichson, 1840)
<i>pueli</i> Fagniez, 1946	Cucujoidea
<i>ragusae</i> Schilsky, 1894	Kateretidae Erichson, 1843
<i>subaeneus</i> Schönherr, 1817	Kateretes Herbst, 1793
<i>aeneus</i> Olivier, 1790	<i>pedicularis</i> (Linnaeus, 1758)
<i>scaber</i> Suffrian, 1843	<i>pusillus</i> (Thunberg, 1794)
<i>virens</i> Marsham, 1802	<i>bipustulatus</i> (Payk., 1798 nec Thunberg, 1781)
<i>flavipes</i> (Olivier, 1790) nec (Fabricius, 1781)	<i>rufilabris</i> (Latreille, 1807)
<i>puncticollis</i> Reitter, 1888 nec Germar, 1855	<i>Heterhelus</i> Jacquel du Val, 1858
<i>tibialis</i> Zetterstedt, 1828 nec Solier, 1849	<i>scutellaris</i> (Heer, 1841)
<i>Dolichosoma</i> Stephens, 1830	<i>Brachypterus</i> Kugelann, 1794
<i>lineare</i> (Rossi, 1792)	<i>glaber</i> (Stephens, 1835)
<i>filiforme</i> Panzer, 1799	<i>urticae</i> (Fabricius, 1792)
<i>Malachiidae</i> Fleming, 1821	<i>erythropus</i> (Marsham, 1802)
<i>Malachiinae</i> Fleming, 1821	<i>Brachypterolus</i> Grouvelle, 1913
<i>Colotini</i> Abeille de Perrin, 1890	<i>linariae</i> (Stephens, 1830)
<i>Charopus</i> Erichson, 1840	<i>cornelli</i> Spornraft, 1966
<i>graminicola</i> (Dejean, 1833)	<i>pulicarius</i> (Linnaeus, 1758)
<i>flavipes</i> (Paykull, 1798 nec Fabricius, 1787)	Nitidulidae Latreille, 1802
<i>Attalini</i> Abeille de Perrin, 1890	<i>Epuraeinae</i> Kirejtshuk, 1986
<i>Hypebaeus</i> Kiesenwetter 1863	<i>Epuraeini</i> Kirejtshuk, 1986
<i>flavipes</i> (Fabricius, 1787)	<i>Epuraea</i> (<i>Epuraea</i>) Erichson, 1843
Note: This species is known only from Pededze river valley Nature protection area (E Latvia). North border of main distribution area.	<i>aestiva</i> (Linnaeus, 1758)
<i>Ebaeus</i> Erichson, 1840	<i>depressa</i> (Illiger, 1798)
<i>lapplandicus</i> Evers, 1993	<i>angustula</i> Sturm, 1844
<i>pedicularius</i> sensu Evers, 1993	<i>biguttata</i> (Thunberg, 1784)
<i>Nepachys</i> Thomson, 1859	<i>unicolor</i> (Olivier, 1790)
<i>cardiacae</i> (Linnaeus, 1761)	<i>x-rubrum</i> J.R.Sahlberg, 1911
Note: Very rare species, known mostly from southern part of Latvia. No confirmed records in the last 50 years.	<i>binotata</i> Reitter, 1872
<i>Axinotarsus</i> Motschulsky, 1853	<i>boreella</i> (Zetterstedt, 1828)
<i>marginalis</i> (Laporte de Castelnau, 1840)	<i>contractula</i> J.R.Sahlberg, 1889
<i>pulicarius</i> (Fabricius, 1775)	<i>helleni</i> Sjöberg, 1939
<i>Malachiini</i> Fleming, 1821	<i>deleta</i> Erichson, 1843
<i>Cordylepherus</i> Evers, 1985	<i>distincta</i> (Grimmer, 1841)
<i>viridis</i> (Fabricius, 1792)	<i>hilleri</i> Reitter, 1877
<i>Malachius</i> Fabricius, 1775	<i>castanea</i> (Duft., 1825 nec R.F.Sahlberg, 1820)
<i>aeneus</i> (Linnaeus, 1758)	<i>fageticola</i> Audisio, 1991
<i>bipustulatus</i> (Linnaeus, 1758)	<i>longula</i> Erichson, 1845
<i>Clanoptilus</i> Motschulsky, 1854	<i>marseuli</i> Reitter, 1872
— <i>elegans</i> (Olivier, 1790)	<i>acuta</i> Biström, 1977
<i>marginellus</i> (Olivier, 1790)	<i>bickhardti</i> Saint-Claire Deville, 1906
	<i>pusilla</i> (Illiger, 1798 nec Thunberg, 1794)
	<i>melina</i> (Erichson, 1843)
	<i>muehli</i> , Reitter, 1908
	<i>oblonga</i> (Herbst, 1793)

- danica* Sjöberg, 1939
opalizans J.R.Sahlberg, 1889
pallescens Stephens, 1832
abietina J.R.Sahlberg, 1889
floreæ Erichson, 1845
placida Mäklin, 1853
pygmaea (Gyllenhal, 1808)
rufomarginata (Stephens, 1830)
silacea (Herbst, 1784)
silesiaca Reitter, 1872
terminalis (Mannerheim, 1843)
adumbrata Mannerheim, 1852
tenenbaumi Sjöberg, 1939
thoracica Tournier, 1872
variegata (Herbst, 1793)
Epuraea (***Dadopora*** Thomson, 1859)
guttata (Olivier, 1811)
Epuraea (***Epuraeanella*** Crotch, 1874)
limbata (Fabricius, 1787)
neglecta (Heer, 1841)
Carpophilinae Erichson, 1843
Carpophilini Erichson, 1843
Carpophilus Stephens, 1830
hemipterus (Linnaeus, 1758)
Meligethinae Thomson, 1859
Pria Stephens, 1829
dulcamarae (Scopoli, 1763)
Meligethes (***Meligethes*** Stephens, 1830)
atratus (Olivier, 1790)
denticulatus denticulatus (Heer, 1841)
hebes Erichson, 1845
flavimanus flavimanus Stephens, 1830
lumbaris Sturm, 1845
Meligethes (***Clypeogethes*** Scholtz, 1932)
aeneus aeneus (Fabricius, 1775)
alpestris Heer, 1841
asperrimus Guillebeau, 1897
australis Küster, 1848
bonvouloiri Brisout de Barnevile, 1872
boops Easton, 1957
brassicae Reitter, 1875
californicus Reitter, 1871
cleominis Easton, 1959
coerulea (Marsham, 1802)
coeruleus Reitter, 1871
florilegulus (Geoffroy, 1785)
gracilis Brisout de Barnevile, 1863
latipes (Marsham, 1802)
minor Rey, 1889
moerens LeConte, 1857
mutatus Harold, 1868
nigricornis Stephens, 1830
nigrina (Marsham, 1802)
peristericus Roubal, 1943
- psyllus* (Herbst, 1784)
pubens Rey, 1889
rotundangulus Ganglbauer, 1899
rubripennis Reitter, 1871
ruficornis LeConte, 1859
rufimanus LeConte, 1857
semiaeneus Ganglbauer, 1899
subtilis Waltl, 1838
urticæ Stephens, 1830
viridipennis Motschulsky, 1866
atramentarius Forster, 1849
bidens Brisout de Barnevile, 1863
brunnicornis Sturm, 1845
— buyssoni Brisout de Barnevile, 1882
carinulatus Forster, 1849
coeruleovirens Forster, 1849
coracinus Sturm, 1845
czwalinai Reitter, 1871
difficilis (Heer, 1841)
egenus Erichson, 1845
exilis Sturm, 1845
gagathinus Erichson, 1845
haemorrhoidalis Forster, 1849
— immundus Kraatz, 1858
kunzei Erichson, 1845
 Note: Imported species probably with no persistent populations in Latvia.
lugubris Sturm, 1845
morosus Erichson, 1845
nigrescens Stephens, 1830
picipes Sturm, 1845
obscurus Erichson, 1843
distinctus Sturm, 1845
ochropus Sturm, 1845
ovatus Sturm, 1845
pedicularis (Gyllenhal, 1808)
viduatus (Heer, 1841)
subaeneus Sturm, 1845
sulcatus Brisout de Barnevile, 1863
sympyti (Heer, 1841)
tenebrosus Forster, 1849
viduatus (Heer, 1841)
tristis Sturm, 1845
umbrosus Sturm, 1845
viridescens (Fabricius, 1787)
Meligethes (***Acanthogethes*** Reitter, 1871)
solidus (Kugelann, 1794)
Meligethes (***Astylogethes*** Kirejtshuk, 1992)
corvinus Erichson, 1845
subrugosus (Gyllenhal, 1808)
caudatus Guillebeau, 1897
Nitidulinae Erichson, 1843

- Nitidulini** Erichson, 1843
Nitidula Fabricius, 1775
bipunctata (Linnaeus, 1758)
carnaria (Schaller, 1783)
rufipes (Linnaeus, 1767)
Omosita Erichson, 1843
colon (Linnaeus, 1758)
depressa (Linnaeus, 1758)
discoidea (Fabricius, 1775)
Soronia Erichson, 1843
grisea (Linnaeus, 1758)
punctatissima (Illiger, 1794)
Amphotis Erichson, 1843
marginata (Fabricius, 1781)
Ipidia (*Ipidia*) Erichson, 1843
binotata Reitter, 1875
 quadrimaculata (Quensel, 1790 nec Scop., 1772)
 quadrinotata (F., 1798 nec L.G.Scriba, 1793)
 quadriplagiata Biström, 1978
Pocadius Erichson, 1843
ferrugineus (Fabricius, 1775)
Thalycra Erichson, 1843
fervida (Olivier, 1790)
Cychramini Lacordaire, 1855
Cychramus Kugelann, 1794
luteus (Fabricius, 1787)
variegatus (Herbst, 1792)
 quadripunctatus (Herbst, 1792)
Strongylini Erichson, 1859
Cylloides Erichson, 1843
ater (Herbst, 1792)
Cryptarchinae Reitter, 1884
Cryptarchini Thomson, 1859
Cryptarcha Shuckard, 1839
strigata (Fabricius, 1787)
undata (Olivier, 1790)
 imperialis Fabricius, 1792
Glischrochilus (*Glischrochilus*) Reitter, 1873
quadripunctatus (Linnaeus, 1758)
Glischrochilus (*Librodor*) Reitter, 1884
grandis Tournier, 1872
hortensis (Geoffroy, 1785)
quadriguttatus (Fabricius, 1776)
Pityophagus Shuckard, 1839
ferrugineus (Linnaeus, 1761)
Cybocephalinae Jacquelín du Val, 1858
Cybocephalus (*Cybocephalus*) Erichson, 1844
politus (Gyllenhal, 1813)
Monotomidae Laporte de Castelnau, 1840
Rhizophaginae Redtenbacher, 1845
Cyanostolus Ganglbauer, 1899
aeneus (Richter, 1820)
Rhizophagus Herbst, 1793
bipustulatus (Fabricius, 1792)
cribratus Gyllenhal, 1827
depressus (Fabricius, 1792)
dispar (Paykull, 1800)
ferrugineus (Paykull, 1800)
grandis Gyllenhal, 1827
nitidulus (Fabricius, 1798)
parvulus (Paykull, 1800)
perforatus Erichson, 1845
picipes (Olivier, 1790)
Monotominae Laporte de Castelnau, 1840
Monotoma Herbst, 1793
angusticollis (Gyllenhal, 1827)
bicolor Villa, 1835
brevicollis Aubé, 1837
conicicollis Aubé, 1837
picipes Herbst, 1793
 brevipennis Kunze, 1839
Silvanidae Kirby, 1837
Silvaninae Kirby, 1837
Airaphilus Redtenbacher, 1858
elongatus (Gyllenhal, 1813)
Ahasverus Des Gozis, 1881
advena (Waltl, 1834)
Oryzaephilus Ganglbauer, 1899
mercator (Fauvel, 1889)
surinamensis (Linnaeus, 1758)
Silvanus Latreille, 1804
bidentatus (Fabricius, 1792)
 affinis Reitter, 1876
 sulcatum (Fabricius, 1792)
unidentatus (Olivier, 1790)
 gratiosus Motschulsky, 1863
 planum (herbst, 1797)
 siculus Stierlin, 1864
Silvanoprus Reitter, 1911
fagi (Guérin-Méneville, 1844)
Brontinae Erichson, 1845
Psammoecus Latreille, 1829
bipunctatus (Fabricius, 1792)
Uleiota Latreille, 1796
planata (Linnaeus, 1761)
Dendrophagus Schönherr, 1809
crenatus (Paykull, 1799)
Note: This species is included on the list of indicator species of natural forest habitats.
Cucujidae Latreille, 1802
Pediacus Shuckard, 1839
fuscus Erichson, 1845

- Cucujus** O.F.Müller, 1764
cinnaberinus (Scopoli, 1763)
 Note: Very rare species with only a few known localities in Western and Eastern Latvia. Officially protected species in Latvia.
- Bothrideridae** Erichson, 1845
Bothriderinae Erichson, 1845
Bothrideres Dejean, 1835
contractus (Geoffroy, 1765)
bipunctatus (Gmelin, 1790)
- Cryptophagidae** Kirby, 1837
Cryptophaginae Kirby, 1837
Telmatophilini Jacquelin du Val, 1858
Telmatophilus Heer, 1841
— brevicollis Aubé, 1862
caricis (Olivier, 1790)
schoenherrii (Gyllenhal, 1808)
typhae (Fallén, 1802)
- Cryptophagini** Kirby, 1837
Henoticus Thomson, 1868
serratus (Gyllenhal, 1808)
- Pteryngium** Reitter, 1887
crenatum (Fabricius, 1798)
- Cryptophagus (Micrambe)** Thomson, 1863
abietis (Paykull, 1798)
bimaculatus (Panzer, 1798)
villosus Heer, 1841
- Cryptophagus (Cryptophagus)** Herbst, 1792
acutangulus Gyllenhal, 1827
angustus Ganglbauer, 1899
parallelus Thomson, 1871 nec Brisout de Barnevile, 1863
- badius* Sturm, 1845
cellaris (Scopoli, 1763)
corticinus Thomson, 1863
dentatus (Herbst, 1793)
distinguendus Sturm, 1845
dorsalis R.F.Sahlberg, 1819
labilis Erichson, 1846
laticollis Lucas, 1846
affinis Sturm, 1845 nec R.F.Sahlberg, 1834
- lycoperdi* (Scopoli, 1763)
micaceus Rey, 1889
pilosus Gyllenhal, 1827
populi Paykull, 1800
pseudodentatus Bruce, 1934
quercinus Kraatz, 1852
saginatus Sturm, 1845
scanicus (Linnaeus, 1758)
schmidti Sturm, 1845
scutellatus Newman, 1834
- subfumatus* Kraatz, 1856
Spavius Motschulsky, 1844
glaber (Gyllenhal, 1808)
Antherophagus Dejean, 1821
nigricornis (Fabricius, 1787)
pallens (Linnaeus, 1758)
- Caenoscelini** Falcoz, 1929
Caenoscelis Thomson, 1863
subdeplanata Brisout de Barnevile, 1882
- Atomariinae** LeConte, 1861
Atomariini LeConte, 1861
Atomaria Stephens, 1830
affinis (R.F.Sahlberg, 1834)
analis Erichson, 1846
apicalis Erichson, 1846
atra (Herbst, 1793)
atrata Reitter, 1875
atricapilla Stephens, 1830
basalis Erichson, 1846
nitidula Erichson, 1846 nec Heer, 1841
- berolinensis* Kraatz, 1853
elongatula Erichson, 1846
- Note: This species is known only from old oak forest in Pededzes lejteces Nature protection area in eastern Latvia.
- fuscata* (Schönherr, 1808)
agnita Kangas, 1961
fusipes (Gyllenhal, 1808)
gutta Newman, 1834
hislopi Wollaston, 1857
impressa Erichson, 1846
lewisi Reitter, 1877
linearis Stephens, 1830
mesomela (Herbst, 1792)
morio Kolenati, 1846
nigrirostris Stephens, 1830
fuscicollis Mannerheim, 1852
- nigriventris* Stephens, 1830
nitidula (Marsham, 1802)
procerula Erichson, 1846
- Note: Insufficiently known species, known only from the area of lake Engures (W Latvia).
- pseudatra* Reitter, 1887
reitteri Lövendal, 1892
pulchra Erichson, 1846
prolixa auct. nec Erichson, 1846
pusilla (Paykull, 1798)
rubella Heer, 1871
berolinensis Kraatz, 1853
bicolor auct. nec Erichson, 1846
rubricollis Brisout de Barnevile, 1863

<i>ruficornis</i> (Marsham, 1802)	<i>tomentosus</i> (DeGeer, 1774)
<i>scutellaris</i> (Motschulsky, 1849)	Laemophloeidae Ganglbauer, 1899
<i>subangulata</i> J.F.Sahlberg, 1926	Laemophloeus Dejean, 1835
<i>testacea</i> (Stephens, 1830)	<i>muticus</i> (Fabricius, 1781)
<i>ruficornis</i> auct. nec (Marsham, 1802)	Cryptolestes Ganglbauer, 1899
<i>turgida</i> Erichson, 1846	<i>abietis</i> (Wankowicz, 1865)
<i>umbrina</i> (Gyllenhal, 1827)	<i>corticinus</i> (Erichson, 1846)
<i>sjoegergi</i> Palm, 1949	Note: This species is known from single locality in Eleja area (S Latvia) situated at the border with Lithuania. Probably should be more widely distributed in pine and mixed forests in the whole of Latvia.
<i>zetterstedti</i> (Zetterstedt, 1838)	<i>ferrugineus</i> (Stephens, 1831)
Ephistemini Leng, 1920	<i>emgei</i> Reitter, 1887
Ootypus Ganglbauer, 1899	Phalacridae Leach, 1815
<i>globosus</i> (Waltl, 1838)	Phalacrinae Leach, 1815
Ephistemus Stephens, 1829	Phalacrus Paykull, 1800
<i>globulus</i> (Paykull, 1798)	<i>brisouti</i> Rye, 1872
Curelius Casey, 1900	<i>hybridus</i> Flach, 1888
<i>exiguus</i> (Erichson, 1846)	<i>caricis</i> Sturm, 1807
Languriidae Crotch, 1873	<i>nigrinus</i> auct. nec Marsham, 1802
Xenoscelinae Ganglbauer, 1899	<i>championi</i> Guillebeau, 1892
Zavaljus Reitter, 1880	<i>suecicus</i> Palm, 1947
<i>brunneus</i> (Gyllenhal, 1808)	<i>substriatus</i> Gyllenhal, 1813
<i>fausti</i> Reitter, 1880	Olibrus Erichson, 1845
Erotylidae Latreille, 1802	<i>aeneus</i> (Fabricius, 1792)
Tritominae Curtis, 1834	<i>affinis</i> (Sturm, 1807)
Tritoma Fabricius, 1775	<i>baudueri</i> Flach, 1888
<i>bipustulata</i> Fabricius, 1775	<i>bicolor</i> (Fabricius, 1792)
<i>subbasalis</i> (Reitter, 1896)	<i>obscurus</i> Guillebeau, 1892
<i>jakowlewi</i> (Semenov, 1898)	<i>bimaculatus</i> Küster, 1848
Triplax (<i>Platichna</i> Herbst, 1793)	<i>corticalis</i> (Panzer, 1797)
<i>aenea</i> (Schaller, 1783)	<i>liquidus</i> Erichson, 1845
<i>russica</i> (Linnaeus, 1758)	<i>millefolii</i> (Paykull, 1800)
Triplax (<i>Platichna</i> Thomson, 1863)	<i>norvegicus</i> Munster, 1901
<i>rufipes</i> (Fabricius, 1775)	<i>pygmaeus</i> (Sturm, 1807)
<i>scutellaris</i> Charpentier, 1825	Stilbus Seidlitz, 1872
Dacninae Gistel, 1856	<i>atomarius</i> (Linnaeus, 1767)
Dacne Latreille, 1796	<i>oblongus</i> (Erichson, 1845)
<i>bipustulata</i> (Thunberg, 1781)	<i>testaceus</i> (Panzer, 1797)
Combocerus Bedel, 1868	Sphindidae Jacquelin du Val, 1860
<i>glaber</i> (Schaller, 1783)	Sphindinae Jacquelin du Val, 1860
Cerylonidae Billberg, 1820	Sphindus Dejean, 1821
Ceryloninae Billberg, 1820	<i>dubius</i> (Gyllenhal, 1808)
Cerylon Latreille, 1802	Arpidiphorus Dejean, 1821
<i>deplanatum</i> Gyllenhal, 1827	<i>orbiculatus</i> (Gyllenhal, 1808)
<i>fagi</i> Brisout de Barnevile, 1867	Corylophidae LeConte, 1852
<i>ferrugineum</i> Stephens, 1830	Corylophinae LeConte, 1852
<i>histeroides</i> (Fabricius, 1792)	Corylophini LeConte, 1852
<i>impressum</i> Erichson, 1845	Corylophus Stephens, 1835
Byturidae Jacquelin du Val, 1858	<i>cassidoides</i> (Marsham, 1802)
Byturinae Jacquelin du Val, 1858	
Byturus Latreille, 1796	
<i>ochraceus</i> (L.G.Scriba, 1790)	
<i>aestivus</i> auct. nec (Linnaeus, 1758)	

- politus* Reitter, 1898
- Orthoperini** Jacquelin du Val, 1857
- Orthoperus** Stephens, 1829
- atomus* (Gyllenhal, 1808)
- atomarius* auct. nec Heer, 1841
 - picea* Stephens, 1829
- brunnipes** (Gyllenhal, 1808)
- kluki* Wankowicz, 1865
- mundus** Matthews, 1885
- improvisus* Bruce, 1946
- nigrescens** Stephens, 1829
- atomus* Stephens, 1829
 - coriaceus* auct. nec Mulsant, Rey, 1861
 - truncatus* Stephens, 1829
- punctatus** Wankowicz, 1865
- punctulatus** Reitter, 1876
- Sericoderinae** Matthews, 1888
- Sericoderus** Stephens, 1829
- lateralis* (Gyllenhal, 1827)
- Endomychidae** Leach, 1815
- Mycetaeinae** Jacquelin du Val, 1857
- Mycetaeini** Jacquelin du Val, 1857
- Mycetaea** Stephens, 1830
- subterranea* (Fabricius, 1801)
- hirta* (Marsham, 1802 nec Schaefer, 1769)
- Leiestini** Thomson, 1863
- Leiesthes** Chevrolat, 1837
- seminigra* (Gyllenhal, 1808)
- Endomychinae** Leach, 1815
- Lycoperdinini** Redtenbacher, 1844
- Lycoperdina** (*Golgia*) Mulsant, 1846
- succincta* (Linnaeus, 1767)
- Mycetina** Mulsant, 1846
- cruciata* (Schaller, 1783)
- Endomychini** Leach, 1815
- Endomychus** Panzer, 1795
- coccineus* (Linnaeus, 1758)
- Coccinellidae** Latreille, 1807
- Epilachninae** Mulsant, 1846
- Madaini** Gordon, 1975
- Subcoccinella** Guérin-Méneville, 1844
- vigintiquatuorpunctata* (Linnaeus, 1758)
- Cynegetis** Dejean, 1836
- impunctata* (Linnaeus, 1767)
- Coccidulinae** Mulsant, 1846
- Coccidulini** Mulsant, 1846
- Coccidula** Gyllenhal, 1827
- rufa* (Herbst, 1783)
- scutellata* (Herbst, 1783)
- Scymninae** Mulsant, 1846
- Scymnini** Mulsant, 1846
- Scymnus** Kugelann, 1794
- abietis* (Paykull, 1798)
- ater* Kugelann, 1794
- auritus* Thunberg, 1795
- bipunctatus* (Kugelann, 1794)
- Note: Record needs confirmation.
- femoralis* Gyllenhal, 1827
- ferrugatus* (Moll, 1785)
- frontalis* (Fabricius, 1787)
- haemorrhoidalis* Herbst, 1797
- jakowlewi* Weise, 1892
- triangularis* J.R.Sahlberg, 1914
- limbatus* *limbatus* Stephens, 1832
- testaceus* auct. nec Motschulsky, 1837
- nigrinus* Kugelann, 1794
- pallipediformis apetzoides* Capra, Fürsch, 1967
- apetzii* auct. nec Mulsant, 1846
- rubromaculatus* (Goeze, 1777)
- suturalis* Thunberg, 1795
- Nephus** Mulsant, 1846
- bipunctatus* (Kugelann, 1794)
- quadrimaculatus* (Herbst, 1783)
- redtenbacheri* Mulsant, 1846
- Stethorini** Dobzhansky, 1924
- Stethorus** Weise, 1885
- punctillum* Weise, 1891
- Hyperaspini** Mulsant, 1846
- Hyperaspis** Dejean, 1836
- campestris* (Herbst, 1783)
- reppensis* (Herbst, 1783)
- Chilocorinae** Mulsant, 1846
- Platynaspini** Mulsant, 1846
- Platynaspis** Redtenbacher, 1843
- luteorubra* (Goeze, 1777)
- Chilocorini** Mulsant, 1846
- Chilocorus** Leach, 1815
- bipustulatus* (Linnaeus, 1758)
- renipustulatus* (L.G.Scriba, 1790)
- Exochomus** Redtenbacher, 1843
- nigromaculatus* (Goeze, 1777)
- flavipes* (Thunberg, 1781)
- quadripustulatus* (Linnaeus, 1758)
- Coccinellinae** Latreille, 1807
- Coccinula** Dobzhansky, 1925
- quatuordeimpustulata* (Linnaeus, 1758)
- Anisosticta** Dejean, 1836
- novemdecimpunctata* (Linnaeus, 1758)
- Tytthaspis** Crotch, 1874
- sedecimpunctata* (Linnaeus, 1761)
- Sospita** Mulsant, 1846
- vigintiguttata* (Linnaeus, 1758)

<i>Myzia</i> Mulsant, 1846	<i>brevicollis</i> (Thomson, 1868)
<i>oblongoguttata</i> (Linnaeus, 1758)	<i>consimilis</i> Mannerheim, 1844
<i>Myrrha</i> Mulsant, 1846	<i>hirtus</i> Gyllenhal, 1827
<i>octodecimguttata</i> (Linnaeus, 1758)	<i>minutus</i> (Linnaeus, 1767)
<i>Propylea</i> Mulsant, 1846	<i>nidicola</i> (Palm, 1944)
<i>quatuordecimpunctata</i> (Linnaeus, 1758)	<i>pseudominutus</i> (Strand, 1958)
<i>Calvia</i> Mulsant, 1846	<i>Enicmus</i> Thomson, 1859
<i>decemguttata</i> (Linnaeus, 1767)	<i>brevicornis</i> (Mannerheim, 1844)
<i>quatuordecimguttata</i> (Linnaeus, 1758)	<i>fungicola</i> Thomson, 1868
<i>quindecimguttata</i> (Fabricius, 1777)	<i>histrio</i> Joy, Tomlin, 1910
<i>Vibidia</i> Mulsant, 1846	<i>planipennis</i> Strand, 1940
<i>duodecimguttata</i> (Poda, 1761)	<i>rugosus</i> (Herbst, 1793)
<i>Halyzia</i> Mulsant, 1846	<i>testaceus</i> (Stephens, 1830)
<i>sedecimguttata</i> (Linnaeus, 1758)	<i>transversus</i> (Olivier, 1790)
<i>Psyllobora</i> Dejean, 1836	<i>Dienerella</i> Reitter, 1911
<i>vigintiduopunctata</i> (Linnaeus, 1758)	<i>argus</i> (Reitter, 1884)
<i>Anatis</i> Mulsant, 1846	<i>costulata</i> (Reitter, 1877)
<i>ocellata</i> (Linnaeus, 1758)	<i>elongata</i> (Curtis, 1830)
<i>Aphidecta</i> Weise, 1899	<i>filiformis</i> (Gyllenhal, 1827)
<i>obliterata</i> (Linnaeus, 1758)	<i>filum</i> (Aubé, 1850)
<i>Hippodamia</i> Mulsant, 1846	<i>Stephostethus</i> LeConte, 1878
<i>notata</i> (Laicharting, 1781)	<i>angusticollis</i> (Gyllenhal, 1827)
<i>septemmaculata</i> (DeGeer, 1775)	<i>angulatus</i> (Mannerheim, 1875)
<i>tredecimpunctata</i> (Linnaeus, 1758)	<i>kokujevi</i> (Semenov, 1898)
<i>undecimnotata</i> (Schneider, 1792)	<i>laticeps</i> (Belon, 1881)
<i>variegata</i> (Goeze, 1777)	<i>lardarius</i> (DeGeer, 1775)
<i>Coccinella</i> Linnaeus, 1758	<i>pandellei</i> (Brisout de Barneville, 1863)
<i>hieroglyphica</i> Linnaeus, 1758	<i>rugicollis</i> (Olivier, 1790)
<i>magnifica</i> Redtenbacher, 1843	<i>Thes</i> Semenov, 1910
<i>distincta</i> Faldermann, 1837 nec Herbst, 1793	<i>bergrothi</i> (Reitter, 1880)
<i>divaricata</i> auct. nec Olivier, 1808	<i>Cartodere</i> (<i>Cartodere</i> Thomson, 1859)
<i>quinquepunctata</i> Linnaeus, 1758	<i>constricta</i> (Gyllenhal, 1827)
<i>saucerottii lutshniki</i> Dobzhansky, 1917	<i>Cartodere</i> (<i>Aridius</i> Motschulsky, 1866)
<i>septempunctata</i> Linnaeus, 1758	<i>nodifer</i> (Westwood, 1839)
<i>trifasciata</i> Linnaeus, 1758	<i>Corticariinae</i> Curtis, 1829
Note: Rare and insufficiently known species	<i>Corticaria</i> Marsham, 1802
with only few records within last 100 years.	<i>abietorum</i> Motschulsky, 1867
<i>undecimpunctata</i> Linnaeus, 1758	<i>abietum</i> Motschulsky, 1867
<i>Oenopia</i> Mulsant, 1850	<i>elongata</i> (Gyllenhal, 1827)
<i>conglobata</i> (Linnaeus, 1758)	<i>ferruginea</i> Marsham, 1802
<i>Adalia</i> Mulsant, 1850	<i>fulva</i> (Comolli, 1837)
<i>bipunctata</i> (Linnaeus, 1758)	<i>impressa</i> (Olivier, 1790)
<i>conglomerata</i> (Linnaeus, 1758)	<i>lateritia</i> Mannerheim, 1844
<i>decempunctata</i> (Linnaeus, 1758)	<i>linearis</i> (Paykull, 1798)
<i>Harmonia</i> Mulsant, 1846	<i>rubripes</i> Mannerheim, 1844
<i>quadripunctata</i> (Pontoppidan, 1763)	<i>longicollis</i> (Zetterstedt, 1838)
<i>Latridiidae</i> Erichson, 1842	<i>longicornis</i> (Herbst, 1793)
<i>Latridiinae</i> Erichson, 1842	<i>polypori</i> J.R.Sahlberg, 1900
<i>Latridius</i> Herbst, 1793	<i>eppelsheimi</i> Reitter, 1886 nec Reitter, 1875
<i>anthracinus</i> Mannerheim, 1844	<i>pubescens</i> (Gyllenhal, 1827)
	<i>saginata</i> Mannerheim, 1844

<i>serrata</i> (Paykull, 1798)						
<i>umbilicata</i> (Beck, 1817)						
<i>Cortinicara</i> Johnson, 1975						
<i>gibbosa</i> (Herbst, 1793)						
<i>transversalis</i> (Gyllenhal, 1827)	1827	nec				
Mannerheim, 1844)						
<i>Corticarina</i> Reitter, 1880						
<i>fuscula</i> (Gyllenhal, 1827)						
<i>trifoveolata</i> (Redtenbacher, 1849)						
<i>lambiana</i> (Sharp, 1910)						
<i>latipennis</i> (J.R.Sahlberg, 1871)						
<i>obfuscata</i> Strand, 1937						
<i>similata</i> (Gyllenhal, 1827)						
<i>truncatella</i> (Mannerheim, 1844)						
<i>Melanophthalma</i> Motschulsky, 1866						
<i>curticollis</i> (Mannerheim, 1844)						
<i>aegyptiaca</i> Otto, 1978						
<i>crocata</i> (Mannerheim, 1844)						
<i>hortensis</i> (Mannerheim, 1844)						
<i>moraviaca</i> Reitter, 1875						
<i>sinica</i> Johnson, 1972						
<i>transversalis</i> (Mannerheim, 1844)	1844	nec				
Gyllenhal, 1827)						
<i>wollastoni</i> (Waterhouse, 1859)						
Tenebrionoidea						
Zopheridae Solier, 1834						
Colydiinae Erichson, 1842						
Colydiini Erichson, 1842						
<i>Colydium</i> Fabricius, 1792						
<i>filiforme</i> Fabricius, 1792						
Synchitini Erichson, 1845						
<i>Synchita</i> Hellwig, 1792						
<i>humeralis</i> (Fabricius, 1792)						
<i>Bitoma</i> Herbst, 1793						
<i>crenata</i> (Fabricius, 1775)						
Orthocerini Blanchard, 1845						
<i>Orthocerus</i> Latreille, 1796						
<i>clavicornis</i> (Linnaeus, 1758)						
— <i>crassicornis</i> (Erichson, 1845)						
Mycetophagidae Leach, 1815						
<i>Triphylus</i> Dejean, 1821						
<i>bicolor</i> (Fabricius, 1776)						
<i>immaculatus</i> Roubal, 1936						
<i>meralis</i> (Marsham, 1802)						
<i>pilosus</i> (Herbst, 1792)						
<i>punctatus</i> (Hellwig, 1792)						
<i>Litargus</i> Erichson, 1846						
<i>connexus</i> (Geoffroy, 1785)						
<i>bifasciatus</i> (Fabricius, 1787)						
<i>lunatus</i> (Fabricius, 1792)						
<i>marginalis</i> (Panzer, 1793)						
<i>mediojunctus</i> Pic, 1903						
<i>signatus</i> (Panzer, 1798)						
<i>Mycetophagus</i> Hellwig, 1792						
<i>ater</i> (Reitter, 1879)						
<i>jaroslawensis</i> (Semenov, 1898)						
Note: This species is only known from Ilgas area in extreme SE of Latvia at the border with Lithuania and Belarus'.						
<i>decempunctatus</i> Fabricius, 1801						
<i>rossicus</i> (Semenov, 1898)						
<i>fulvicollis</i> Fabricius, 1792						
<i>deubeli</i> (Mehely, 1890)						
<i>sexmaculatus</i> Ragusa, 1891						
— <i>irroratus</i> Reitter, 1879						
<i>multipunctatus</i> Fabricius, 1792						
<i>similis</i> (Marsham, 1802)						
<i>piceus</i> (Fabricius, 1776)						
<i>bosnicus</i> Apfelbeck, 1911						
<i>brunneus</i> Panzer, 1798						
<i>flavotinctus</i> Roubal, 1931						
<i>histrio</i> (C.R.Sahlberg, 1837)						
<i>humeralis</i> (Schilsky, 1888)						
<i>hungaricus</i> Papp, 1946						
<i>lunaris</i> Fabricius, 1801						
<i>punctulatus</i> (Schilsky, 1888)						
<i>undulatus</i> (Marsham, 1802)						
<i>variabilis</i> Hellwig, 1792						
<i>varius</i> (Marsham, 1802)						
<i>populi</i> Fabricius, 1798						
<i>quadriguttatus</i> P.W.J.Müller, 1821						
<i>bipustulatus</i> Melsheimer, 1846						
<i>bisignatus</i> Melsheimer, 1856						
<i>disjunctemaculatus</i> Roubal, 1936						
<i>pubescens</i> Stephens, 1830						
<i>variegatus</i> C.R.Sahlberg, 1837						
<i>quadripustulatus</i> (Linnaeus, 1761)						
<i>antemacularis</i> (Dalla Torre, 1879)						
<i>bipustulatus</i> (Schilsky, 1888)						
<i>boleti</i> (Herbst, 1783)						
<i>impustulatus</i> (Schilsky, 1888)						
<i>quadrimaculatus</i> (Schaller, 1783)						
<i>ruficollis</i> (Schilsky, 1889)						
<i>winteri</i> Reitter, 1911						
Note: This species is included on the list of indicator species of natural forest habitats.						
<i>salicis</i> Brisout de Barneville, 1862						
<i>tschitscherini</i> (Reitter, 1897 nec Semenov, 1898)						
<i>tschitscherini</i> (Semenov, 1898 nec Reitter, 1897)						
<i>Typhaea</i> Stephens, 1829						
<i>haagi</i> Reitter, 1874						
Note: This species is known by small serie from single locality in Vārkava area (E Latvia).						
<i>stercorea</i> (Linnaeus, 1758)						
<i>crenata</i> Melsheimer, 1846						
<i>fumata</i> (Linnaeus, 1767)						
<i>testacea</i> (Hellwig, 1792)						
<i>tomentosa</i> Stephens, 1830						
<i>variabilis</i> (Herbst, 1792)						

Ciidae Leach, 1819	<i>conjuncta</i> Pic, 1901
Ciinae Leach, 1819	<i>obscurior</i> Pic, 1901
Cis Laterille, 1796	<i>perrisi</i> Houlbert, 1934
<i>alter</i> Silfverberg, 1991	<i>pfefferi</i> Roubal, 1930
<i>nitidus</i> auct. nec (Fabricius, 1792)	
<i>bidentatus</i> (Olivier, 1790)	<i>fungorum</i> Fabricius, 1790
<i>boleti</i> (Scopoli, 1763)	<i>bicolor</i> (Marsham, 1802)
<i>castaneus</i> Mellie, 1848	<i>collaris</i> (Quensel, 1790)
<i>fuscoclavis</i> Nyholm, 1954	<i>dermestoides</i> Herbst, 1792
<i>comptus</i> Gyllenhal, 1827	
<i>dentatus</i> Mellie, 1848	Eustrophinae Gistel, 1856
<i>glabratus</i> Mellie, 1848	Eustrophus Latreille, 1809
<i>hanseni</i> Strand, 1965	<i>dermestoides</i> (Fabricius, 1792)
<i>hispidus</i> (Paykull, 1798)	<i>siculus</i> (Baudi di Selve, 1892)
<i>jacquemartii</i> Mellie, 1848	Hallomeninae Mulsant, 1856
<i>lineatocribratus</i> Mellie, 1848	Hallomenus Panzer, 1794
<i>micans</i> (Fabricius, 1792)	<i>axillaris</i> (Illiger, 1807)
<i>nitidus</i> (Fabricius, 1792)	<i>dimidiatus</i> Stephens, 1832
<i>punctulatus</i> (Gyllenhal, 1827)	<i>dorsalis</i> (Beck, 1817)
<i>quadridens</i> Mellie, 1848	<i>fuscus</i> Gyllenhal, 1810
<i>rugulosus</i> Mellie, 1848	<i>piceus</i> (Marsham, 1802)
<i>setiger</i> Mellie, 1848	<i>scapulatus</i> Fairmaire, 1881
<i>petropolitanus</i> Jakobson, 1895	<i>binotatus</i> (Quensel, 1790)
Ennearthron Mellie, 1847	<i>bipunctatus</i> Paykull, 1799
<i>cornutum</i> (Gyllenhal, 1827)	<i>boleti</i> (Kugelann, 1792)
<i>laricinum</i> (Mellie, 1848)	<i>elateroides</i> (Kugelann, 1792)
Orthocis Casey, 1898	<i>humeralis</i> Panzer, 1794
<i>alni</i> (Gyllenhal, 1813)	
<i>festivus</i> (Panzer, 1793)	Melandryidae Leach, 1815
<i>linearis</i> (J.R.Sahlberg, 1900)	Melandryinae Leach, 1815
<i>perrisi</i> auct. nec (Abeille de Perrin, 1874)	Orchesia Latreille, 1807
<i>pygmaeus</i> (Marsham, 1802)	<i>fasciata</i> (Illiger, 1798)
<i>oblongus</i> (Mellie, 1848)	<i>laterufa</i> Pic, 1927
<i>rhododactylus</i> (Marsham, 1802)	<i>trifasciata</i> (Zetterstedt, 1840)
<i>vestitus</i> (Mellie, 1848)	
Hadreule Thomson, 1859	<i>fusiformis</i> Solsky, 1871
<i>elongatula</i> (Gyllenhal, 1827)	<i>acicularis</i> Reitter, 1886
Sulcacis Dury, 1917	<i>nadeshdae</i> Semenov, 1898
<i>affinis</i> (Gyllenhal, 1827)	<i>picea</i> Seidlitz, 1891
<i>fronticornis</i> (Panzer, 1809)	<i>luteipalpis</i> Mulsant, Guillebeau, 1857
Orophiinae Thomson, 1863	<i>micans</i> (Panzer, 1794)
Ropalodontus Mellie, 1847	<i>abeillei</i> Guillebeau, 1888
<i>perforatus</i> (Gyllenhal, 1813)	<i>australis</i> Guillebeau, 1887
<i>strandii</i> Lohse, 1969	<i>clavicornis</i> (Latreille, 1804)
Octotemnus Mellie, 1847	<i>corsica</i> Guillebeau, 1888
<i>glabriculus</i> (Gyllenhal, 1827)	<i>rey</i> Guillebeau, 1887
<i>mandibularis</i> (Gyllenhal, 1813)	<i>subimpressa</i> Guillebeau, 1887
Tetratomidae Billberg, 1820	
Tetratominae Billberg, 1820	<i>minor</i> Walker, 1837
Tetratoma Fabricius, 1790	<i>sepicola</i> Rosenhauer, 1847
<i>ancora</i> Fabricius, 1790	<i>tetratoma</i> Thomson, 1864
	<i>undulata</i> Kraatz, 1853
	<i>fasciata</i> (Gyllenhal, 1810)
	Abdera Stephens, 1832
	<i>affinis</i> (Paykull, 1799)
	<i>flexuosa</i> (Paykull, 1799)
	<i>undata</i> (Panzer, 1799)
	Wanachia Schulze, 1912
	<i>triguttata</i> (Gyllenhal, 1810)
	<i>scutellaris</i> (Mulsant, 1856)
	<i>suturalis</i> (Baudi di Selve, 1877)

Dircaeaa Fabricius, 1798***— australis*** Fairmaire, 1856*imitata* Olexa, 1955*quadriguttata* Duftschmid, 1812 nec Paykull, 1798*quadrimaculata* (Illiger, 1798) in part*quadrisignata* Mulsant, 1856***quadriguttata*** Paykull, 1798 nec Duftschmid, 1812*quadrimaculata* (Illiger, 1798) in part*guttata* Stierlin, 1898***Phloiotrya*** Stephens, 1832***rufipes*** (Gyllenhal, 1810)*testaceipennis* (Pic, 1900)

Note: This species is known from one locality only (Slīteres National park, NW Latvia) but should be more widely distributed in western and eastern parts of Latvia.

Serropalpus Sellenius, 1786***barbatus*** (Schaller, 1783)*biguttatus* (Schellenberg, 1802)*niponicus* Lewis, 1895*striatus* Hellenius, 1786***Xylita*** Paykull, 1798***laevigata*** (Hellenius, 1786)*buprestoides* (Fabricius, 1793)*decolorata* (Randall, 1838)*discolor* (Fabricius, 1798)*lividipennis* Reitter, 1911*robusta* Motschulsky, 1872***Rushia*** Forel, 1893***parreyssi*** (Mulsant, 1856)*ferruginea* (Gredler, 1866)*mollis* (Graëlls, 1858)*paganettii* (Obenberger, 1917)*revelierei* (Mulsant, 1859)

Note: This species is known from one locality only (Slīteres National park, NW Latvia) but should be more widely distributed in western and eastern parts of Latvia.

Hypulus Paykull, 1798***quercinus*** (Quensel, 1790)*blekingensis* (Schneider, 1792)*dubius* (Fabricius, 1792)*undulatus* Pic, 1925***Zilora*** Mulsant, 1856***obscura*** (Fabricius, 1794)*sericea* (Sturm, 1807)***ferruginea*** (Paykull, 1798)

Note: Note: This species is known from single locality only (Cekule & Bunči areas, Central Latvia), but is no doubt more widely distributed in young and middle-aged pine forests of Latvia.

Melandrya Fabricius, 1801***caraboides*** (Linnaeus, 1761)*coerulescens* Petri, 1912*luridipes* Pic, 1907*rufibarbis* (Schaller, 1783)*serrata* (Fabricius, 1775)***dubia*** (Schaller, 1783)*canaliculata* (Fabricius, 1787)*caraboides* Latreille, 1818*rufipes* Chevrolat, 1844

Note: This species is included on the list of indicator species of natural forest habitats.

Phryganophilus R.F.Sahlberg, 1833***ruficollis*** (Fabricius, 1798)*analis* (Fabricius, 1792)*paradoxus* (Paykull, 1799)

Note: Very rare species. Officially protected species in Latvia.

Osphyinae Mulsant, 1856***Conopalpus*** Gyllenhal, 1810***testaceus*** (Olivier, 1790)*flavicollis* Gyllenhal, 1810 nec Peletier, Audinet-Serville, 1825*flavicollis* Peletier, Audinet-Serville, 1825 nec Gyllenhal, 1810*nigricornis* Germar, 1821*obscuriceps* Pic, 1925*thoracicus* Baudi di Selve, 1877*vigorsi* Stephens, 1832

Note: This species is known from single locality in Vīdale area (Slīteres National park, NW Latvia).

Incertae sedis***Dolotarsus*** Jacquelin du Val, 1863***lividus*** (R.F.Sahlberg, 1833)*ephippius* (Schaum, 1859)*rufipes* Jacquelin du Val, 1863*suturus* (Gredler, 1866)***Mordellidae*** Latreille, 1802***Mordellinae*** Latreille, 1802***Mordellini*** Latreille, 1802***Tomoxia*** Costa, 1854***bucephala*** Costa, 1854*biguttata* (Gyllenhal, 1827 nec Rossi, 1794)***Variimorda*** Méquignon, 1946***briantea*** (Comolli, 1837)***villosa*** (Schrank, 1781)*fasciata* (Fabricius, 1775 nec Forster, 1771)***Mordella*** Linnaeus, 1758***aculeata*** Linnaeus, 1758***holomelaena*** Apfelbeck, 1914***Hoshikananomia*** Kono, 1935***perlata*** (Sulzer, 1776)***Curtimorda*** Méquignon, 1946

maculosa (Naezen, 1794)

Mordellistenini Ermisch, 1941

Mordellistena Costa, 1854

brevicauda (Boheman, 1849)

humeralis (Linnaeus, 1758)

- *parvula* (Gyllenhal, 1827)

pseudodonana Ermisch, 1956

nana Ermisch, 1956 nec Motschulsky, 1860

pumila (Gyllenhal, 1810)

- *rufifrons* Schilsky, 1894

Note: The Type specimen of *M. rufifrons* Schilsky is identical with *M. humeralis* (L.) and has a label "*Mordellistena humeralis*, det. Horak". This synonymy is still unpublished and the right to publish it for the first time remains with Mr. J.Horak. I exclude this species from the Latvian beetle list.

variegata (Fabricius, 1798)

Mordellochroa Emery, 1876

abdominalis (Fabricius, 1775)

Rhipiphoridae Laporte de Castelnau, 1840

Pelecotominae Seidlitz, 1875

Pelecotoma Fischer von Waldheim, 1809

fennica (Paykull, 1799)

Note: Scarce species known only from single locality (Slīteres National park, NW Latvia) and with no confirmed records in the last 100 years.

Gemminger, Harold, 1870

Metocus Dejean, 1834

paradoxus (Linnaeus, 1761)

Stenotrachelidae Thomson, 1859

Stenotrachelinae Thomson, 1859

Scotodes Eschscholtz, 1818

annulatus Eschscholtz, 1818

Note: Rare and insufficiently known species.

Meloidae Gyllenhal, 1810

Melinae Gyllenhal, 1810

Lyttini LeConte, 1862

Lytta Fabricius, 1775

vesicatoria (Linnaeus, 1758)

Note: Very rare species, known from two localities in Central (Sigulda, Gaujas National park) and extreme SW (Papes ezers Nature park) Latvia. North border of main distribution area. Officially protected species in Latvia.

Alosimus Mulsant, 1857

- *syriacus* (Linnaeus, 1758)

Cerocomini Semenov, 1895

Cerocoma Geoffroy, 1762

schaefferi (Linnaeus, 1758)

Note: Scarce species with no confirmed records in the last 100 years. North border of main distribution area.

Meloini Blanchard, 1845

Meloe Linnaeus, 1758

brevicollis Panzer, 1793

proscarabaeus Linnaeus, 1758

violaceus Marsham, 1802

Nemognathinae Laporte de Castelnau, 1840

Sitarini Mulsant, 1857

Apalus Fabricius, 1775

bimaculatus (Linnaeus, 1761)

Note: Scarce species with no confirmed records in the last 50 years.

Oedemeridae Latreille, 1810

Nacerdinae Mulsant, 1858

Ditylini Mulsant, 1858

Ditylus Fischer von Waldheim, 1817

laevis (Fabricius, 1787)

helopoides Fischer von Waldheim, 1817

Chrysanthia W.Schmidt, 1844

geniculata (W.Schmidt, 1846)

cupreomicans Westhoff, 1881

nigricornis Westhoff, 1881

saturata Westhoff, 1881

viridis W.Schmidt, 1846 nec (DeGeer, 1775)

viridissima (Linnaeus, 1758)

korbi Reitter, 1894

thalassina (Fabricius, 1775)

viridis (DeGeer, 1775) nec W.Schmidt, 1846

Nacerdini Mulsant, 1858

Nacerdes (*Nacerdes*) Dejean, 1834

melanura (Linnaeus, 1758)

acuta (Marsham, 1802)

analis (Olivier, 1811)

apicalis (Say, 1835)

erminea (Germar, 1817)

italica (Chevrolat, 1877)

lepturoides (Thunberg, 1784)

nigripes (Fabricius, 1781)

notata (Fabricius, 1792)

particularis (Pic, 1924)

rufa (Brullé, 1838)

sardea W.Schmidt, 1846

sardoa Heyden, Reitter, Weisse, 1906

testacea (Geoffroy, 1785)

Anogcodes Dejean, 1834

rufiventris (Scopoli, 1763)

bipartitus (Schrank, 1776)

dorsalis (Olivier, 1811)

melanocephalus (Fabricius, 1794)

ustulatus (Scopoli, 1763)

fasciata (Villers, 1789)

- melanura* (Fabricius, 1787)
rufus (Fischer von Waldheim, 1820)
scutellaris (Waltl, 1839)
- Calopodinae** A.Costa, 1852
- Calopodini** A.Costa, 1852
- Calopus* Fabricius, 1775
serraticornis (Linnaeus, 1758)
pretneri G.Müller, 1929
- Oedemerinae** Latreille, 1810
- Asclerini** Gistel, 1856
- Ischnomera* Stephens, 1832
sanguinicollis (Fabricius, 1787)
flavicollis (Panzer, 1794)
sanguineicollis Gistel, 1856
- Oedemerini** Latreille, 1810
- Oedemera* (*Oedemera*) Olivier, 1789
croceicollis (Gyllenhal, 1827)
sarmatica Morawitz, 1861
femorata (Scopoli, 1763)
flavescens (Linnaeus, 1767)
pratterana (Schrank, 1781)
similis W.Schmidt, 1846
- flavipes* (Fabricius, 1792)
aenea (Olivier, 1795)
clavipes (Fabricius, 1801)
pusilla A.Costa, 1852
- lurida* (Marsham, 1802)
viridula Seidlitz, 1899
- podagrariae* (Linnaeus, 1767)
fulva (Geoffroy, 1785)
sebastiani Pic, 1901
sericans Mulsant, 1858
testacea (Fabricius, 1792)
- Note: Very rare species with only one record in the last 100 years. North border of main distribution area.
- virescens* (Linnaeus, 1767)
striata (Herbst in Fuessly, 1784)
- Pythidae** Blanchard, 1845
- Pythinae** Blanchard, 1845
- Pytho* Laterille, 1796
- depressus* (Linnaeus, 1767)
atriceps Pic, 1912
basipennis Pic, 1912
castaneus (Fabricius, 1792)
coeruleus (Herbst in Fuessly, 1782)
coeruleus (Fabricius, 1792)
festivus (Fabricius, 1792)
lignarius (DeGeer, 1775)
planus (Herbst, 1799)
- Pyrochroidae** Latreille, 1807
- Pyrochroa* Geoffroy 1762
coccinea (Linnaeus, 1761)
kabyliana Pic, 1898
purpurata Schrank, 1781
- rubra* DeGeer, 1775 (female)
serraticornis (Scopoli, 1763)
kiesenwetteri Fairmaire, 1849
purpurata O.F.Müller, 1788 nec Schrank, 1781
rubens (Schaller, 1783)
ruberrima Geoffroy, 1785
rutilans Voet, 1806
satrapa Schrank, 1781
tauricola Pic, 1912
- Schizotus** Newman, 1838
pectinicornis (Linnaeus, 1758)
rubra (DeGeer, 1775) (male)
- Scriptiidae** Mulsant, 1856
- Anaspinae** Mulsant, 1856
- Cyrtanaspis* Emery, 1876
phalerata (Germar, 1831)
- Anaspis* (*Anaspis*) Geoffroy, 1762
arctica Zetterstedt, 1828
bohemica Schilsky, 1898
norvegica Munster, 1924
- frontalis* (Linnaeus, 1758)
garneysi Fowler, 1889
lurida Stephens, 1832
marginicollis Lindberg, 1925
ruficollis (Fabricius, 1792)
bickhardti Schilsky, 1906
- thoracica* (Linnaeus, 1758)
- Anaspis* (*Nassipa*) Emery, 1876
flava (Linnaeus, 1758)
rufilabris (Gyllenhal, 1827)
- Anaspis* (*Silaria*) Mulsant, 1856
brunnipes (Mulsant, 1856)
- Scriptiinae** Mulsant, 1856
- Scraptia* Latreille, 1807
fuscula P.W.J.Müller, 1821
innata (Kangas, 1959)
- Salpingidae** Leach, 1815
- Salpinginae** Leach, 1815
- Sphaeriestes* Stephens, 1829
bimaculatus (Gyllenhal, 1810)
castaneus (Panzer, 1796)
achilleae Bonelli, 1812
brunnescens Pic, 1892
immaculatus Stephens, 1831
piceae Germar, 1824
- stockmanni* (Biström, 1977)
ater (Paykull, 1798 nec DeGeer, 1774)
- Rabocerus** Mulsant, 1859
foveolatus (Ljungh, 1823)
bishopi Sharp, 1909
impressithorax Pic, 1903
mutilatus (Champion, 1886)
- gabrieli* (Gerhardt, 1901)

foveolatus (Ljungh, 1823) in part
Colposis Mulsant, 1859
mutilatus (Beck, 1817)
maritimus Pic, 1903
virescens Mulsant, 1859
Note: North border of main distribution area.
Cariderus Mulsant, 1859
aeneus (Olivier, 1807)
megricus Iablokoff-Khnzorian, 1956
numidicus Pic, 1904
ornithorrhynchus Abeille de Perrin, 1874
planirostris (Fabricius, 1787)
fulvirostris (Fabricius, 1787)
luteonitens Fairmaire, 1879
quadriimpressus Desbrochers des Loges, 1900
spinolae Costa, 1847
Rhinosimus Latreille, 1802
ruficollis (Linnaeus, 1761)
bicolor Stephens, 1831
coeruleocephalus Latreille, 1804
oboris (Fabricius, 1787)
ostratus (DeGeer, 1775)
viridicollis Pic, 1892
Mycteridae Blanchard, 1845
Mycterinae Blanchard, 1845
Mycterus Clairville, 1798
curculioides (Fabricius, 1781)
gracilior Fairmaire 1874
griseus (Gistel 1856)
griseus Clairville 1798
Note: Scarce species with no confirmed records in the last 50 years.
Boridae Thomson, 1859
Borinae Thomson, 1859
Boros Herbst, 1797
schneideri (Panzer, 1795)
boros (Fabricius, 1801)
corticalis Paykull, 1798
elongatus Herbst, 1797
Note: Very rare species recently known only from a few localities in old broad-leaved forests of Eastern and Northern Latvia. Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.
Anthicidae Latreille, 1819
Anthicinae Latreille, 1819
Notoxini Stephens, 1829
Notoxus Geoffroy, 1762
monoceros *monoceros* (Linnaeus, 1760)
biconjunctus Pic, 1900
biinterruptus Pic, 1900
cucullatus Geoffroy, 1785
immaculatus Pic, 1899

integer LaFerté-Sénectère, 1849a
integer LaFerté-Sénectère, 1849b
latemaculatus Pic, 1892
mongolicus Pic, 1914
obscuricollis Pic, 1916
platycerus LaFerté-Sénectère, 1849a
platycerus LaFerté-Sénectère, 1849b
septemmaculata Roubal, 1933
subjuncta Pic, 1911
subobliteratus Pic 1903
unicornis Westhoff, 1882
Formicomini Bonadona, 1974
Anthelephila Hope, 1834
pedestris (Rossi, 1790)
atratulus (Reitter, 1889)
brevipilis (Desbrochers des Loges, 1875)
cursor (LaFerté-Sénectère, 1849a)
cursor (LaFerté-Sénectère, 1849b)
equestris (Panzer, 1800)
fusca (Geoffroy, 1799)
nobilis (Faldermann, 1837)
sareptanus (Desbrochers des Loges, 1875)
thomsoni (Pic, 1901)
thoracicus (Panzer, 1794)
Note: This species is known from a single individual only (Cekule area, Central Latvia). North border of main distribution area.
Anthicini Latreille, 1819
Omonadus Mulsant, Rey, 1866
floralis (Linnaeus, 1758)
basalis (Villa, Villa, 1838)
basillaris (Say, 1824)
breviculus (Philippi, 1864)
calycinus (Panzer, 1793)
fallax (Broun, 1893)
formicarius (Olivier, 1795)
formicoides (Geoffroy, 1785)
fusca (Marsham, 1802)
massauensis (Pic, 1900)
myrmicocephalus (Rossi, 1792)
pedicularius (Schrank, 1781)
reducteapicalis (Pic, 1915)
semirufus (Fairmaire, Germain, 1860)
syriacus (Baudi di Selve, 1881)
formicarius *formicarius* (Goeze, 1777)
enodis (Casey, 1895)
picianus (Koch, 1931)
quisquilius (Thomson, 1864)
rixator (Casey, 1895)
scenicus (Casey, 1895)
semirufus (Fairmaire, 1896)
Cordicomus Pic, 1894
sellatus (Panzer, 1796)
subfumosus (Reitter, 1911)
Stricticomus Pic, 1894
tobias (Marseul, 1879)
binhanus (Pic, 1927)

corporaali (Pic, 1923)
mauritiensis (Pic, 1898)
mundulus (Sharp, 1885)
parisiensis (Saint-Albin, 1952)
postoculatus (Fairmaire, 1896)
tanakai (Nomura, 1960)
turanicus (Reitter, 1889)

Hirticomus Pic, 1894

hispidus (Rossi, 1792)
bicolor (Olivier, 1795)
hirtellus (Panzer, 1796)

Note: This species is known only from the Abava river valley Nature protection area (Central Latvia), but it should be more widely distributed in Western and South-Western parts of Latvia. Northeast border of main distribution area.

Anthicus Paykull, 1798

antherinus (Linnaeus, 1761)
astrachanicus Csiki, 1901
cinctellus (Rossi, 1792)
semitestaceus Pic, 1892
syriae Pic, 1892
tripustulatus (Fabricius, 1792)
valens Pic, 1896
ater ater (Thunberg, 1787 nec Panzer, 1796)
ater (Panzer, 1796 nec Thunberg, 1787)
axillaris Schmidt, 1842
inflatus Kolenati, 1846
varians Kolenati, 1846
bimaculatus (Illiger, 1801)
dauricus Motschulsky, 1845
fasciatus Schilsky, 1888
pallens Schilsky, 1888
pallescens Pic, 1894
sagitta Krynicki, 1829
schilskyi Pic, 1894
flavipes flavipes (Panzer, 1796)
flavescens Pic, 1899
nigriceps Mannerheim, 1843
obscurus Küster, 1848
rufipes Paykull, 1800
scoticus Rye, 1872

Note: A. *scoticus* Rye is a newly recognised and currently unpublished synonym of *A. flavipes flavipes* (Pz.). The synonymy will be published by D.Telnov in Chandler D., Nardi G. & Telnov D. "Nomenclatorial Notes on the Palaearctic Anthicidae(Coleoptera)" (in print).

— *inaequalis* Marseul, 1879

umbrinus LaFerté-Sénectère, 1849a
setulosus Boheman, 1851
umbrinus LaFerté-Sénectère, 1849b

Note: This species is known from only a single

record (Rīga city, Central Latvia), but it should be more widely distributed in northern and eastern parts of the country. South-west border of main distribution area.

Aderidae Winkler, 1927

Aderinae Winkler, 1927

Phytobaenini Baguena, 1948

Phytobaenus R.F.Sahlberg, 1834

amabilis R.F.Sahlberg, 1834

Aderini Winkler, 1927

Aderus Westwood, 1829

populneus (Creutzer, 1796)

Otoleus Mroczkowski, 1987

— *pruinosus* (Kiesenwetter, 1861)

Euglenesini Seidlitz, 1875

Euglenes Westwood, 1830

oculatus (Paykull, 1798)

pygmaeus (DeGeer, 1775)

nitidifrons (Thomson, 1886)

Anidorus Mulsant, Rey, 1866

nigrinus (Germar, 1831)

Tenebrionidae Latreille, 1802

Lagriinae Latreille, 1825

Lagria Fabricius, 1775

hirta (Linnaeus, 1758)

depilis Mulsant, 1856

flava Geoffroy, 1758

glabrata Fabricius, 1775

lurida Krynicki, 1832

pubescens Linnaeus, 1767

nudipennis Mulsant, 1856

pontica Motschulsky, 1860

seminuda Reitter, 1889

spadicea Scopoli, 1763

villosa DeGeer, 1775

Alleculinae Mulsant, 1854

Alleculini Mulsant, 1854

Allecula Fabricius, 1801

morio (Fabricius, 1787)

Prionychus Solier, 1835

ater (Fabricius, 1775)

Note: This species is included on the list of indicator species of natural forest habitats.

Hymenalia Mulsant, 1851

rufipes (Fabricius, 1792)

Mycetocharini Mulsant, 1856

Mycetochara (*Mycetochara* Berthold, 1827)

axillaris (Paykull, 1799)

flavipes (Fabricius, 1792)

humeralis (Fabricius, 1787)

linearis (Illiger, 1794)

obscura (Zetterstedt, 1838)

Gonoderini Seidlitz, 1896**Pseudocistela** Crotch, 1873*cerambooides* (Linnaeus, 1758)

Note: This species is included on the list of indicator species of natural forest habitats.

Isomira Mulsant, 1856*murina* (Linnaeus, 1758)**Omophlinae** Mulsant, 1856**Omophlini** Mulsant, 1856**Cteniopus (Cteniopus)** Solier, 1835*sulphureus* (Linnaeus, 1758)*flavus* (Scopoli, 1763)*luteus* (Geoffroy, 1785)

Note: Scarce and insufficiently known species, with no confirmed records in the last 50 years.

Omophlus (Paromophlus) Znojko, 1950*rufitarsis* (Leske, 1785)*amerinae* Kirsch, 1869*armeriae* Curtis, 1836*betulae* Bach, 1856*frigidus* Mulsant, 1856*pinicola* Redtenbacher, 1849*pubescens* Mulsant, 1856

Note: Very rare and insufficiently known species.

Tenebrioninae Latreille, 1802**Alphitobiini** Reitter, 1917**Alphitobius** Stephens, 1829*diaperinus* (Panzer, 1797)*longipennis* Walker, 1858*opatroides* Brullé, 1838*ovatus* (Herbst, 1799)*rufipes* Walker, 1858**Blaptini** Leach, 1815**Blaps** Fabricius, 1775*— lethifera* Marsham, 1802*brevis* Fischer von Waldheim, 1844*carinula* Seidlitz, 1893*convexa* Fischer von Waldheim, 1844*damascena* Fischer von Waldheim, 1844*dorsata* Fischer von Waldheim, 1844*fatidica* Sturm, 1807*mortisaga* (Linnaeus, 1758)*acuminata* DeGeer, 1775*brevicornis* Seidlitz, 1893*canaliculata* Faldermann, 1837*carbo* Steven, 1829*confusa* Fischer von Waldheim, 1844*dahli* Solier, 1848*elongata* Ménétriés, 1832*ensifera* Motschulsky, 1845*reflexicollis* Fischer von Waldheim, 1844

Note: Rare species, known mostly from the eastern part of Latvia.

Bolitophagini Kirby, 1837**Bolitophagus** Illiger, 1798*reticulatus* (Linnaeus, 1767)*boleti* (Piller, Mitterpacher, 1783)*cornutus* (Thunberg, 1784)*crenatus* (Fabricius, 1792)*gibbus* (Panzer, 1789)**Eledona** Latreille, 1796*agricola* (Herbst, 1783)*sulcata* Thunberg, 1784**Helopini** Latreille, 1802**Stenomax** Allard, 1876*aeneus* (Scopoli, 1763)*arboreus* (Schrank, 1781)*incurvus* (Küster, 1850)

Note: Very rare and insufficiently known species.

Nalassus Mulsant, 1854*dermestoides* (Illiger, 1798)

Note: Very rare and insufficiently known species.

— gilvipes (Ménétriés, 1849)*brevicollis* (Küster, 1850)**Opatrini** Hope, 1840**Opatrum** Fabricius, 1775*riparium* W.Scriba, 1865*sabulosum* (Linnaeus, 1761)*intermedium* Fischer von Waldheim, 1844*rugosum* DeGeer, 1775*tricarinatum* Motschulsky, 1860**Melanimon** Steven, 1829*tibiale* (Fabricius, 1781)**Phylan** Dejean, 1821*gibbus* (Fabricius, 1775)*arenosus* Marsham, 1802*gemellatus* Marsham, 1802*maritimus* Leach, 1815*paludicola* Chevrier, 1865*pilipes* Herbst, 1797

Note: Very rare species known only from SW part of Latvia (Baltic Sea shore S of Liepāja) and with no confirmed records in the last 50 years.

Ulomini Blanchard, 1845**Gnatocerus** Thunberg, 1814*cornutus* (Fabricius, 1798)*cucullatus* Montrouzier, 1860*laeviusculus* Stephens, 1832*maxillaris* (Beauvois, 1805)*ruber* Thunberg, 1814**Uloma** Dejean, 1821*culinaris* (Linnaeus, 1758)*cucullata* Fischer von Waldheim, 1832

- rufa* (Piller, Mitterpacher, 1783)
perroudi Mulsant, Guillebeau, 1855
- Tenebrionini** Latreille, 1802
- Neatus* LeConte, 1862
- picipes* (Herbst, 1797)
badius (Say, 1823)
- Tenebrio** Linnaeus, 1758
- molitor* (Linnaeus, 1758)
laticollis Stephens, 1832
- obscurus* Fabricius, 1792
pollens Casey, 1924
- Triboliini** Mulsant, 1854
- Palorus* Mulsant, 1854
depressus (Fabricius, 1790)
unicolor (Olivier, 1790)
- Tribolium** McLeay, 1825
castaneum (Herbst, 1797)
bifoveolatus (Duftschmid, 1812)
cinnamomea (Herbst, 1792)
ferruginea (Herbst, 1797)
navale (Herbst, 1792)
rubens Laporte de Castenau, 1840
testaceum (Fabricius, 1798)
- confusum* Jacquelin du Val, 1863
destructor Uyttenboogart, 1934
madens (Charpentier, 1825)
obscurus Redtenbacher, 1842
- Diaperinae** Latreille, 1802
- Crypticini** Mulsant, 1854
- Crypticus* Reitter, 1817
quisquilius (Linnaeus, 1761)
glaber (Fabricius, 1775)
laevis (Gmelin, 1788)
laticollis (Panzer, 1798)
luctuosus (Schrank, 1781)
niger (Geoffroy, 1785)
pusillus (Herbst, 1799)
opacus Brancsik, 1899
- Diaperini** Latreille, 1802
- Alphitophagus* Stephens, 1832
bifasciatus (Say, 1823)
pictus Ménétriés, 1832
populi Redtenbacher, 1849
quadripustulatus Stephens, 1832
unifasciatus Donisthorpe, 1925
- Diaperis** O.F.Müller, 1764
boleti (Linnaeus, 1758)
fasciata (Scopoli, 1763)
fungi Motschulsky, 1845
quercana Baudi di Selve, 1876
interrupta Heyden, 1890
morio Baudi di Selve, 1876
posticalis Heyden, 1890
- Neomida** Latreille, 1829
haemorrhoidalis (Fabricius, 1787)
- cornifrons* (Rossi, 1792)
- Note: This species is included on the list of indicator species of natural forest habitats.
- Platydema** Laporte de Castelnau, Brullé, 1831
violaceum (Fabricius, 1790)
dytiscoides (Rossi, 1790)
- Note: This species is included on the list of indicator species of natural forest habitats.
- Pentaphyllus** Dejean, 1821
testaceus (Hellwig, 1792)
- Hypophlaeini** Shuckard, 1840
- Corticeus* Piller, Mitterpacher, 1783
bicolor (Olivier, 1790)
fraxini (Kugelann, 1794)
ferrugineus Mulsant, 1854
- linearis* (Fabricius, 1790)
pallidus Hellén, 1903
perrisi Pic, 1917
- longulus* (Gyllenhal, 1827)
pini (Panzer, 1799)
leonardi Reitter, 1906
nocivus Wollaston, 1862
- suturalis* (Paykull, 1800)
bivittatus Reitter, 1875
- unicolor* Piller, Mitterpacher, 1783
castaneus Fabricius, 1790
cimeterius Herbst, 1783
pseudocastaneus Reitter, 1906
purkynei Roubal, 1932
taxicornis Rossi, 1790
- Note: This species is included on the list of indicator species of natural forest habitats.
- Myrmecixeninae** Jacquelin du Val, 1858
- Myrmecixenus** Chevrolat, 1835
subterraneus Chevrolat, 1835
- Scaphidemini** Reitter, 1922
- Scaphidema** Redtenbacher, 1849
metallicum (Fabricius, 1792)
aeneus Panzer, 1793
bicolor Fabricius, 1798
sulcipenne Petri, 1926
- Coelometopinae** Lacordaire, 1859
- Upis* Fabricius, 1792
ceramboides (Linnaeus, 1758)
reticulata Say, 1824
variolosa (DeGeer, 1775)
- Note: Scarce species with no confirmed records in the last 50 years.
- Chrysomeloidea**
- Cerambycidae** Latreille, 1802
- Prioninae** Latreille, 1802
- Ergates* Audinet-Serville, 1832
faber *faber* (Linnaeus, 1761)

<p><i>alkani</i> Demelt, 1968</p> <p>Note: Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.</p> <p>Prionus Geoffroy, 1762</p> <p><i>coriarius</i> (Linnaeus, 1758)</p> <p>Note: Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.</p> <p>Tragosoma Audinet-Serville, 1832</p> <p><i>depsarium</i> (Linnaeus, 1767)</p> <p>Note: Very rare and insufficiently known species with only a few actual records from NW and NE parts of Latvia. Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.</p> <p>Lepturinae Latreille, 1802</p> <p>Rhamnusium Latreille, 1829</p> <p><i>bicolor</i> (Schrank, 1781)</p> <ul style="list-style-type: none"> <i>etruscum</i> (Rossi, 1790) <i>glaucopterum</i> (Schaller, 1783) <i>gracilicorne</i> Théry, 1894 <i>rubroviolaceus</i> (Villers, 1789) <i>rufivcollis</i> (Herbst, 1784) <i>salicis</i> (Fabricius, 1787) <p>Note: Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.</p> <p>Rhagium (<i>Rhagium</i> Fabricius, 1775)</p> <p><i>inquisitor</i> <i>inquisitor</i> (Linnaeus, 1758)</p> <p>Rhagium (<i>Hagrium</i> Villiers, 1978)</p> <p>— <i>bifasciatum</i> Fabricius, 1775</p> <p>Rhagium (<i>Megarhagium</i> Reitter, 1912)</p> <p><i>mordax</i> (DeGeer, 1775)</p> <p><i>sycophanta</i> (Schrank, 1781)</p> <p>Note: Scarce and insufficiently known species.</p> <p>Oxymirus Mulsant, 1862</p> <p><i>cursor</i> (Linnaeus, 1758)</p> <p>Stenocorus (<i>Stenocorus</i> Geoffroy, 1762)</p> <p><i>meridianus</i> (Linnaeus, 1758)</p> <p>Note: Officially protected species in Latvia.</p> <p>Pachyta Dejean, 1821</p> <p><i>lamed</i> <i>lamed</i> (Linnaeus, 1758)</p> <p><i>quadrimaculata</i> (Linnaeus, 1758)</p> <p>Brachyta Fairmaire, 1864</p> <p><i>interrogationis</i> <i>interrogationis</i> (Linnaeus, 1758)</p> <p>Evodinus LeConte, 1850</p> <p><i>borealis</i> (Gyllenhal, 1827)</p> <p>Note: This species is known only from single area in Ropaži area (Central Latvia). South</p>	<p>border of main distribution area.</p> <p>Gaurotes (<i>Carilia</i> Mulsant, 1863)</p> <p><i>virginea</i> <i>virginea</i> (Linnaeus, 1758)</p> <ul style="list-style-type: none"> <i>thalassina</i> (Schrank, 1781) <p>Acmaeops LeConte, 1850</p> <p><i>marginata</i> (Fabricius, 1781)</p> <p><i>pratensis</i> (Laicharting, 1784)</p> <p>Dinoptera Mulsant, 1863</p> <p><i>collaris</i> (Linnaeus, 1758)</p> <p>Cortodera Mulsant, 1863</p> <p><i>femorata</i> (Fabricius, 1787)</p> <p>Grammoptera Audinet-Serville, 1835</p> <p>— <i>abdominalis</i> (Stephens, 1831)</p> <ul style="list-style-type: none"> <i>variegata</i> (Germar, 1824) <p><i>ruficornis</i> <i>ruficornis</i> (Fabricius, 1781)</p> <ul style="list-style-type: none"> <i>atra</i> (Fabricius, 1775) <i>holomelina</i> Pool, 1905 <p>— <i>ustulata</i> (Schaller, 1783)</p> <p>Pidonia Mulsant, 1863</p> <p><i>lurida</i> (Fabricius, 1792)</p> <p>Etorufus Matsushita, 1933</p> <p><i>pubescens</i> (Fabricius, 1787)</p> <p>Lepturobosca Reitter, 1912</p> <p><i>virens</i> (Linnaeus, 1758)</p> <p>Macroleptura Nakane, Ohbayashi, 1957</p> <p><i>thoracica</i> (Creutzer, 1799)</p> <p>Note: Very rare species, known from only a few individuals from western and eastern Latvia. This species is included on the list of indicator species of natural forest habitats.</p> <p>Leptura Linnaeus, 1758</p> <p><i>aethiops</i> Poda, 1761</p> <p><i>annularis</i> <i>annularis</i> (Fabricius, 1801)</p> <ul style="list-style-type: none"> <i>arcuata</i> Panzer, 1793 <p><i>quadrifasciata</i> <i>quadrifasciata</i> Linnaeus, 1758</p> <p>Lepturalia Reitter, 1912</p> <p><i>nigripes</i> (DeGeer, 1775)</p> <p>Note: This species is included on the list of indicator species of natural forest habitats.</p> <p>Anastrangalia Casey, 1924</p> <p>— <i>dubia</i> <i>dubia</i> (Scopoli, 1763)</p> <p>Note: All previous references are invalid and relate to <i>A. reyi</i> (Heyden).</p> <p><i>reyi</i> (Heyden, 1889)</p> <ul style="list-style-type: none"> <i>inexpectata</i> (Jansson, Sjöberg, 1928) <i>ochracea</i> (Rey, 1885) <p><i>sanguinolenta</i> (Linnaeus, 1761)</p> <ul style="list-style-type: none"> <i>sandoensis</i> (Palm, 1953) <p>Stictoleptura Casey, 1924</p> <p><i>rubra</i> <i>rubra</i> (Linnaeus, 1758)</p> <p><i>scutellata</i> (Fabricius, 1781)</p>
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Note: This species is known from one locality only (Slīteres National park, NW Latvia). North border of main distribution area.

variicornis (Dalman in Schönherr, 1817)

Note: This species is known from single locality only (Slīteres National park, NW Latvia).

Paracorymbia (*Paracorymbia*) Miroshnikov, 1998)

—*fulva* (DeGeer, 1775)

maculicornis maculicornis (DeGeer, 1775)

Anoplodera Mulsant, 1839

rufipes rufipes (Schaller, 1783)

krueperi (Ganglbauer, 1882)

sexguttata (Fabricius, 1775)

Note: Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.

Vadonia Mulsant, 1863

—*unipunctata* (Fabricius, 1787)

occidentalis (K.Daniel, J.Daniel, 1891)

Judolia Mulsant, 1863

sexmaculata (Linnaeus, 1758)

Pachytodes Pic, 1891

cerambyciformis (Schrank, 1781)

quadrimaculatus (Scopoli, 1763)

Note: Very rare species.

Alosterna Mulsant, 1863

ingrica (Baeckmann, 1902)

tabacicolor tabacicolor (DeGeer, 1775)

bicoloripes Pic, 1914

Pseudovadonia Lobanov, Danilevsky, Murzin, 1981

livida (Fabricius, 1776)

pecta (K.Daniel, J.Daniel, 1891)

Strangalia Audinet-Serville, 1835

attenuata (Linnaeus, 1758)

Note: This species is included on the list of indicator species of natural forest habitats.

Rutpela Nakane, Ohbayashi, 1959

maculata (Poda, 1761)

Stenurella Villiers, 1974

bifasciata bifasciata (O.F.Müller, 1776)

melanura (Linnaeus, 1758)

melanurella (Reitter, 1901)

semicrassa (Pic, 1901)

nigra (Linnaeus, 1758)

Necydalinae Latreille, 1825

Necydalis (*Necydalis*) Linnaeus, 1758)

major Linnaeus, 1758

Note: Officially protected species in Latvia;

also included on the list of indicator species of natural forest habitats.

ulmi (Chevrolat, 1836)

annulatus (Petagna, 1819)

Note: Scarce species with no confirmed records in the last 100 years. North border of main distribution area.

Spondylidinae Audinet-Serville, 1832

Spondylis Fabricius, 1758

buprestoides (Linnaeus, 1758)

Nothorhina Redtenbacher, 1845

muricata (Dalman, 1817)

punctata (Fabricius, 1798)

scabricolle (Redtenbacher, 1842)

Note: Officially protected species in Latvia; also included on the list of indicator species of natural forest habitats.

Asemum Eschscholtz, 1830

striatum (Linnaeus, 1758)

Tetropium Kirby, 1837

castaneum (Linnaeus, 1758)

fuscum (Fabricius, 1787)

Arhopalus Audinet-Serville, 1834

ferus (Mulsant, 1839)

polonicum (Motschulsky, 1845)

rusticus (Linnaeus, 1758)

lugubris (Gmelin, 1790)

triste (Fabricius, 1787)

Cerambycinae Latreille, 1802

Stromatium Audinet-Serville, 1834

Stromatium unicolor (Olivier, 1795)

fulvus (Villers, 1789)

inermis Tournier, 1872)

pallidum (Zoubkoff, 1833)

platyfemur (Chevrolat, 1882)

streps (Fabricius, 1798)

Note: Probably introduced species, known only from Ķemeri area (Central Latvia).

Cerambyx (*Cerambyx*) Linnaeus, 1758)

cerdo cerdo Linnaeus, 1758

acuminatus (Motschulsky, 1852)

heros Scopoli, 1763

pfisteri Stierlin, 1864

Note: Scarce species, known mostly by unconfirmed records. North border of main distribution area. Officially protected species in Latvia.

—*scopolii scopolii* Fuesslins, 1775

cerdo Poda, 1761 nec Linnaeus, 1758

Gracilia Audinet-Serville, 1834

—*minuta* (Fabricius, 1781)

albanica Csiki, 1931

Obrium Dejean, 1821

brunneum (Fabricius, 1792)
cantharinum cantharinum (Linnaeus, 1767)
ferrugineum (Fabricius, 1781)
fuscicornis (Gmelin, 1790)

Nathrius Bréthes, 1916
— *brevipennis* (Mulsant, 1839)
porteri Bréthes, 1916
rufipennis (Dufour, 1851)

Molorchus Fabricius, 1792
minor minor (Linnaeus, 1758)

Glypha (*Glypha*) Newman, 1840
umbellatarum umbellatarum (Schreber, 1759)
depressa (Motschulsky, 1845)
diversipes (Pic, 1897)
minima (Scopoli, 1763)
obscuripes (P.W.J.Müller, 1948)

Aromia Audinet-Serville, 1833
moschata moschata (Linnaeus, 1758)
thea Reitter, 1894

Ropalopus (*Ropalopus*) Mulsant, 1839
clavipes (Fabricius, 1775)
nigroplanum (DeGeer, 1775)
femoratus (Linnaeus, 1758)
punctatus (Fabricius, 1798)
— *macropus* (Germar, 1824)

Hylotrupes Audinet-Serville, 1834
bajulus (Linnaeus, 1758)

Semanotus Mulsant, 1839
undatus (Linnaeus, 1758)

Callidium (*Callidium*) Fabricius, 1775
violaceum (Linnaeus, 1758)

Callidium (*Palaeocallidium*) Plavilstshikov, 1940
coriaceum Paykull, 1800

Callidium (*Callidostola*) Reitter, 1912
aeneum aeneum (DeGeer, 1775)

Pyrrhidium Fairmaire, 1864
— *sanguineum* (Linnaeus, 1758)

Leioderus Redtenbacher, 1849
kollari Redtenbacher, 1849

Phymatodes (*Phymatodes*) Mulsant, 1839
testaceus (Linnaeus, 1758)
italicum (Gmelin, 1790)
ruficolle (Fabricius, 1781)
violaceum (Rossi, 1790)

Poecilium (*Poecilium*) Fairmaire, 1864
alni alni (Linnaeus, 1767)
— *fasciatum* (Villers, 1789)
unifasciatum (Olivier, 1790)

Xylotrechus (*Xylotrechus*) Chevrolat, 1860
arvicola (Olivier, 1795)
heydeni (Stierlin, 1864)
rusticus (Linnaeus, 1758)

Clytus Laicharting, 1784
arietis arietis (Linnaeus, 1758)
Cyrtoclytus Ganglbauer, 1882
capra (Germar, 1824)
Plagionotus Mulsant, 1842
arcuatus (Linnaeus, 1758)
detritus detritus (Linnaeus, 1758)
convertinii (Petagna, 1819)

Note: This species is recently distributed mostly in eastern part of Latvia.

floralis (Pallas, 1773)
abruptus (Kraatz, 1870)
puinosus (Kraatz, 1870)

Chlorophorus Chevrolat, 1863
figuratus (Scopoli, 1763)
herbstii (Brahm, 1790)
— *sartor* (O.F.Müller, 1766)
massiliensis (Linnaeus, 1767)

Anaglyptus Mulsant, 1839
mysticus (Linnaeus, 1758)

Note: This species is known from one locality only (Slīteres National park, NW Latvia).

Lamiinae Latreille, 1825

Iberodorcadion (*Iberodorcadion*) Breuning, 1943
fuliginator fuliginator (Linnaeus, 1758)

Note: This scarce species is known by single individual from Kandava area (Central Latvia). North border of main distribution area.

Mesosa (*Mesosa*) Latreille, 1829
— *curculionoides* (Linnaeus, 1761)

Mesosa (*Aphelocnemia*) Stephens, 1831
myops (Dalman, 1817)

Note: Very rare and insufficiently known species with only two confirmed records in the last 80 years. Officially protected species in Latvia.

Agapanthia (*Agapanthia*) Aud.-Serville, 1835
villosoviridescens (DeGeer, 1775)
— *violacea* (Fabricius, 1775)

Lamia Fabricius, 1775
textor (Linnaeus, 1758)

Monochamus Dejean, 1821
galloprovincialis pistor (Germar, 1818)
rosenmuelleri (Cederhjelm, 1798)
rosenmuelleri Jakobson, 1910
urussovi (Fischer von Waldheim, 1806)

Note: West border of main distribution area. This species is included on the list of indicator species of natural forest habitats.

— *saltuarius* Gebler, 1830

- sutor* (Linnaeus, 1758)
Anaesthetis Dejean, 1835
testacea (Fabricius, 1781)
Pogonocherus (*Pogonocherus* Dejean, 1821)
hispidulus (Piller, Mitterpacher, 1783)
hispidus (Linnaeus, 1758)
Pogonocherus (*Pityphilus* Mulsant, 1863)
decoratus Fairmaire, 1855
fasciculatus fasciculatus (DeGeer, 1775)
ovatus (Goeze, 1777)
ovalis (Gmelin, 1790)
Acanthocinus Dejean, 1821
aedilis (Linnaeus, 1758)
griseus griseus (Fabricius, 1792)
Leiopus Audinet-Serville, 1835
nebulosus (Linnaeus, 1758)
punctulatus (Paykull, 1800)
Exocentrus Dejean, 1835
lusitanus (Linnaeus, 1767)
Note: Very rare and insufficiently known species. Should be widely distributed in broad leaved forests of Latvia.
Aegomorphus Haldeman, 1847
clavipes (Schrantz, 1781)
varius (Fabricius, 1787)
Oplosia Mulsant, 1862
cinerea (Mulsant, 1839)
fennica (Paykull, 1800)
Saperda (*Anaarea* Mulsant, 1839)
carcharias (Linnaeus, 1758)
similis (Laicharting, 1784)
Note: Insufficiently known species, actually recorderd from few localities mostly in central and eastern Latvia.
Saperda (*Saperda* Fabricius, 1775)
perforata (Pallas, 1773)
Note: This species is included on the list of indicator species of natural forest habitats.
scalaris scalaris (Linnaeus, 1758)
hieroglyphica (Pallas, 1773)
xantha Demelt, 1960
Saperda (*Lopezcolonia* Alonso-Zarazaga, 1998)
punctata (Linnaeus, 1767)
Saperda (*Compsidia* Mulsant, 1839)
populnea (Linnaeus, 1758)
Menesia Mulsant, 1856
bipunctata (Zoubkoff, 1829)
Note: Very rare and insufficiently known species.
Stenostola Dejean, 1835
dubia (Laicharting, 1784)

- alboscutellata* Kraatz, 1862
Opsilia Mulsant, 1862
— *coerulescens* (Scopoli, 1763)
Phytoecia (*Phytoecia* Dejean, 1835)
— *caerulea* (Scopoli, 1772)
coelestis (Townson, 1797)
coerulea auct.
flavimana (Creutzer in Panzer, 1796)
rufimana (Fabricius, 1801)
cylindrica (Linnaeus, 1758)
fuliginosa (Scopoli, 1786)
kammereri Schatzmayr, 1928
simplonica Stierlin, 1879
nigricornis (Fabricius, 1781)
caroni Mulsant, Godart, 1876
julii Mulsant, 1862
solidaginis Bach, 1856
virgula (Charpentier, 1825)
Note: This species is known from many localities in SW Latvia (Daugavpils area). North border of main distribution area.
Oberea (*Oberea* Dejean, 1835)
linearis (Linnaeus, 1761)
oculata (Linnaeus, 1758)
pupillata (Gyllenhal, 1817)
Tetrops Stephens, 1829
praeustus (Linnaeus, 1758)
nigra Kraatz, 1859
Bruchidae Latreille, 1802
Bruchinae Latreille, 1802
Bruchini Latreille, 1802
Bruchus O.F.Müller, 1764
affinis affinis Frölich, 1799
flavimanus Boheman, 1833
obscuritarsis Motschulsky, 1873
atomarius (Linnaeus, 1761)
fahraei Gyllenhal, 1839
granarius Linnaeus, 1767
seminarius Bach, 1854
troglodytes Fahraeus, 1839
varicornis Brullé, 1832
viciae Motschulsky, 1854
laticollis Boheman, 1833
loti Paykull, 1800
lathyri Stephens, 1831
muelleri Schilsky, 1905
oxytropis Gebler, 1830
wasastjernii Fahraeus, 1839
pisorum (Linnaeus, 1758)
cruciger Geoffroy, 1785
intermedius Motschulsky, 1854
lunaris Rey, 1893
pisii Linnaeus, 1767
sparsus Fabricius, 1801
unifasciatus Mulsant, Rey, 1858

rufimanus rufimanus Boheman, 1833
fabae Motschulsky, 1854
granarius (Marsham, 1802)

Acanthoscelidini Bridwell, 1946

Acanthoscelides Schilsky, 1905

obtectus (Say, 1831)
breweri (Crotch, 1867)
irresectus (Fahraeus, 1839)
obsoletus (Say, 1831)
pallidipes (Fahraeus, 1839)
subellipticus (Wollaston, 1854)
varicornis (Motschulsky, 1873)

Bruchidius Schilsky, 1905

marginalis (Fabricius, 1776)
marginellus (Fabricius, 1792)

Amblycerinae Bridwell, 1932

Spermophagus Schönherr, 1833

sericeus (Geoffroy, 1785)
cardui (Boheman, 1829)
cisti (Olivier, 1790)
euphorbiae Küster, 1849
sulcifrons Küster, 1849

Note: This species is known by single record from extreme SW of Latvia only (North env. of Liepāja), but it should be more widely distributed in SW part of Latvia. North border of main distribution area.

Chrysomelidae Latreille, 1802

Donaciinae Kirby, 1837

Macroplea Samouelle, 1819

appendiculata (Panzer, 1794)
baltica Seidlitz, 1875
gyllenhalii Lacordaire, 1845
kraussei Reitter, 1910
sahlbergii Lacordaire, 1845
schioedtei Guérin-Méneville, 1844
zosterae (Fabricius, 1801)

mutica (Fabricius, 1798)
equiseti (Fabricius, 1798)
japana Jacoby, 1885
mosellae Bellevoye, 1870
mucronata (Hoppe, 1795)

Plateumaris Thomson, 1859

braccata (Scopoli, 1772)
abdominalis (Olivier, 1795)
nigra (Fabricius, 1792)
palustris (Herbst, 1784)
violacea (Pallas, 1773)

sericea (Linnaeus, 1761)
aenea (Hoppe, 1795)
asiatica (Faldermann, 1837)
comari (Suffrian, 1846)
discolor (Panzer, 1795)
geniculata (Thomson, 1866)
palustris (Schilling, 1837)
proteus (Kunze, 1818)

rustica (Kunze, 1818)
affinis Kunze, 1818
sulcifrons Weise, 1900

Donacia Fabricius, 1775

antiqua Kunze, 1818
gracilis Suffrian, 1845
simplicifrons Lacordaire, 1845

aquatica (Linnaeus, 1758)
coccineofasciata Harrer, 1784
dentipes Fabricius, 1792
fasciata Herbst, 1784
latonae Gistel, 1857
vittata Olivier, 1791

bicolora Zschach, 1788
aurea Hoppe, 1795
sagittariae Fabricius, 1792

brevicornis Ahrens, 1810
platysterna Thomson, 1866

brevitarsis Thomson, 1884

crassipes Fabricius, 1775
micans Hoppe, 1795
spinosa Thomson, 1866
striata Panzer, 1793

dentata Hoppe, 1795
melanocephala Marsham, 1802

fennica (Paykull, 1800)
flavidula Reitter, 1913
ochroleuca Weise, 1912

impressa (Paykull, 1799)

marginata Hoppe, 1795
lateralis Bonelli, 1812
lemnæ Fabricius, 1810
limbata Panzer, 1796
trojana Goecke, 1948
vittata Panzer, 1796

obscura Gyllenhal, 1813
splendens Jakobson, 1925

semicuprea Panzer, 1796
aenea Gistel, 1857

simplex Fabricius, 1775
linearis Hoppe, 1795

sparganii Ahrens, 1810

thalassina Germar, 1811

versicolorea (Brahm, 1790)
bidens Olivier, 1791
cincta Germar, 1810

vulgaris Zschach, 1788
typhae Ahrens, 1810

Donaciella Reitter, 1920

cinerea (Herbst, 1784)
hydrocharis (Fabricius, 1792)
hydrochoeridis (Fabricius, 1801)
tarsata (Panzer, 1796)

clavipes (Fabricius, 1793)
aerea (Schrink, 1798)
glabrata (Solsky, 1872)

- menyanthidis* (Gyllenhal, 1827)
tomentosa (Ahrens, 1810)
Orsodacninae Thomson, 1859
Orsodacne Latreille, 1802
cerasi (Linnaeus, 1758)
 chlorotica (Olivier, 1791)
 fulvicollis (Fabricius, 1792)
 glabrata (Fabricius, 1798)
Zeugophorinae Böving, Craighead, 1931
Zeugophora Kunze, 1818
flavicollis (Marsham, 1802)
frontalis Suffrian, 1840
subspinosa (Fabricius, 1781)
turneri Power, 1863
 rufotestacea Kraatz, 1871
Criocerinae Latreille, 1807
Criocerini Latreille, 1807
Lilioceris Reitter, 1912
lili (Scopoli, 1763)
 amurensis Pic, 1916
 liliorum (Thomson, 1866)
merdigera (Linnaeus, 1758)
 brunnea (Fabricius, 1792)
Crioceris O.F.Müller, 1764
asparagi asparagi (Linnaeus, 1758)
 campestris Linnaeus, 1767
duodecimpunctata (Linnaeus, 1758)
Lemini Gyllenhal, 1813
Lema Fabricius, 1798
cyanella (Linnaeus, 1758)
 puncticollis Curtis, 1830
 rugicollis Suffrian, 1841
Oulema Gozis, 1886
erichsonii (Suffrian, 1841)
 weisei Seidlitz, 1891
duftschmidi (Redtenbacher, 1874)
gallaeciana (Heyden, 1870)
 lichenis (Voet, 1806)
melanopus (Linnaeus, 1758) s. l.
 melanopa (Linnaeus, 1761)
Clytrinae Kirby, 1837
Labidostomis Germar, 1822
humeralis (Schneider, 1792)
 impressihumera Lacordaire, 1848
lepidus Lefevre, 1872
Note: North border of main distribution area.
longimana (Linnaeus, 1761)
 glycyrrhizae Becker, 1880
 pallida (Geoffroy, 1785)
 uralensis Lacordaire, 1848
lucida (Germar, 1824)
 albipennis Lacordaire, 1848
 fulgida Lacordaire, 1848

- Note: North border of main distribution area.
tridentata (Linnaeus, 1758)
 cyanicollis Lacordaire, 1848
 kluczickii Nowicki, 1872
 leithneri Redtenbacher, 1874
 viridicollis Lacordaire, 1848
Clytra Laicharting, 1781
laeviuscula (Ratzeburg, 1837)
 quadripuncta Laicharting, 1781
quadripunctata (Linnaeus, 1758)
 quadrisignata Märkel, 1842
Smaragdina Chevrolat, 1837
affinis (Illiger, 1794)
flavicollis Charpentier, 1825
salicina (Scopoli, 1763)
 cyanea (Fabricius, 1775)
Cptocephala Chevrolat, 1837
unifasciata (Scopoli, 1763)
 femoralis Küster, 1845
Cryptocephalinae Gyllenhal, 1813
Pachybrachini Chapuis, 1874
Pachybrachis Chevrolat, 1837
hieroglyphicus *hieroglyphicus* (Laicharting, 1781)
 atomarius (Gebler, 1830)
Cryptocephalini Gyllenhal, 1813
Cryptocephalus O.F.Müller, 1764
androgyne Marseul, 1875
 coerulescens R.F.Sahlberg, 1839
 flavilabris Gyllenhal, 1813
 matsumurai Takizava, 1971
 pelleti Marseul, 1875
 similis L.Medvedev, 1956
– *anticus* Suffrian, 1848
 nigerrimus Ballion, 1878
 octacosmus Bedel, 1891
 octoguttatus Schneider, 1792
 sexpustulatus Rossi, 1790
aureolus Suffrian, 1847 nec Seidlitz, 1875
biguttatus (Scopoli, 1763)
 bipustulatus Fabricius, 1775
bilineatus (Linnaeus, 1767)
 armeniacus (Faldermann, 1837)
 partitus Jacoby, 1885
 spitzyi Suffrian, 1848
bipunctatus (Linnaeus, 1758)
 dispar Paykull, 1799
 intemeratus Gozis, 1907
 lineola Fabricius, 1792
 paradoxus Suffrian, 1848
chrysopus (Gmelin, 1788)
 biguttatus (Schaller, 1783)
 huebneri Fabricius, 1792
 haemorrhoidalis Schneider, 1792
cordiger (Linnaeus, 1758)

<i>coryli</i> (Linnaeus, 1758)	<i>ochropezzus</i> Suffrian, 1853
<i>rubroscutellatus</i> Burlini, 1967	
<i>vitis</i> (Fabricius, 1775)	
<i>decemmaculatus</i> (Linnaeus, 1758)	<i>octopunctatus</i> (Scopoli, 1763)
<i>betulaenanae</i> Schill, 1837	<i>apotmetus</i> Jakobson, 1900
<i>bothnicus</i> (Linnaeus, 1758)	<i>hongyuanensis</i> Tan, 1992
<i>decempunctatus</i> (Linnaeus, 1761)	<i>potaissiae</i> Csiki, 1953
<i>frenatus</i> Laicharting, 1781 nec Fabricius, 1792	<i>variabilis</i> Schneider, 1791
<i>frenatus</i> Fabricius, 1792 nec Laicharting, 1781	
<i>pisanus</i> Burlini, 1967	
<i>trilineatus</i> Fabricius, 1798	
<i>distinguendus</i> Schneider, 1792	<i>pallifrons</i> Gyllenhal, 1813
<i>alni</i> Ménétriés, 1836	<i>epipleuralis</i> Weise, 1898
<i>ongudajensis</i> Jakobson, 1900	<i>furcifrons</i> Mannerheim, 1844
<i>variegatus</i> Gyllenhal, 1813	<i>insignis</i> Paykull, 1799
<i>elegantulus</i> Gravenhorst, 1807	<i>pallidifrons</i> Clavareau, 1913
<i>elongatulus</i> Olivier, 1808	
<i>jucundus</i> Faldermann, 1837	
<i>kuatunensis</i> Pic, 1954	
<i>negrei</i> Burlini, 1967	
<i>tesselatus</i> Germar, 1813	
<i>yangweii</i> Chen, 1942	
<i>exiguus</i> Schneider, 1792	<i>parvulus</i> O.F.Müller, 1776
<i>adocetus</i> Jakobson, 1901	<i>amatus</i> Baly, 1873
<i>amiculus</i> Baly, 1873	<i>barbareae</i> Stephens, 1834
<i>kiyosatonus</i> Kimoto, 1964	<i>coeruleus</i> Olivier, 1790
<i>wasasternii</i> Gyllenhal, 1827	<i>consolanus</i> Baly, 1874
<i>flavipes</i> Fabricius, 1781	<i>flavilabris</i> Fabricius, 1787
<i>dispar</i> Weise, 1882	<i>fulcratus</i> Germar, 1824
<i>wydleri</i> Faldermann, 1837	<i>inurbanus</i> Harold, 1874
<i>frontalis</i> Marsham, 1802	<i>livens</i> Gmelin, 1788
<i>marginellus</i> Gravenhorst, 1807	<i>obliquostriatus</i> Motschulsky, 1866
<i>fulvus</i> Goeze, 1777	<i>nigrocoeruleus</i> Goeze, 1777
<i>ochraceus</i> Stephens, 1829	<i>nitens</i> Rossi, 1792
<i>signaticollis</i> Suffrian, 1848	<i>permodesmus</i> Baly, 1873
<i>hypocoeridis</i> (Linnaeus, 1758)	<i>violaceus</i> Geoffroy, 1785
<i>auratus</i> Gerhard, 1909	
<i>auricollis</i> Clavareau, 1913	<i>pini</i> (Linnaeus, 1758)
<i>bicolor</i> Gerhard, 1909	<i>abietis</i> Redtenbacher, 1874
<i>cristatus</i> Dufour, 1843 nec Weise, 1882	<i>laevigatus</i> Suffrian, 1847
<i>cristatus</i> Weise, 1882 nec Dufour, 1843	<i>porrii</i> Suffrian, 1847
<i>cyanescens</i> Weise, 1893	
<i>similis</i> Stephens, 1834	<i>populi</i> Suffrian, 1848
<i>violaceus</i> Gerhard, 1909	<i>brachialis</i> Mulsant, 1859
<i>labiatus</i> (Linnaeus, 1761)	<i>punctiger</i> Paykull, 1799
<i>longicornis</i> Thomson, 1866	
<i>moraei</i> (Linnaeus, 1758)	<i>pusillus</i> Fabricius, 1777
<i>cruciatus</i> Marseul, 1875	<i>gracilis</i> Stephens, 1834
<i>nitidulus</i> Fabricius, 1787	<i>marginella</i> Donovan, 1801
<i>ochrostoma</i> Harold, 1872	<i>marginellus</i> Marsham, 1802
<i>nitidus</i> (Linnaeus, 1758)	<i>minutus</i> (Herbst, 1783)
<i>flavifrons</i> Fabricius, 1792	<i>raphaelensis</i> Gautier, 1861
<i>nitens</i> (Linnaeus, 1761)	<i>sonchi</i> Schrank, 1798
<i>ocellatus</i> Drapiez, 1819	<i>verticalis</i> Boheman, 1851
<i>corsicus</i> Pic, 1909	
<i>geminus</i> Gyllenhal, 1827	<i>pygmaeus</i> Fabricius, 1792
<i>hummleri</i> Pic, 1909	— <i>quadripustulatus</i> Gyllenhal, 1813
	<i>apicalis</i> Marseul, 1875
	<i>bisignatus</i> Suffrian, 1847
	<i>quercerti</i> Suffrian, 1848
	<i>rufipes</i> (Goeze, 1777)
	<i>gracilis</i> Fabricius, 1792
	<i>minutus</i> Fabricius, 1792
	<i>septimaniensis</i> Pic, 1905
	<i>vitticollis</i> Weise, 1891
	<i>sericeus</i> (Linnaeus, 1758)
	<i>aureolus</i> Seidlitz, 1875 nec Suffrian, 1847
	<i>bidens</i> Thomson, 1868
	<i>coeruleus</i> Weise, 1882
	<i>intrusus</i> Weise, 1881
	<i>mirandus</i> Burlini, 1967
	<i>robustus</i> Suffrian, 1853
	<i>sexpunctatus</i> (Linnaeus, 1758)

signatifrons Suffrian, 1847

vittatus Fabricius, 1775

digonensis Pic, 1900

Lamprosomatinae Lacordaire, 1849

Oomorphus Curtis, 1831

concolor (Sturm, 1807)

maritima (Stephens, 1829)

unicolor Brullé, 1834

Synetinae Edwards, 1953

Syneta Lacordaire, 1845

betulae (Fabricius, 1792)

Note: This species is only known from northern parts of Latvia. South border of main distribution area.

Eumolpinae Hoppe, 1840

Eumolpini Hoppe, 1840

Eumolpus Illiger, 1798

asclepiadeus (Pallas, 1776)

praetiosus (Schneider, 1791)

Note: North border of main distribution area.

Adoxini Baly, 1863

Pachnephorus Redtenbacher, 1845

pilosus (Rossi, 1790)

aeneus (Schneider, 1792)

arenarius (Panzer, 1797)

carbonarius (Grimmer, 1841)

lepidopterus Küster, 1846

tessellatus (Duftschmid, 1825)

Bromius Chevrolat, 1837

obscurus (Linnaeus, 1758)

Chrysomelinae Latreille, 1802

Chrysomelini Latreille, 1802

Leptinotarsa Chevrolat, 1837

decemlineata (Say, 1824)

Chrysolina Motschulsky, 1860

americana (Linnaeus, 1758)

barbarica (Gmelin, 1790)

decemstriata (Goeze, 1777)

nitidula (Geoffroy, 1785)

Note: Very rare species with no confirmed records in the last 50 years. Introduced species.

analis (Linnaeus, 1767)

schach Tenenbaum, 1927

carnifex (Fabricius, 1792)

cerealis (Linnaeus, 1767)

— *chalcites* (Germar, 1824)

morio (Krynicki, 1832)

fastuosa speciosa (Linnaeus, 1767)

fuliginosa (Olivier, 1807)

geminata (Paykull, 1799)

approximata (Zenker, 1815)

bifoveolata Brancsik, 1910

graminis (Linnaeus, 1758)

artemisiae Motschulsky, 1860

gypsophilae (Küster, 1845)

rufomarginata (Suffrian, 1851)

haemoptera (Linnaeus, 1758)

acuticollis Fairmaire, 1877

atra (Goeze, 1777)

hottentotta (Fabricius, 1792)

hyperici (Forster, 1771)

fucata Fabricius, 1781

gemellata Geoffroy, 1785

kuesteri (Helliesen, 1912)

limbata (Fabricius, 1775)

marginata (Linnaeus, 1758)

menthae (Schrank, 1776)

opulenta (Suffrian, 1851)

pachysoma Hubenthal, 1911

rufa (Duftschmid, 1825)

oricalcia (O.F.Müller, 1776)

politata (Linnaeus, 1758)

quadrigemina (Suffrian, 1851)

alternata (Suffrian, 1851)

indigena Weise, 1884

normandi Bechyné, 1949

Locality in Latvia is the single one in Baltia

— *reitteri reitteri* (Weise, 1884)

saxonica Silfverberg, 1977

sanguinolenta (Linnaeus, 1758)

marginalis (Duftschmid, 1825)

staphylea (Linnaeus, 1758)

cuprea (DeGeer, 1775)

rubrocuprea (Geoffroy, 1785)

subferruginea (Suffrian, 1851)

sturmi (Westhoff, 1882)

diversipes Bedel, 1892

violacea Weise, 1916

— *susterai* Bechyne, 1950

morio (Krynicki, 1832)

varians (Schaller, 1783)

incerta Meissner, 1907

margarita (Olivier, 1798)

marshami (Donovan, 1790)

ooensis Weise, 1916

subaenaea (Suffrian, 1851)

Colaphus Dahl, 1823

sophiae (Schaller, 1783)

erythropus (Gmelin, 1790)

Note: North border of main distribution area.

Gastrophysa Chevrolat, 1837

polygoni (Linnaeus, 1758)

elongata Jolivet, 1951

viridula (DeGeer, 1775)

raphani (Herbst, 1783)

Phaedon Latreille, 1829

armoraciae (Linnaeus, 1758)

<i>betulae</i> Küster, 1846	<i>atrovirens</i> (Cornelius, 1857)
cochleariae (Fabricius, 1792)	<i>laticollis</i> (Suffrian, 1851)
<i>egenus</i> (Gyllenhal, 1827)	<i>cavifrons</i> Thomson, 1866
<i>galeopsis</i> Letzner, 1849	<i>polaris</i> (Schneider, 1886)
<i>grammicus</i> Suffrian, 1851	Note: South border of main distribution area.
<i>omissus</i> R.F.Sahlberg, 1838	<i>tibialis</i> (Suffrian, 1851)
<i>regnianum</i> Tottenham, 1941	<i>vitellinae</i> (Linnaeus, 1758)
<i>laevigatus</i> (Duftschmid, 1825)	<i>angusticollis</i> Motschulsky, 1860
<i>gramicus</i> (Duftschmid, 1825)	<i>latipennis</i> Motschulsky, 1860
<i>sabulicola</i> Suffrian, 1851	<i>vulgatissima</i> (Linnaeus, 1758)
Hydrothassa Thomson, 1866	Timarcha Latreille, 1829
<i>glabra</i> (Herbst, 1783)	<i>goettingensis</i> (Linnaeus, 1758)
<i>hannoveriana</i> (Fabricius, 1775)	— <i>pratensis</i> (Duftschmid, 1825)
<i>ranunculi</i> (Herbst, 1783)	<i>tenebricosa</i> (Fabricius, 1775)
<i>marginella</i> (Linnaeus, 1758)	Entomoscelis Chevrolat, 1837
Prasocuris Latreille, 1802	<i>adonidis</i> (Pallas, 1771)
<i>junci</i> (Brahm, 1790)	<i>trilineata</i> (Fabricius, 1777)
<i>beccabungae</i> (Illiger, 1794)	Note: Very rare and insufficiently known
<i>phellandrii</i> (Linnaeus, 1758)	species with no confirmed records in the last
Plagiodesma Chevrolat, 1837	50 years.
<i>versicolora</i> (Laicharting, 1781)	Galerucinae Latreille, 1802
Chrysomela Linnaeus, 1758	Galerucini Latreille, 1802
<i>collaris</i> Linnaeus, 1758	Galerucella Crotch, 1873
<i>escheri</i> Heer, 1836	<i>calmariensis</i> (Linnaeus, 1767)
<i>cuprea</i> Fabricius, 1775	<i>grisescens</i> (Joannis, 1866)
<i>ruficaudis</i> DeGeer, 1775	<i>reducta</i> Chen, 1942
<i>lapponica</i> Linnaeus, 1758	<i>vittaticollis</i> Baly, 1874
<i>populi</i> Linnaeus, 1758	<i>lineola</i> (Fabricius, 1781)
<i>saliceti</i> Suffrian, 1849	<i>verna</i> (Laicharting, 1781)
<i>tremula</i> Fabricius, 1787	<i>nymphaeae</i> (Linnaeus, 1758) s. l.
<i>tremulae</i> Paykull, 1799	<i>pusilla</i> (Duftschmid, 1825)
<i>vigintipunctata</i> (Scopoli, 1763)	<i>tenella</i> (Linnaeus, 1761)
Linaeidea Motschulsky, 1860	<i>parva</i> (Herbst, 1783)
<i>aenea</i> (Linnaeus, 1758)	<i>semenovi</i> Rybakov, 1889
<i>coeruleoviolacea</i> (DeGeer, 1775)	Xanthogaleruca Laboissière, 1934
Gonioctena Chevrolat, 1837	<i>luteola</i> (O.F.Müller, 1766)
<i>decemnotata</i> (Marsham, 1802)	Pyrrhalta Joannis, 1866
<i>rufipes</i> (DeGeer, 1775)	<i>viburni</i> (Paykull, 1799)
<i>sexpunctata</i> (Fabricius, 1787)	Lochmaea Weise, 1883
<i>flavicornis</i> (Suffrian, 1851)	<i>caprea</i> (Linnaeus, 1758)
<i>intermedia</i> (Helliesen, 1913)	<i>crataegi</i> (Forster, 1771)
— <i>kaufmanni</i> (Miller, 1881)	<i>suturalis</i> (Thomson, 1866)
<i>linnaeana</i> (Schrink, 1781)	Galeruca O.F.Müller, 1764
<i>triandrae</i> (Suffrian, 1851)	<i>dahli</i> (Joannis, 1866)
<i>pallida</i> (Linnaeus, 1758)	<i>interrupta</i> (Illiger, 1802)
<i>frontalis</i> (Olivier, 1807)	<i>hungarica</i> I.Frivaldszky von Frivald, 1876
<i>quinquepunctata</i> (Fabricius, 1787)	<i>laticollis</i> R.F.Sahlberg, 1837
<i>viminalis</i> (Linnaeus, 1758)	<i>confinis</i> Gebler, 1848
<i>baaderi</i> (Panzer, 1794)	<i>flava</i> Küster, 1855
<i>collaris</i> Grndl, 1882	<i>fontinalis</i> Boheman, 1851
<i>decempunctata</i> (Linnaeus, 1758)	<i>pomonae</i> (Scopoli, 1763)
<i>fridrichsdalensis</i> (O.F.Müller, 1764)	<i>dispar</i> Joannis, 1866
<i>haemorrhoidalis</i> (Scopoli, 1763)	
Phratora Chevrolat, 1837	

- rustica* Schaller, 1883
tanaceti (Linnaeus, 1758)
bonvouloiri Joannis, 1866
cicatricosa Chevrolat, 1872
declivis Joannis, 1866
tristis Scopoli, 1863
- Sermylassini** Mroczkowski, 1991
Sermylassa Reitter, 1912
halensis (Linnaeus, 1767)
nigricornis (Fabricius, 1775)
- Agelasticini** Chevrolat, 1875
Agelastica Chevrolat, 1837
alni (Linnaeus, 1758)
violacea Laicharting, 1781
- Luperini** Chapuis, 1875
Phyllobrotica Chevrolat, 1837
quadrimaculata (Linnaeus, 1758)
bimaculata (Panzer, 1795)
melanogaster (Gmelin, 1790)
- Calomicrus** Stephens, 1831
pinicola (Duftschmid, 1825)
Luperus Geoffroy, 1762
flavipes (Linnaeus, 1767)
carniolicus Kiesenwetter, 1861
ochropus Gmelin, 1790
- longicornis* (Fabricius, 1781)
diniensis Bellier, 1870
- luperus* (Sulzer, 1776)
dispar Redtenbacher, 1874
erythromelas Gmelin, 1790
lyperus Bedel, 1897
niger Goeze, 1777
rufipes Goeze, 1777
- Note: North border of main distribution area.
- Alticinae** Newman, 1834
Phyllotreta Chevrolat, 1837
armoraciae (Koch, 1803)
atra (Fabricius, 1775)
aethiopissa (Schrank, 1789)
atterima (Schrank, 1798)
punctata (Marsham, 1802)
- cruciferae* (Goeze, 1777)
colorea Foudras, 1860
elongata (Stephens, 1834)
obscurella (Illiger, 1807)
poeciloceras (Comolli, 1837)
- exclamationis* (Thunberg, 1784)
brassicae (Olivier, 1789)
quadripustulata (Paykull, 1799)
- flexuosa* (Illiger, 1794)
fallax Allard, 1860
- nemorum* (Linnaeus, 1758)
nigripes (Fabricius, 1775)
lens Thunberg, 1874
lepidii (Koch, 1803)
- nigroaenea* Roger, 1856
talassicola Heikertinger, 1944
ochripes (Curtis, 1837)
caucasica Harold, 1875
excisa Redtenbacher, 1849
- striolata*** (Fabricius, 1803)
vittata auct.
- tetrastigma*** (Comolli, 1837)
undulata Kutschera, 1860
australis Blackburn, 1890
blackburni Bryant, 1925
- vittula* (Redtenbacher, 1849)
rivularis Motschulsky, 1849
- Aphthona*** Chevrolat, 1837
abdominalis (Duftschmid, 1825)
finitima Ogloblin, 1927
- *aeneomicans* Allard, 1875
metallica Weise, 1893
- cyparissiae* (Koch, 1803)
erichsoni (Zetterstedt, 1838)
curvifrons Bach, 1859
nigritarsis Motschulsky, 1845
- euphorbiae*** (Schrank, 1781)
atratula Allard, 1859
hilaris (Stephens, 1831)
punctigera Mulsant, Rey, 1874
subaptera Mulsant, Rey, 1874
- *flaviceps* Allard, 1859
— *gracilis* Faldermann, 1837
- lutescens* (Gyllenhal, 1813)
sternalis Mulsant, Rey, 1874
- *nigriceps* (Redtenbacher, 1842)
decorata Kutschera, 1861
sicula Foudras, 1860
- nonstriata* (Goeze, 1777)
coerulea Geoffroy, 1785
plenifrons Wollaston, 1864
pseudoacori Marsham, 1802
thoracica Ogloblin, 1917
- pallida* Bach, 1856
- pygmaea* (Kutschera, 1861)
violacea (Koch, 1803)
ciscaucasica L.Medvedev, 1962
sublaevis Boheman, 1851
tbilisiensis Král, 1969
- Longitarsus*** Berthold, 1827
absynthii Kutschera, 1862
aeneicollis (Faldermann, 1837)
rubenticollis Allard, 1860
suturalis auct.
- *aeruginosus* (Foudras, 1859)
anchusae (Paykull, 1799)
apicalis (Beck, 1817)
atricillus (Linnaeus, 1761)
ballotae (Marsham, 1802)

<i>canescens</i> Foudras, 1860	<i>lateralis</i> (Illiger, 1807)
<i>brunneus</i> (Duftschmid, 1825)	<i>longipes</i> Bach, 1859
<i>castaneus</i> (Duftschmid, 1825)	<i>marginatus</i> (Geoffroy, 1785)
<i>curtus</i> (Allard, 1860)	<i>saltatrix</i> Wollaston, 1854
<i>liliputanus</i> Allard, 1866	<i>trilineolatus</i> Foudras, 1860
<i>papaveris</i> Allard, 1866	<i>obliteratus</i> (Rosenhauer, 1847)
<i>viduus</i> Allard, 1866	<i>consociatus</i> Forster, 1849
<i>exoletus</i> (Linnaeus, 1758)	<i>ochroleucus</i> (Marsham, 1802)
<i>boppardiensis</i> Bach, 1856	<i>albellus</i> (Duméril, 1816)
<i>exoletus</i> (Linnaeus, 1761)	<i>cognatus</i> Wollaston, 1860
<i>femoralis</i> (Marsham, 1802)	<i>parvulus</i> (Paykull, 1799)
<i>lividus</i> Fauvel, 1888	<i>maderensis</i> Allard, 1863
<i>ferrugineus</i> (Foudras, 1860)	<i>pumilis</i> (Illiger, 1807)
<i>dlabolai</i> Král, 1964	<i>pellucidus</i> (Foudras, 1860)
<i>menthae</i> Bedel, 1898	<i>ionius</i> Mohr, 1962
<i>mentaphagus</i> Gentner, 1926	<i>paleaceus</i> Mulsant, Rey, 1874
<i>waterhousei</i> Kutschera, 1864	<i>testaceus</i> Allard, 1860
<i>fulgens</i> (Foudras, 1860)	<i>pratensis</i> (Panzer, 1794)
<i>fuscoaeneus</i> Redtenbacher, 1849	<i>funereus</i> Mulsant, Rey, 1874
<i>ganglbaueri</i> Heikertinger, 1912	<i>medicaginis</i> Allard, 1860
<i>senecionis</i> Brisout de Barneville, 1873	<i>obsoletus</i> Mulsant, Rey, 1874
<i>gracilis</i> Kutschera, 1864	<i>pusillus</i> (Gyllenhal, 1813)
<i>helvolus</i> Kutschera, 1863	<i>quadriguttatus</i> (Pontoppidan, 1765)
<i>holsaticus</i> (Linnaeus, 1758)	<i>cynoglossi</i> (Marsham, 1802)
<i>haemorrhoidalis</i> Jacoby, 1855	<i>quadrimaculatus</i> (Koch, 1803)
<i>jacobaeae</i> (Waterhouse, 1861)	<i>quadrinotatus</i> (Gmelin, 1790)
<i>longipennis</i> Kutschera, 1863	<i>quadripunctatus</i> (Geoffroy, 1785)
<i>longisetata</i> Weise, 1889	<i>quadripunctulatus</i> Stephens, 1839
<i>clarus</i> Allen, 1967	<i>quadripustulatus</i> (Fabricius, 1775)
<i>kwangsiensis</i> Chen, 1939	<i>quatuorpustulatus</i> (Olivier, 1798)
<i>luridus</i> (Scopoli, 1763)	<i>rubiginosus</i> (Foudras, 1860)
<i>amurensis</i> Weise, 1887	<i>scutellaris</i> (Mulsant, Rey, 1874)
<i>castaneus</i> (Duftschmid, 1825)	<i>succineus</i> (Foudras, 1860)
<i>fuscescens</i> Stephens, 1831	<i>amiculus</i> Baly, 1874
<i>fusculus</i> Kutschera, 1864	<i>suturellus</i> (Duftschmid, 1825)
<i>syriacus</i> Allard, 1866	<i>fuscicollis</i> Stephens, 1831
<i>lycopi</i> (Foudras, 1860)	<i>senecionis</i> Motschulsky, 1851
<i>limnophilus</i> Abeille, 1907	<i>thoracicus</i> Allard, 1860
<i>seriatus</i> Kutschera, 1863	— <i>symphyti</i> Heikertinger, 1912
<i>melanocephalus</i> (DeGeer, 1775)	<i>tabidus</i> (Fabricius, 1775)
<i>atricapillus</i> (Duftschmid, 1825)	<i>jaceae</i> (Panzer, 1809)
<i>crassicornis</i> Foudras, 1860	<i>verbasci</i> (Panzer, 1794)
<i>piciceps</i> Stephens, 1831	<i>Altica</i> O.F.Müller, 1764
— <i>membranaceus</i> (Foudras, 1860)	<i>aenescens</i> (Weise, 1888)
<i>teucrii</i> Allard, 1860	<i>weisei</i> Kemner, 1919
<i>minusculus</i> (Foudras, 1860)	<i>ampelophaga</i> Guérin-Méneville, 1858
<i>monticola</i> Kutschera, 1863	<i>convexipennis</i> Ogloblin, 1921
<i>nanus</i> (Foudras, 1860)	<i>hispana</i> Allard, 1869
<i>brevicollis</i> Kutschera, 1863	Note: Introduced species.
<i>vitreus</i> Kutschera, 1863	<i>brevicollis</i> Foudras, 1860 s. str.
<i>nasturtii</i> (Fabricius, 1792)	<i>coryli</i> Allard, 1860
<i>niger</i> (Koch, 1803)	<i>carduorum</i> Guérin-Méneville, 1858
<i>elongatus</i> Bach, 1859	<i>basalis</i> Allard, 1859
<i>nigerrimus</i> (Gyllenhal, 1827)	<i>cirsii</i> Israelson, 1956
<i>nigrofasciatus</i> (Goeze, 1777)	<i>carinthiacea</i> (Weise, 1888)
	<i>daghestanica</i> Shapiro, 1970

- issykensis* Ogloblin, 1921
chamaenerii (H.Lindberg, 1926)
 — *deserticola* (Weise, 1889)
engstroemi (J.R.Sahlberg, 1894)
helianthemi (Allard, 1859)
 cognata Kutschera, 1864
 potentillae Allard, 1859
 pusilla Duftschmid, 1825
longicollis (Allard, 1860)
 britteni Sharp, 1914
 sandini Kemner, 1919
lythri Aubé, 1843
 epilobii Allard, 1859
oleracea (Linnaeus, 1758)
 alchemillae Palij, 1968
palustris (Weise, 1888)
quercetorum Foudras, 1860
tamaricis Schrank, 1785
 articulata Beck, 1817
 azurea Knoch, 1843
 consobrina Duftschmid, 1825
 erucae Fabricius, 1792
Batophila Foudras, 1859
fallax Weise, 1888
rubi (Paykull, 1799)
Lythraria Bedel, 1897
salicariae (Paykull, 1800)
Ochrosis Foudras, 1859
ventralis (Illiger, 1807)
Neocrepidodera Heikertinger, 1911
 — *crassicornis* (Faldermann, 1837)
 — *femorata* (Gyllenhal, 1813)
 femoralis (Duftschmid, 1825)
ferruginea (Scopoli, 1763)
 exoleta (Linnaeus, 1761)
 similis (Stephens, 1831)
interpunctata (Motschulsky, 1859)
motschulskii Konstantinov, 1991
nigritula (Gyllenhal, 1813)
 ovulum (Duftschmid, 1825)
transversa (Marsham, 1802)
Derocrepis Weise, 1886
rufipes (Linnaeus, 1758)
Hippuriphila Foudras, 1859
modeeri (Linnaeus, 1761)
 chrysopygis (Beck, 1817)
 mancula LeConte, 1861
Crepidodera Chevrolat, 1837
aurata (Marsham, 1802)
 versicolor Kutschera, 1860
aurea (Geoffroy, 1785)
 cyanea (Marsham, 1802)
 gaudens (Stephens, 1831)
 metallica (Duftschmid, 1825)
- fulvicornis* (Fabricius, 1792)
 smaragdina Foudras, 1860
lamina (Bedel, 1901)
nitidula (Linnaeus, 1758)
plutus (Latreille, 1804)
 chloris Foudras, 1860
Epithrix Foudras, 1859
pubescens (Koch, 1803)
Podagriva Chevrolat, 1837
 — *fuscicornis* (Linnaeus, 1766)
Mantura Stephens, 1831
chrysanthemi (Koch, 1803)
 reclairei Everts, 1921
obtusata (Gyllenhal, 1813)
 pallidicornis Waltl, 1839
 subobtusata Jansson, 1943
rustica (Linnaeus, 1767)
 semiaenea (Fabricius, 1792)
Chaetocnema Stephens, 1831
aerosa (Letzner, 1846)
arida Foudras, 1860
aridula (Gyllenhal, 1827)
 — *breviuscula* (Faldermann, 1837)
compressa (Letzner, 1847)
 tarda Bach, 1859
concinna (Marsham, 1802)
 dentipes (Koch, 1803)
confusa (Bohemian, 1851)
hortensis (Geoffroy, 1785)
 aridella (Paykull, 1799)
 scabricollis Allard, 1860
laevicollis (Thomson, 1866)
 heikertingeri Ljubitschev, 1963
mannerheimi (Gyllenhal, 1827)
obesa (Boieldieu, 1859)
 jurassica Pic, 1915
 meridionalis Foudras, 1860
sahlbergi (Gyllenhal, 1827)
semicoerulea (Koch, 1803)
 meridionalis Allard, 1859
subcoerulea (Kutschera, 1864)
 punctulata Mulsant, Rey, 1874
tibialis (Illiger, 1807)
 caesaraugustana Fuente, 1909
 obscuripes Pic, 1909
 pumila Allard, 1859
Dibolia Latreille, 1829
cryptocephala (Koch, 1803)
cynoglossi (Koch, 1803)
 marrubii Hering, 1924
depressiuscula Letzner, 1846
 laevicollis Foudras, 1860
occultans (Koch, 1803)

<i>timida</i> (Illiger, 1807)	<i>sophiae</i> Heikertinger, 1914 nec (Illiger, 1807)
<i>eryngii</i> Bach, 1859	
<i>Sphaeroderma</i> Stephens, 1831	Cassidinae Stephens, 1831
<i>rubidum</i> (Graëlls, 1858)	<i>Pilemostoma</i> Desbrochers des Loges, 1891
<i>nitidum</i> Portevin, 1934	<i>fastuosa</i> (Schaller, 1783)
<i>ocularium</i> Allard, 1860	<i>ocellata</i> (Herbst, 1799)
<i>testaceum</i> (Fabricius, 1775)	Hypocassida Weise, 1893
<i>cardui</i> (Gyllenhal, 1813)	<i>subferruginea</i> (Schrank, 1776)
<i>Argopus</i> Fischer, 1824	<i>cornea</i> (Marseul, 1869)
<i>ahrensii</i> (Germar, 1817)	<i>costata</i> (Boheman, 1856)
<i>hemisphaericus</i> Duftschmid, 1825	<i>ferruginea</i> (Fabricius, 1781)
<i>Psylliodes</i> Berthold, 1827	<i>fusca</i> (Laicharting, 1781)
<i>affinis</i> (Paykull, 1799)	Cassida Linneus, 1758
<i>attenuatus</i> (Koch, 1803)	<i>atrata</i> Fabricius, 1787
<i>vicus</i> Redtenbacher, 1849	<i>denticollis</i> Suffrian, 1844
<i>chalcomerus</i> (Illiger, 1807)	<i>ferruginea</i> Goeze, 1777
<i>allardi</i> Bach, 1866	<i>thoracica</i> Geoffroy, 1785
<i>brunnipes</i> (Duftschmid, 1825)	<i>tincta</i> Weise, 1891
<i>unimaculatus</i> Stephens, 1831	<i>flaveola</i> Thunberg, 1794
<i>chrysocephalus</i> (Linnaeus, 1758)	<i>exsculpta</i> Charpentier, 1825
<i>anglicanus</i> (Gmelin, 1790)	<i>obsoleta</i> Illiger, 1798
<i>atlanticus</i> Fauvel, 1897	<i>hemisphaerica</i> Herbst, 1799
<i>cyanopterus</i> (Illiger, 1807)	<i>anglica</i> Curtis, 1826
<i>nigricollis</i> (Marsham, 1802)	<i>concinna</i> Steffens, 1832
<i>nuceus</i> (Illiger, 1807)	<i>corii</i> Fairmaire, 1876
<i>sophiae</i> (Illiger, 1807)	<i>rossii</i> Wollaston, 1857
<i>cucullatus</i> <i>cucullatus</i> (Illiger, 1807)	<i>lineola</i> Creutzer, 1799
<i>cupreus</i> (Koch, 1803)	<i>bicostata</i> Fischer von Waldheim, 1842
<i>cypricolor</i> Allard, 1860	<i>nigroguttata</i> Gorham, 1885
<i>foudrasi</i> Bach, 1866	<i>signata</i> Herbst, 1799
<i>herbaceus</i> Foudras, 1859	<i>suturalis</i> Fischer von Waldheim, 1842
<i>obscurus</i> (Duftschmid, 1825)	<i>margaritacea</i> Schaller, 1783
<i>politus</i> (Gravenhorst, 1807)	<i>mutabilis</i> Villers, 1789
<i>cupreatus</i> (Duftschmid, 1825)	<i>spergulae</i> Marsham, 1802
– <i>deplanatus</i> L.Medvedev, 1962	<i>superba</i> Gmelin, 1787
<i>dulcamarae</i> (Koch, 1803)	<i>murraea</i> Linnaeus, 1767
<i>hyoscyami</i> (Linnaeus, 1758)	<i>subobliterata</i> Pic, 1902
<i>coeruleus</i> (Paykull, 1799)	<i>varia</i> Latreille, 1804
<i>luteolus</i> (O.F.Müller, 1776)	<i>variegata</i> Geoffroy, 1785
<i>nana</i> (Duftschmid, 1825)	<i>nebulosa</i> Linnaeus, 1758
<i>pallida</i> Stephens, 1831	<i>affinis</i> Fabricius, 1775
<i>propinqua</i> Redtenbacher, 1849	<i>tigrina</i> DeGeer, 1775
<i>marcidus</i> (Illiger, 1807)	<i>nobilis</i> Linnaeus, 1758
<i>operosa</i> Foudras, 1860	<i>laevis</i> Herbst, 1799
<i>napi</i> (Fabricius, 1792)	<i>pulchella</i> Panzer, 1797
<i>ecalcarata</i> Redtenbacher, 1849	<i>splendidula</i> Marsham, 1802
<i>lunariae</i> Jacquet, 1877	<i>urticae</i> Brahm, 1790
<i>rapae</i> (Illiger, 1807)	<i>viridula</i> Paykull, 1798
<i>picinus</i> (Marsham, 1802)	– <i>pannonica</i> Suffrian, 1844
<i>apicalis</i> Stephens, 1831	<i>panzeri</i> Weise, 1907
<i>melanophthalmus</i> (Duftschmid, 1825)	<i>prasina</i> Illiger, 1798
<i>piceus</i> Redtenbacher, 1849	<i>chloris</i> Suffrian, 1844
<i>rufilabris</i> (Koch, 1803)	<i>viridana</i> Herbst, 1799
<i>reitteri</i> Weise, 1888	<i>rubiginosa</i> O.F.Müller, 1776
– <i>tricolor</i> Weise, 1888	<i>graeca</i> Kraatz, 1874
<i>cyanopterus</i> Redtenbacher, 1849 nec (Ill., 1807)	<i>melanoscelis</i> Schrank, 1798
	<i>similis</i> Marsham, 1802

- singularis* Stephens, 1832
sanguinolenta O.F.Müller, 1776
 cruentata Donovan, 1793
sanguinosa Suffrian, 1844
 languida Cornelius, 1851
seladonia Gyllenhal, 1827
stigmatica Suffrian, 1844
subreticulata Suffrian, 1844
vibex Linnaeus, 1767
 angusticollis Habenbach, 1822
 dorsalis Herbst, 1799
 liriophora Kirby, 1797
viridis Linnaeus, 1758
 equestris Fabricius, 1787
vittata Villers, 1789
 atlantica Escalera, 1914
 oblonga Illiger, 1798
 salicorniae Curtis, 1826
- Curculionoidea**
- Nemonychidae** Bedel, 1883
- Doydirhynchinae** Pierce, 1916
- Cimberidini** Des Gozis, 1882
- Cimberis** Des Gozis, 1882
- attelaboides* (Fabricius, 1787)
 canescens Semenov, 1900
 rhinomacer (Paykull, 1792)
- Doydirhynchini** Pierce, 1916
- Doydirhynchus** Dejean, 1821
- austriacus* (Olivier, 1807)
 castaneus Germar, 1833
 fulvipennis Reitter, 1916
 karamani Stierlin, 1886
 lutescens Schilsky, 1903
 pallidicolor Pic, 1905
 testaceus Voss, 1932
- Anthribidae** Billberg, 1820
- Anthribinae** Billberg, 1820
- Gonotropis** LeConte, 1876
- dorsalis* *dorsalis* (Thunberg, 1796)
- Rhaphitropis** Reitter, 1916
- marchicus* (Herbst, 1797)
 cinctus (Bach, 1854)
 maculosus (Mulsant, 1859)
 reyi (Des Gozis, 1882)
 scriptus (Thunberg, 1815)
- Platyrhinus** Clairville, Schellenberg, 1798
- resinosus* (Scopoli, 1763)
 ater (Geoffroy, 1762)
 costirostris Clairville, Schellenberg, 1798
 flavifrons (Fuessly, 1775)
 latirostris (Fabricius, 1792)
 oblingus (Sulzer, 1776)
- Note: This species is included on the list of indicator species of natural forest habitats.
- Dissoleucas** Jordan, 1925
- niveirostris* (Fabricius, 1798)
 dubius (Ponza, 1805)
- Platystomus** Schneider, 1791
- albinus* (Linnaeus, 1758)
 desertus Schilsky, 1907
 thierieti Viturat, 1895
 uniformis Reitter, 1916
- Anthribus** O.F.Müller, 1764
- fasciatus* Forster, 1770
 marmoratus Geoffroy, 1785
 rubripes (Rey, 1893)
 rufipes Schilsky, 1907
 scabrosus Fabricius, 1775
 ventralis (Rey, 1893)
- nebulosus* Forster, 1771
 clathratus (Herbst in Fuessly 1786)
 gavoyi Chobaut, 1922
 kuesteri Reitter, 1916
 nebulosus (Küster, Bose-Gutf., 1859)
 variegatus Geoffroy, 1785
 varius (Fabricius, 1787)
- scapularis* Gebler, 1833
 constrictus (Stierlin, 1867)
 gebleri (Reitter, 1916)
 rosae (Reitter, 1916)
- Choraginae** Kirby, 1819
- Choragus** Kirby, 1819
- horni* Wolfrum, 1930
- Rhynchitidae** Gistel, 1848
- Rhynchititae** Gistel, 1848
- Isotheini** Scudder, 1893
- Deporaus** (*Deporaus*) Samouelle, 1819
- betulae* (Linnaeus, 1758)
 excordatoniger (DeGeer, 1775)
 fagi (Scopoli, 1763)
 femoralis (Latreille, 1804)
 femoratus (Olivier, 1789)
 populi (Scopoli, 1763)
- Caenorhinus** (*Caenorhinus*) Thomson, 1859
- mannerheimi* (Hummel, 1823)
 alliarie (Stephens, 1831)
 constrictus (Gyllenhal, 1839)
 cyaneipennis (Stephens, 1831)
 laevicollis (Stephens, 1831)
 megacephalus (Germar, 1824)
 planipennis Roelofs, 1874
- Rhynchitini** Gistel, 1848
- Temnocerus** Thunberg, 1815
- caeruleus* (Fabricius, 1798)
 tomentosus (Gyllenhal, 1839)
 uncinatus (Thomson, 1865)
- longiceps* (Thomson, 1888)
 harwoodi (Joy, 1911)
- nanus* (Paykull, 1792)

<i>angustifrons</i> (Voss, 1932)	<i>violaceus</i> (Poda, 1761)
<i>cylindricus</i> (Stephens, 1831)	<i>viridis</i> (Geoffroy, 1785)
<i>planirostris</i> (Fabricius, 1801)	<i>viridulus</i> (Westhoff, 1882)
<i>Neocoenorrhinus</i> Voss, 1952	<i>populi</i> (Linnaeus, 1758)
<i>germanicus</i> (Herbst, 1797)	<i>cupreus</i> (Wasmann, 1887)
<i>fauconneti</i> (Pic, 1926)	<i>cuprifer</i> Schilsky, 1903
<i>rugipennis</i> (Stephens, 1831)	<i>nigripes</i> Faust, 1882
<i>Neocoenorhinidius</i> Legalov, 2003	<i>nigritulus</i> Schilsky, 1889
<i>pauxillus</i> (Germar, 1824)	<i>parvulus</i> Sharp, 1889
<i>alliariae</i> (Rossi, 1790 nec Fabricius, 1775)	<i>tartaricus</i> Faust, 1882
<i>atrococeruleus</i> (Stephens, 1831)	
<i>livescens</i> (Voss, 1929)	Attelabidae Billberg, 1820
<i>pauculus</i> (Voss, 1933)	Attelabinae Billberg, 1820
<i>persicus</i> (Gyllenhal, 1839)	Attelabini Billberg, 1820
<i>Tatianaerhynchites</i> (<i>Tatianaerhynchites</i> Legalov, 2002)	<i>Attelabus</i> (<i>Attelabus</i> Linnaeus, 1758)
<i>aequatus</i> (Linnaeus, 1767)	<i>nitens</i> (Scopoli, 1763)
<i>bicolor</i> (Rossi, 1790)	<i>ater</i> Fleischer, 1887
<i>creticus</i> (Voss, 1955)	<i>atricornis</i> Mulsant, Guillebeau, 1855
<i>fulgidus</i> (Geoffroy, 1785)	<i>balcanicus</i> K.Daniel, J.Daniel, 1898
<i>interstitialis</i> (Desbrochers des Loges, 1869)	<i>coccineus</i> (Geoffroy, 1785)
<i>paykulli</i> (Schilsky, 1903)	<i>coryli</i> O.F.Müller, 1776
<i>phryganophilus</i> (Iablokoff-Khnzorian, 1956)	<i>curculionides</i> Linnaeus, 1767
<i>purpureus</i> (Goeze, 1777)	<i>curculioniformis</i> Schrank, 1798
<i>ruber</i> (Geoffroy, 1785)	<i>fascicollis</i> Reitter, 1916
<i>ruber</i> (Fairmaire, 1859)	<i>hispanicus</i> Jaekel, 1860
<i>semiruber</i> (Stierlin, 1863)	<i>maculipes</i> Rondani, 1833
<i>Epirhynchites</i> (<i>Tshernyshevius</i> Legalov, 2003)	<i>marginalis</i> Pic, 1898 nec Reitter, 1916
<i>auratus</i> (Scopoli, 1763)	<i>marginalis</i> Reitter, 1916 nec Pic, 1898
<i>angorensis</i> (Pic, 1926)	<i>marginatus</i> Wasmann, 1887
<i>armeniacus</i> (Zaitzev, 1937)	<i>obscurior</i> Pic, 1898
<i>chamaecerasi</i> (Ter-Minassian, 1950)	<i>obsidianus</i> A.Costa, 1886
<i>ferghanensis</i> (Nevsky, 1928)	<i>pseudosuturalis</i> Ter-Minassian, 1950
<i>microcarpae</i> (Ter-Minassian, 1944)	<i>pulvinicollis</i> Jaekel, 1860
<i>minor</i> (Schilsky, 1903)	<i>rubricollis</i> Reitter, 1916
<i>theressae</i> (Pic, 1926)	<i>rufipes</i> Schilsky, 1903
<i>Involvulus</i> Schrank, 1798	<i>suturalis</i> Jacobs, 1926
<i>cupreus</i> (Linnaeus, 1761)	Apoderinae Jaekel, 1860
<i>aeneus</i> (Latreille, 1804)	Apoderini Jaekel, 1860
<i>dybofskyi</i> (Faust, 1882)	<i>Compsapoderus</i> (<i>Compsapoderus</i> Voss, 1927)
<i>metallicus</i> Schrank, 1798	<i>erythropterus</i> (Gmelin, 1790)
<i>punctatus</i> (Herbst, 1797)	<i>atricolor</i> (Faust, 1887)
<i>Byctiscini</i> Voss, 1923	<i>bicolor</i> (Redtenbacher, 1868)
<i>Byctiscus</i> Thomson, 1859	<i>frontalis</i> (Faust, 1882)
<i>betulae</i> (Linnaeus, 1758)	<i>intermedius</i> (Illiger, 1794)
<i>alni</i> (O.F.Müller, 1776)	<i>politus</i> (Gebler, 1825)
<i>betuleti</i> (Fabricius, 1792)	<i>Apoderus</i> (<i>Apoderus</i> Olivier, 1807)
<i>bispinus</i> (O.F.Müller, 1776)	<i>coryli</i> (Linnaeus, 1758)
<i>destitulus</i> Voss, 1930	<i>alni</i> Chinaglia, 1908
<i>cuprinus</i> Schilsky, 1903	<i>atricollis</i> Gortaniet, Grandi, 1904
<i>inermis</i> (O.F.Müller, 1776)	<i>avellanae</i> (Linnaeus, 1767)
<i>intermedius</i> Voss, 1930	<i>carnicus</i> Gortaniet, Grandi, 1904
<i>nitens</i> (Marsham, 1802)	<i>collaris</i> (Scopoli, 1763)
<i>populi</i> (Paykull, 1792)	<i>dauricus</i> (Laxmann, 1770)
<i>unispinus</i> (O.F.Müller, 1776)	<i>denigratus</i> (Gmelin, 1790)
<i>violacea</i> (Donovan, 1794)	<i>dubius</i> Gortaniet, Grandi, 1904
	<i>excoriatoruber</i> (DeGeer, 1775)
	<i>fiorii</i> Gortaniet, Grandi, 1904
	<i>gibbicollis</i> Faust, 1882
	<i>homalinus</i> Voss, 1927

- kamtschaticus* Motschulsky, 1845
lineatus Gortaniet, Grandi, 1904
morio (Bonelli, 1812)
niger Gortaniet, Grandi, 1904
nigricollis Faust, 1882
nigrifrons Della Beffa, 1912
nigripes Rey, 1893
rubricollis Gortaniet, Grandi, 1904
rubripes Chinaglia, 1908
ruficeps Rey, 1893
superans Faust, 1882
superbus Schilsky, 1903
- Apionidae** Schönherr, 1823
- Apioninae** Schönherr, 1823
- Ceratapiini** Alonso-Zarazaga, 1990
- Taphrotopium** Reitter, 1916
sulcifrons (Herbst, 1797)
- Omphalapion** Schilsky, 1901
dispar (Germar, 1817)
hookerorum (Kirby, 1808)
hookeri (Kirby, 1808)
- Diplapion** Reitter, 1916
confluens (Kirby, 1808)
stolidum (Germar, 1817)
- Ceratapion (Acanephodus)** Alonso-Zarazaga, 1990
basicorne (Illiger, 1807)
alliariae auct. nec (Linnaeus, 1758)
onopordi onopordi (Kirby, 1808)
- Ceratapion (Echinostroma)** Alonso-Zarazaga, 1990
penetrans penetrans (Germar, 1817)
- Ceratapion (Ceratapion)** Schilsky, 1901
armatum (Gerstaecker, 1854)
austriacum (Wagner, 1904)
gibbirostre (Gyllenhal, 1813)
carduorum (Gyllenhal, 1813 nec Kirby, 1808)
- Kalcapiini** Alonso-Zarazaga, 1990
- Melanapion (Melanapion)** Wagner, 1930
minimum (Herbst, 1797)
- Squamapion** Bokor, 1923
atomarium (Kirby, 1808)
elongatum (Germar, 1817)
flavimanum (Gyllenhal, 1833)
- Taeniapion** Schilsky, 1906
urticarium urticarium (Herbst, 1784)
- Kalcapion** Schilsky, 1906
pallipes (Kirby, 1808)
- Aspidapiini** Alonso-Zarazaga, 1990
- Aspidapion (Aspidapion)** Schilsky, 1901
radiolus radiolus (Marsham, 1802)
- Ixapiini** Alonso-Zarazaga, 1990
- Trichopterapion** Wagner, 1930
holosericeum (Gyllenhal, 1833)
- Piezotrachelini** Voss, 1959
- Pseudoprotapion** Ehret, 1990
astragali astragali (Paykull, 1800)
- Protapion** Schilsky, 1908
apricans (Herbst, 1797)
fagi (Kirby, 1808)
flavifemoratum (Kirby, 1808) in part
flavipes (Panzer, 1795 nec Paykull, 1792 nec Fabricius, 1775)
ochropus (Gmelin, 1777)
tubicen (Wencker, 1864) in part (female)
- assimile* (Kirby, 1808)
bohemani (Bedel, 1885 nec Thomson, 1865)
incertum (Desbrochers des Loges, 1894-95)
propinquum (Desbr. des Loges, 1908 nec Hartmann, 1907)
ryei (Blackburn, 1874)
- dissimile* (Germar, 1817)
adjectum (Desbrochers des Loges, 1894-95)
heterocerum (Thomson, 1865)
- filiostre* (Kirby, 1808)
brevirostre (Schilsky, 1902 nec Gerstaecker, 1854)
cantianum (Wagner, 1906-08)
morio (Germar, 1817)
- fulvipes* (Geoffroy, 1785)
bergrothi (Desbrochers des Loges, 1894-95)
burgodionum (Schubert, 1957)
dichroum (Bedel, 1885)
flavipes (Paykull, 1792 nec Fabricius, 1775)
rosai (Pesarini, 1957)
- interjectum interjectum* (Desbrochers des Loges, 1895)
oblitum (Desbrochers des Loges, 1894-95 nec Smith, 1884)
- nigritarse* (Kirby, 1808)
concoloripes (Desbrochers des Loges, 1908)
waterhousei (Bohemian, 1839)
- ononidis* (Gyllenhal, 1827)
bohemani (Thomson, 1865 nec Bedel, 1885)
ononicola (Bach, 1854)
- ruficrus* (Germar, 1817)
- trifolii* (Linnaeus, 1768)
aestivum (Germar, 1817)
flavifemoratum (Kirby, 1808)
flavipes (Laicharting, 1871)
leachi (Stephens, 1831)
linderi (Wencker, 1858)
- Aplemonini** Kissinger, 1968
- Pseudostenapion** Wagner, 1930
simum (Germar, 1817)
- Aizobius** Alonso-Zarazaga, 1990
sedi (Germar, 1817)
- Perapion (Perapion)** Wagner, 1907

<i>affine</i> (Kirby, 1808)	<i>cerdo</i> (Gerstaecker, 1854)
<i>curtirostre</i> (Germar, 1817)	<i>craccae</i> (Linnaeus, 1767)
<i>marchicum</i> (Herbst, 1797)	<i>opeticum</i> (Bach, 1854)
<i>oblongum</i> (Gyllenhal, 1839)	<i>pomonae</i> (Fabricius, 1798)
<i>violaceum violaceum</i> (Kirby, 1808)	<i>subulatum</i> (Kirby, 1808)
Apionini Schönherr, 1823	Eutrichapion (<i>Eutrichapion</i> Reitter, 1916)
Apion Herbst, 1797	<i>ervi</i> (Kirby, 1808)
<i>cruentatum</i> Walton, 1844	<i>viciae</i> (Paykull, 1800)
<i>frumentarium</i> (Linnaeus, 1758)	Eutrichapion (<i>Psilocalymma</i> Alonso-Zarazaga, 1990)
<i>miniatum</i> Germar, 1833	<i>facetum</i> (Gyllenhal, 1839)
<i>sanguineum</i> (DeGeer, 1775)	Eutrichapion (<i>Cnemapion</i> Bokor, 1923)
<i>haematodes</i> Kirby, 1808	<i>vorax</i> (Herbst, 1797)
<i>frumentarium</i> (Paykull, 1792 nec Linnaeus, 1758)	Nanophyinae Gistel, 1856
<i>rubens</i> Stephens, 1839	Nanophyes Schönherr, 1838
<i>rubiginosum</i> Grill, 1893	<i>marmoratus marmoratus</i> (Goeze, 1777)
<i>sanguineum</i> Grill, 1893 nec (DeGeer, 1775)	Nanomimus Alonso-Zarazaga, 1989
Oxystomatini Alonso-Zarazaga, 1990	<i>circumscriptus</i> (Aubé, 1864)
Catapion Schilsky, 1906	Curculionidae Latreille, 1802
<i>pubescens</i> (Kirby, 1811)	Otiorhynchinae Schönherr, 1826
<i>seniculus</i> (Kirby, 1808)	Otiorhynchini Schönherr, 1826
Betulapion Ehret, 1994	Otiorhynchus Germar, 1824
<i>simile simile</i> (Kirby, 1811)	— <i>aurosparsus</i> Germar, 1824
Stenopterapion (<i>Stenopterapion</i> Bokor, 1923)	<i>conspersus</i> (Herbst, 1795)
<i>meliloti</i> (Kirby, 1808)	<i>desertus</i> Rosenhauer, 1847
<i>tenue</i> (Kirby, 1808)	<i>pellucidus</i> J.R.Sahlberg, 1900
Ischnopterapion (<i>Ischnopterapion</i> Bokor, 1923)	<i>ligustici</i> (Linnaeus, 1758)
<i>loti</i> (Kirby, 1808)	<i>lugdunensis</i> (Boheman, 1843)
Ischnopterapion (<i>Chlorapion</i> Györffy, 1956)	<i>nodosus</i> (O.F.Müller, 1764)
<i>virens</i> (Herbst, 1797)	<i>dubius</i> (Ström, 1783)
Synapion Schilsky, 1902	<i>maurus</i> (Gyllenhal, 1813)
<i>ebeninum</i> (Kirby, 1808)	<i>ovatus ovatus</i> (Linnaeus, 1758)
Holotrichapion (<i>Apiops</i> Alonso-Zarazaga, 1990)	<i>raucus raurus</i> (Fabricius, 1776)
<i>pisi</i> (Fabricius, 1801)	<i>rugifrons</i> (Gyllenhal, 1813)
<i>pullum</i> (Gyllenhal, 1833)	<i>scaber</i> (Linnaeus, 1758)
<i>aestimatum</i> (Faust, 1890)	<i>setosus</i> (Fabricius, 1801)
Holotrichapion (<i>Legaricaption</i> Ehret, 1990)	<i>singularis</i> (Linnaeus, 1767)
<i>aethiops</i> (Herbst, 1797)	<i>picipes</i> (Fabricius, 1776)
Hemitrichapion (<i>Dimesomyops</i> Alonso-Zarazaga 1990)	<i>smreczynskii</i> Cmoluch, 1968
<i>pavidum</i> (Germar, 1817)	Note: An introduced species.
Cyanapion (<i>Cyanapion</i> Bokor, 1923)	<i>sulcatus</i> (Fabricius, 1775)
<i>spencii</i> (Kirby, 1808)	<i>tristis</i> (Scopoli, 1763)
Cyanapion (<i>Bothyorrhynchapion</i> Bokor, 1923)	<i>nigrata</i> (Fabricius, 1776)
<i>affer</i> (Gyllenhal, 1833)	Trachyphloeus Germar, 1817
<i>platalea</i> (Gyllenhal, 1833 nec Germar, 1817)	<i>angustisetulus</i> Hansen, 1915
<i>gyllehalii</i> (Kirby, 1808)	<i>aristatus</i> (Gyllenhal, 1827)
Oxystoma Duméril, 1806	<i>bifoveolatus</i> (Beck, 1817)
	<i>scaber</i> auct. nec (Linnaeus, 1758)
	<i>digitalis</i> Gyllenhal, 1827
	<i>spinimanus</i> Germar, 1824
	Omiini Shuckard, 1840

- Omiamima*** Silfverberg, 1977
 — *concinna* (Bohemian, 1843)
 — *mollina* (Bohemian, 1834)
- Phyllobiini*** Schönherr, 1826
- Phyllobius*** Germar, 1824
arborator (Herbst, 1797)
argentatus (Linnaeus, 1758)
betulinus (Bechstein, Scharfenberg, 1805)
 betulae (Fabricius, 1801 nec Linnaeus, 1758)
calcaratus (Fabricius, 1792)
 glaucus (Scopoli 1763)
maculicornis Germar, 1824
oblongus (Linnaeus, 1758)
pomaceus Gyllenhal, 1834
 alneti (Fabricius, 1792 nec Schrank, 1781)
 urticæ (DeGeer, 1775 nec Scopoli, 1763)
pyri (Linnaeus, 1758)
 — *roboretanus* Gredler, 1882
 parvulus sensu (Olivier, 1807)
- vespertinus* (Fabricius, 1792)
 mali (Paykull, 1792) in part
- virideaeris* (Laicharting, 1781)
 pomonae Olivier, 1807
- viridicollis* (Fabricius, 1792)
 cloropus (Linnaeus, 1758)
- Brachyderini*** Schönherr, 1826
- Polydrusus*** Germar, 1817
cervinus (Linnaeus, 1758)
corruscus Germar, 1824
flavipes (DeGeer, 1775)
fulvicornis (Fabricius, 1792)
 ruficornis (Bonsdorff, 1785 nec Linnaeus, 1758)
 — *impar* Des Gozis, 1882
impressifrons Gyllenhal, 1834
- Note: A species increasing its distribution.
- mollis* (Ström, 1768)
 micans (Fabricius, 1792)
pallidus (Gyllenhal, 1834)
 atomarius (Olivier, 1807 nec Linnaeus, 1761)
- pilosus* Gredler, 1866
 binotatus Thomson, 1868
- pterygomalis* Boheman, 1840
- undatus* (Fabricius, 1781)
 tereticollis auct. nec (DeGeer, 1775)
- Eusomus*** Germar, 1824
 — *ovulum* Germar, 1824
- Sciaphilus*** Schönherr, 1823
- asperatus* (Bonsdorff, 1785)
 muricatus (Fabricius, 1792)
- Brachysomus*** Schönherr, 1823
- echinatus* (Bonsdorff, 1785)
- Barypeithes*** Jacquelin du Val, 1854
- pellucidus* (Bohemian, 1834)
 gabrieli Penecke, 1928
- trichopterus* (Gautier, 1863)
 violatus Seidlitz, 1868
- Brachyderes*** Schönherr, 1823
incanus (Linnaeus, 1758)
- Strophosomini*** Gistel, 1856
- Strophosoma*** Billberg, 1820
capitatum (DeGeer, 1775)
 rufipes Stephens, 1831
- faber* (Herbst, 1785)
fulvicorne Walton, 1848
 curvipes Thomson, 1865
- melanogrammum* (Forster, 1771)
- Cneorhinini*** Lacordaire, 1863
- Liophloeus*** Germar, 1817
tessulatus (O.F.Müller, 1776)
 maurus (Marsham, 1802)
 nubilis (Fabricius, 1776)
- Philopedon*** Stephens, 1831
plagiatus (Schaller, 1783)
 geminatus (Fabricius, 1787)
- Barynotini*** Lacordaire, 1863
- Barynotus*** Germar, 1817
moerens (Fabricius, 1792)
 elevatus (Marsham, 1802)
- obscurus* (Fabricius, 1775)
 murinus (O.F.Müller, 1776)
- Tropiphorini*** Leng, 1920
- Tropiphorus*** Schönherr, 1842
elevatus (Herbst, 1795)
- Tanymecini*** Lacordaire, 1863
- Chlorophanus*** C.R.Sahlberg, 1823
graminicola (Schönherr, 1832)
- Note: This species is known from extreme South of Latvia only (Skaistkalne area).
- viridis* (Linnaeus, 1758)
- Tanymecus*** Germar, 1817
palliatus (Fabricius, 1787)
- Sitoninae*** Gistel, 1856
- Sitonini*** Gistel, 1856
- Sitona*** Germar, 1817
ambiguus Gyllenhal, 1834
 lineellus auct. nec (Bonsdorff, 1785)
- cylindricollis* Fahraeus, 1840
- griseus* (Fabricius, 1775)
- hispidulus* (Fabricius, 1776)
- humeralis* Stephens, 1831
- inops* Gyllenhal, 1832
- lepidus* Gyllenhal, 1834
 flavescens (Marsham, 1802 nec Fabricius, 1787)
- lineatus* (Linnaeus, 1758)

<i>lineellus</i> (Bonsdorff, 1785)	<i>dealbatus</i> (Gmelin, 1790)
<i>decipiens</i> H.Lindberg, 1933	<i>marmoratus</i> (Fabricius, 1792)
<i>longulus</i> Gyllenhal, 1834	<i>tigrinus</i> (Panzer, 1789 nec Geoffroy, 1785)
<i>macularius</i> (Marsham, 1802)	Cleonis Dejean, 1821
<i>crinitus</i> (Herbst, 1795 nec Gmelin, 1790)	<i>pigra</i> (Scopoli, 1763)
<i>ononidis</i> Sharp, 1866	<i>sulcirostris</i> (Linnaeus, 1767)
<i>puncticollis</i> Stephens, 1831	Erirhinae Schönherr, 1825
<i>striatellus</i> Gyllenhal, 1834	Rhytirrhinini Lacordaire, 1863
<i>tibialis</i> (Herbst, 1795 nec Sparrman, 1785)	Gronops Schönherr, 1823
<i>sulcifrons</i> (Thunberg, 1798)	<i>inaequalis</i> Boheman, 1842
<i>suturalis</i> Stephens, 1831	<i>lunatus</i> (Fabricius, 1775)
<i>elegans</i> Gyllenhal, 1834	Phytonomini Gistel, 1856
Cleoninae Schönherr, 1826	Hypera Germar, 1817
Lixini Schönherr, 1826	<i>adspersa</i> (Fabricius, 1792)
Larinus Schönherr, 1823	<i>alternans</i> Stephens, 1834
<i>jaceae</i> (Fabricius, 1775)	<i>julini</i> R.F.Sahlberg, 1843
<i>planus</i> (Fabricius, 1792)	<i>pollux</i> (Fabricius, 1801)
<i>carlinae</i> (Olivier, 1807)	<i>arator</i> (Linnaeus, 1758)
<i>sturnus</i> (Schaller, 1783)	<i>polygoni</i> (Linnaeus, 1761)
Lixus Fabricius, 1801	<i>arundinis</i> (Paykull, 1792)
<i>bardanae</i> (Fabricius, 1787)	<i>contaminata</i> (Herbst, 1795)
<i>cylindrus</i> (Fabricius, 1787 nec Herbst, 1783)	<i>dauci</i> (Olivier, 1807)
<i>iridis</i> Olivier, 1807	<i>fasciulata</i> (Herbst, 1795 nec DeGeer, 1775)
<i>turbatus</i> Gyllenhal, 1827	<i>diversipunctata</i> (Schrank, 1798)
<i>paraplecticus</i> (Linnaeus, 1758)	<i>elongata</i> (Paykull, 1792 nec Fabricius, 1775)
— <i>punctiventris</i> Boheman, 1836	<i>variegata</i> (Bach, 1854)
Rhinocyllini Lacordaire, 1863	<i>fuscocinerea</i> (Marsham, 1802)
Rhinocyllus Germar, 1819	<i>murina</i> (Fabricius, 1792 nec O.F.Müller, 1764)
<i>conicus</i> (Frölich, 1792)	<i>meles</i> (Fabricius, 1792)
Note: This species is known from single locality in Slītere area (Slīteres National park, NW Latvia), but it should be more widely distributed in the whole country. A species increasing its distribution.	<i>trifolii</i> (Herbst, 1795)
Cleonini Schönherr, 1826	<i>nigrirostris</i> (Fabricius, 1775)
Conioleononus Motschulsky, 1860	<i>variabilis</i> (Fabricius, 1776)
<i>hollbergi</i> (Fahraeus, 1842)	<i>plantaginis</i> (DeGeer, 1775)
<i>glaucus</i> (Fabricius, 1787 nec Scopoli, 1763)	<i>olivieri</i> Capiomont, 1868
<i>turbatus</i> (Fahraeus, 1842)	<i>postica</i> (Gyllenhal, 1813)
<i>nebulosus</i> (Linnaeus, 1758)	<i>variabilis</i> (Herbst, 1795 nec Fabricius, 1776)
— <i>nigrosuturatus</i> (Goeze, 1777)	<i>rumicis</i> (Linnaeus, 1758)
Stephanocleonus Motschulsky, 1860	<i>suspiciosa</i> (Herbst, 1795)
<i>tetragrammus</i> (Pallas, 1781)	<i>meles</i> (Gyllenhal, 1813)
<i>concinnus</i> (Gyllenhal, 1834)	<i>pedestris</i> (Paykull, 1792 nec Poda, 1761)
Chromoderus Motschulsky, 1860	<i>venusta</i> (Fabricius, 1781)
<i>affinis</i> (Schrank, 1781)	<i>trilineata</i> (Marsham, 1802)
<i>albidus</i> (Fabricius, 1787)	<i>viciae</i> (Gyllenhal, 1813)
<i>fasciatus</i> (O.F.Müller, 1776 nec Scopoli, 1763)	<i>zoilus</i> (Scopoli, 1763)
Mecaspis Schönherr, 1823	<i>medius</i> (Marsham, 1802)
— <i>striatellus</i> (Fabricius, 1792)	<i>punctata</i> (Fabricius, 1775)
Cyphocleonus Motschulsky, 1860	Erirhinini Schönherr, 1825
	Grypus Germar, 1817
	<i>brunnirostris</i> (Fabricius, 1792)
	<i>equiseti</i> (Fabricius, 1775)
	Thryogenes Bedel, 1884
	<i>festucae</i> (Herbst, 1795)
	<i>nereis</i> (Paykull, 1800)

- scirrhosus* (Gyllenhal, 1836)
brunneirostris (Olivier, 1807)
- Notaris** Germar, 1817
acridulus (Linnaeus, 1758)
aethiops (Fabricius, 1792)
bimaculatus (Fabricius, 1787)
maerkeli (Bohemian, 1843)
rhamni Herbst, 1795
scirpi (Fabricius, 1787 nec Rossi 1790)
- Dorytomus** Germar, 1817
dejeani Faust, 1882
costirostris Gyllenhal, 1835
dorsalis (Linnaeus, 1758)
validirostris Gyllenhal, 1836
edoughensis Desbrochers des Loges, 1875
affinis (Paykull, 1800 nec Schrank, 1781)
- Note: This species has no valid name. *D. edoughensis* is an African species and does not occur in the Baltic region. Specimens of *Dorytomus* arranged to “*edoughensis*” from Scandinavia and Baltic States should be further observed as sensu Silfverberg, 1992 “*Enumeratio Coleopterorum Fennoscandiae, Daniae et Baltiae*”.
- ictor* (Herbst, 1795)
longimanus (Forster, 1771)
vorax (Fabricius, 1792)
- majalis* (Paykull, 1792)
melanophthalmus (Paykull, 1792)
punctator (Herbst, 1795)
- minutus* (Gyllenhal, 1836)
nebulosus (Gyllenhal, 1836)
nordenskioldi Faust, 1882
ocallescens (Gyllenhal, 1836)
reussi Formanek, 1908
rufatus (Bedel, 1888)
pectoralis Gyllenhal, 1836
- salicinus* (Gyllenhal, 1827)
salicis Walton, 1851
schoenherri Faust, 1882
suratus (Gyllenhal, 1836)
flavipes (Panzer, 1797 nec DeGeer, 1775 nec Fabricius, 1775)
- taeniatus* (Fabricius, 1781)
costirostris R.F.Sahlberg, 1835
- tortrix* (Linnaeus, 1761)
tremulae (Fabricius, 1787)
variegatus Gyllenhal, 1836
- Pseudostyphlus** Tournier, 1874
pillumus (Gyllenhal, 1836)
- Cioninae** Schönherr, 1825
Cionini Schönherr, 1825
- Cionus** Clairville, Schellenberg, 1798
hortulanus (Geoffroy, 1785)
longicollis Brisout de Barneville, 1863
montanus Wingelmüller, 1914
nigritarsis Reitter, 1904
olivieri Rosenschoeld, 1838
thapsus Olivier, 1807
scrophulariae (Linnaeus, 1758)
tuberculosus (Scopoli, 1763)
scrophulariae Latreille, 1807 nec (L., 1758)
verbasci Fabricius, 1807
- Stereonychus** Suffrian, 1854
fraxini (DeGeer, 1775)
- Cleopus** Dejean, 1821
pulchellus (Herbst, 1795)
solani (Fabricius, 1792)
- Tychiinae** Thomson, 1859
Tychiini Thomson, 1859
Tychus Germar, 1817
aureolus Kiesenwetter, 1851
breviusculus Desbrochers des Loges, 1873
haematopus auct. nec Gyllenhal, 1836
micaceus Rey, 1895
- junceus* (Reich, 1797)
lineatulus Stephens, 1831
medicaginis Brisout de Barneville, 1862
meliloti Stephens, 1831
picirostris (Fabricius, 1787)
polylineatus (Germar, 1824)
quinquepunctatus (Linnaeus, 1758)
schneideri (Herbst, 1795)
squamulatus Gyllenhal, 1836
flavicollis auct. nec Stephens, 1831
- stephensi* Gyllenhal, 1836
tomentosus (Herbst, 1795 nec Olivier, 1790)
- Sibinia** Germar, 1817
nigritarsis (Desbrochers der Loges, 1870)
pellucens (Scopoli, 1772)
primita (Herbst, 1795)
pyrrhocodactyla (Marsham, 1802)
potentillae Germar, 1824
- unicolor* (Fahraeus, 1843)
viscariae (Linnaeus, 1761)
- Ellescini** Thomson, 1859
Ellescus Dejean, 1821
bipunctatus (Linnaeus, 1758)
infirmus Herbst, 1795
scanicus (Paykull, 1792)
- Acalyptinae** Thomson, 1859
Acalyptus Schönherr, 1833
carpini (Fabricius, 1792)

Curculioninae Latreille, 1802	
Anthonomini Thomson, 1859	
<i>Anthonomus</i> Germar, 1817	
<i>brunnipennis</i> Curtis, 1840	
<i>conspersus</i> Desbrochers des Loges, 1868	
<i>humeralis</i> (Panzer, 1795)	
<i>pedicularius</i> (Linnaeus, 1758)	
<i>phyllocola</i> (Herbst, 1795)	
<i>pinivorax</i> Silfverberg, 1977	
<i>pubescens</i> (Paykull, 1792 nec Fabricius, 1775)	
<i>piri</i> Kollar, 1837	
<i>pomorum</i> (Linnaeus, 1758)	
<i>rubi</i> (Herbst, 1795)	
<i>sorbi</i> Germar, 1821	
<i>ulmi</i> (DeGeer, 1775)	
<i>inversus</i> Bedel, 1884	
<i>Furcipes</i> Desbrochers des Loges, 1868	
<i>rectirostris</i> (Linnaeus, 1758)	
Brachonyx Schönherr, 1825	
<i>pineti</i> (Paykull, 1792)	
Bradybatus Germar, 1824	
<i>kellneri</i> Bach, 1854	
Curculionini Latreille, 1802	
Curculio Linnaeus, 1758	
<i>betulae</i> (Stephens, 1831)	
<i>cerasorum</i> Paykull, 1792 nec Fabricius, 1775	
<i>crux</i> Fabricius, 1776	
<i>glandium</i> Marsham, 1802	
<i>nucum</i> Linnaeus, 1758	
<i>salicivorus</i> Paykull, 1792	
<i>pyrrhoceras</i> Marsham, 1802	
<i>venosus</i> (Gravenhorst, 1807)	
<i>villosus</i> Fabricius, 1781	
Rhynchaeninae Thomson, 1859	
Rhynchaenus Clairville, Schellenberg, 1798	
<i>alni</i> (Linnaeus, 1758)	
<i>saltator</i> (Geoffroy, 1785)	
<i>calceatus</i> (Germar, 1821)	
<i>decoratus</i> (Germar, 1821)	
<i>fagi</i> (Linnaeus, 1758)	
<i>foliorum</i> (O.F.Müller, 1764)	
<i>saliceti</i> (Paykull, 1792)	
<i>iota</i> (Fabricius, 1787)	
<i>lonicerae</i> (Herbst, 1795)	
<i>pilosus</i> (Fabricius, 1781)	
<i>populicola</i> Silfverberg, 1977	
<i>populi</i> (Fabricius, 1792 nec Linnaeus, 1758)	
<i>pratensis</i> (Germar, 1821)	
<i>pseudostigma</i> Tempere, 1982	
<i>quercus</i> (Linnaeus, 1758)	
<i>rufitarsis</i> (Germar, 1821)	
	<i>rufus</i> (Schrank, 1781)
	<i>rusci</i> (Herbst, 1795)
	<i>salicis</i> (Linnaeus, 1758)
	<i>signifer</i> (Creutzer, 1799)
	<i>avellanae</i> (Donovan, 1797 nec Paykull, 1792)
	<i>stigma</i> (Germar, 1821)
	<i>testaceus</i> (O.F.Müller, 1776)
	Rhamphus Clairville, Schellenberg, 1798
	<i>oxyacanthalae</i> (Marsham, 1802)
	<i>pulicarius</i> (Herbst, 1795)
	Gymnetrinae Thomson, 1859
	Gymnetrini Thomson, 1859
	Gymnetron Schönherr, 1825
	<i>antirrhini</i> (Paykull, 1800)
	<i>beccabungae</i> (Linnaeus, 1761)
	<i>collinum</i> (Gyllenhal, 1813)
	<i>labile</i> (Herbst, 1795)
	<i>linariae</i> (Panzer, 1793)
	<i>melanarium</i> (Germar, 1821)
	<i>pascuorum</i> (Gyllenhal, 1813)
	<i>rostellum</i> (Herbst, 1795)
	<i>stimulosum</i> (Germar, 1821)
	<i>veronicae</i> (Germar, 1821)
	<i>vilosolum</i> Gyllenhal, 1838
	Mecinini Gistel, 1856
	Mecinus Germar, 1817
	<i>collaris</i> Germar, 1821
	<i>heydeni</i> Wencker, 1866
	<i>ianthinus</i> Germar, 1821
	<i>pyraster</i> (Herbst, 1795)
	Miarini Pierce, 1919
	Miarus Schönherr, 1825
	<i>campanulae</i> (Linnaeus, 1767)
	<i>frigidus</i> Franz, 1947
	<i>distinctus</i> (Bohemian, 1845)
	<i>longirostris</i> auct. nec (Gyllenhal, 1813)
	<i>graminis</i> (Gyllenhal, 1813)
	<i>micros</i> (Germar, 1821)
	Anoplinae Bedel, 1884
	Anoplus Germar, 1820
	<i>plantaris</i> (Naezen, 1794)
	<i>roboris</i> Suffrian, 1840
	Bagoinae Thomson, 1859
	Bagous Germar, 1817
	<i>alismatis</i> (Marsham, 1802)
	<i>tibialis</i> (Bohemian, 1845)
	<i>binodulus</i> (Herbst, 1795)
	<i>atrirostris</i> (Olivier, 1808)
	<i>glabrirostris</i> (Herbst, 1795)
	<i>lutosus</i> (Gyllenhal, 1813)
	<i>caudatus</i> Thomson, 1868

- | | |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <i>cruentatus</i> J.R.Sahlberg 190X
<i>lутulentus</i> (Gyllenhal, 1813)
<i>nigritarsis</i> Thomson, 1865
<i>lutulosus</i> (Gyllenhal, 1827)
<i>petro</i> (Herbst, 1795)
<i>tempestivus</i> (Herbst, 1795)
<i>angustulus</i> Thomson, 1868
<i>sjoeborgi</i> Bruce, 1968
<i>thomsoni</i> Bruce, 1968
<i>tubulus</i> Caldara, O'Brien, 1994
<i>angustus</i> Silfverberg, 1977
<i>cylindrus</i> (Paykull, 1800 nec Fabricius, 1781)
Tanysphyrinae Gistel, 1856
Tanysphyrus Germar, 1817
<i>lemnæ</i> (Paykull, 1792)
<i>callæ</i> Voss, 1943
Cossoninae Schönherr, 1825
Dryophthorini Schönherr, 1825
Dryophthorus Schönherr, 1825
<i>corticalis</i> (Paykull, 1792)
Cotasterini Faust, 1886
Pselactus Broun, 1886
<i>spadix</i> (Herbst, 1795)
Cossonini Schönherr, 1825
Cossonus Clairville, Schellenberg, 1798
<i>cylindricus</i> R.F.Sahlberg, 1835
<i>parallelepipedus</i> (Herbst, 1795)
Rhyncolini Gistel, 1856
Rhyncolus Germar, 1817
<i>ater</i> (Linnaeus, 1758)
<i>chloropus</i> auct. nec (Linnaeus, 1758)
<i>elongatus</i> (Gyllenhal, 1827)
<i>reflexus</i> Boheman, 1838
<i>sculpturatus</i> Waltl, 1839
<i>nitidipennis</i> Thomson, 1868
Phloeophagus Schönherr, 1838
<i>lignarius</i> (Marsham, 802)
<i>turbatus</i> Schönherr, 1845
Brachytemnus Wollaston, 1873
<i>porcatus</i> (Germar, 1824)
Stereocorynes Wollaston, 1873
<i>truncorum</i> (Germar, 1824)
Magdalinae LeConte, 1876
Magdalis Germar, 1817
<i>armigera</i> (Geoffroy, 1785)
<i>aterrima</i> (Fabricius, 1775 nec Linnaeus 1758)
<i>barbicornis</i> (Latreille, 1804)
<i>mixta</i> (Desbrochers des Loges, 1870)
<i>carbonaria</i> (Linnaeus, 1758)
<i>atramentaria</i> Germar, 1824
<i>caucasica</i> (Tournier, 1872)
<i>duplicata</i> Germar, 1819 | <i>weisei</i> Schreiner, 1882
<i>frontalis</i> (Gyllenhal, 1827)
<i>linearis</i> (Gyllenhal, 1827)
<i>memnonia</i> (Gyllenhal, 1837)
<i>nitida</i> (Gyllenhal, 1827)
<i>ruficornis</i> (Linnaeus, 1758)
<i>pruni</i> (Linnaeus, 1767)
<i>violacea</i> (Linnaeus, 1758)
Molytinae Schönherr, 1823
Lepyrini Kirby, 1837
Lepyrus Germar, 1817
<i>capucinus</i> (Schaller, 1783)
<i>bimaculatus</i> (Olivier 1808)
<i>binotatus</i> (Fabricius, 1793)
<i>palustris</i> (Scopoli, 1763)
<i>bipunctatus</i> (Geoffroy, 1785)
<i>colon</i> (Linnaeus, 1771)
Hylobiini Kirby, 1837
Hylobius Germar, 1817
<i>abietis</i> (Linnaeus, 1758)
<i>excavatus</i> (Laicharting, 1781)
<i>piceus</i> (DeGeer, 1775 nec Pallas, 1771 nec Scopoli, 1763)
<i>pinastri</i> (Gyllenhal, 1813)
Molytini Schönherr, 1823
Liparus Olivier, 1807
<i>coronatus</i> (Goeze, 1777)
Pissodini Gistel, 1856
Pissodes Germar, 1817
<i>castaneus</i> (DeGeer, 1775)
<i>notatus</i> (Fabricius, 1787 nec Bonsdorff, 1785)
<i>harcyniae</i> (Herbst, 1795)
<i>piceae</i> (Illiger, 1807)
<i>pini</i> (Panzer, 17XX nec Linnaeus, 1758)
<i>pini</i> (Linnaeus, 1758)
<i>piniphilus</i> (Herbst, 1797)
<i>validirostris</i> (R.F.Sahlberg, 1834)
<i>piniphilus</i> sensu Hansen, 1918
Trachodini LeConte, 1866
Trachodes Germar, 1824
<i>hispidus</i> (Linnaeus, 1758)
Cryptorhynchinae Schönherr, 1825
Cryptorhynchus Illiger, 1807
<i>lapathi</i> (Linnaeus, 1758)
Acalles Schönherr, 1825
<i>camelus</i> (Fabricius, 1792)
<i>echinatus</i> (Germar, 1824)
<i>— lemur</i> (Germar, 1824)
Baridinae Schönherr, 1839
Baris Germar, 1817
<i>artemisiae</i> (Herbst, 1795)
<i>carbonaria</i> (Bohemian, 1836) |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

dalmatina Brisout de Barneville, 1862
laticollis (Marsham, 1802)
lepidii Germar, 1824
timida (Rossi, 1792)
Limnobaris Bedel, 1885
dolorosa (Goeze, 1777)
 pillistriata (Stephens, 1831)
pusio (Boheman, 1843)
t-album atriplicis (Fabricius, 1776)
 reitteri Munster, 1928
t-album t-album (Linnaeus, 1758)
Ceutorhynchinae Gistel, 1856
Mononychini LeConte, 1876
Mononychus Germar, 1824
punctumalbum (Herbst, 1784)
Phytobiini Gistel, 1856
Eubrychius Thomson, 1859
velutus (Beck, 1817)
Litodactylus Redtenbacher, 1849
leucogaster (Marsham, 1802)
Pelenomus Thomson, 1859
canaliculatus (Fahraeus, 1843)
comari (Herbst, 1795)
quadricorniger (Colonelli, 1986)
 quadricornis (Gyllenhal, 1813 nec Paykull, 1792)
velaris (Gyllenhal, 1827)
waltoni (Boheman, 1843)
Neophytobius Wagner, 1936
quadrinodosus (Gyllenhal, 1813)
Rhinoncus Schönherr, 1826
albicinctus Gyllenhal, 1837
bruchoides (Herbst, 1784)
castor (Fabricius, 1792)
inconspicetus (Herbst, 1795)
 gramineus (Fabricius, 1792 nec Gmelin, 1790)
pericarpius (Linnaeus, 1758)
perpendicularis (Reich, 1797)
Cnemogonini Colonnelli, 1979
Auleutes Dietz, 1896
epilobii (Paykull, 1800)
Scleropterini Schultze, 1902
Rutidosoma Stephens, 1831
fallax (Otto, 1897)
globulus (Herbst, 1795)
Scleropterus Schönherr, 1825
serratus (Germar, 1824)
Ceutorhynchini Gistel, 1856
Amalus Schönherr, 1825
scortillum (Herbst, 1795)
 haemorrhous (Herbst, 1795 nec Gmelin, 1790)

Poophagus Schönherr, 1837
sisymbrii (Fabricius, 1775)
Tapinotus Schönherr, 1826
sellatus (Fabricius, 1794)
Coeliodes Schönherr, 1837
dryados (Gmelin, 1790)
 querqus (Fabricius, 1787 nec Linnaeus, 1758)
rubicundus (Herbst, 1795)
trifasciatus Bach, 1854
Thamiocolus Thomson, 1859
sahlbergi (R.F.Sahlberg, 1845)
 lamii R.F.Sahlberg, 1835
viduatus (Gyllenhal, 1813)
Micrelus Thomson, 1859
ericae (Gyllenhal, 1813)
Zacladus Reitter, 1913
geranii (Paykull, 1800)
 affinis (Paykull, 1792 nec Schrank, 1781)
Ceutorhynchus Germar, 1824
abbreviatulus (Fabricius, 1792)
Note: North border of main distribution area.
arcuatus (Herbst, 1795)
asperifoliarum (Gyllenhal, 1813)
assimilis (Paykull, 1792)
barbareae Suffrian, 1847
campestris Gyllenhal, 1837
chalybaeus Germar, 1824
 moguntiagus Schultze, 1895
 timidus Weise, 1883
cochleariae (Gyllenhal, 1813)
contractus (Marsham, 1802)
 micros Schultze, 1897
cruciger (Herbst, 1784)
 crucifer (Olivier, 1807)
distinctus Brisout de Barneville, 1870
 marginatus (Paykull, 1792 nec Olivier, 1790)
 simillimus Edwards, 1911
dubius Brisout de Barneville, 1883
 berteroae Penecke, 1928
erysimi (Fabricius, 1787)
euphorbiae Brisout de Barneville, 1866
fennicus Faust, 1894
 gammeli Hajoss, 1929
figuratus Gyllenhal, 1837
 chrysanthemi auct. nec Germar, 1824
floralis (Paykull, 1792)
gallorhenanus Hoffmann, 1954
geographicus (Goeze, 1777)
Note: This species is only known from single locality in Randu pļavas coastal meadows in Kuivīži-Aināži area (Rīga Gulf shore, N Latvia). Probably should be more widely

distributed in whole of Latvia. North border of main distribution area.

hampei Brisout de Barneville, 1869
hirtulus Germar, 1824
ignitus Germar, 1824
inaffectatus Gyllenhal, 1837
javeti Brisout de Barneville, 1869
litura (Fabricius, 1775)
obstrictus (Marsham, 1802)
 assimilis auct. nec (Paykull, 1792)
pallidactylus (Marsham, 1802)
 quadridens (Panzer, 1795 nec Fabricius, 1775)
pectoralis Weise, 1883
pervicax Weise, 1883
pleurostigma (Marsham, 1802)
 sulcicollis Gyllenhal, 1836
plumbeus Brisout de Barneville, 1869
pollinarius (Forster, 1771)
posthumus Germar, 1824
pulvinatus Gyllenhal, 1837
 convexus Voss, 1959
 satrapa Voss, 1967
punctiger (R.F.Sahlberg, 1835)
rapae Gyllenhal, 1837
rhenanus Schultze, 1895
rugulosus (Herbst, 1795)
 rubiginosus Schultze, 1896
sulcicollis (Paykull, 1800)
suturalis (Fabricius, 1775)
sympyti Bedel, 1885
syrites Germar, 1824
triangulum Boheman, 1845
Sirocalodes Voss, 1958
depressicollis (Gyllenhal, 1813)
 nigrinus (Marsham, 1802 nec Herbst, 1795)
quercicola (Paykull, 1792)
Calosirus Thomson, 1859
terminatus (Herbst, 1795)
Trichosirocalus Colonnelli, 1979
barnevillei (Grenier, 1866)
troglodytes (Fabricius, 1787)
Nedyus Schönherr, 1825
quadrimaculatus (Linnaeus, 1758)
Coeliastes Weise, 1883
lamii (Fabricius, 1792)
Zygopinae Lacordaire, 1866
Coryssomerini Thomson, 1859
Coryssomerus Schönherr, 1825
capucinus (Beck, 1817)
Euryommatus Roger, 1857
mariae Roger, 1857

Ithyporinae Lacordaire, 1866
Orobitini Thomson, 1859
Orobitis Germar, 1817
cyanus (Linnaeus, 1758)
Rhynchophorinae Schönherr, 1833
Sitophilus Schönherr, 1838
granarius (Linnaeus, 1758)
muricatus (Fabricius, 1801)
oryzae (Linnaeus, 1763)
Scolytidae Latreille, 1807
Hylesininae Erichson, 1836
Hylastini LeConte, 1876
Hylurgops LeConte, 1876
glabratus (Zetterstedt, 1828)
 crenatus (Panzer, 1813)
 decumanus (Erichson, 1836)
 paykulli (Duftschmid, 1825)
 tenebrosus (R.F.Sahlberg, 1836)
palliatus (Gyllenhal, 1813)
 abitiperda (Bechstein, 1818)
 fuscus (Duftschmid, 1825)
 helpferi (Villa, 1835)
 marginatus (Duftschmid, 1825)
 piceus (Marsham, 1802)
 rufescens (Stephens, 1830)
 rufus (Marsham, 1802)
Hylastes Erichson, 1836
angustatus (Herbst, 1793)
 graphus (Duftschmid, 1825)
 scandinavicus Lekander, 1965
ater (Paykull, 1800)
 angusticollis Eggers, 1929
 chloropus (Duftschmid, 1825)
 pinicola (Bedel, 1888)
attenuatus Erichson, 1836
brunneus Erichson, 1836
 aterrimus Eggers, 1933
 rotundicollis Reitter, 1894
 subalpinus Eggers, 1940
cunicularius Erichson, 1836
 starki Eggers, 1933
opacus Erichson, 1836
 simplex Rey, 1892
Hylesinini Erichson, 1836
Hylesinus Fabricius, 1801
crenatus (Fabricius, 1787)
fraxini (Panzer, 1799)
 haemorrhoidalis Marsham, 1802
 henscheli Knotek, 1894
 melanocephalus Fabricius, 1801
 picipennis Stephens, 1836
 rufescens Marsham, 1802

Note: Probably correct name for this species is *H. varius* (Fabricius).

toranio (Danhoine, 1788)
bicolor Brullé, 1832
essau Gredler, 1866
oleiperda (Fabricius, 1792)
scaber Marsham, 1802
suturalis Redtenbacher, 1842

Hylastinus Bedel, 1888

obscurus (Marsham, 1802)
crenatus (Olivier, 1795)
crenulatus (Duftschmid, 1825)
fuscencens (Stephens, 1830)
piceus (Stephens, 1830)
pilosus Eggers, 1944
trifolii (P.W.J.Müller, 1803)

Pteleobius Bedel, 1888

vittatus (Fabricius, 1787)
furcatus (Marsham, 1802)
sericeus (Stephens, 1832)

Note: North border of main distribution area.

Tomicini Thomson, 1859

Xylechinus Chapuis, 1869

pilosus (Ratzeburg, 1837)

Hylurgus Latreille, 1807

ligniperda (Fabricius, 1787)
elongatus (Herbst, 1793)
flavipes (Panzer, 1795)

Tomicus Latreille, 1802-03

minor (Hartig, 1834)
corsicus (Eggers, 1911)

piniperda (Linnaeus, 1758)
analogus (LeConte, 1868)
major (Eggers, 1943)
testaceus (Fabricius, 1787)

Dendroctonus Erichson, 1836

micans (Kugelann, 1794)
abietinus (Fabricius, 1792)
ligniperda (Paykull, 1800)

Phloeotribini Chapuis, 1869

Phloeotribus Latreille, 1796

rhododactylus (Marsham, 1802)
retamae (Peris, 1864)
spartii (Nördlinger, 1847)
tarsalis (Forster, 1849)

spinulosus (Rey, 1883)

chapuisi (Blandford, 1891)
perrisi (Guillebeau, 1894)
crenatus (Guillebeau, 1893)

Polygraphini Chapuis, 1869

Polygraphus Erichson, 1836

— *poligraphus* (Linnaeus, 1758)
griseus Eggers, 1923
pubescens (Fabricius, 1792)

Note: All previous references are invalid and relate to *P. punctifrons* Thomson.

punctifrons Thomson, 1886

subopacus Thomson, 1871
nanus Schedl, 1955

Scolytinae Latreille, 1807

Scolytini Latreille, 1807

Scolytus Geoffroy, 1762

carpini (Ratzeburg, 1837)
balcanicus (Eggers, 1911)
peregrinus (Eggers, 1908)

intricatus (Ratzeburg, 1837)

lenkoranus Eggers, 1942
penicillatus Reitter, 1913
picicolor (Stephens, 1830)
simmeli Eggers, 1923
tiburtinus Claus, 1958

laevis Chapuis, 1869

loevendali (Eggers, 1912)
pomonacearum Butovitsch, 1929

mali (Bechstein, 1805)

bicallosus Eggers, 1933
castaneus (Ratzeburg, 1837)
nitidulus Chapuis, 1869
pruni (Ratzeburg, 1837)
pyri (Ratzeburg, 1837)

multistriatus (Marsham, 1802)

abhorrens (Wichmann, 1913)
flavicornis (Chevrolat, 1829)
nodifer Reitter, 1913
therondi Hoffmann, 1939
ulmi Redtenbacher, 1849

ratzeburgi Janson, 1856

amurensis (Eggers, 1908)
destructor (Ratzeburg, 1837)
lineatus Kurenzov, 1941
sahlbergi (Eggers, 1912)
sibiricus (Eggers, 1922)

rugulosus (P.W.J.Müller, 1818)

assimilis Boheman, 1858
haemorrhous Schmiedeberger, 1837
mangliensis Lezhava, 1940
mediterraneus (Eggers, 1922)
taxicola Lezhava, 1943

scolytus (Fabricius, 1775)

destructor Olivier, 1795
fuchsii Reitter, 1913
geoffroi (Goeze, 1777)
niger Geoffroy, 1785
punctatus O.F.Müller, 1776

Ipini Bedel, 1888

Pityogenes Bedel, 1888

bidentatus (Herbst, 1784)

ater (Fabricius, 1792)
bidens (Fabricius, 1792)
bispinus (Guyon, 1855)
obtusus Eggers, 1932

chalcographus (Linnaeus, 1761)

bicolor (Chevrolat, 1838)
sexdentatus (Olivier, 1795)

spinosus (DeGeer, 1775)
xylographus (R.F.Sahlberg, 1836)
quadridens (Hartig, 1834)
trepanatus (Nordlinger, 1848)
austriacus (Wachtl, 1887)
elongatus (Loevendal, 1889)
Pityokteines Fuchs, 1911
curvidens (Germar, 1824)
calligraphus (Duftschmid, 1825)
orthographus (Duftschmid, 1825)
psilonotus (Germar, 1824)
 Note: North border of main distribution area.
Orthotomicus Ferrari, 1867
laricis (Fabricius, 1792)
chalcographus (Olivier, 1795)
denticulatus (Sturm, 1826)
longicollis (Gyllenhal, 1827)
oblitus (Peris, 1862)
proximus (Eichhoff, 1867)
feiferi (Keler, 1925)
omissus (Eichhof, 1871)
suturalis (Gyllenhal, 1827)
nigritus (Gyllenhal, 1827)
Ips DeGeer, 1775
acuminatus (Gyllenhal, 1827)
geminatus (Zetterstedt, 1828)
iconographus (Kugelann, 1837)
porographus (Eschscholtz, 1837)
quadridentatus (Sturm, 1826)
amitinus (Eichhoff, 1871)
duplicatus (R.F.Sahlberg, 1836)
infuscatus (Eichhoff, 1877)
judeichi (Kirsch, 1870)
rectangulus (Ferrari, 1867)
sexdentatus (Börner, 1767)
pinastri (Bechstein, 1818)
stenographus (Duftschmid, 1825)
typographus (Linnaeus, 1758)
octodentatus (Paykull, 1800)
Dryocoetini Lindemann, 1876
Lymantor Loevendal, 1889
aceris (Lindemann, 1875)
coryli (Perris, 1855)
sepicola Loevendal, 1889
Dryocoetes Eichhoff, 1864
alni (Georg, 1856)
marshami (Rey, 1868)
similis Eggers, 1911
autographus (Ratzeburg, 1837)
americanus (Hopkins, 1915)
artepunctatus Eggers, 1914
pseudotsugae Swaine, 1915
semicastaneus (Mannerheim, 1852)
septentrionis (Mannerheim, 1843)
suecicus Eggers, 1932
victoris (Mulsant, Rey, 1853)

hectographus Reitter, 1913
villosus (Fabricius, 1792)
eichhoffi Ferrari, 1867
histerinus (Dufour, 1843)
pilosus (Dejean, 1837)
Crypturgini LeConte, 1876
Crypturgus Erichson, 1836
cinereus (Herbst, 1793)
apfelbecki Eggers, 1940
atticus Eggers, 1911
corsicus Eggers, 1911
minutus (Duftschmid, 1828)
subcribrosus Eggers, 1933
tenerrimus (R.F.Sahlberg, 1836)
hispidulus Thomson, 1870
maulei Roubal, 1910
pusillus (Gyllenhal, 1813)
aphodioides (Villa, Villa, 1833)
atomus LeConte, 1868
danicus Eggers, 1922
minimus (Stebbing, 1903)
Xyloterini Lindemann, 1876
Trypodendron Stephens, 1830
domesticum (Linnaeus, 1758)
limbatus (Herbst, 1784)
laeve Eggers, 1939
—lineatum (Olivier, 1795)
bivittata (Kirby, 1837)
borealis Swaine, 1917
cavifrons (Mannerheim, 1843)
meridionale Eggers, 1940
 Note: All previous references are invalid and relate to *T. leave* Eggers.
signatum (Fabricius, 1787)
limbatus (Paykull, 1800)
quinquelineatus (Adams, 1827)
waringi (Curtis, 1840)
Xyleborini LeConte, 1876
Xyleborus Eichhoff, 1864
cryptographus (Ratzeburg, 1837)
villosus (Ratzeburg, 1837)
dispar (Fabricius, 1792)
aequalis (Reitter, 1913)
brevis (Panzer, 1793)
pyri (Peck, 1817)
ratzeburgi (Kolenati, 1846)
swainei (Drake, 1921)
thoracicus (Panzer, 1793)
tachygraphus (R.F.Sahlberg, 1834)
monographus (Fabricius, 1792)
tuberculatus (Herbst, 1793)
Cryphalini Lindemann, 1876
Trypophloeus Fairmaire, 1868
asperatus (Gyllenhal, 1813)
berezinae Stark, 1952

Note: This species is referred in the “World Catalogue of Scolytidae” by Wood and Bright (1992) as *binodulus* (Ratzeburg, 1837). The question about the priority in name is unclear and hence the most commonly name for the species used in Europe is conserved here.

bispinus Eggers, 1927

palmi Hansen, 1955

Ernoporus Thomson, 1859

tiliae (Panzer, 1793)

lederi (Reitter, 1889)

ratzeburgi (Ferrari, 1867)

Cryphalus Erichson, 1836

abietis (Ratzeburg, 1837)

Note: This species is referred in the “World Catalogue of Scolytidae” by Wood and Bright (1992) as *asperatus* (Gyllenhal, 1813). The question about the priority in name is unclear and hence the most commonly name for the species used in Europe is conserved here.

piceae (Ratzeburg, 1837)

orientalis Eggers, 1911

subdepressus Eggers, 1940

Note: North-western border of main distribution area.

Corthylini LeConte, 1876

Pityophthorus Eichhoff, 1864

lichtensteinii (Ratzeburg, 1837)

rossicus Eggers, 1915

micrographus (Linnaeus, 1758)

fennicus Eggers, 1914

morosovi Spessivtseff, 1926

Platypodidae Latreille, 1802

Platypodinae Strohmeyer, 1911

Platypodini Schedl, 1938

Platypus Herbst, 1793

cylindrus (Fabricius, 1792)

cylindriformis (Reitter, 1894)

flavicornis (Kugelann, 1792)

platypus (Duftschmid, 1825)

Note: This species is known from single locality situated in extreme SE of Latvia at the border with Lithuania and Belarus'. North border of main distribution area.

INDEX TO FAMILY NAMES

Family	Page	Family	Page
Sphaeriusidae	4; 6	Trogossitidae	5; 67
Trachypachidae	4; 6	Cleridae	5; 68
Carabidae	4; 6	Dasytidae	5; 68
Gyrinidae	4; 28	Malachiidae	5; 69
Haliplidae	4; 29	Kateretidae	5; 69
Noteridae	4; 29	Nitidulidae	5; 69
Dytiscidae	4; 29	Monotomidae	5; 71
Helophoridae	4; 35	Silvanidae	5; 71
Hydrophilidae	4; 35	Cucujidae	5; 71
Hydrochidae	4; 38	Bothrideridae	5; 72
Spercheidae	4; 38	Cryptophagidae	5; 72
Georissidae	4; 38	Languriidae	5; 73
Sphaeritidae	4; 38	Erotylidae	5; 73
Histeridae	4; 38	Cerylonidae	5; 73
Hydraenidae	4; 39	Byturidae	5; 73
Ptiliidae	4; 39	Laemophloeidae	5; 73
Leiodidae	4; 40	Phalacridae	5; 73
Colonidae	4; 41	Sphindidae	5; 73
Cholevidae	4; 41	Corylophidae	5; 73
Platypsyllidae	4; 41	Endomychidae	5; 74
Scydmaenidae	4; 42	Coccinellidae	5; 74
Agyrtidae	4; 42	Latridiidae	5; 75
Silphidae	4; 42	Zopheridae	5; 76
Staphylinidae	4; 42	Mycetophagidae	5; 76
Scaphidiidae	4; 51	Ciidae	5; 77
Pselaphidae	4; 52	Tetratomidae	5; 77
Lucanidae	4; 52	Melandryidae	5; 77
Trogidae	4; 53	Mordellidae	5; 78
Geotrupidae	4; 53	Rhipiphoridae	5; 79
Scarabaeidae	4; 53	Stenotrachelidae	5; 79
Clambidae	4; 58	Meloidae	5; 79
Eucinetidae	4; 58	Oedemeridae	5; 79
Scirtidae	4; 58	Pythidae	5; 80
Dascillidae	4; 58	Pyrochroidae	5; 80
Buprestidae	4; 58	Scriptiidae	5; 80
Byrrhidae	4; 61	Salpingidae	5; 80
Elmidae	4; 61	Mycteridae	5; 81
Dryopidae	4; 62	Boridae	5; 81
Limnichidae	4; 62	Anthicidae	5; 81
Heteroceridae	4; 62	Aderidae	5; 82
Psephenidae	4; 62	Tenebrionidae	5; 82
Elateridae	4; 62	Cerambycidae	5; 84
Eucnemidae	4; 64	Bruchidae	5; 88
Throscidae	4; 64	Chrysomelidae	5; 89
Lycidae	4; 64	Nemonychidae	5; 98
Drilidae	4; 65	Anthribidae	5; 98
Lampyridae	4; 65	Rhynchtidae	5; 98
Cantharidae	4; 65	Attelabidae	5; 99
Dermestidae	4; 65	Apionidae	5; 100
Bostrichidae	4; 66	Curculionidae	5; 102
Lyctidae	4; 66	Scolytidae	5; 109
Anobiidae	5; 66	Platypodidae	5; 111
Ptinidae	5; 67		
Lymexylidae	5; 67		


NOTES

THE ENTOMOLOGICAL SOCIETY OF LATVIA

The Entomological Society of Latvia (LEB in official abbreviation) has united entomologists, arachnologists and other invertebrate biologists from Latvia and the World since 1951. LEB is a discussion club for scientists, applied biologists and wildlife managers. It publishes scientific journals, principally "Latvijas Entomologs", and distributes knowledge to a wider audience through mass media.

Your LEB membership gives you the opportunity to serve Science and extend your professional development and recognition by colleagues, to publish papers in our journals "Latvijas entomologs" and "Acta coleopterologica latvica", to get the latest information on our activities and to participate in our projects, seminars and expeditions. The membership fee is EUR 10.- per year (including subscription to all our issues).

APPLICATION FORM



LATVIJAS ENTOMOLOGIJAS BIEDRĪBA

Pasts: Latvijas Universitātēs Bioloģijas Fakultāte, Kronvalda bulv. 4, LV-1586, Rīga, Latvija.
Tel.: 9191557, 7034880 ; fakss: 8830891 ; e-pasts: adalia@lanet.lv ; URL: <http://invertebrata.from.lv>

Please, admit me to the Entomological Society of Latvia (LEB). Membership fee is EUR 10.- including all subscriptions.

First name (-s): _____ Name: _____

Birth date: _____

Education: _____

Title (-s): _____

Specialisation in entomology: _____

Affiliation, position: _____

Correspondence address: _____

Phone: _____ Fax: _____

E-mail: _____ URL: _____

Submission date: _____ Signature: _____

_____ admitted to the Entomological Society of Latvia by decision of Council of LEB No ____ Date ____

1st Chairman: _____

ENTOMOLOGICAL SOCIETY OF LATVIA

c/o Faculty of Biology, University of Latvia, Kronvalda Blvd. 4, LV-1586 Rīga, Latvia / Lettland
Tel.: (+371) 7034880 ; Fax: (+371) 8830891 ; E-mail: adalia@lanet.lv ; URL: <http://invertebrata.from.lv>