

Errata

A number of important errors and omissions from this Atlas have been drawn to the authors' attention:

There are twelve species of Coccinellidae that have been ascribed a conservation designation. In the atlas only eleven are mentioned (page 30). The additional species is:

Hippodamia variegata (Nationally notable B)

Additionally *Coccinella magnifica* is listed in table 4 as **nationally notable B** and should be **nationally notable A**. *Hyperaspis pseudopustulata* should be listed as **nationally notable B**.

Please refer to <http://jncc.defra.gov.uk/page-3408> for updates.

480 records kindly provided by Martin Harvey were regretfully omitted from the atlas. Martin should be listed in the atlas for submitting more than 100 records. All the data provided by him will be available through the NBN Gateway (UK Ladybird Survey). The authors would like to express their sincere apologies to Martin. The table below lists the records from Martin that would constitute new 10-km squares:

Species	New 10-km
<i>Adalia bipunctata</i>	SU57, SU70, SZ39
<i>Adalia decempunctata</i>	SU57
<i>Anatis ocellata</i>	SP41, SU57
<i>Aphidecta oblitterata</i>	SP81, SX67
<i>Calvia quattuordecimguttata</i>	SU57, SU78
<i>Chilocorus renipustulatus</i>	SU47, SU55, SU57, SU66, SU88
<i>Coccinella hieroglyphica</i>	SU47, SU56
<i>Coccinella septempunctata</i>	SO22
<i>Coccinella undecimpunctata</i>	SU57
<i>Exochomus quadripustulatus</i>	SU57, SU78, SU79
<i>Halyzia sedecimguttata</i>	SK15, SP72, SU10, SU46, SU49, SU57, SU62
<i>Harmonia quadripunctata</i>	SU57
<i>Myrrha octodecimguttata</i>	SU88
<i>Propylea quattuordecimpunctata</i>	SU36
<i>Psyllobora vigintiduopunctata</i>	SU57
<i>Rhyzobius chrysomeloides</i>	TQ08
<i>Tytthaspis sedecimpunctata</i>	SO87, SU47, SU57, SU68, SU78

***Henosepilachna argus* (page 114)** - The southern-most record mapped should be deleted.

***Nephus quadrimaculatus* (page 156)** - A record from Trevor James (10-km TL23, post 1990) was not mapped and the authors apologise to Trevor for this omission.

***Cynegetis impunctata* (page 162)** - The reference cited, Thomas *et al.* (2008), in support of the first record of *Cynegetis impunctata* is not in press and, as such, the record must stand as unconfirmed.

***Exochomus nigromaculatus* (page 163)** - The 80+ records mentioned as being from Peter Hodge are for *Exochomus quadripustulatus* and not *Exochomus nigromaculatus* as stated. The authors apologise to Peter for this error. The text should therefore read as follows:

Exochomus nigromaculatus is a black ladybird (with two orange-red markings on the pronotum, and orange-red legs) about the same size as its closest relation in Britain, the pine ladybird. *Exochomus nigromaculatus* is a species of dry heathland in continental Europe. The species had not been recorded in Britain since the 1830s until it was found near Doncaster in 1967 (Skidmore, 1985).

The authors would welcome information relating to any other errors and omissions that may come to light.