

NEW AND RARE BEETLE (COLEOPTERA) SPECIES FROM CURONIAN SPIT (LITHUANIA)

RADVILĖ NAGROCKAITĖ¹, BRIGITA TAMUTĖ^{1,2,3}, VYTAUTAS TAMUTIS^{2,3}

¹Department of Biology, Vytautas Magnus University, Vileikos 8, LT-44248 Kaunas, Lithuania.

²Department of Entomology, Kaunas T.Ivanauskas zoological museum, Laisvės al. 106, LT-44253 Kaunas, Lithuania.

³Department of Biology and Plant Protection, Aleksandras Stulginskis University, Studentų 11, LT-53361 Akademija, Lithuania.

E-mail: b.tamute@gmail.com;

Introduction

The Curonian Spit is surrounded by the Baltic Sea and the Curonian Lagoon. It is unique of which is comprised of relief created by the sea and wind, the highest spit in the whole Northern Europe, Lagoon marl prints, remains of former Lagoon's forest and soil brought by the wind and constant interesting eolic processes. Very specific climate conditions determine unique vegetation cover in the Curonian Spit that is critical for the diversity of insects and their population size.

The fauna of Coleoptera is investigated already more than 130 years in the Curonian Spit. However, data on many species is fragmentary. The first data on Coleoptera fauna in the Curonian Spit present F. L. Lentz (1897). After many years new data on some Coleoptera species distribution were published (Pileckis, 1963, 1968; Sharova, Grūntal, 1973; Bercio, Folwaczny, 1979). During the last decade there were many reports on the Coleoptera fauna from the Curonian Spit (Šablevičius, Ferenca, 1995; Barševskis, 2001; Šablevičius, 2003; Ferenca, 2004; Tamutis, 2005; Žiogas, Zolubas, 2005; Ferenca et al., 2006; Ivinskis et al., 2009).

The field practices of students are important both to study process and research experience. In many cases their collected material is very important for faunistic and ecological investigations of insects in Lithuania. In this report we present the records on new and rare beetle species found in the Curonian spit during the field practices time of students of Vytautas Magnus University in the last decade of May, 2011.

Material and Methods

The material was collected in the environs of Juodkrantė during transects expeditions and using sweep net. The stereomicroscope SMZ - 168 and various guides (Freude et al. 1967, 1979; Hůrka, 1996) have been used for species identification. The beetles were collected by Radvilė Nagrockaitė (R.N.) and Eglė Žilinskytė (E.Ž.).

In this report we followed the classification of the order Coleoptera accepted by Bouchard et al. (2011) and used the genera and species names accepted in the Catalogue of Lithuanian beetles (Tamutis et al. 2011). The material is deposited in the Kaunas T. Ivanauskas Zoological Museum.

List of localities

Neringa t.	Juodkrantė env. (1)	55°32'57,3"N, 21°07'11,2"E
	Juodkrantė env. (2)	55°33'20,6"N, 21°06'21,3"E
	Juodkrantė env. (3)	55°32'35,9"N, 21°05'56,4"E

Results

List of species

CARABIDAE

Calosoma inquisitor (Linnaeus, 1758)

Juodkrantė env. (2), 27 05 2011, 1 spec. (E.Ž.).

CLERIDAE

Opilo mollis (Linnaeus, 1758)

Juodkrantė env. (3), 28 05 2011, 1 spec. (R.N.).

COCCINELIDAE

Harmonia axyridis (Pallas, 1773)

Juodkrantė env. (1), 26 05 2011, 1 spec. (R.N.).

Discussion

Calosoma inquisitor (Linnaeus, 1758) was mentioned twice in Curonian Spit (Ferenca, 2004, 2006). Our specimen was found on the beach of the sea. This species is included into Lithuanian Red Data book since 1990 (Red Data Book of Lithuania, 1992).

Opilo mollis (Linnaeus, 1758) has been noted for Lithuania by Löbl *et al.* (2007) in Catalogue of Palaearctic Coleoptera. However, this notification was not confirmed by local entomologists till now. *O. mollis* is considered as very rare species in some neighboring countries (Burakowski *et al.*, 1986; Telnov, 2004; Lundberg & Gustafsson, 1995), not yet found in Belarus and Kaliningrad Region.

Harmonia axyridis (Pallas, 1773) is new alien species in Lithuanian fauna. The presumed native distribution of *H. axyridis* extends from Altai Mountains in the west to the Pacific Coast in the east, and from southern Siberia in the north to southern China in the south. The first established population was documented in 1988 in North America. After this initial detection, it spread rapidly across North America (Koch, 2003). *H. axyridis* has also been released in Europe (Katsoyannos *et al.*, 1997; Iperti & Bertand, 2001). *H. axyridis* occurs in many color forms. Adults are strongly oval and convex, about 6 mm long and 5 mm wide. *H. axyridis* have a mix of individuals ranging in color from pale yellow-orange to bright red-orange, with or without black spots on the wing covers. The head, antennae and mouthparts are generally straw-yellow but are sometimes tinged with black. *H. axyridis* was found in the wood lot on the grass near Juodkrantė. It was typical red color with 19 black spots (Figure 1.). This species was recently found in Poland (Przewozny *et al.*, 2007) and Latvia (Barševskis, 2009) as well.

Acknowledgement

We warmly thank Romas Ferenca for original and nice photograph which was used for illustrations in current paper.

References

- Barševskis A. 2001. New and rare species of beetles (Insecta: Coleoptera) in the Baltic countries and Byelorussia. *Baltic Journal of Coleopterology* 1 (1–2): 3–18.
- Barševskis A. 2009. Multicoloured Asian lady beetle (*Harmonia axyridis* (Pallas, 1773) (Coleoptera: Coccinellidae) for the first time in the fauna of Latvia. *Baltic Journal of Coleopterology* 9 (2): 135–138.
- Bercio H., Folwaczny B. 1979. *Verzeichnis der Käfer Preussens* [The check-list of beetles of Prussia]. Parzeller & Co, Fulda, Germany, 369 pp.
- Bouchard P., Bousquet Y., Davies A. E., Alonso-Zarazaga M. A., Lawrence J. F., Lyal C. H. C., Newton A. F., Reid C. A. M., Schmitt M., Ślipiński S. A., Smith A. B. T. 2011. Family-group names in Coleoptera (Insecta). *ZooKeys* 88: 1–972.
- Burakowski B., Mroczkowski M., Stefańska J. 1986. *Katalog fauny Polski, Tom. 11: Chrząszcze – Coleoptera. Dermestoidea, Bostrichoidea, Cleroidea i Lymexyloidea* [Catalogue of Polish fauna, Vol. 11: Beetles – Coleoptera. Dermestoidea, Bostrichoidea, Cleroidea and Lymexyloidea]. Warzawa, Poland, 243 pp.
- Ferenca R. 2004. New and rare for Lithuania beetles (Coleoptera) species registered in 1978–2004. *New and Rare for Lithuania Insect Species* 16: 11–22.
- Ferenca R. 2006. Fauna and ecology of beetles (insecta, Coleoptera) in Kuršių Nerija national park [abstract]. *6th Symposium of Baltic coleopterologists „Communities of coleoptera in agricultural and forest ecosystems“*, 7–9 September, 2006, Kaunas, Lithuania, 35–36.
- Ferenca R., Ivinskis P., Tamutis V. 2006. New and rare for Lithuania species of beetles (Coleoptera). *New and Rare for Lithuania Insect Species* 17: 11–21.
- Freude H., Harde K. W., Lohse G. A. (eds). 1967. *Die Käfer Mitteleuropas, Band 7: Clavicornia*. Krefeld: Goecke & Evers.
- Freude H., Harde K. W., Lohse G. A. (eds). 1979. *Die Käfer Mitteleuropas, Band 6: Diversicornia*. Krefeld: Goecke & Evers.
- Hůrka, K. 1996. *Carabidae of the Czech and Slovak Republics. Illustrated key*. Zlín: Kabourek.
- Iberti G., Bertand E. 2001. Hibernation of *Harmonia axyridis* (Coleoptera: Coccinellidae) in South-Eastern France. *Acta Societas Zoologicae Bohemicae* 65: 207–210.
- Ivinskis P., Meržijevskis A., Rimšaitė J. 2009. Data on new and rare for the Lithuanian fauna species of Coleoptera. *New and Rare for Lithuania Insect Species* 21: 45–63.
- Katsoyannos P., Kontodimas D. C., Stathas G. J., Tsartsalis C. T. 1997. Establishment of *Harmonia axyridis* on citrus and some data on its phenology in Greece. *Phytoparasitica* 25: 183–191.
- Koch R. L. 2003. The multicolored Asian lady beetle, *Harmonia axyridis*: A review of its biology, uses in biological control, and non-target impacts. *Journal of Insect Science* 3 (32): 1–16.

- Lentz F. L. 1879. Catalog der Preussischen Käfer [Catalogue of Prussian beetles]. *Beiträge zur Naturkunde Preussens* 4: 1–64.
- Löbl I., Rolčík J., Kolibáč J., Gerstmeier R. 2007. Cleridae. In: Löbl I., Smetana A. (Eds) *Catalogue of Palaearctic Coleoptera, Vol. 4: Elateroidea – Cucuoidea*. Apollo Books, Stenstrup, Denmark, 367–384.
- Lundberg S., Gustavsson B. 1995. *Catalogus Coleopterorum Sueciae*. Natural History Museum, Stockholm, 302 pp.
- Pileckis S. 1963. Naujos vabalų (Coleoptera) rūšys Lietuvos TSR [New species of beetles (Coleoptera) in Lithuanian SSR]. *LŽŪA mokslo darbai* 10 (19): 53–64.
- Pileckis S. 1968. Naujos ir mažai žinomos vabalų (Coleoptera) rūšys Lietuvos TSR faunoje [New and little known species of beetles (Coleoptera) in the fauna of Lithuanian SSR]. *LŽŪA mokslo darbai* 15 (2): 29–37.
- Przewozny M., Barłożek T., Bunalski M. 2007. *Harmonia axyridis* (Pallas, 1773) (Coleoptera:Coccinellidae) new species of ladybird beetles for Polish fauna. *Polish Journal of Entomology* 76: 177–182.
- Red Data Book of Lithuania* 1992. Vilnius. Environmental Protection Department of the Republic of Lithuania, 87–119.
- Šablevičius B., Ferenca R. 1995. 14 new and 3 rare for Lithuania species of Coleoptera found in 1987–1994. In: Jonaitis V. (Ed) *New and Rare for Lithuania Insect Species. Records and Descriptions of 1994–1995*, Vilnius, 145–147.
- Šablevičius B. 2003. New and rare for Lithuania beetle (Coleoptera) species. *New and Rare for Lithuania Insect Species* 15: 11–24.
- [Sharova I. H., Grüntal S. J.] Шарова И. Х., Грюнталль С. Ю. 1973. К изучению жужелиц (Coleoptera, Carabidae) заповедника “Жувитас” и косы Куршю-Нярия [Carabidae Beetles (Coleoptera, Carabidae) of the reservation “Žuvintas” and of the Curonian Spit]. *Acta Entomologica Lituanica* 2: 63–73.
- Tamutis V. 2005. Beetles (Insecta, Coleoptera) of the Red Data Book of Lithuania. In: Skłodowski J., Huruk S., Barševskis A., Tarasiuk S. (Eds) *Protection of Coleoptera in the Baltic Sea Region*. Warsaw, Poland, 23–35.
- Tamutis V., Tamutė B., Ferenca R. 2011. A. Catalogue of Lithuanian beetles (Insecta: Coleoptera). *ZooKeys* 121:263–264.
- Telnov D. 2004. Checklist of Latvian Beetles (Insecta: Coleoptera). In: Telnov D. (Ed) *Compendium of Latvian Coleoptera* 1: 1–113.
- Žiogas A., Zolubas P. 2005. Rare and protected forest beetle species in the national parks of Lithuania. In: Skłodowski J., Huruk S., Barševskis A., Tarasiuk S. (Eds) *Protection of Coleoptera in the Baltic Sea Region*. Warsaw, Poland, 37–46.

Naujos ir retos vabalų (Coleoptera) rūšys iš Kuršių nerijos (Lietuva)

R.NAGROCKAITĖ, B. TAMUTĖ, V. TAMUTIS

Santrauka

Straipsnyje pateikiama informacija apie 1 naują Lietuvai svetimžemę boružių rūšį *Harmonia axyridis* (Pallas, 1773) ir 2 retas vabalų rūšis aptiktas Juodkrantės apylinkėse Kuršių nerijoje 2011 metais. Taip pat apžvelgiamas šių rūsių paplitimas kaimyninėse valstybėse. Pirmą kartą skelbiami tikri *Opilo mollis* (L.) radvietės duomenys.



Figure 1. The specimen of *Harmonia axyridis* found in Curonian Spit, 2011.

Received: October 14, 2011