

## Culture collection of parasitic protists at the Zoological Institute RAS (CCPP ZIN RAS)

---

Marina N. Malysheva<sup>1</sup>, Maria A. Mamkaeva<sup>2</sup>,  
Alexei Yu. Kostygov<sup>1,3</sup>, Alexander O. Frolov<sup>1</sup> and  
Sergey A. Karpov<sup>1,2</sup>

<sup>1</sup> Zoological Institute, Russian Academy of Sciences, St. Petersburg, Russia

<sup>2</sup> St. Petersburg State University, St. Petersburg, Russia

<sup>3</sup> Life Science Research Centre, Faculty of Science, University of Ostrava, Ostrava, Czech Republic

### Summary

Establishment of the bank of living cultures of trypanosomatids at the Laboratory of Protozoology, Zoological Institute of the Russian Academy of Sciences, St. Petersburg, started in 1979. The first edition of the collection catalogue (Malysheva et al., 2001) included 33 strains of trypanosomatids. The present, second edition contains descriptions of 52 strains of trypanosomatids and 28 strains of aphelids and parasitic chytridiomycetes. The strains of aphelids and parasitic chytridiomycetes were isolated and initially cultivated at the Center for Culturing Collection of Microorganisms of the Research park of St. Petersburg State University. This collection is named the Culture Collection of Parasitic Protists of the Zoological Institute RAS (CCPP ZIN RAS).

**Key words:** Aphelididae, bibliography, culture collection, GenBank, Gromochytriaceae, living cultures, Mesochytriaceae, parasitic protists, Trypanosomatidae

### Introduction

Study of biodiversity of parasitic protists is one of the “hot spots” in modern biology. Many of these organisms cause dangerous infections of humans, animals and plants, and their genetic peculiarities are the unique examples of evolutionary changes in eukaryotes. The establishment, support and accession of such culture collections are very important tasks for study of biodiversity, cell biology, and parasite-host interactions. A

depositarium of parasitic protozoans was established at the Laboratory of Protozoology of the Zoological Institute RAS in 1979. The first edition of the collection catalogue was published in 2001 and included 33 strains of trypanosomatids (Malysheva et al., 2001). The present, second edition contains descriptions of 52 strains of trypanosomatids augmented with 28 strains of aphelids and parasitic chytridiomycetes. The structure of the catalogue is generally the same as in the first edition (Malysheva et al., 2001); however, it was modified in two aspects:

1) the deposition GenBank Numbers were included, and 2) a new section “List of the lost strains” was added.

## Cultivating methods

Axenic trypanosomatid cultures have been maintained in the liquid nutrient medium BHI (Difco) with addition of hemin (25 mg/ml) and inoculated once a month. The cultures were placed in the fridge (5°C) after inoculation and subsequent incubation during three days at 25°C.

For cultivation of the *Blastocrithidia* flagellates, a two-phase medium has been used: the blood agar as a dense phase, and a BHI medium with the addition of serum (5-10%) as a liquid phase. For cultivation of the *Trypanosoma* flagellates, a slightly modified two-phase medium has been used: the blood agar as a dense phase, and a physiological solution with the addition of serum (5-10%) as a liquid phase.

Accumulating cultures of the aphelids and chytrids were maintained in culture on *Tribonema gayanum* Pasch. (strain 20 CALU) as the host cultivated in the mineral medium: K<sub>2</sub>HPO<sub>4</sub> (66.7 mg/l); MgSO<sub>4</sub> × 7H<sub>2</sub>O (33 mg/l); KNO<sub>3</sub> (100 mg/l); microelements (1 mg/l) of stock solution: ZnSO<sub>4</sub> × 7H<sub>2</sub>O (0.22 g/l), MgSO<sub>4</sub> (1.81 g/l), CuSO<sub>4</sub> × 5H<sub>2</sub>O (0.079 g/l), NaBO<sub>3</sub> × 4H<sub>2</sub>O (2.63 mg/l), (NH<sub>4</sub>)<sub>6</sub>Mo<sub>7</sub>O<sub>2</sub> × 4H<sub>2</sub>O (1 mg/l), FeSO<sub>4</sub> × 7H<sub>2</sub>O (9.3 mg/l), CaCl<sub>2</sub> (1.2 mg/l), Co(NO<sub>3</sub>)<sub>2</sub> × H<sub>2</sub>O (0.02 mg/l), EDTA (10 mg/l), pH 7.0-7.2 (Gromov and Titova, 1991) in the 12 ml glass tubes containing 3-5 ml of the medium at room temperature in the presence of light (24 h). The inoculation was performed twice a month.

## Catalogue of living cultures

Class Kinetoplastea Honigberg 1963 emend. Vickerman 1976

Subclass Metakinetoplastina Vickerman 2004

Order Trypanosomatida Kent 1880 stat. nov. Hollande 1952

Family Trypanosomatidae Doflein 1951

Genus *Angomonas* Souza and Corte-Real, 1991

*A. deanei* (Carvalho, 1973)

Synonyms: *Crithidia deanei* Carvalho, 1973; *Critidinia roitmani*, Fiorini et al., 1989; *Herpetomonas roitmani* (Fiorini et al., 1989)

**1 Strain TCC080E (=ATCC 50305)**

**Depositor:** S.A. Podlipaev

**History:** 1990, received from E.P. Camargo, São Paulo University, Brazil. Isolated in Brazil.

**Host:** *Ornidia obesa* (Fabricius, 1775) (Diptera).

**GenBank:** AF038023, AF038026, EU079130, EU095978, HM593012

**Bibliography:** Fiorini et al., 1989; Faria e Silva et al., 1991; Hollar et al., 1998; Yurchenko et al., 2009, Teixeira et al., 2011.

Genus *Crithidia* Léger, 1902

*C. brevicula* Frolov et Malysheva, 1989

Synonyms: *Crithidia allae* Frolov et Malysheva, 1989 (NBrA); *Wallaceina brevicula* Frolov et Malysheva, 1989 (NBr, NBrB); *Wallaceina inconsitans* Podlipaev, Frolov et Kolesnikov, 1990 (GK); *Wallaceina vicina* Malysheva et Frolov, 2004 (WG); *Leptomonas peterhoffi* Podlipaev, 1985 (in part) (101); *Wallaceina podlipaevi* Malysheva et Frolov, 2009 (101); *Blastocrithidia gerricola* Podlipaev, 1985 (in part) (KV1); *Blastocrithidia miridarum* Podlipaev et Frolov, 1987 (in part) (BM33); *Leptomonas* sp. (F2); *Leptomonas* sp. (F5); *Leptomonas* sp. (F6); *Leptomonas* sp. (F7); *Leptomonas* sp. (F8); *Leptomonas* sp. (CL8).

**2 Strain NBr**

**Depositor:** A.O. Frolov

**History:** 1986, isolated by A.O. Frolov in the vicinity of the Lyady village, Pskov Province, Russia.

**Host:** *Nabis (Nabis) brevis* Scholtz, 1847 (Heteroptera: Nabidae).

**GenBank:** AF153045, AF316620, AJ401357, AJ401378, EU088277- EU088279, KJ443335- KJ443337, KJ443350, KJ474892- KJ474894, KJ474931, KJ474932, Z32854,

**Bibliography:** Frolov and Malysheva, 1989a, 1989b; 1990; Podlipaev et al., 1991; Malysheva and Frolov, 1995; Podlipaev et al., 1998; Podlipaev and Frolov, 2000; Yurchenko et al., 2000; Merzlyak et al., 2001a, 2001b; Yurchenko et al., 2006; Yurchenko et al., 2009; Kostygov et al., 2014

**3 Strain BrB**

**Depositor:** A.O. Frolov

**History:** Clonal culture. 1986, isolated from the culture of *C. brevicula* (NBr) in the Laboratory of Protozoology ZIN RAS, St. Petersburg, Russia.

**Host:** *Nabis (Nabis) brevis* Scholtz, 1847 (Heteroptera: Nabidae).

**Bibliography:** Frolov and Malysheva, 1989a, 1989b; Kolesnikov et al., 1990; Skarlato et al., 1990; Podlipaev et al., 1991; Malysheva and Frolov, 1995.

**4 Strain WG**

**Depositor:** M.N. Malysheva

**History:** 2000, isolated by M.N. Malysheva in Komarovo, Leningrad Province, Russia.

**Host:** *Limnoporus rufoscutellatus* (Latreille,

1807) (Heteroptera: Gerridae).

**GenBank:** KJ443338-KJ443340, KJ443354, KJ474895-KJ474897, KJ474907-KJ474909

**Bibliography:** Kostygov et al., 2004; Malysheva and Frolov, 2004; Kostygov et al., 2014.

#### 5 Strain 101

**Depositor:** S.A. Podlipaev

**History:** 1980, isolated by S.A. Podlipaev in Peterhof, St. Petersburg, Russia.

**Host:** *Nabis (Nabicula) flavomarginatus* Scholtz, 1847 (Heteroptera: Nabidae).

**GenBank:** JN944133, AF153039, AF322390, EU088283-EU088285, KJ443308-KJ443311, KJ474867, KJ474935-KJ474937, KJ443306, KJ443307, KJ474866, KJ474933, KJ474934

**Bibliography:** Podlipaev, 1985; Kolesnikov et al., 1990; Marakhova et al., 1991; Podlipaev et al., 1991; Podlipaev et al., 1998; Podlipaev and Frolov, 2000; Merzlyak et al., 2001a; Kostygov et al., 2004; Yurchenko et al., 2006; Malysheva and Frolov, 2009; Yurchenko et al., 2009; Gerasimov et al., 2012a; Kostygov et al., 2014.

#### 6 Strain KV1

**Synonyms:** *Blastocritidia gerricola* Podlipaev, 1985 (in part).

**Depositor:** S.A. Podlipaev

**History:** 1981, isolated by S.A. Podlipaev at the Curonian Spit, Kaliningrad Province, Russia.

**Host:** *Gerris lacustris* (Linnaeus, 1758) (Heteroptera).

**GenBank:** AF153036, AF322391, AJ401379, AY547459; KJ443332-KJ443334, KJ443351, KJ474889-KJ474891, KJ474928-KJ474930.

**Bibliography:** Podlipaev, 1985; Kolesnikov et al., 1990; Podlipaev et al., 1991; Podlipaev et al., 1998; Podlipaev and Frolov, 2000; Merzlyak et al., 2001a, 2001b; Podlipaev et al., 2004; Yurchenko et al., 2006; Kostygov et al., 2014.

#### 7 Strain BM33

**Depositor:** A.O. Frolov

**History:** 1985, isolated by A.O. Frolov in the Berezitsy village, Pskov Province, Russia.

**Host:** *Leptopterna dolobrata* (Linnaeus, 1758) (Heteroptera: Miridae).

**GenBank:** AY547462, AY547483, KJ443312, KJ443313, KJ443355, KJ474868- KJ474870, KJ474916-KJ474918

**Bibliography:** Podlipaev and Frolov, 1987; Podlipaev et al., 2004; Kostygov et al., 2014.

#### 8 Strain F2

**Depositor:** S.A. Podlipaev

**History:** 1986, isolated by S.A. Podlipaev at the Cape Kartesh (White Sea), Karelia, Russia.

**Host:** *Nabis (Nabicula) flavomarginatus* Scholtz, 1847 (Heteroptera: Nabidae).

**GenBank:** AF375664, DQ140170, KJ443317-KJ443319, KJ443344, KJ474874-KJ474876, KJ474913-KJ474915,

**Bibliography:** Podlipaev and Rokitskaya, 1999; Kostygov et al., 2004; Yurchenko et al., 2006; Kostygov et al., 2014.

#### 9 Strain F5

**Depositor:** S.A. Podlipaev

**History:** 1986, isolated by S.A. Podlipaev at the Cape Kartesh (White Sea), Karelia, Russia.

**Host:** *Nabis (Nabicula) flavomarginatus* Scholtz, 1847 (Heteroptera: Nabidae).

**GenBank:** AY547460, KJ443320-KJ443322, KJ443345, KJ474877-KJ474879, KJ474923, KJ474924

**Bibliography:** Podlipaev et al., 1991; Podlipaev and Rokitskaya, 1999; Kostygov et al., 2004; Podlipaev et al., 2004; Kostygov et al., 2014.

#### 10 Strain F7

**Depositor:** S.A. Podlipaev

**History:** 1986, isolated by S.A. Podlipaev at the Cape Kartesh (White Sea), Karelia, Russia.

**Host:** *Nabis (Nabicula) flavomarginatus* Scholtz, 1847 (Heteroptera: Nabidae).

**GenBank:** KJ443326-KJ443328, KJ443347, KJ474883-KJ474885, KJ474910-KJ474912

**Bibliography:** Kostygov et al., 2004; Kostygov et al., 2014.

#### 11 Strain CL8

**Depositor:** A.O. Frolov

**History:** 1984, isolated by A.O. Frolov in Peterhof, St. Petersburg, Russia.

**Host:** *Nabis (Dolichonabis) limbatus* Dahlbom, 1851 (Heteroptera: Nabidae).

**GenBank:** KJ443314-KJ443316, KJ443349, KJ474871-KJ474873, KJ474925-KJ474927

**Bibliography:** Podlipaev and Rokitskaya, 1999; Kostygov et al., 2004; Kostygov et al., 2014.

#### 12 Strain GER4

**Depositor:** M.N. Malysheva

**History:** 2013, isolated by M.N. Malysheva in Komarovo, Leningrad Province, Russia.

**Host:** *Limnoperus rufoscutellatus* (Latreille, 1807) (Heteroptera: Gerridae)

*C. fasciculata* Léger, 1902

#### 13 Strain Cf-C1

**Depositor:** S.A. Podlipaev

**History:** 1992, received from D. Maslov, University of California-Riverside, USA.

**Host:** unknown.

**GenBank:** AF124228, AODS00000000, CFU 05816, CFU12625, CFU30216- CFU30223, D13007-D13036, EU714010, JX271578- JX271580, K03444- K03446, K03448- K03452, L18916, M19066, M19266, M20252, M22852, U05816, U30216- U30223, X03047, X04483, X04760,

**Bibliography:** Birkenmeyer et al., 1985; Ray et al., 1986; Feagin et al., 1988; Maslov et al., 1988; Sugisaki and Ray, 1988; Hughes et al., 1989; van der Spek et al., 1991; Campbell, Ray, 1993; Malysheva and Frolov, 1993; Sugisaki and Takanami, 1993; Maslov et al., 1994; Yasuhira and Simpson, 1995; Podlipaev et al., 1998; Engel and Ray, 1999; Podlipaev and Rokitskaya, 1999; Podlipaev and Frolov, 2000; Lye et al., 2010; Malysheva et al., 2010; Arendt, 2013.

*C. guilhermei* Soares et al., 1986

#### 14 Strain Cg

**Depositor:** S.A. Podlipaev

**History:** 1990, received from E.P. Camargo, São Paulo University, Brazil, isolated in Rio de Janeiro, Brazil.

**Host:** *Lucilia cuprina* (Wiedemann, 1830) (Diptera: Calliphoridae).

**GenBank:** HQ834477

**Bibliography:** Soares et al., 1986; Faria e Silva, et al., 1991; Branquinha et al., 1996; Podlipaev and Rokitskaya, 1999.

*Critchidia* spp.

#### 15 Strain C4

**Depositor:** S.A. Podlipaev

**History:** 1987, isolated by S.A. Podlipaev in the Nizhnesvirsky Nature Reserve, Leningrad Province, Russia.

**Host:** *Limnoperus rufoscutellatus* (Latreille, 1807) (Heteroptera: Gerridae).

**GenBank:** AY547463, AY547482

**Bibliography:** Podlipaev et al., 1998; Skarlato et al., 1998; Podlipaev and Frolov, 2000; Podlipaev et al., 2004.

#### 16 Strain Cfm9

**Depositor:** A.O. Frolov

**History:** 1983, isolated by A.O. Frolov in Peterhof, St. Petersburg, Russia.

**Host:** *Nabis (Nabicula) flavomarginatus* Scholtz, 1847 (Heteroptera: Nabidae).

**GenBank:** AF153041, AF320820, AY547465, AY547491

**Bibliography:** Merzlyak et al., 2001a; Podlipaev et al., 2004; Yurchenko et al., 2006.

#### 17 Strain CL6

**Depositor:** A.O. Frolov

**History:** 1983, isolated by A.O. Frolov in the Berezitsy village, Pskov Province, Russia.

**Host:** *Nabis (Nabicula) limbatus* Dahlbom, 1851.

**Bibliography:** Podlipaev and Rokitskaya, 1999.

Genus *Leptomonas* Kent, 1880

*Leptomonas pyrrhocoris* (Zotta, 1912)

#### 18 Strain LP

**Depositor:** M.N. Malysheva

**History:** Isolated by M.N. Malysheva, Krasnodar Region, Russia.

**Host:** *Pyrrhocoris apterus* Linnaeus, 1758 (Heteroptera: Pyrrhocoridae).

**GenBank:** AY029072, DQ140169

**Bibliography:** Yurchenko et al., 2006; Votýpka et al., 2012.

#### 19 Strain K1

**Depositor:** S.A. Podlipaev

**History:** 2004, isolated by S.A. Podlipaev in Burkina Faso, West Africa.

**Host:** *Dysdercus* sp. (Heteroptera: Pyrrhocoridae).

#### 20 Strain D1

**Depositor:** S.A. Podlipaev

**History:** 2004, isolated by S.A. Podlipaev in Burkina Faso, West Africa.

**Host:** *Dysdercus* sp. (Heteroptera: Pyrrhocoridae).

#### 21 Strain PP1

**Depositors:** A.O. Frolov, M.N. Malysheva

**History:** 2014, isolated in Pskov Province, Russia.

**Host:** *Pyrrhocoris apterus* Linnaeus, 1758 (Heteroptera: Pyrrhocoridae).

**Bibliography:** Frolov et al., 2014.

#### 22 Strain PP2

**Depositors:** A.O. Frolov, M.N. Malysheva

**History:** 2014, isolated in Pskov Province, Russia.

**Host:** *Pyrrhocoris apterus* Linnaeus, 1758 (Heteroptera: Pyrrhocoridae).

**Bibliography:** Frolov et al., 2014.

#### 23 Strain PP3

**Depositors:** A.O. Frolov, M.N. Malysheva

**History:** 2014, isolated in Pskov Province, Russia.

**Host:** *Pyrrhocoris apterus* Linnaeus, 1758 (Heteroptera: Pyrrhocoridae).

**Bibliography:** Frolov et al., 2014.

**24 Strain PP1 cl 10**

**Depositor:** M.N. Malysheva

**History:** Clonal culture. 2014, isolated from the culture of *L. pyrrhocoris* (Strain PP1) in the Laboratory of Protozoology ZIN RAS, St. Petersburg, Russia.

**25 Strain PP1 cl 11**

**Depositor:** M.N. Malysheva

**History:** Clonal culture. 2014, isolated from the culture of *L. pyrrhocoris* (Strain PP1) in the Laboratory of Protozoology ZIN RAS, St. Petersburg, Russia.

**26 Strain PP2 cl 2**

**Depositor:** M.N. Malysheva

**History:** Clonal culture. 2014, isolated from the culture of *L. pyrrhocoris* (Strain PP2) in the Laboratory of Protozoology ZIN RAS, St. Petersburg, Russia.

**27 Strain PP2 cl. 3**

**Depositor:** M.N. Malysheva

**History:** Clonal culture. 2014, isolated from the culture of *L. pyrrhocoris* (Strain PP2) in the Laboratory of Protozoology ZIN RAS, St. Petersburg, Russia.

**28 Strain PP3 cl. 4**

**Depositor:** M.N. Malysheva

**History:** Clonal culture. 2014, isolated from the culture of *L. pyrrhocoris* (Strain PP3) in the Laboratory of Protozoology ZIN RAS, St. Petersburg, Russia.

**29 Strain PP3 cl. 5**

**Depositor:** M.N. Malysheva

**History:** Clonal culture. 2014, isolated from the culture of *L. pyrrhocoris* (Strain PP3) in the Laboratory of Protozoology ZIN RAS, St. Petersburg, Russia.

**30 Strain PP3 cl. 6**

**Depositor:** M.N. Malysheva

**History:** Clonal culture. 2014, isolated from the culture of *L. pyrrhocoris* (Strain PP3) in the Laboratory of Protozoology ZIN RAS, St. Petersburg, Russia.

**31 Strain 25EC**

**Depositor:** S.A. Podlipaev

**History:** 2003, received from D. Maslov, University of California-Riverside, USA. 2003, isolated by D. Maslov, El Ceibo, Costa Rica.

**Host:** nymph *Dysdercus* sp. (Heteroptera: Pyrrhocoridae).

**GenBank:** JF937081

**Bibliography:** Votýpka et al., 2012.

**32 Strain 27EC**

**Depositor:** S.A. Podlipaev

**History:** 2003, received from D. Maslov, University of California-Riverside, USA. 2003, isolated by D. Maslov, El Ceibo, Costa Rica.

**Host:** nymph *Dysdercus* sp. (Heteroptera: Pyrrhocoridae).

**33 Strain 28EC**

**Depositor:** S.A. Podlipaev

**History:** 2003, received from D. Maslov, University of California-Riverside, USA. 2003, isolated by D. Maslov, El Ceibo, Costa Rica.

**Host:** nymph *Dysdercus* sp. (Heteroptera: Pyrrhocoridae).

**GenBank:** JF937082-JF937084

**Bibliography:** Votýpka et al., 2012.

**34 Strain 29EC**

**Depositor:** S.A. Podlipaev

**History:** 2003, received from D. Maslov, University of California-Riverside, USA. 2003, isolated by D. Maslov, El Ceibo, Costa Rica.

**Host:** undetermined nymph Pyrrhocoridae (Heteroptera).

**35 Strain 30EC**

**Depositor:** S.A. Podlipaev

**History:** 2003, received from D. Maslov, University of California-Riverside, USA. 2003, isolated by D. Maslov, El Ceibo, Costa Rica.

**Host:** nymph *Dysdercus* sp. (Heteroptera: Pyrrhocoridae).

**36 Strain 31EC**

**Depositor:** S.A. Podlipaev

**History:** 2003, received from D. Maslov, University of California-Riverside, USA. 2003, isolated by D. Maslov, El Ceibo, Costa Rica.

**Host:** nymph *Dysdercus* sp. (Heteroptera: Pyrrhocoridae).

**37 Strain 32EC**

**Depositor:** S.A. Podlipaev

**History:** 2003, received from D. Maslov, University of California-Riverside, USA. 2003, isolated by D. Maslov, El Ceibo, Costa Rica.

**Host:** nymph *Dysdercus* sp. (Heteroptera: Pyrrhocoridae).

**38 Strain 12VL**

**Depositor:** S.A. Podlipaev

**History:** 2003, received from D. Maslov, University of California-Riverside, USA. 2003, isolated by D. Maslov, Carrara, Costa Rica.

**Host:** *Dysdercus obliquus* (Heteroptera: Pyrrhocoridae).

**GenBank:** JF937077

**Bibliography:** Votýpka et al., 2012.

*Leptomonas podlipaevi* Yurchenko, Lukeš, Xu, Maslov, 2006

**39 Strain UCR5**

**Depositor:** S.A. Podlipaev

**History:** 2003, received from D. Maslov, University of California-Riverside, USA. 2002, isolated by D. Maslov, California, USA.

**Host:** *Boisea rubrolineata* (Barber, 1956) (Heteroptera: Rhopalidae).

**GenBank:** DQ019000; DQ140174, DQ140175, DQ383649

**Bibliography:** Yurchenko et al., 2006.

*L. seymouri* Wallace, 1977

**40 Strain ATCC 30220**

**Depositor:** S.A. Podlipaev

**History:** 1992, received from D. Maslov, University of California-Riverside, USA. 1959, isolated by F.G. Wallace, Florida, USA.

**Host:** *Dysdercus suturellus* (Herrick-Schaeffer, 1842) (Heteroptera: Pyrrhocoridae).

**GenBank:** AF047495, AF053738, AF153040, AF338253, AH015821, AJ243569, AJ245951, AJ313532, AJ318494-AJ318497, AJ401380, AJ511865, AY160127, AY160128, AY547477, DQ239758, DQ239760, DQ239761, EF421167-EF421174, EU623433, GQ153669-GQ153671, JN944129, S78663, U23406, X07487, X07488, X07716, X14005, X78552, X82217

**Bibliography:** Wallace, 1977; Camargo et al., 1978a, 1978b; Galinari and Camargo, 1978; Goncalves de Lima et al., 1979; Fish et al., 1981; Freymuller and Camargo, 1981; Alfieri and Camargo, 1982; Camargo et al., 1982; Dollahon et al., 1983; Kim and Ray, 1985; Bellofatto et al., 1988; Bellofatto and Cross, 1988; Conchon et al., 1989; Teixeira and Camargo, 1989; Gazzinelli et al., 1991; Palfi et al., 1994; Xu et al., 1994; Hartree and Bellofatto, 1995; He and Bellofatto, 1995; Teixeira et al., 1995; Branquinha et al., 1996; Hannaert et al., 1998; Bell and Bindereif, 1999; Podlipaev and Rokitskaya, 1999; Bell et al., 2000; Gilinger and Bellofatto, 2001; Merzlyak et al., 2001a,b; Podlipaev et al., 2004; Bessolitsina et al., 2005; Das and Bellofatto, 2005; Flegontov et al., 2006a,b; Nasereddin et al., 2008; Jackson, 2010; Gerasimov et al., 2012a.

*Leptomonas* spp.

**41 Strain LS**

**Depositor:** S.A. Podlipaev

**History:** 1992, received from D. Maslov,

University of California-Riverside, USA, isolated in USA.

**Host:** unknown.

**GenBank:** J03814, L19409, X53914

**Bibliography:** Lake et al., 1988; Fernandes et al., 1993; Podlipaev and Rokitskaya, 1999.

**42 Strain 24EC**

**Depositor:** S.A. Podlipaev

**History:** 2003, received from D. Maslov, University of California-Riverside, USA. 2003, isolated by D. Maslov, El Ceibo, Costa Rica.

**Host:** undetermined nymph of Pyrrhocoridae (Heteroptera).

Genus *Blastocrithidia* Laird, 1959

*Blastocrithidia* sp.

**43 Strain BLP**

**Depositor:** M.N. Malysheva

**History:** 2015, isolated in Pskov Province, Russia.

**Host:** *Pyrrhocoris apterus* Linnaeus, 1758 (Heteroptera: Pyrrhocoridae).

**Remark:** non-axenic.

Genus *Herpetomonas* Kent, 1880

*H. muscarum* (Leidy, 1856)

Synonym: *H. megaseliae* Daggett et al. 1972

**44 Strain:** ATCC 30260

**Depositor:** S.A. Podlipaev

**History:** 1992, received from D. Maslov, University of California-Riverside, USA. 1968, isolated by F.G. Wallace in Minneapolis, Minnesota, USA.

**Host:** *Musca domestica* Linnaeus, 1758 (Diptera: Muscidae).

**GenBank:** AY547484, AUXJ00000000, AY547484, DQ092548, EU095983, EU095984, FJ807255, JF756655, JN944119, JQ359731, JQ359748, JQ406959-JQ406961, KC503350-KC503410, KF160206-KF160242, KP689382, KP689382, KP689391, KP689399, L10849, L10850, L18872, U01007, U01011, U01015, U05812

**Bibliography:** Rogers and Wallace, 1971; Galinari and Camargo, 1978; Yoshida, 1978; Fish et al., 1981; Freymuller and Camargo, 1981; Coombs et al., 1982; Alfieri and Camargo, 1982; Van der Ploeg et al., 1984; Teixeira and Camargo, 1989; Faria e Silva et al., 1991; Landweber and Gilbert, 1993; Maslov et al., 1994; Lopes et al., 1997; Redman and Coombs, 1997; Podlipaev et al., 1998; Podlipaev and Rokitskaya, 1999; Podlipaev and Frolov, 2000; Podlipaev et al., 2004; Yurchenko et al., 2006; Gile et al., 2009; Yurchenko et al., 2009; Rocha et al.,

2010; Alves et al., 2011; Gerasimov et al., 2012a; Alves et al., 2013; Borghesan et al., 2013; Klein et al., 2013.

*H. samuelpessoai* (Galvao et al., 1970).

Synonyms: *Leptomonas pessoai* Galvao et al., 1970, *H. pessoai* sensu Levine, 1978.

**45 Strain:** ATCC 30252

**Depositor:** S.A. Podlipaev

**History:** 1992, received from D. Maslov, University of California-Riverside, USA, isolated in Brazil.

**Host:** *Zelus leucogrammus* (Perty, 1833) (Heteroptera: Reduviidae).

**GenBank:** AF047494, AF064359-AF064399, AJ401381, AJ493125, FJ807256, JN944120, JQ359716, JQ359743, JQ406989-JQ406991, L10851, L10852, L38605, U01008, U01012, U01016, X62331

**Bibliography:** Galvão et al., 1970; Roitman et al., 1972; Roitman and Roitman, 1972; Camargo et al., 1978; Galinari and Camargo, 1978; Yoshida, 1978; Goncalves de Lima et al., 1979; Fish et al., 1981; Freymuller and Camargo, 1981; Alfieri and Camargo, 1982; Fish et al., 1982; Pinto et al., 1982; Kim and Ray, 1985; Palatnik et al., 1987; Jansen et al., 1988; Conchon et al., 1989; Teixeira and Camargo, 1989; Faria e Silva et al., 1991; Aksoy, 1992; Gazzinelli et al., 1992; Hupperich et al., 1992; Landweber and Gilbert, 1993; Landweber and Gilbert, 1994; Hannaert et al., 1998; Podlipaev et al., 1998; Fu et al., 1999; Podlipaev and Rokitskaya, 1999; Podlipaev and Frolov, 2000; Merzlyak et al., 2001; Gile et al., 2009; Gerasimov et al., 2012a; Borghesan et al., 2013.

*H. nabiculae* (Podlipaev 1985)

Synonyms: *Leptomonas nabiculae* Podlipaev 1985, *L. peterhoffi* Podlipaev 1985, *L. occidentalis* Podlipaev 1985, *Leptomonas* sp. Nfm.

**46 Strain:** Nfm2

**Depositor:** A.O. Frolov

**History:** 1991, isolated by A.O. Frolov in Berezitsy village, Pskov Province, Russia.

**Host:** *Nabis (Nabicia) flavomarginatus* Scholtz, 1847 (Heteroptera: Nabidae).

**GenBank:** JN624300

**Bibliography:** Podlipaev et al., 1998; Podlipaev and Frolov, 2000; Merzlyak et al., 2001a; Podlipaev et al., 2004; Kostygov et al., 2011.

Genus *Wallacemonas* Kostygov et Yurchenko, 2014

*W. collosoma* (Wallace, Clark, Dyer et Collins, 1960)

Synonym: *Leptomonas collosoma* Wallace, Clark, Dyer et Collins, 1960.

**47 Strain:** ATCC 30261

**Depositor:** S.A. Podlipaev

**History:** 1992, received from D. Maslov, University of California-Riverside, USA. 1960, isolated by F.G. Wallace in Minneapolis, Minnesota, USA.

**Host:** *Limnopus dissortis* (Drake and Harris, 1930) (Heteroptera: Gerridae).

**GenBank:** AF006632, AF006750, AF050095, AF126283, AF153038, AF204671, AF331656, AF351198, AH015822, AY046598, AY145448, AY495451, AY495452, AY594669, DQ239762-DQ239765, EU084898, EU084899, HG323624, JF946515, JN582046, JN944123, K02633, L32919, S76723, X56453-X56455, X79014

**Bibliography:** Wallace et al., 1960; Camargo et al., 1978a,b; Galinari and Camargo, 1978; Fish et al., 1981; Freymuller and Camargo, 1981; Alfieri and Camargo, 1982; Camargo et al., 1982; Dollahon et al., 1983; Conchon et al., 1989; Teixeira et al., 1989; Gazzinelli et al., 1991; Ben-Shlomo et al., 1997; Xu et al., 1997; Levitan et al., 1998; Podlipaev et al., 1998; Goncharov et al., 1999; Podlipaev and Rokitskaya, 1999; Li et al., 2000; Podlipaev and Frolov, 2000; Liang et al., 2001; Merzlyak et al., 2001a; Liu et al., 2003; Liang et al., 2004; Yurchenko et al., 2009; Gerasimov et al., 2012a,b; Kostygov et al., 2014.

*W. rigidus* (Podlipaev, Malysheva et Kolesnikov, 1991)

Synonyms: *Leptomonas rigidus*, *Leptomonas* sp. PL11, *Leptomonas* sp. Sld

**48 Strain:** PL11

**Depositor:** S.A. Podlipaev

**History:** 2001, isolated by P. Merkulov and E. Slisarenko at the Cape Kartesh (White Sea), Karelia, Russia.

**Host:** *Salda littoralis* (Linnaeus, 1758) (Heteroptera: Saldidae).

**GenBank:** JN582047, JN944127, JN944128

**Bibliography:** Kostygov et al., 2004; Podlipaev et al., 2004; Gerasimov et al., 2012b.

**49 Strain:** Sld

**Depositor:** S.A. Podlipaev

**History:** 2000, isolated by S.A. Podlipaev at the Cape Kartesh (White Sea), Karelia, Russia.

**Host:** *Saldua pallipes* (Fabricius, 1794) (Heteroptera: Saldidae).

**GenBank:** JF946516, JN582048

**Bibliography:** Kostygov et al., 2004; Podlipaev et al., 2004, Gerasimov et al., 2012a, 2012b.

*Wallacemonas* sp.

**50 Strain Wsd**

**Depositor:** S.A. Podlipaev

**History:** 2001, isolated by P. Merkulov and E. Slisarenko at the Cape Kartesh (White Sea), Karelia, Russia.

**Host:** *Salda littoralis* (Linnaeus, 1758) (Heteroptera: Saldidae).

**GenBank:** AY547488, JF946514, JF946517, JN582045, JN944131, JN944132, JN944134, JN944135

**Bibliography:** Kostygov et al., 2004; Podlipaev et al., 2004, Gerasimov et al., 2012a, 2012b.

**51 Strain TrypX**

**Depositor:** S.A. Podlipaev

**History:** 1997, received from J. Lukeš, Institute of Parasitology, Czech Republic. 1983, isolated in Alexandria, Egypt.

**Host:** *Rattus norvegicus* (Berkenhout, 1769) (Rodentia: Muridae).

**Bibliography:** Morsy et al., 1988; Podlipaev et al., 1998; Podlipaev and Frolov, 2000; Podlipaev et al., 2004.

Genus *Trypanosoma* Gruby, 1843

*Trypanosoma* sp.

**52 Strain TPE**

**Depositor:** M.N. Malysheva

**History:** 2015, isolated by M.N. Malysheva in Pskov Province, Russia.

**Host:** *Pelophylax esculetus* (Linnaeus, 1758).

**Remark:** non-axenic.

Superphylum Opisthosporidia

Phylum Aphelida

Class Aphelidea

Order Aphelidida Gromov, 2000

Family Aphelidiidae Gromov, 2000

Genus *Aphelidium* Zopf, 1885

*Aphelidium* aff. *melosirae*

**53 Strain X-101**

**Depositor:** M.A. Mamkaeva

**History:** 2012, isolated by M.A. Mamkaeva in the vicinity of Sergievka park, Peterhof, St. Petersburg, Russia. Sample P-1.

**Host:** *Tribonema gayanum* Pash.

**GenBank:** KJ566931

**Bibliography:** Karpov et al. 2014.

**54 Strain X-102**

**Depositor:** M.A. Mamkaeva

**History:** 2012, isolated by M.A. Mamkaeva in the vicinity of the Kuty village, Kingisepp District, Leningrad Province, Russia. Sample P-2.

**Host:** *Tribonema gayanum* Pash.

Aphelinida spp.

**55 Strain X-103**

**Depositor:** M.A. Mamkaeva

**History:** 2014, isolated by M.A. Mamkaeva in the vicinity of the Zaborka village, Novgorod Province, Russia. Sample 19.

**Host:** *Tribonema gayanum* Pash.

**56 Strain X-104**

**Depositor:** M.A. Mamkaeva

**History:** 2014, isolated by M.A. Mamkaeva in the vicinity of the Kurakino village, Novgorod Province, Russia. Sample 21.

**Host:** *Tribonema gayanum* Pash.

**57 Strain X-105**

**Depositor:** M.A. Mamkaeva

**History:** 2014, isolated by M.A. Mamkaeva in the vicinity of the Kurakino village, Novgorod Province, Russia. Sample 23.

**Host:** *Tribonema gayanum* Pash.

**58 Strain X-106**

**Depositor:** M.A. Mamkaeva

**History:** 2014, isolated by M.A. Mamkaeva in the vicinity of Peterhof, St. Petersburg, Russia. Sample 24.

**Host:** *Tribonema gayanum* Pash.

**59 Strain X-107**

**Depositor:** M.A. Mamkaeva

**History:** 2015, isolated by M.A. Mamkaeva in the vicinity of the Okulovka town, Novgorod Province, Russia. Sample K-4.

**Host:** *Tribonema gayanum* Pash.

**60 Strain X-108**

**Depositor:** M.A. Mamkaeva

**History:** 2014, isolated by M.A. Mamkaeva from a roadside ditch in the vicinity of the Krestzy village, Novgorod Province, Russia. Sample A-1.

**Host:** *Tribonema gayanum* Pash.

**61 Strain X-109**

**Depositor:** M.A. Mamkaeva

**History:** 2014, isolated by M.A. Mamkaeva from a roadside ditch in the vicinity of the Krestzy village, Novgorod Province, Russia. Sample A-2.

**Host:** *Tribonema gayanum* Pash.

**62 Strain X-110**

**Depositor:** M.A. Mamkaeva

**History:** 2014, isolated by M.A. Mamkaeva from the pond of Krestzy village, Novgorod Province, Russia. Sample A-3.

**Host:** *Tribonema gayanum* Pash.

**63 Strain X-111**

**Depositor:** M.A. Mamkaeva

**History:** 2014, isolated by M.A. Mamkaeva from a roadside ditch in the vicinity of the Zaytzevo village, Novgorod Province, Russia. No specimen: A-6.

**Host:** *Tribonema gayanum* Pash.

**64 Strain X-112**

**Depositor:** M.A. Mamkaeva

**History:** 2014, isolated by M.A. Mamkaeva in the vicinity of Vyborg, Leningrad Province, Russia. Sample V-1.

**Host:** *Tribonema gayanum* Pash.

Class Chytridiomycetes

Genus "Amoeboradix"

"Amoeboradix" sp.

**65 Strain X-113**

**Depositor:** M.A. Mamkaeva

**History:** 2014, isolated by M.A. Mamkaeva from a roadside ditch in the vicinity of Sergievka park, Peterhof, St. Petersburg, Russia. Sample K-1.

**Host:** *Tribonema gayanum* Pash.

**66 Strain X-114**

**Depositor:** M.A. Mamkaeva

**History:** Isolated in the vicinity of the Opolie village, Kingisepp District, Leningrad Province, Russia. Sample K-2.

**Host:** *Tribonema gayanum* Pash.

Genus *Polyphagus* Nowak.

*P. parasiticus* Nowak.

**67 Strain X-115**

**Depositor:** M.A. Mamkaeva

**History:** 2014, isolated by M.A. Mamkaeva from a roadside ditch in the vicinity of the Zaborka village, Novgorod Province, Russia. Sample P.p.

**Host:** *Tribonema gayanum* Pash.

**68 Strain X-116**

**Depositor:** M.A. Mamkaeva

**History:** 2015, isolated by M.A. Mamkaeva in the vicinity of the Zaytzevo village, Novgorod Province, Russia. Sample P-6.

**Host:** *Tribonema gayanum* Pash.

Genus *Chytridium* Braun

*Chytridium* sp.

**69 Strain X-117**

**Depositor:** M.A. Mamkaeva

**History:** 2015, isolated by M.A. Mamkaeva in the vicinity of the Hartchevnia village, Novgorod Province, Russia. Sample H-1.

**Host:** *Tribonema gayanum* Pash.

**70 Strain X-118**

**Depositor:** M.A. Mamkaeva

**History:** isolated by M.A. Mamkaeva from a roadside ditch road, Leningrad Province, Russia. Sample CH-4.

**Host:** *Tribonema gayanum* Pash.

**71 Strain X-119**

**Depositor:** M.A. Mamkaeva

**History:** 2015, isolated by M.A. Mamkaeva in the vicinity of the Zaytzevo village, Novgorod Province, Russia. Sample CH-6.

**Host:** *Tribonema gayanum* Pash.

Genus *Rhizophydium* Schenk

*Rhizophydium* sp.

**72 Strain X-120**

**Depositor:** M.A. Mamkaeva

**History:** 2015, isolated by M.A. Mamkaeva in the vicinity of the Tregubovo village, Novgorod Province, Russia. Sample H-3

**Host:** *Tribonema gayanum* Pash.

**73 Strain X-121**

**Depositor:** M.A. Mamkaeva

**History:** 2015, isolated by M.A. Mamkaeva in the Kuty village, Kingisepp District, Leningrad Province, Russia. Sample KU-1.

**Host:** *Tribonema gayanum* Pash.

**74 Strain X-122**

**Depositor:** M.A. Mamkaeva

**History:** 2015, isolated by M.A. Mamkaeva in the Kuty village, Kingisepp District, Leningrad Province, Russia. Sample KU-2.

**Host:** *Tribonema gayanum* Pash.

Order Mesochytriales Karpov et Aleoshin, 2014

Family Mesochytriaceae Karpov et Aleoshin,

2014

Genus *Mesochytrium* Gromov et al. 2000

*Mesochytrium* sp.

**75 Strain X-123**

**Depositor:** M.A. Mamkaeva

**History:** 2012, isolated by M.A. Mamkaeva in the vicinity of Stary Peterhof, St. Petersburg, Russia. Sample 10.

**Host:** *Tribonema gayanum* Pash.

**GenBank:** FJ804149, FJ804153

**Bibliography:** Karpov et al. 2010.

Order Gromochytriales Karpov et Aleoshin, 2014

Family Gromochytriaceae Karpov et Aleoshin, 2014

Genus *Gromochytrium* Karpov et Aleoshin, 2014

*Gromochytrium* sp.

### 76 Strain X-124

**Depositor:** M.A. Mamkaeva

**History:** 2012, isolated by M.A. Mamkaeva in the Kuty village, Kingisepp District, Leningrad Province, Russia. Sample X-22.

**Host:** *Tribonema gayanum* Pash.

### 77 Strain X-125

**Depositor:** M.A. Mamkaeva

**History:** 2013, isolated by M.A. Mamkaeva in the Zaborka village, Novgorod Province, Russia. Sample V-1.

**Host:** *Tribonema gayanum* Pash.

Class Monoblepharidomycetes

### 78 Strain X-126

**Depositor:** M.A. Mamkaeva

**History:** 2014, isolated by M.A. Mamkaeva in the vicinity of Kotka, Finland. Sample F-1.

**Host:** *Tribonema gayanum* Pash.

### 79 Strain X-127

**Depositor:** M.A. Mamkaeva

**History:** 2014, isolated by M.A. Mamkaeva from a roadside ditch in the vicinity of the Zaborka village, Novgorod Province, Russia. Sample PP1.

**Host:** *Tribonema gayanum* Pash.

### 80 Strain X-128

**Depositor:** M.A. Mamkaeva

**History:** 2014, isolated by M.A. Mamkaeva from a pond in the Zaytzevo village, Novgorod Province, Russia. Sample R.m.

**Host:** *Tribonema gayanum* Pash.

## List of lost cultures

Genus *Critidia* Léger, 1902

*C. brevicula* Frolov et Malyshova, 1989

### 1 Strain NBrA

**Depositor:** A.O. Frolov

**History:** Isolated from culture *C. brevicula* (NBr) in the Laboratory of Protozoology ZIN RAS, St. Petersburg, Russia.

**Host:** *Nabis (Nabis) brevis* (Scholtz, 1847) (Heteroptera).

**Bibliography:** Frolov and Malyshova, 1989a, 1990; Kolesnikov et al., 1990; Marakhova et al., 1991; Podlipaev et al., 1991.

### 2 Strain ZK

**Depositor:** A.O. Frolov

**History:** 1986, isolated by A.O. Frolov in the village Berezitsy, Pskov Province, Russia.

**Host:** *Grypocoris sexguttatus* (Fabricius, 1777) (Heteroptera: Miridae).

**GenBank:** AF121798, AF124056, AF153044, AJ401356, AJ401382, EU076608, EU088280-EU088282, KJ443341-KJ443343, KJ443353, KJ474898-KJ474900, KJ474904-KJ474906

**Bibliography:** Podlipaev and Frolov, 1987; Frolov and Skarlato, 1990; Kolesnikov et al., 1990; Podlipaev et al., 1990; Marakhova et al., 1991; Podlipaev et al., 1991; Marakhova et al., 1993; Podlipaev et al., 1998; Podlipaev and Rokitskaya, 1999; Podlipaev and Frolov, 2000; Yurchenko et al., 2000; Merzlyak et al., 2001a, 2001b; Yurchenko et al., 2009; Kostygov et al., 2014.

### 3 Strain F6

**Depositor:** S.A. Podlipaev

**History:** 1986, isolated by S.A. Podlipaev at the Cape Kartesh (White Sea), Karelia, Russia.

**Host:** *Nabis (Nabicula) flavomarginatus* Scholtz, 1847 (Heteroptera: Nabidae).

**GenBank:** AF153042, AY029073, AY547461, AY547478, KJ443323-KJ443325, KJ443346, KJ474880-KJ474882, KJ474901-KJ474903

**Bibliography:** Kolesnikov et al., 1990; Podlipaev et al., 1991; Merzlyak et al., 2001a; Kostygov et al., 2004; Podlipaev et al., 2004; Kostygov et al., 2014.

### 4 Strain F8

**Depositor:** S.A. Podlipaev

**History:** 1986, isolated by S.A. Podlipaev at the Cape Kartesh (White Sea), Karelia, Russia.

**Host:** *Nabis (Nabicula) flavomarginatus* Scholtz, 1847 (Heteroptera: Nabidae).

**GenBank:** KJ443329-KJ443331, KJ443348, KJ474886-KJ474888, KJ474919-KJ474922

**Bibliography:** Podlipaev and Rokitskaya, 1999; Kostygov et al., 2004; Kostygov et al., 2014.

*C. dedva* Kostygov et al., 2011

### 5 Strain D2

**Depositor:** S.A. Podlipaev

**History:** 1982, isolated by S.A. Podlipaev in the village Beloostrov, Leningrad Province, Russia.

**Host:** *Nabis (Nabicula) flavomarginatus* Scholtz, 1847 (Heteroptera).

**GenBank:** JN624299

**Bibliography:** Podlipaev, 1985; Kolesnikov et al., 1990; Podlipaev et al., 1991; Podlipaev et al., 1998; Kostygov et al., 2004, Kostygov et al., 2011.

*Crithidia* sp.

### 6 Strain P

**Depositor:** S.A. Podlipaev

**History:** 1988, isolated by S.A. Podlipaev in the village Beloostrov, Leningrad Province, Russia.

**Host:** *Panorpa communis* Linnaeus, 1758 (Mecoptera).

**Bibliography:** Podlipaev et al., 1998; Bulat et al., 1999; Podlipaev and Rokitskaya, 1999; Podlipaev and Frolov, 2000; Kostygov et al., 2004; Podlipaev et al., 2004.

Genus *Wallaceonas* Kostygov et Yurchenko, 2014

*W. rigidus* (Podlipaev, Malysheva et Kolesnikov, 1991)

Synonyms: *Leptomonas rigidus* Podlipaev, Malysheva et Kolesnikov, 1991; *Wallaceina rigida*.

### 7 Strain 1b

**Depositor:** S.A. Podlipaev

**History:** 1987, isolated by S.A. Podlipaev at the Cape Kartesh (White Sea), Karelia, Russia.

**Host:** *Salda littoralis* (Linnaeus, 1758) (Heteroptera).

**GenBank:** JN582049

**Bibliography:** Podlipaev et al., 1991, Kolesnikov et al., 1990.

## Acknowledgements

Support for this work was provided by RFBR grant No. 15-29-02734. The strains were isolated and initially cultivated at the Center for Culturing Collection of Microorganisms of the Research park of St. Petersburg State University.

## References

- Aksoy S. 1992. Spliced leader RNA and 5S rRNA genes in *Herpetomonas* spp. are genetically linked. *Nucleic Acids Res.* 20 (4), 913.
- Alfieri S.C. and Camargo E.P. 1982. Trypanosomatidae: isoleucine requirement and threonine deaminase in species with and without endosymbionts. *Exp. Parasitol.* 53 (3), 371–380.
- Alves J.M., Voegtly L., Matveyev A.V., Lara A.M., da Silva F.M., Serrano M.G., Buck G.A., Teixeira M.M. and Camargo E.P. 2011. Identification and phylogenetic analysis of heme synthesis genes in trypanosomatids and their bacterial endosymbionts. *PLoS One.* 6 (8), e23518.
- Alves J.M., Klein C.C., da Silva F.M., Costa-Martins A.G., Serrano M.G., Buck G.A., Vasco-ncelos A.T., Sagot M.F., Teixeira M.M., Motta M.C. and Camargo E.P. 2013. Endosymbiosis in trypanosomatids: the genomic cooperation between bacterium and host in the synthesis of essential amino acids is heavily influenced by multiple horizontal gene transfers. *BMC Evol. Biol.* 13, 190.
- Arendt C.S. 2013. *Crithidia fasciculata* adenosine transporter 1 (CfAT1), a novel high-affinity equilibrative nucleoside transporter specific for adenosine. *Mol. Biochem. Parasitol.* 191 (2), 75–79.
- Bell M. and Bindereif A. 1999. Cloning and mutational analysis of the *Leptomonas seymouri* U5 snRNA gene: function of the Sm site in core RNP formation and nuclear localization. *Nucleic Acids Res.* 27 (20), 3986–3994.
- Bell M., Wöhner R. and Bindereif A. 2000. U4 small nuclear RNA genes of trypanosomes: cloning of the *Leptomonas seymouri* gene and mutational analysis of core snRNP assembly. *Gene.* 247 (1-2), 77–86.
- Bellofatto V. and Cross G.A. 1988. Characterization of RNA transcripts from the alpha tubulin gene cluster of *Leptomonas seymouri*. *Nucleic Acids Res.* 16 (8), 3455–3469.
- Bellofatto V., Cooper R. and Cross G.A. 1988. Discontinuous transcription in *Leptomonas seymouri*: presence of intact and interrupted mini-exon gene families. *Nucleic Acids Res.* 16 (15), 7437–7456.
- Ben-Shlomo H., Levitan A., Béja O. and Michaeli S. 1997. The trypanosomatid *Leptomonas collosoma* 7SL RNA gene. Analysis of elements controlling its expression. *Nucleic Acids Res.* 25 (24), 4977–4984.
- Bessolitsina E.A., Fedyakov A.V., Merzlyak E.M. and Kolesnikov A.A. 2005. The ATPase subunit 6 gene of *Leptomonas seymouri* (Trypanosomatidae) is transcribed and edited as a polycistronic mRNA. *Mol. Biol.* 39 (1), 53–57.
- Birkenmeyer L., Sugisaki H. and Ray D.S. 1985. The majority of minicircle DNA in *Crithidia fasciculata* strain CF-C1 is of a single class with nearly homogeneous DNA sequence. *Nucleic Acids Res.* 13 (19), 7107–7118.
- Borghesan T.C., Ferreira R.C., Takata C.S., Campaner M., Borda C.C., Paiva F., Milder R.V., Teixeira M.M. and Camargo E.P. 2013. Molecular phylogenetic redefinition of *Herpetomonas* (Kinetoplastea, Trypanosomatidae), a genus of insect parasites associated with flies. *Protist.* 164 (1), 129–152.
- Branquinha M.H., Vermelho A.B., Goldenberg S. and Bonaldo M.C. 1996. Ubiquity of cysteine-

- and metalloproteinase activities in a wide range of trypanosomatids. *J. Eukaryot. Microbiol.* 43 (2), 131–135.
- Bulat S.A., Mokrousov I.V. and Podlipaev S.A. 1999. Classification of trypanosomatids from insects and plants by the UP-PCR (universally primed PCR) technique and cross dot blot hybridization of PCR products. *Europ. J. Protistol.* 35, 319–327.
- Camargo E.P., Itow S. and Alfieri S.C. 1978a. Proteolytic activities in cell extracts of trypanosomatids. *J. Parasitol.* 64, 1120–1121.
- Camargo E.P., Coelho J.A., Moraes G. and Figueiredo E.N. 1978b. *Trypanosoma* spp., *Leishmania* spp. and *Leptomonas* spp.: enzymes of ornithine-arginine metabolism. *Exp. Parasitol.* 46, 141–144,
- Camargo E.P., Mattei D.M., Barbieri C.L. and Morel C.M. 1982. Electrophoretic analysis of endonuclease-generated fragments of k-DNA, of esterase isoenzymes, and of surface proteins as aids for species identification of insect trypanosomatids. *J. Protozool.* 29, 25–258.
- Campbell A.G. and Ray D.S. 1993. Functional complementation of an *Escherichia coli* ribonuclease H mutation by a cloned genomic fragment from the trypanosomatid *Crithidia fasciculata*. *Proc. Natl Acad. Sci. USA.* 90 (20), 9350–9354.
- Cho J. and Eichinger D. 1998. *Crithidia fasciculata* induces encystation of *Entamoeba invadens* in a galactose-dependent manner. *J. Parasitol.* 84 (4), 705–710.
- Conchon I., Campaner M., Sbravate C. and Camargo E.P. 1989. Trypanosomatids, other than *Phytomonas* spp., isolated and cultured from fruit. *J. Protozool.* 36, 412–414.
- Coombs G.H. 1982. Proteinases of *Leishmania mexicana* and other flagellate protozoa. *Parasitology*. 84 (1), 149–155.
- Da Silva J.B. and Roitman I. 1982. Effect of temperature and osmolarity on growth of *Crithidia fasciculata*, *Crithidia hutneri*, *Crithidia thermophila*, and *Herpetomonas samuelpessoai*. *J. Eukaryot. Microbiol.* 29 (2), 269–272,
- Das A. and Bellofatto V. 2003. RNA polymerase II-dependent transcription in trypanosomes is associated with a SNAP complex-like transcription factor. *Proc. Natl Acad. Sci. USA.* 100 (1), 80–85.
- Dollahon N.R., Hager R. and Hua T.C. 1983. *Herpetomonas megaseliae*, *Crithidia fasciculata*, and *Leptomonas collosoma* (Kinetoplastida, Trypanosomatidae) in feces of lizards fed culture forms of the flagellates. *J. Protozool.* 30, 58–62.
- Engel M.L. and Ray D.S. 1999. The kinetoplast structure-specific endonuclease I is related to the 5' exo/endonuclease domain of bacterial DNA polymerase I and colocalizes with the kinetoplast topoisomerase II and DNA polymerase beta during replication. *Proc. Natl Acad. Sci. USA.* 96 (15), 8455–8460.
- Faria e Silva P.M., Solý-Cava A.M., Soares M.J., Motta M.C., Fiorini J.E. and de Souza W. 1991. *Herpetomonas roitmani* (Fiorini et al., 1989) n. comb.: a trypanosomatid with a bacterium-like endosymbiont in the cytoplasm. *J. Protozool.* 38 (5), 489–494.
- Feagin J.E., Shaw J.M., Simpson L. and Stuart K. 1988. Creation of AUG initiation codons by addition of uridines within cytochrome b transcripts of kinetoplastids. *Proc. Natl Acad. Sci. USA.* 85 (2), 539–543. Erratum in: *Proc. Natl Acad. Sci. USA.* 85 (10), 3540.
- Fernandes A.P., Nelson K. and Beverley S.M. 1993. Evolution of nuclear ribosomal RNAs in kinetoplastid protozoa: perspectives on the age and origins of parasitism. *Proc. Natl Acad. Sci. USA.* 90 (24), 11608–11612.
- Fiorini J.E., de Faria e Silva P.M., Soares M.J. and Brazil R.P. 1989. Três novas espécies de tripanosomatídeos de insetos isolados em Alfenas, Minas Gerais, Brasil. *Mem. Inst. Oswaldo Cruz.* 84 (1), 69–74.
- Fish W.R., Holz G.G. Jr., Beach D.H., Owen E. and Anekwe G.E. 1981. The cyclopropane fatty acid of trypanosomatids. *Mol. Biochem. Parasitol.* 3, 103–115.
- Fish W.R., Holz G.G. Jr. and Beach D.H. 1982. Some *Phytomonas* and *Herpetomonas* species form unique iso-branched polyunsaturated fatty acids. *Mol. Biochem. Parasitol.* 5 (1), 1–18.
- Flegontov P.N., Strelkova M.V. and Kolesnikov A.A. 2006a. The *Leishmania major* maxicircle divergent region is variable in different isolates and cell types. *Mol. Biochem. Parasitol.* 146 (2), 173–179.
- Flegontov P.N., Guo Q., Ren L., Strelkova M.V. and Kolesnikov A.A. 2006b. Conserved repeats in the kinetoplast maxicircle divergent region of *Leishmania* sp. and *Leptomonas seymouri*. *Mol. Genet. Genomics.* 276 (4), 322–333.
- Freymuller E. and Camargo E. P. 1981. Ultrastructural differences between species of trypanosomatids with and without endosymbionts. *J. Protozool.* 28, 175–182.
- Frolov A.O. and Malysheva M.N. 1989a. Description of *Crithidia allae* sp. n. and *Crithidia brevicula* sp. n. (Protozoa, Trypanosomatidae) from the predator bug *Nabis brevis* Scholtz (Hemiptera,

- Miridae). Zool. Zh. 68 (7), 5–10 (in Russian with English summary).
- Frolov A.O. and Malysheva M.N. 1989b. The developmental cycle of *Crithidia brevicula* (Trypanosomatidae) in the laboratory culture. Tsitologiya. 31, 971–975 (in Russian with English summary).
- Frolov A.O. and Malysheva M.N. 1990. Electron microscopical study of the endomastigotes of *Crithidia allae* (Trypanosomatidae). Zool. Zh. 69 (6), 5–10 (in Russian with English summary).
- Frolov A.O. and Skarlato S.O. 1990. The structure of rosette-like cell associations in the lower trypanosomatids. Tsitologiya. 32(5), 455–461 (in Russian with English summary).
- Frolov A.O. and Malysheva M.N. 1992. The endomastigotes, a special type of the transmission stages of trypanosomatids of the genus *Proteomonas*. Parazitologiya. 26, 351–353 (in Russian with English summary).
- Fu G., Lambson B. and Barker D. 1999. Characterisation of kinetoplast DNA minicircles from *Herpetomonas samuelpessoai*. FEMS Microbiol. Lett. 172 (1), 65–71.
- Galinari S. and Camargo E. P. 1978. Trypanosomatid protozoa: survey of acetylornithinase and ornithine acetyltransferase. Exp. Parasitol. 46, 277–282.
- Galvão A.B., Oliveira R.L., Carvalho A.L.M. and Veiga G.P. 1970. *Leptomonas pessoai* sp. n. (Trypanosomatidae, Kinetoplastida, Protozoa). Rev. Goiana Med. 16, 229–236.
- Gazzinelli R.T., Romanha A.J., Fontes G., Chiari E., Gazzinelli G. and Brener Z. 1991. Distribution of carbohydrates recognized by the lectins *Euonymus europaeus* and concanavalin A in monoxenic and heteroxenic trypanosomatids. J. Protozool. 38, 320–325.
- Gerasimov E.S., Efimova N.S. and Kolesnikov A.A. 2012a. Three patterns of trypanosomatid cryptogene structural organization. Molecular Biology. 46 (4), 548–556.
- Gerasimov E.S., Kostygov A.Y., Yan S. and Kolesnikov A.A. 2012b. From cryptogene to gene? ND8 editing domain reduction in insect trypanosomatids. Europ. J. Protistol. 48 (3), 185–193.
- Gile G.H., Faktorová D., Castlejohn C.A., Burger G., Lang B.F., Farmer M.A., Lukes J. and Keeling P.J. 2009. Distribution and phylogeny of EFL and EF-1alpha inrr5 Euglenozoa suggest ancestral co-occurrence followed by differential loss. PLoS One. 4 (4), e5162.
- Gilinger G. and Bellofatto V. 2001. Trypanosome spliced leader RNA genes contain the first identified RNA polymerase II gene promoter in these organisms. Nucleic Acids Res. 29 (7), 1556–1564.
- Goncalves de Lima V.M., Roitman I. and Kilgour V. 1979. Five trypanosomatid species of insects distinguished by isoenzymes. J. Protozool. 26 (4), 648–652.
- Goncharov I., Palfi Z., Bindereif A. and Michaleli S. 1999. Purification of the spliced leader ribonucleoprotein particle from *Leptomonas collosoma* revealed the existence of an Sm protein in trypanosomes. Cloning the SmE homologue. J. Biol. Chem. 274 (18), 12217–12221.
- Gromov B.V. and Titova N.N. 1991. CALU-collection of algal cultures in the laboratory of microbiology of Biological Institute of Sankt-Petersburg University. In: Catalogue of Microalgal Cultures in the Collections of the USSR (Ed. Semenenco V.S.). IPPAS, Moskow, pp. 76–125.
- Gromov B.V., Mamkaeva K.A. and Pljusch A.V. 2000. *Mesochytrium penetrans* gen. et sp. nov. (Chytridiales) – a parasite of the green algae *Chlorococcum minutum* (Chlorococcales), with an unusual behaviour of the sporangia. Nova Hedwigia. 71, 151–160.
- Hannaert V., Opperdoes F.R. and Michels P.A. 1998. Comparison and evolutionary analysis of the glycosomal glyceraldehyde-3-phosphate dehydrogenase from different Kinetoplastida. J. Mol. Evol. 47 (6), 728–738.
- Hartree D. and Bellofatto V. 1995. Essential components of the mini-exon gene promoter in the trypanosomatid *Leptomonas seymouri*. Mol. Biochem. Parasitol. 71 (1), 27–39.
- He P. and Bellofatto V. 1995. Structure of the *Leptomonas seymouri* trans-spliceosomal U2 snRNA-encoding gene; potential U2-U6 snRNA interactions conform to the cis-splicing counterpart. Gene. 165 (1), 131–135.
- Hollar L., Lukeľ J. and Maslov D.A. 1998. Monophyly of endosymbiont containing trypanosomatids: phylogeny versus taxonomy. J. Eukaryot. Microbiol. 45 (3), 293–297.
- Hughes D.E., Shonekan O.A. and Simpson L. 1989. Structure, genomic organization and transcription of the bifunctional dihydrofolate reductase-thymidylate synthase gene from *Crithidia fasciculata*. Mol. Biochem. Parasitol. 34 (2), 155–166.
- Hupperich K., Camargo E.P. and Milder R. 1992. Ultrastructural study of the host-parasite

- relationship of trypanosomatids in the housefly. Parasitol. Res. 78 (1), 48–55.
- Jackson A.P. 2010. The evolution of amastin surface glycoproteins in trypanosomatid parasites. Mol. Biol. Evol. 27 (1), 33–45.
- Jansen A.M., Carreira J.C. and Deane M.P. 1988. Infection of a mammal by monogenetic insect trypanosomatids (Kinetoplastida, Trypanosomatidae). Mem. Inst. Oswaldo Cruz. 83 (3), 271–272.
- Karpov S.A., Letcher P.M., Mamkaeva M.A. and Mamkaeva K.A. 2010. Phylogenetic position of the genus *Mesochytrium* (Chytridiomycota) based on zoospore ultrastructure and 18S and 28S rRNA gene sequences. Nova Hedwigia. 90 (1-2), 81–94.
- Karpov S.A., Mamkaeva M.A., Benzerara K., Moreira D. and López-García P. 2014. Molecular phylogeny and ultrastructure of *Aphelidium* aff. *melosirae* (Aphelida, Opisthosporidia). Protist. 165, 512–526.
- Kim R. and Ray D.S. 1985. Conservation of a 29-base-pair sequence within maxicircle ARS fragments from six species of trypanosomes. Gene. 40 (2-3), 291–299.
- Klein C.C., Alves J.M., Serrano M.G., Buck G.A., Vasconcelos A.T., Sagot M.F., Teixeira M.M., Camargo E.P. and Motta M.C. 2013. Biosynthesis of vitamins and cofactors in bacterium-harbouring trypanosomatids depends on the symbiotic association as revealed by genomic analyses. PLoS One. Nov 19; 8 (11), e79786.
- Kolesnikov A.A., Maslov D.A. and Podlipaev S.A. 1990. Comparative restriction enzyme cleavage analysis of kinetoplast DNA from the lower trypanosomatids isolated in the North-West region of the USSR. Arch. Protistenkd. 138, 239–250.
- Kostygov A.Y., Sliarenko E.A., Merkulov P.A. and Podlipaev S.A. 2004. Genetic diversity of insect trypanosomatids from subarctic and North-West Russia revealed by UP-PCR typing. Protistology. 3 (4), 257–264.
- Kostygov A.Y., Malysheva M.N. and Frolov A.O. 2011. Investigation of causes of the conflict between taxonomy and molecular phylogeny of trypanosomatids by the example of *Leptomonas nabiculae* Podlipaev, 1987. Parazitologiya. 45 (6), 409–424 (in Russian with English summary).
- Kostygov A.Y., Grybchuk-Leremenko A., Malysheva M.N., Frolov A.O. and Yurchenko V. 2014. Molecular revision of the genus *Wallaceina*. Protist. 165, 594–604.
- Lake J.A., de la Cruz V.F., Ferreira P.C., Morel C. and Simpson L. 1988. Evolution of parasitism: kinetoplastid protozoan history reconstructed from mitochondrial rRNA gene sequences. Proc. Natl Acad. Sci. U.S.A. 85 (13), 4779–4783.
- Landweber L.F. and Gilbert W. 1993. RNA editing as a source of genetic variation. Nature. 363 (6425), 179–182.
- Landweber L.F. and Gilbert W. 1994. Phylogenetic analysis of RNA editing: a primitive genetic phenomenon. Proc. Natl. Acad. Sci. USA. 91 (3), 918–921.
- Levitin A., Xu Y., Ben-Dov C., Ben-Shlomo H., Zhang Y. and Michaeli S. 1998. Characterization of a novel trypanosomatid small nucleolar RNA. Nucleic Acids Res. 26 (7), 1775–1783.
- Li L., Otake L.R., Xu Y. and Michaeli S. 2000. The trans-spliceosomal U4 RNA from the monogenetic trypanosomatid *Leptomonas collosoma*. Cloning and identification of a transcribed tRNA-like element that controls its expression. J. Biol. Chem. 275 (4), 2259–2264.
- Liang X.H., Liu L. and Michaeli S. 2001. Identification of the first trypanosome H/ACA RNA that guides pseudouridine formation on rRNA. J. Biol. Chem. 276 (43), 40313–40318.
- Liang X.H., Ochaion A., Xu Y.X., Liu Q. and Michaeli S. 2004. Small nucleolar RNA clusters in trypanosomatid *Leptomonas collosoma*. Genome organization, expression studies, and the potential role of sequences present upstream from the first repeated cluster. J. Biol. Chem. 279 (7), 5100–5119.
- Liu L., Ben-Shlomo H., Xu Y.X., Stern M.Z., Goncharov I., Zhang Y. and Michaeli S. 2003. The trypanosomatid signal recognition particle consists of two RNA molecules, a 7SL RNA homologue and a novel tRNA-like molecule. J. Biol. Chem. 278 (20), 18271–18280.
- Lopes A.H., Dutra P.M., Rodrigues C.O., Soares M.J., Angluster J. and Cordeiro R.S. 1997. Effect of platelet-activating factor on the process of cellular differentiation of *Herpetomonas muscarum muscarum*. J. Eukaryot. Microbiol. 44 (4), 321–325.
- Lye L.F., Owens K., Shi H., Murta S.M., Vieira A.C., Turco S.J., Tschudi C., Ullu E. and Beverley S.M. 2010. Retention and loss of RNA interference pathways in trypanosomatid protozoans. PLoS Pathog. 6 (10), e1001161.
- Malysheva M.N. and Skarlato S.O. 1989. Studies on the ultrastructural organization of *Leptomonas peterhoffi* (Trypanosomatidae) grown on liquid and solid culture media. Tsitologiya. 31 (3): 267–272 (in Russian with English summary).
- Malysheva M.N. and Frolov A.O. 1993. Free-living disseminating instars of trypanosomatids from

- water insects. *Parazitologiya*. 27 (4), 337–340 (in Russian with English summary).
- Malysheva M.N. and Frolov A.O. 1995. The developmental cycle of *Proteomonas breviculus* (Trypanosomatidae) in the predatory bugs of the family Nabidae (Hemiptera). *Parazitologiya*. 29 (4), 289–297 (in Russian with English summary).
- Malysheva M.N., Frolov A.O. and Podlipaev S.A. 1995. Two stages life cycle of *Proteomonas brevicula* (Mastigophora: Trypanosomatidae): isolation from host and their investigations *in vitro*. Abstr. 2nd Europ. Congr. Protistology, Clermont-Ferrand. P. 66.
- Malysheva M.N., Podlipaev S.A. and Frolov A.O. 2001. List of living cultures of homoxenous trypanosomatids in the collection of the Laboratory of Protozoology, Zoological Institute, St. Petersburg (Kinetoplastida). *Zoosystematica Rossica*. 9, 245–246.
- Malysheva M.N. and Frolov A.O. 2004. The description of *Wallaceina vicina* sp. n. (Kinetoplastida: Trypanosomatidae) from the water strider *Gerris rufoscutellatus* Latreille, 1807 (Hemiptera: Gerridae). *Parazitologiya*. 38 (5), 470–476 (in Russian with English summary).
- Malysheva M.N. and Frolov A.O. 2009. On the problem of identification of homoxenous trypansome cultures with the description of a new species *Wallaceina podlipaevi* sp. n. (Kinetoplastida: Trypanosomatidae). *Parazitologiya*. 43 (6), 502–515 (in Russian with English summary).
- Malysheva M.N., Danilov K.L., Kalninsh K.K., Lavrik N.L., Fokin G.A., Frolov A.O. and Skarlato S.O. 2010. Influence of thermoconditioned types of water on the proliferative activity of flagellates *Crithidia fasciculata*. *Water: Technology and Ecology*. 3, 20–29 (in Russian with English summary).
- Marakhova N.V., Skarlato S.O., Frolov A.O. and Tsouladze A.M. 1991. The molecular karyotype polymorphism in the lower trypanosomatids. *Tsitologiya*. 33 (10), 59–64 (in Russian with English summary).
- Marché S., Roth C., Philippe H., Dollet M. and Baltz T. 1995. Characterization and detection of plant trypanosomatids by sequence analysis of the small subunit ribosomal RNA gene. *Mol. Biochem. Parasitol.* 71 (1), 15–26.
- Maslov D.A., Rezekina L.A. and Kolesnikov A.A. 1988. Some sequence elements of minicircle DNA from the kinetoplast of the trypanosomatid *Crithidia fasciculata*. *Molecular Biology (Moscow)* 22, 1482–1490.
- Maslov D.A., Avila H.A., Lake J.A. and Simpson L. 1994. Evolution of RNA editing in kinetoplastid protozoa. *Nature*. 368 (6469), 345–348.
- Merzlyak E., Yurchenko V., Kolesnikov A.A., Alexandrov K., Podlipaev S.A. and Maslov D.A. 2001a. Diversity and phylogeny of insect trypanosomatids based on small subunit rRNA genes: polyphyly of *Leptomonas* and *Blastocrithidia*. *J. Eukaryot. Microbiol.* 48 (2), 161–169.
- Merzlyak E.M., Zakharova M.Yu. and Kolesnikov A.A. 2001b. Monogenetic trypanosomatids: comparison of the ND8 editing gene. *Eur. J. Protistol.* 37, 233–239.
- Morsy T.A., Schnur L.F., Feinsod F.M., Michael S.A., Saah A., Salama M.M. and Wahba M.M. 1988. The discovery and preliminary characterization of a novel trypanosomatid parasite from *Rattus norvegicus* and stray dogs from Alexandria, Egypt. *Ann. Trop. Med. Parasitol.* 82, 437–444.
- Nasereddin A., Bensoussan-Hermano E., Schönian G., Baneth G. and Jaffe C.L. 2008. Molecular diagnosis of Old World cutaneous leishmaniasis and species identification by use of a reverse line blot hybridization assay. *J. Clin. Microbiol.* 46 (9), 2848–2855.
- Palatnik C.B., Previato J.O., Gorin P.A. and Mendonça-Previato L. 1987. *Leptomonas samueli* glycoconjugates. Comparison with *Herpetomonas samuelssoai*. *Comp. Biochem. Physiol. B*. 86 (3), 593–599.
- Palfi Z., Xu G.L. and Bindereif A. 1994. Spliced leader-associated RNA of trypanosomes. Sequence conservation and association with protein components common to trans-spliceosomal ribonucleoproteins. *J. Biol. Chem.* 269 (48), 30620–30625.
- Pinto A.S., Pinto A.C., De Souza W. and Angluster J. 1982. Fatty acid composition in *Herpetomonas samuelssoai*. Influence of growth conditions. *Comp. Biochem. Physiol.* 73, 351–356.
- Podlipaev S.A. 1985. New species of lower trypanosomatids from true bugs (Heteroptera) of families Gerridae and Nabidae: life cycle stages in nature and during laboratory cultivation. *Proc. Zool. Inst. Ac. Sci USSR*. 129, 35–47 (in Russian with English summary).
- Podlipaev S.A. and Frolov A.O. 1987. Description and laboratory cultivation of *Blastocrithidia miridarum* sp. n. (Mastigophora, Trypanosomatidae). *Parazitologiya*. 21 (4), 545–552 (in Russian with English summary).
- Podlipaev S.A., Frolov A.O. and Kolesnikov A.A. 1990. *Proteomonas inconstans* n. gen., n. sp.

- (Kinetoplastida: Trypanosomatidae), a parasite of the bug *Calocoris sexguttatus* (Hemiptera: Miridae). *Parazitologiya*. 24 (4), 339–346 (in Russian with English summary).
- Podlipaev S.A., Filatov M.V. and Pantina K.A. 1991. A comparative determination of DNA content in the lower trypanosomatids from Hemiptera of the north-west of the USSR. *Parazitologiya*. 25 (3), 250–257 (in Russian with English summary).
- Podlipaev S.A., Malysheva M.N. and Kolesnikov A.A. 1991. *Leptomonas rigidus* sp. n. (Trypanosomatidae) – a parasite of *Salda littoralis* L. (Hemiptera: Heteroptera). *Acta Protozool.* 30, 121–127.
- Podlipaev S.A., Mokrousov I.V. and Bulat S.A. 1998. Towards molecular systematics of trypanosomatids from insects by means of the polymerase chain reaction with universal primers (UP-PCR). *Parazitologiya*. 32 (4), 317–326 (in Russian with English summary).
- Podlipaev S.A. and Rokitskaya T.A. 1999. Classification of isolates of insect's trypanosomatids: isoenzyme analysis. *Parazitologiya*. 33 (4), 350–357 (in Russian with English summary).
- Podlipaev S.A. and Frolov A.O. 2000. Phylogeny of Trypanosomatids: molecular and morphological researches. *Parazitologiya*. 34 (3), 169–182 (in Russian with English summary).
- Podlipaev S.A., Sturm N.R., Fiala I., Fernandes O., Westenberger S.J., Dollet M., Campbell D.A. and Lukeš J. 2004. Diversity of insect trypanosomatids assessed from the spliced leader RNA and 5S rRNA genes and intergenic region. *J. Eukaryot. Microbiol.* 51 (3), 283–290.
- Ray D.S., Hines J.C., Sugisaki H. and Sheline C. 1986. kDNA minicircles of the major sequence class of *C. fasciculata* contain a single region of bent helix widely separated from the two origins of replication. *Nucleic Acids Res.* 14 (20), 7953–7965.
- Redman C.A. and Coombs G.H. 1997. The products and pathways of glucose catabolism in *Herpetomonas muscarum ingenoplastis* and *Herpetomonas muscarum muscarum*. *J. Eukaryot. Microbiol.* 44 (1), 46–51.
- Rocha L. de S., dos Santos C.B., Falqueto A., Grimaldi G.Jr. and Cupolillo E. 2010. Molecular biological identification of monoxenous trypanosomatids and *Leishmania* from anthropophilic sand flies (Diptera: Psychodidae) in Southeast Brazil. *Parasitol. Res.* 107 (2), 465–468.
- Rogers W.E. and Wallace F.G. 1971. Two new subspecies of *Herpetomonas muscarum* (Leidy, 1856) Kent, 1880. *J. Protozool.* 18 (4), 645–654.
- Roitman I. and Roitman C. 1972. Increase growth inhibition of trypanosomatidal drugs at 37 °C. *J. Protozool.* 19, 59.
- Roitman C., Roitman I. and Azevedo H.P. 1972. Growth of an insect trypanosomatid at 37 °C in a defined medium. *J. Protozool.* 19, 346–349.
- Shaw J.M., Feagin J.E., Stuart K. and Simpson L. 1988. Editing of kinetoplastid mitochondrial mRNAs by uridine addition and deletion generates conserved amino acid sequences and AUG initiation codons. *Cell*. 53 (3), 401–411.
- Skarlato S.O., Parshkova T.A. and Frolov A.O. 1990. Electron microscope and molecular biology investigation of the nucleus in the lower trypanosomatids, *Blastocrithidia miridarum* and *Crithidia brevicula*. *Tsitologiya*. 32 (4), 317–324 (in Russian with English summary).
- Skarlato S.O., Somova N.V., Rozanov Yu.M., Kudryavtsev B.N., Bobyleva N.N. and Frolov A.O. 1998. Characteristic organization of the chromosomal apparatus of *Crithidia* sp. *Tsitologiya*. 40 (11), 991–1005 (in Russian with English summary).
- Soares M.J., Brazil R.P., Tanuri A. and de Souza W. 1986. Some ultrastructural aspects of *Crithidia guilhermei* n. sp. isolated from *Phaenicia cuprina* (Diptera: Calliphoridae). *Can. J. Zool.* 64 (12), 2837–2842.
- Sugisaki H. and Ray D.S. 1987. DNA sequence of *Crithidia fasciculata* kinetoplast minicircles. *Mol. Biochem. Parasitol.* 23 (3), 253–263.
- Sugisaki H. and Takanami M. 1993. The 5'-terminal region of the apocytochrome b transcript in *Crithidia fasciculata* is successively edited by two guide RNAs in the 3' to 5' direction. *J. Biol. Chem.* 268 (2), 887–891.
- Teixeira M.M. and Camargo E.P. 1989. Monoclonal antibodies for the identification of trypanosomatids of the genus *Phytomonas*. *J. Protozool.* 36, 262–264.
- Teixeira M.M.G., Campaner M. and Camargo E.P. 1995. Characterization of the target antigens of *Phytomonas*-specific monoclonal antibodies. *J. Eukaryot. Microbiol.* 42 (3), 232–237.
- Van der Ploeg L.H., Cornelissen A.W., Barry J.D. and Borst P. 1984. Chromosomes of kinetoplastida. *EMBO J.* 3 (13), 3109–3115.
- Van der Spek H., Arts G.J., Zwaal R.R., van den Burg J., Sloof P. and Benne R. 1991. Conserved genes encode guide RNAs in mitochondria of *Crithidia fasciculata*. *EMBO J.* 10 (5), 1217–1224.
- Votýpková J., Klepetková H., Yurchenko V., Horák A., Lukeš J. and Maslov D. 2012. Cosmopolitan

- distribution of a trypanosomatid *Leptomonas pyrrhocoris*. *Protist.* 163, 616–631.
- Wallace F.G., Clark T.B., Dyer M.I. and Collins T. 1960. Two new species of flagellates cultivated from insects of the genus *Gerris*. *J. Protozool.* 7, 390–393.
- Wallace F.G. 1977. *Leptomonas seymouri* sp. n. from the cotton stainer *Dysdercus suturellus*. *J. Protozool.* 24, 483–484.
- Xu G. L., Wieland B. and Bindereif A. 1994. Trans-spliceosomal U6 RNAs of *Crithidia fasciculata* and *Leptomonas seymouri*: deviation from the conserved ACAGAG sequence and potential base pairing with spliced leader RNA. *Mol. Cell Biol.* 14 (7), 4565–4570.
- Xu Yx, Ben-Shlomo H. and Michaeli S. 1997. The U5 RNA of trypanosomes deviates from the canonical U5 RNA: the *Leptomonas collosoma* U5 RNA and its coding gene. *Proc. Natl. Acad. Sci. USA.* 94 (16), 8473–8478.
- Xu Y, Liu L, Lopez-Estraco C. and Michaeli S. 2001. Expression studies on clustered trypanosomatid box C/D small nucleolar RNAs. *J. Biol. Chem.* 276 (17), 14289–14298.
- Yasuhira S. and Simpson L. 1995. Minicircle-encoded guide RNAs from *Crithidia fasciculata*. *RNA.* 1 (6), 634–643.
- Yoshida N. 1978. *Herpetomonas mariadeanei* sp. n. (Protozoa, Trypanosomatidae) from *Muscina stabulans* (Fallen, 1816) (Diptera, Muscidae). *J. Protozool.* 25 (4), 421–425.
- Yurchenko V., Kolesnikov A. A. and Lukeš J. 2000. Phylogenetic analysis of Trypanosomatina (Protozoa: Kinetoplastida) based on minicircle conserved regions. *Folia Parasitol.* 47, 1–5.
- Yurchenko V., Lukeš J., Xu X. and Maslov D.A. 2006. An integrated morphological and molecular approach to a new species description in the Trypanosomatidae: the case of *Leptomonas podlipaevi* n. sp., a parasite of *Boisea rubrolineata* (Hemiptera: Rhopalidae). *J. Eukaryot. Microbiol.* 53 (2), 103–111.
- Yurchenko V.Y., Lukeš J., Jirku M. and Maslov D. A. 2009. Selective recovery of the cultivation-prone components from mixed trypanosomatid infections: a case of several novel species isolated from Neotropical Heteroptera. *Int. J. Syst. Evol. Microbiol.* 59 (4), 893–909.

**Address for correspondence:** Alexander O. Frolov. Angliyskiy prosp. 32, 190121 Saint-Petersburg, Russia; e-mail: [frolal@yandex.ru](mailto:frolal@yandex.ru)