

Review of tintinnid species (Ciliophora: Tintinnina) from the Barents Sea

Olga Chovgan

Murmansk Marine Biological Institute, Kola Scientific Center of the Russian Academy of Sciences, Murmansk, Russia

| Submitted June 18, 2019 | Accepted July 01, 2019 |

Summary

The paper presents a list of tintinnid species (Tintinnina Kofoid et Campbell, 1929) in the Barents Sea according to records from the XX through the XXI century. The research also includes original data (2012–2017), which allowed identifying three new tintinnid species in recent years: *Ormosella hackeli* Kofoid et Campbell, 1929, *Parundella caudata* (Ostenfeld, 1899) Jörgensen, 1924, and *Amphorellopsis tetragona* (Jörgensen, 1924) Kofoid et Campbell, 1929. These three alien species of marine ciliates belong to the warm-water fauna; they were recorded in the western part of the Barents Sea and are inherited to the North Cape Current. The research includes a brief biogeographic characteristic and photographic material of alien tintinnids in the Barents Sea.

Key words: Barents Sea, ciliates, distribution, plankton, Tintinnina

Introduction

Currently the total number of tintinnid species (suborder Tintinnina Kofoid et Campbell, 1929) in the Barents Sea remains unknown. There are heterogeneity and inconsistency of taxonomy information in plankton records (Lifshits, 1978; Burkovsky, 1974; Dolan et al., 2017). The morphological variability of lorica complicates distinguishing between typical forms of tintinnids, varieties and different stages of life cycle (Rossolimo, 1927; Lifshits, 1977; Dolan et al., 2017). In addition, records of the species composition of ciliates in the Western Arctic that allow to reliably assess the present ciliate diversity in the Barents Sea are limited

and episodic (Meunier, 1910; Burkovsky et al., 1974; Burkovsky, 1976; Lifshits, 1977, 1978; Lifshits et al., 1978; Dolan et al., 2017).

In recent decades, the thermal and ice regimes have been changed in the Barents Sea (Polyakov et al., 2017; Aksenov and Ivanov, 2018). This makes a comparative analysis of the tintinnid ciliates from the Barents Sea in the XX century and at present an ecologically relevant issue. The Atlantic currents provide more than 50% of the warm water contribution to the Barents Sea, so “atlantisation” is one of the key causes of water warming in the Western Arctic (Aksenov and Ivanov, 2018). Thus, there appears a possibility of natural invasion of the Barents Sea by the warm-water tintinnid species.




Fig. 1. General view of the Barents Sea tintinnids: *Amphorellopsis tetragona* (A), *Ormosella haeckeli* (B), *Parundella caudata* (C).

Material and methods

To analyze the species composition of tintinnids, we summarized the published data from the XX century until present. In addition, this research bases on the original data from the plankton samples of the Murmansk Marine Biological Institute collected in 2012–2017. The plankton material was collected in the autumn-winter and spring seasons; the sampling area is located in the western part of the Barents Sea: the standard oceanographic section “Kola meridian” (69°30′–77° N, 33°30′ E). Sampling was carried out using a bathometer and plankton net with a mesh size of 30 μm; the obtained samples were concentrated by gentle inverse filtering through a lavsan filter and were fixed in buffered formaldehyde (1% final concentration). A Nageotte counting chamber was used to examine ciliate assemblages. In order to clarify the tissues of the organisms, a hypochlorite solution was applied, followed by heating.

Results and discussion

According to the references, the taxonomic composition of tintinnid ciliates is represented by 20 genera, including 73 species (Table 1). Original data display first appearance of three alien tintinnid species in the Barents Sea region. In 2012–2017, the presence of *Parundella caudata* (Ostenfeld, 1899) Jörgensen, 1924 (Fig. 1C) and *Ormosella haeckeli* Kofoid et Campbell, 1929 (Fig. 1B) (Makarevich and Oleinik, 2017) was confirmed. *Amphorellopsis tetragona* (Jörgensen, 1924) Kofoid et Campbell, 1929 (Fig. 1A) has not been previously noted in the Barents Sea and adjacent waters (Meunier, 1910;

Rossolimo, 1927; Burkovsky et al., 1974; Burkovsky, 1976; Lifshits, 1977; Dolan et al., 2017).

***Amphorellopsis tetragona* (Jörgensen, 1924) Kofoid and Campbell, 1929 (= *Amphorella tetragona* Jörgensen, 1924).**

The species was described in the Mediterranean Sea (Brandt, 1906; Jörgensen, 1924; Zakaria et al., 2007; Dolan, 2016). *A. tetragona* was noted in the Eastern Pacific (Kofoid and Campbell, 1929; Gomez, 2007), in the Marmara (Toklu-Aliçli et al., 2010; Durmus and Balkis, 2014) and the Adriatic (Monti et al., 2012) seas.

In the original description (Kofoid, Campbell, 1929), the genus *Amphorellopsis* included 7 species, which differed mainly in the number and degree of development of the facets and plicae. A key feature that differentiates *A. tetragona* from other species of genus *Amphorellopsis* is four alary plicae and more flaring mouth above fusiform bowl (Kofoid, Campbell, 1929).

***Ormosella haeckeli* Kofoid and Campbell, 1929.**

For the first time, tintinnids of this species were found in the Pacific Ocean: Peruvian and Equatorial Counter Current (Kofoid and Campbell, 1929); date on *O. haeckeli* was included to ciliate fauna of the South Atlantic (Alder, 1999; Dolan and Pierce, 2014). Previously this species was listed for the Barents Sea as *Ormosella* aff. *cornucopia* (Makarevich and Oleinik, 2017).

In the first description, the genus *Ormosella* included 7 species, of which 3 are characterized by common features: contracted bell-shaped lorica; wall of lorica is hyaline without agglutinates; the collar is clearly defined, in the shape of a truncated cone; the aboral part is sharply narrowed. The

Table 1. (Continuation).

Species (according to the a World Register of Marine Species "WoRMS")	Literature source†												Original name in the literature source	
	1	2	3	4	5	6	7	8	9	10	11	12		
<i>Ptychocylis drygalskii</i> Brandt, 1896	+						+							
<i>Ptychocylis minor</i> Gruber, 1879*							+							
<i>Ptychocylis obtusa</i> Brandt, 1906		+		+	+	+	+	+	+	+	+			
<i>Ptychocylis urnula</i> (Claparède et Lachmann, 1858)							+		+		+			
<i>Salpingacantha ampla</i> Kofoid et Campbell, 1929							+							
<i>Salpingacantha undata</i> (Jørgensen, 1899)							+							
<i>Salpingacantha unguiculata</i> (Brandt, 1906) Kofoid et Campbell, 1929							+	+						<i>Salpingella</i> cf. <i>ungiculata</i> (8)
<i>Salpingella acuminata</i> (Claparède et Lachmann, 1858) Jørgensen, 1924	+	+	+	+	+	+	+		+	+				
<i>Salpingella decurtata</i> Jørgensen, 1924							+							<i>Tintinnus acuminatus</i> (1, 9), <i>Salpingella</i> aff. <i>acuminate</i> (10)
<i>Salpingella minutissima</i> Kofoid et Campbell, 1929							+							
<i>Salpingella rotundata</i> (Kofoid et Campbell, 1929)**							+							
<i>Salpingella secata</i> (Brandt, 1896)							+	+						
<i>Steenstrupiella steenstrupii</i> (Claparède et Lachmann, 1858) Kofoid et Campbell, 1929							+							
<i>Stenosemella oliva</i> (Meunier, 1910)		+	+	+	+		+	+						<i>Stenostomella</i> cf. <i>oliva</i> (8)
<i>Stenosemella robusta</i> *		+	+	+	+									
<i>Stenosemella steinii</i> Jørgensen, 1912									+		+			
<i>Stenosemella ventricosa</i> (Claparède et Lachmann, 1858) Jørgensen, 1924							+		+		+			
<i>Tintinnopsis acuta</i> *	+													
<i>Tintinnopsis angusta</i> Meunier, 1910	+													
<i>Tintinnopsis baltica</i> Brandt, 1896							+							
<i>Tintinnopsis beroidea</i> Stein, 1867		+	+		+		+		+		+			
<i>Tintinnopsis campanula</i> Ehrenberg, 1840			+		+				+		+			
<i>Tintinnopsis cylindrica</i> Daday, 1887	+								+					
<i>Tintinnopsis fimbriata</i> Meunier, 1919		+	+		+		+		+					
<i>Tintinnopsis fucus</i> *									+					
<i>Tintinnopsis japonica</i> Hada, 1932		+			+									
<i>Tintinnopsis karajacensis</i> Brandt, 1896							+		+					
<i>Tintinnopsis lata</i> Meunier, 1910	+						+							
<i>Tintinnopsis major</i> Meunier, 1910	+													
<i>Tintinnopsis meunieri</i> Kofoid et Campbell, 1929			+				+							
<i>Tintinnopsis nana</i> Lohmann, 1908			+		+		+		+					
<i>Tintinnopsis nitida</i> Brandt, 1986							+		+					
<i>Tintinnopsis oliva</i> *	+													
<i>Tintinnopsis parvata</i> (Joergensen, 1924)*									+					

Table 1. (Continuation).

Species (according to the a World Register of Marine Species "WoRMS")	Literature source†												Original name in the literature source
	1	2	3	4	5	6	7	8	9	10	11	12	
<i>Tintinnopsis parvula</i> Jörgensen, 1912		+	+		+		+			+			
<i>Tintinnopsis patula</i> Meunier, 1910	+												
<i>Tintinnopsis rapa</i> Meunier, 1910							+						
<i>Tintinnopsis sacculus</i> Brandt, 1896									+		+		
<i>Tintinnopsis sinuata</i> Brandt, 1896							+						
<i>Tintinnopsis tubulosa</i> Levander, 1900									+		+		
<i>Tintinnopsis turbo</i> Meunier, 1919													
<i>Tintinnopsis undella</i> Meunier, 1910		+			+		+						

† Literature sources: 1 – Meunier, 1910; 2 – Burkovsky, 1976; 3 – Lifshits, 1977; 4 – Lifshits, 1978; 5 – Lifshits et al., 1978; 6 – Boltovskoy et al., 1991; 7 – Matishov et al, 2000; 8 – Jensen and Hansen, 2000; 9 – Dvoretzkiy, 2010; 10 – Makarevich and Oleynik, 2017; 11 – Dvoretzkiy and Dvoretzkiy, 2018; 12 – this work. * Species is absent in base of "WoRMS"; ** "nomen dubium"; *** "taxon inquirendum".

researched tintinnids were identified as *O. haeckeli* Kofoid and Campbell, 1929 by 12 longitudinal equal facets on the bowl and pedicel; large collar, clearly prominent from a bowl with an inner nuchal edge (Kofoid and Campbell, 1929).

***Parundella caudata* (Ostenfeld, 1899) Jörgensen, 1924** (= *Tintinnus caudatus* Ostenfeld, 1899; = *Undella* (*Parundella*) *caudata* Jörgensen, 1899).

A typical representative of the ciliate fauna, inhabiting the waters of the South Atlantic (Fernandes, 2004; Thompson and Alder, 2005; Kršinić and Grbec, 2006) and the northern Indian Ocean (Asha Devi et al., 2018); one was found in the Pacific Ocean, the coast of the Philippines and Chile (Fonda Umani et al., 2011; Santiago, 2018). The nearest location to the Barents Sea is North Atlantic, Norwegian, North and Baltic seas (Marshall, 1969; Cordeiro and Sassi, 1997). The earlier information for the presence *P. caudata* in the Barents Sea (Makarevich and Oleinik, 2017) is confirmed by this report.

The first description (Kofoid and Campbell, 1929) included 22 species of the genus *Parundella*, most of which only slightly differed in size, proportions, and general shape of lorica. The investigated tintinnids were assigned to the *Parundella caudata* (Ostenfeld, 1899) Jörgensen, 1924 by a combination indistinct signs (lorica expands slightly below the oral rim, reaching a maximum width in the lower part, then narrows sharply, turning into a long and thin pedicel).

Thus, the found species of tintinnids belong to warm-water fauna. Probably the reason of the atypical finds in the western part of the Barents

Sea is natural invasion through the North Cape Current under the present conditions of temperature fluctuations in this water area. Thereby there is area expansion of ciliate species *A. tetragona*, *O. haeckeli*, *P. caudata* are new for this region in recent years and have a warm-water genesis.

Conclusion

1. The total list of tintinnid composition of the Barents Sea consists of 73 species belongs to 20 genera.

2. *Amphorellopsis tetragona* (Jörgensen, 1924) Kofoid and Campbell, 1929, *Ormosella haeckeli* Kofoid and Campbell, 1929, *Parundella caudata* (Ostenfeld, 1899) Jörgensen, 1924 are new for the Barents Sea and have a warm-water genesis.

Acknowledgements

Thank researchers of MMBI plankton laboratory: A.A. Oleinik and E.V. Druzhkova for significant assistance during this project. The research was carried out within the state assignment of MMBI KSC RAS (No. 0228-2019-0003).

References

Aksenov P.V. and Ivanov V.V. 2018. "Atlantification" as a possible cause for reducing of the Sea-Ice Cover in the Nansen Basin in winter. Arctic and Antarctic research. 64, 42–54 (in Russian with English summary).

- Alder V.A. 1999. Tintinnoinea. In: South Atlantic zooplankton. (Ed: Boltovskoy D.) Backhuys publishers, Leiden, pp. 321–384.
- Asha Devi C.R., Vimal Kumar K.G., Gupta G.V.M., Anilkumar V. and Sudhakar M. 2018. Implications of stratification on the heterotrophic plankton community (micro- and meso zooplankton) in the South Eastern Arabian Sea. *Regional Studies in Marine Science*. 24, 66–81.
- Boltovskoy D., Vivequin S.M. and Swanberg N.R. 1991. Vertical distribution of tintinnids and associated microplankton in the upper layer of the Barents Sea. *Sarsia*. 76, 141–151.
- Brandt K. 1906. Die Tintinnodeen der Plankton-Expedition. Tafeler klärungen nebst kurzer Diagnose der neuen Arten. Ergebnisse der Atlantic Plankton-Expedition, 3 (L.a.) (in German).
- Burkovsky I.V. 1976. New data on tintinnids (Ciliata) of the Arctic and revision of fauna. *Zool. Zh. (Moscow)*. 55, 325–336 (in Russian with English summary).
- Burkovsky I.V., Zamyshlyak E.A. and Poskryakova, N.P. 1974. Revision of the fauna of tintinnida (Ciliata) of the White Sea. *Zool. Zh. (Moscow)*. 53, 1757–1764 (in Russian with English summary).
- Cordeiro T.A. and Sassi R. 1997. Tintinnina (Ciliophora, Protista) of the North Sea during the spring of 1986. *Helgol. Meeresunters.* 51, 155–172.
- Dvoretzky V.G. and Dvoretzky A.G. 2010. Checklist of fauna found in zooplankton samples from the Barents Sea. *Polar. Biol.* 33, 991–1005.
- Dvoretzky V.G. and Dvoretzky A.G. 2018. Composition and distribution of summer zooplankton of polar front (the Barents Sea). *Herald of the Kola Science Centre of the RAS*. 2, 25–37 (in Russian with English summary).
- Dolan J.R. 2017. Historical trends in the species inventory of tintinnids (ciliates of the microzooplankton) in the Bay of Villefranche (NW Mediterranean Sea): shifting baselines. *Eur. J. Protistol.* 57, 16–25.
- Dolan J.R. and Pierce R.W. and Yang E.J. 2017. Tintinnid ciliates of the marine microzooplankton in Arctic seas: a compilation and analysis of species records. *Polar. Biol.* 40, 1247–1260.
- Dolan J.R. and Pierce R.W. 2014. Chapter 6.1. Tintinnid ciliates of the marine plankton. In: *Biogeographic atlas of the Southern Ocean* (Eds: De Broyer C., Koubbi P., Griffiths H.J., Raymond B., d'Udekem d'Acoz C., et al.). SCAR, Cambridge, pp. 254–259.
- Durmus T. and Balkis N. 2014. Tintinnid (Protozoa: Ciliophora) species in the Gulf of Gemlik and some ecological properties of the environment. *Fresenius Environ. Bull.* 23, 2991–2997.
- Fernandes L.F. 2004. Tintinnos (Ciliophora, Tintinnina) de aguas subtropicas na regio Sueste-Sul do Brasil. I: Familias Codonellidae, Codonellopsidae, Coxliellidae, Cyttarocilidae, Epplocilidae, Petalotrichidae, Ptychocylidae, Tintinnidae e Undellidae. *Revista Brasileira de Zoologia*. 21, 521–576 (in Portuguese).
- Fonda Umani S., Monti M., Cataletto B. and Budillon G. 2011. Tintinnid distributions in the Strait of Magellan (Chile). *Polar. Biol.* 34, 1285–1299.
- Gomez F. 2007. Trends on the distribution of ciliates in the open Pacific Ocean. *Acta Oecologia*. 32, 188–202.
- Jørgensen E.G. 1899. Protophyten und Protozoën im Plankton aus der norwegische Westküste. *Bergens Museums Aarbog*. 6, 1–112 (in German).
- Jørgensen E.G. 1924. Mediterranean Tintinnidae. Report on the Danish oceanographical expedition 1908–1910 to the Mediterranean and adjacent seas. Copenhagen. 2. J.3 (Biology).
- Jensen F. and Hansen B.W. 2000. Ciliates and heterotrophic dinoflagellates in the marginal ice zone of the central Barents Sea during spring. *J. Mar. Biol. Assoc. UK*. 80, 45–54.
- Kofoid C.A. and Campbell A.S. 1929. A conspectus of the marine and freshwater ciliata belonging to the suborder Tintinnoidea, with descriptions of new species principally from the Agassiz Expedition to the eastern tropical Pacific 1904–1905. *Univ. Calif. Publ. Zool.* 34, 1–403.
- Kršinić F. and Grbec B. 2006. Horizontal distribution of tintinnids in the open waters of the south Adriatic (Eastern Mediterranean). *Sci. Mar.* 70, 77–88.
- Lifshic A.V. 1977. Investigation of shell ciliates in the coastal zone of the Barents Sea. In: *Biology of the North Seas of the European part of the USSR*. pp. 15–20 (in Russian).
- Lifshits A.V. 1978. Ecological features of tintinnids (Tintinnida, Ciliata) in the coastal zone of the Barents Sea. In: *Patterns of bioproduction processes in the Barents Sea*. pp. 53–63 (in Russian).
- Lifshits, A.V., Fomin, O.K. and Khromov V.M. 1978. Seasonal species tintinnid complexes (Tintinnida, Ciliata) of the coast of the Barents Sea. In: *Patterns of bioproduction processes in the Barents Sea*. pp. 64–71 (in Russian).
- Makarevich P.R. and Oleinik. A.A. 2017. Microplankton of the Barents Sea: current composition and structure on the eve of the winter. *Vestnik of*

MSTU. 20, 316–325 (in Russian with English summary).

Matishov G., Makarevich P., Timofeev S., Kuznetsov L., Druzhkov N., Larionov V., Golubev V., Zuyev A., Adrov N., Denisov V., Iliyn G., Kuznetsov A., Denisenko S., Savinov V., Shavikyn A., Smolyar I., Levitus S., O'Brien T. and Baranova O. 2000. Biological atlas of the Arctic Seas 2000: Plankton of the Barents and Kara Seas. National Oceanographic Data Center / NOAA. Silver Spring, MD, USA.

Marshall S.M. 1969. Order: Tintinnida, Conseil International Pour L'exploration de la Mer. Zooplankton Sheet, pp. 117–127.

Meunier A. 1910. Microplankton des Mers de Barents et de Kara. Duc d'Orleans, 'Campagne Arctique de 1907'. Charles. Bulens, Bruxelles.

Monti M., Minocci M., Milani L. and Fonda Umani S. 2012. Seasonal and interannual dynamics of microzooplankton abundances in the Gulf of Trieste (Northern Adriatic Sea, Italy). Estuarine, Coastal Shelf Sci. 115, 149–157.

Ostenfeld C. 1899. Über CoccoSPAera und einige Tintinniden im Plankton des nördlichen Atlantischen Oceans. Zool. Anz. 22, 433–439.

Polyakov I.V., Pnyushkov A.V., Alkire M.B., Ashik I.M., Baumann T.M., Carmack E.C., et al. 2017. Greater role for Atlantic inflows on sea-ice loss in the Eurasian Basin of the Arctic Ocean. Science. 356, 285–291.

Rossolimo L.L. 1927. Planktonic ciliates of the Kara Sea. Tr. Plavuch. morsk. in-ta. 2, 63–77 (in Russian with English summary).

Santiago J.A. and Lagman M.C. 2018. Species check-list for Tintinnids of the Philippines Archipelago (Protozoa, Ciliophora). ZooKeys. 771, 1–14.

Thompson G.A. and Alder V.A. 2005. Patterns in tintinnid species composition and abundance in relation to hydrological conditions of the Southwestern Atlantic during austral spring. Aquat. Microb. Ecol. 40, 85–101.

Toklu-Aliçli B., Balkis N., Toklu A.S. and Andhöbek A. 2010. First record of *Amphorellopsis tetragona* (Protozoa: Ciliophora: Tintinnina) from the Sea of Marmara. IUFS Journal of Biology. 69, 103–110.

Zakaria H.Y., Radwan A.A. and Said M.A. 2007. Influence of salinity variations on zooplankton community in El-Mex Bay, Alexandria. Egypt J. Aquat. Res. 33, 52–67.

Address for correspondence: Olga Chovgan. Murmansk Marine Biological Institute KSC RAS, Vladimirskaia St. 17, Murmansk, 183010, Russia; e-mail: chovgan@mmbi.info.